

Local extrema and mean value theorems

Analysis 1

S.-J. Kimmerle

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Functions

Differentiation in 1d

Derivatives

Local extrema and mean value theorems

Convex and concave

Excursion: unrestricted optimization in 1d

Integration in 1d

Summary - outlook and review

We consider

Problem (Optimization problem (OP))

Minimize $f(x)$

subject to the constraint $x \in X \subseteq \mathbb{R}$.

*Thereby let $f : X \rightarrow \mathbb{R}$ an at least **1x continuously differentiable** function.*

If $X = \mathbb{R}$, then we call this an unrestricted optimization problem.



Definition (Minima)

- $\hat{x} \in X$ is called a **global minimum** of (OP), if

$$f(\hat{x}) \leq f(x) \quad \text{for all } x \in X \subseteq \mathbb{R}. \quad (2)$$

- $\hat{x} \in X$ is called a **strict global minimum** of (OP), if in (2) “ $<$ ” for all $x \in X$ holds except for $x = \hat{x}$.

- $\hat{x} \in X$ is called a **local minimum** of (OP), if there exists for a $\varepsilon > 0$ a neighbourhood

$$U_\varepsilon(\hat{x}) := \{x \in \mathbb{R}^n \mid |x - \hat{x}| < \varepsilon\}$$

with

$$f(\hat{x}) \leq f(x) \quad \text{für alle } x \in X \cap U_\varepsilon(\hat{x}). \quad (3)$$

- $\hat{x} \in X$ is called a **strict local minimum** of (OP), if in (3) “ $<$ ” for all $x \in X \cap U_\varepsilon(\hat{x})$ holds except for $x = \hat{x}$.

A strict minimum is also called an isolated minimum.
Analogously corresponding maxima are defined.

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Functions

Differentiation in 1d

Derivatives

Local extrema and mean value theorems

Convex and concave

Excursion: unrestricted optimization in 1d

Integration in 1d

Summary - outlook and review



Extrema: example - to be completed in the lecture

Analysis 1

S.-J. Kimmerle

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Functions

Differentiation in 1d

Derivatives


Local extrema and mean value theorems

Convex and concave

Excursion: unrestricted optimization in 1d

Integration in 1d

Summary - outlook and review



A twice continuously differentiable function $f : \mathbb{R} \rightarrow \mathbb{R}$ exhibits a local minimum at \hat{x} ,

then:

- $f'(\hat{x}) = 0$ (necessary condition of 1st order)
- $f''(\hat{x}) \geq 0$ (necessary condition of 2nd order)

Vice versa let $f'(\hat{x}) = 0$ and $f''(\bar{x}) > 0$,

then \hat{x} is a local minimum of f .

(sufficient condition)

A point \hat{x} where $f'(\hat{x}) = 0$, is called a **stationary point**.



$$f : (-1, 1) \rightarrow \mathbb{R}, x \mapsto x^3$$

$$f : [-1, 1] \rightarrow \mathbb{R}, x \mapsto x$$

Introduction

Basics (sets,
mappings, and
numbers)

Proof techniques

Sequences and
series

Functions

Differentiation in
1d

Derivatives


Local extrema and mean
value theorems

Convex and concave

Excursion: unrestricted
optimization in 1d

Integration in 1d

Summary - outlook
and review




Theorem (Rolle)

*Let $a < b$ and $f : [a, b] \rightarrow \mathbb{R}$ a continuous function
that is continuous differentiable in (a, b)
with $f(a) = f(b)$,*

then there exists a $\xi \in (a, b)$ such that

$$f'(\xi) = 0.$$



Corollary (Mean value theorem)

Let $a < b$ and $f : [a, b] \rightarrow \mathbb{R}$ a continuous function that is continuously differentiable in (a, b) ,

then there exists a $\xi \in (a, b)$ such that

$$f'(\xi) = \frac{f(b) - f(a)}{b - a}.$$

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Functions

Differentiation in 1d

Derivatives


Local extrema and mean value theorems

Convex and concave

Excursion: unrestricted optimization in 1d

Integration in 1d

Summary - outlook and review



Mean value theorem - geometrical interpretation

Analysis 1

S.-J. Kimmerle

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Functions

Differentiation in 1d

Derivatives

Local extrema and mean value theorems

Convex and concave

Excursion: unrestricted optimization in 1d

Integration in 1d

Summary - outlook and review

Geometrically the mean value theorem means that the slope of the secant at the graph of f at the points $(a, f(a))$ and $(b, f(b))$ is equal to the slope of the tangent line to the graph of f at a point $(\xi, f(\xi))$.



Theorem (Generalized mean value theorem)

Let $a < b$ and $f, g : [a, b] \rightarrow \mathbb{R}$ continuous functions that are continuously differentiable in (a, b) , and $g'(x) \neq 0$ for all $x \in (a, b)$, then there exists a $\xi \in (a, b)$ such that

$$\frac{f'(\xi)}{g'(\xi)} = \frac{f(b) - f(a)}{g(b) - g(a)}.$$

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Functions

Differentiation in 1d

Derivatives

Local extrema and mean value theorems

Convex and concave

Excursion: unrestricted optimization in 1d

Integration in 1d

Summary - outlook and review

