

We are going to introduce the trigonometric functions \sin and \cos by the **Euler formula**

$$\exp(ix) = \cos(x) + i \sin(x), \quad x \in \mathbb{R}, \quad i \text{ the imaginary unity.}$$

This requires complex numbers \mathbb{C} and the exponential function for complex arguments.

Another important motivation is the solution of equations of the type:

$$x^2 = -a, \quad a \in \mathbb{R}^+.$$

The fundamental theorem of algebra says any polynomial of degree n has n complex roots (if counted properly)
 \rightsquigarrow Linear Algebra.



Definition (Complex numbers)

The set $\mathbb{R} \times \mathbb{R}$ of (ordered) pairs together with the operations

$$(x_1, y_1) + (x_2, y_2) := (x_1 + x_2, y_1 + y_2),$$

$$(x_1, y_1) \cdot (x_2, y_2) := (x_1 x_2 - y_1 y_2, x_1 y_2 + y_1 x_2)$$

is called the **complex numbers**, abbreviated with \mathbb{C} .

The neutral element of the addition is $(0, 0)$,
of the multiplication $(1, 0)$.

The inverse of (x, y) w.r.t. the addition is $(-x, -y)$,
w.r.t. the multiplication (where $(x, y) \neq (0, 0)$) is

$$\left(\frac{x}{x^2 + y^2}, \frac{-y}{x^2 + y^2} \right).$$

[Introduction](#)[Basics \(sets, mappings, and numbers\)](#)[Proof techniques](#)[Sequences and series](#)[Functions](#)[Continuity](#)[Applications for continuous functions](#)[Inverse functions](#)[Exponentiation and logarithm](#)[Complex numbers](#)[Trigonometric functions](#)[Differentiation in 1d](#)[Integration in 1d](#)[Summary - outlook and review](#)

Theorem

Field \mathbb{C} The set of complex numbers \mathbb{C} is a field.

But \mathbb{C} is no ordered field.

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Functions

Continuity

Applications for continuous functions

Inverse functions

Exponentiation and logarithm

Complex numbers

Trigonometric functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review



We identify

- $(1, 0)$ with the unity 1 (of real numbers),
- $(0, 1)$ with the **imaginary unity** i

yielding the standard notation

$$z = a + bi \quad \text{for } (a, b)$$

We call $\operatorname{Re}(z) := a$ the **real part of** z ,
and $\operatorname{Im}(z) := b$ the **imaginary part of** z .

Cartesian complex plane,
similar to \mathbb{R}^2 ,
but with a different structure



Let $x_1, x_2, y_1, y_2, \lambda \in \mathbb{R}$.

Then

- $(x_1 + y_1 i) + (x_2 + y_2 i) = (x_1 + x_2) + (y_1 + y_2) i$
- $(x_1 + y_1 i) \cdot (x_2 + y_2 i) = (x_1 x_2 - y_1 y_2) + (x_1 y_2 + x_2 y_1) i$

In particular:

- $i^2 = -1$
- $\lambda(x_1 + y_1 i) = \lambda x_1 + \lambda y_1 i$
- $(x_1 + y_1 i) \cdot (x_1 - y_1 i) = x_1^2 + y_1^2$

We call $\bar{z} := x_1 - y_1 i$ the **complex conjugate** of $z = x_1 + y_1 i \in \mathbb{C}$.



Geometric interpretation of addition in \mathbb{C}

Analysis 1

S.-J. Kimmerle

Introduction

Basics (sets,
mappings, and
numbers)

Proof techniques

Sequences and
series

Functions

Continuity

Applications for continuous
functions

Inverse functions

Exponentiation and
logarithm

Complex numbers

Trigonometric functions

Differentiation in
1d

Integration in 1d

Summary - outlook
and review



Polar coordinate system for a point $P = (x|y)$:

uses distance $r = \sqrt{x^2 + y^2}$ to the origin 0
and angle ϕ between the positive x -axis and $0P$

For complex numbers:

radius $r = |z| = |x + yi| = \sqrt{z\bar{z}}$, the absolute value in \mathbb{C}

$$\phi = \arg(x + yi) := \begin{cases} \arccos\left(\frac{a}{r}\right) & ; b \geq 0 \\ -\arccos\left(\frac{a}{r}\right) & ; b < 0 \end{cases}$$

Polar complex plane

ϕ is not uniquely defined,

here $\phi \in (-\pi, \pi]$



We consider convergence of the real and imaginary part.

Concepts as (Cauchy) sequences, series, limits ... may be transferred directly from real to complex numbers!

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Functions

Continuity

Applications for continuous functions

Inverse functions

Exponentiation and logarithm

Complex numbers

Trigonometric functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review

