

Theorem (Limits of inequalities)

Consider two convergent real sequences

$\{a_n\}_{n \geq n_0}$ with $\lim_{n \rightarrow \infty} a_n = a$ and

$\{b_n\}_{n \geq n_0}$ with $\lim_{n \rightarrow \infty} b_n = b$.

If $a_n \leq b_n$ for all $n \geq n_0$,
then

$$a = \lim_{n \rightarrow \infty} a_n \leq \lim_{n \rightarrow \infty} b_n = b.$$

Corollary:

$$\alpha \leq a_n \leq A$$

$$\lim_{n \rightarrow \infty}$$

$$\alpha \leq \underbrace{\lim_{n \rightarrow \infty} a_n}_{= a} \leq A$$

Introduction

Basics (sets,
mappings, and
numbers)

Proof techniques

Sequences and
series

Convergence

Series

Functions

Differentiation in
1d

Integration in 1d

Summary - outlook
and review

Definition (Definite divergence of a sequence)

A real sequence $\{a_n\}_{n \leq n_0}$ is called

- **definitely (or certainly) divergent to $+\infty$,**
if there exists for any $k_1 \in \mathbb{R}$ a $N(k_1) \in \mathbb{N}$ s.t.

$$a_n > k_1 \text{ for all } n \geq N(k_1),$$

- **definitely (or certainly) divergent to $-\infty$,**
if there exists for any $k_2 \in \mathbb{R}$ a $N(k_2) \in \mathbb{N}$ s.t.

$$a_n < k_2 \text{ for all } n \geq N(k_2).$$

Otherwise a divergent sequence is called **indefinitely divergent**.

We write

$$\lim_{n \rightarrow \infty} a_n = \pm\infty.$$

Introduction

Basics (sets,
mappings, and
numbers)

Proof techniques

Sequences and
series

Convergence

Series

Functions

Differentiation in
1d

Integration in 1d

Summary - outlook
and review

Cauchy sequence

Analysis 1


S.-J. Kimmerle

Definition (Cauchy sequence)

A real sequence $\{a_n\}_{n \leq n_0}$ is called a **Cauchy sequence**, if for any $\varepsilon > 0$ there exists a natural number $N(\varepsilon)$ s.t.

$$|a_n - a_m| < \varepsilon \quad \text{for all } n, m \geq N(\varepsilon).$$

Every Cauchy sequence in the real numbers has a limit, this is equivalent to (or also one definition of) the completeness of the real numbers.



Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Convergence

Series

Functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review



Equivalence of convergent and Cauchy sequences

Analysis 1

S.-J. Kimmerle

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Convergence

Series

Functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review

Theorem (Cauchy criterion for the convergence of sequences)

A real sequence $\{a_n\}_{n \geq n_0}$ is convergent, iff $\{a_n\}_{n \geq n_0}$ is a Cauchy sequence.

Proof:

$\{a_n\}_{n \geq n_0}$ is convergent $\Leftrightarrow \{a_n\}_{n \geq n_0}$ Cauchy sequence



Definition (Subsequence)

Let $\{a_n\}_{n \geq n_0}$ be a sequence of numbers and $n_1 < n_2 < \dots$ a strictly increasing sequence of natural numbers.

Then $\{a_{n_k}\}$ is called a subsequence of $\{a_n\}$.

not a subsequence: $\{a_1, a_1, a_1, a_3, a_2, a_2, \dots\}$

$$a_n := (-1)^n \quad \{a_1, a_2, a_3, a_4, \dots\} \\ = \{-1, 1, -1, 1, \dots\}$$

$$a_{2n} = (-1)^{2n} = ((-1)^2)^n = 1^n = 1$$

$$\text{subsequence} \rightarrow \{a_2, a_4, a_6, a_8, \dots\} \\ = \{1, 1, 1, 1, \dots\}$$

$$\downarrow \\ a_{2n+1} = (-1)^{2n+1} = ((-1)^2)^n (-1) = -1$$

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Convergence

Series

Functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review



Bolzano-Weierstrass theorem

Analysis 1

S.-J. Kimmerle

Weierstraß

Theorem (Bolzano-Weierstrass)

Any bounded real sequence $\{a_n\}_{n \geq n_0}$
has a convergent subsequence.

n

Example:

$a_n = (-1)^n$ bounded ✓
not definitely divergent

convergent subsequences

$\{1, 1, 1, \dots\}$ with acc.pt. 1
or $\{-1, -1, -1, \dots\}$ with acc.pt. -1
or $\{-1, -1, 1, 1, 1, \dots\}$ with acc.pt. 1
= 2 exceptions

Introduction

Basics (sets,
mappings, and
numbers)

Proof techniques

Sequences and
series

Convergence

Series

Functions

Differentiation in
1d

Integration in 1d

Summary - outlook
and review



Accumulation point

Definition

Theorem (Accumulation point)

A real number a is called an **accumulation point** (or **cluster point** or **limit point**) of a real sequence $\{a_n\}_{n \geq n_0}$, if there exists a subsequence of $\{a_n\}_{n \geq n_0}$ that converges to a .

$$b_n := \frac{1}{n} \longrightarrow 0 \text{ as } n \rightarrow +\infty$$

↑ limit

$$\{ \underline{1}, \underline{\frac{1}{2}}, \underline{\frac{1}{3}}, \underline{\frac{1}{4}}, \underline{\frac{1}{5}}, \dots \}$$

- * any limit is an accumulation point
- * if a sequence converges, the limit is the only accumulation point



Definition (Limes superior, limes inferior)

Let $\{a_n\}_{n \geq n_0}$ be a sequence of real numbers, then we define

- the **limes superior** of the sequence $\{a_n\}_{n \geq n_0}$ as

$$\limsup_{n \rightarrow \infty} a_n := \lim_{n \rightarrow \infty} \left(\sup_{k \geq n} a_k \right),$$

- the **limes inferior** of the sequence $\{a_n\}_{n \geq n_0}$ as

$$\liminf_{n \rightarrow \infty} a_n := \lim_{n \rightarrow \infty} \left(\inf_{k \geq n} a_k \right).$$

Remark: This definition is later important for a general approach to integrals required for stochastics.

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Convergence


Series

Functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review



Definition (Monotone real sequences)

A sequence $\{a_n\}_{n \geq n_0}$ of real numbers is called

- **monotonically increasing**,
if $a_n \leq a_{n+1}$ for all $n \geq n_0$,
- **strictly monotonically increasing**,
if $a_n < a_{n+1}$ for all $n \geq n_0$,
- **monotonically decreasing**,
if $a_n \geq a_{n+1}$ for all $n \geq n_0$,
- **strictly monotonically decreasing**,
if $a_n > a_{n+1}$ for all $n \geq n_0$.

$$a_n := n$$
$$b_n := \frac{1}{n}$$

$$c_n := 1 - \frac{1}{n} \quad \left| \quad e_n := (-1)^n \right.$$
$$d_n := \frac{(-1)^n}{2^n} \quad \left| \quad \text{not monotone} \right.$$
$$\left\{ -\frac{1}{2}, \frac{1}{4}, -\frac{1}{8}, \dots \right\} \quad \text{not monotone}$$

Introduction

Basics (sets,
mappings, and
numbers)

Proof techniques

Sequences and
series

Convergence

Series

Functions

Differentiation in
1d

Integration in 1d

Summary - outlook
and review


Properties of monotone sequences

Analysis 1

S.-J. Kimmerle

Theorem (Convergence of bounded monotone sequences)

Any bounded monotone real sequence $\{a_n\}_{n \geq n_0}$ converges.



An illustration of the statement of the theorem.

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Convergence

Series

Functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review



Definition (Countable set)

A non-empty set M is called **countable**, if there exists a sequence $\{a_n\}_{n \geq n_0}$ with

$$\mathbb{S} \subseteq M = \{a_n \mid n \in \mathbb{N}\}.$$

A non-empty set M is called **uncountable**, if it is not countable.

Evidently, a subset of a countable set is countable.

A finite set is countable. Equivalently, an uncountable set has not a finite number of elements.

A countable set may have a finite or infinite number of elements.

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Convergence

Series

Functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review



Theorem (Countability of the union of sets)

The union of a countable number of countable sets M_n , $n \in \mathbb{N}$, i.e.

$$\bigcup_{n \in \mathbb{N}} M_n = \mathcal{M}$$

is again countable.

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Convergence

Series

Functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review



Countability of rational numbers

Analysis 1

S.-J. Kimmerle

Theorem (Countability of \mathbb{Q})

The set \mathbb{Q} of rational numbers is countable.


Proof: by
Cantor's
diagonal seq.

$$\mathbb{Q} = \left\{ \frac{p}{q} \mid p \in \mathbb{Z}, q \in \mathbb{Z} \setminus \{0\} \right\}$$

We consider $\left\{ \frac{p}{q} \mid p \in \mathbb{N}, q \in \mathbb{N} \right\}$

w.l.o.g. (see the last theorem)

bijection
to \mathbb{N}



Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Convergence

Series

Functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review

