

Theorem (Properties of the absolute value)

For any $w, z \in \mathbb{C}$ there holds:

$$1)a) z \geq 0 \quad 1)b) |z| = 0 \Leftrightarrow z = 0$$

$$2) |z + w| \leq |z| + |w| \quad (\text{triangle inequality})$$

$$3) |z \cdot w| = |z||w| \quad (\text{multiplicativity})$$

Remark: A field, for whose elements a mapping with the properties of an absolute value exists, is called a **valued field**.



We extend the exponential function into the complex plane.

Definition (Complex exponential function)

For $z \in \mathbb{C}$ we define

$$\exp(z) := \sum_{k=0}^{\infty} \frac{1}{k!} z^k = 1 + z + \frac{z^2}{2} + \frac{z^3}{6} + \dots$$

Properties:

- 1 $\exp(z) \neq 0$ for any $z \in \mathbb{C}$
- 2 $\exp(z)$ is continuous
- 3 $\exp(\bar{z}) = \overline{\exp(z)}$
- 4 Let $z = x + yi$,
then $|\exp(z)| = |\exp(x)|$.



Definition (Cosine and sine)

For $x \in \mathbb{R}$ we define:

$$\cos(x) := \operatorname{Re}(e^{ix})$$

$$\sin(x) := \operatorname{Im}(e^{ix})$$

We see that the Euler formula holds:

$$\exp(ix) = \cos(x) + i \sin(x), \quad x \in \mathbb{R}$$

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Functions

Continuity

Applications for continuous functions

Inverse functions

Exponentiation and logarithm

Complex numbers

Trigonometric functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review



Geometric interpretation of the Euler formula

Analysis 1

S.-J. Kimmerle

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Functions

Continuity

Applications for continuous functions

Inverse functions

Exponentiation and logarithm

Complex numbers

Trigonometric functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review

The range of values is the unit circle:

$$\begin{aligned} |\exp(ix)| &= \left(\exp(ix) \overline{\exp(ix)} \right)^{1/2} \\ &= (\exp(ix) \exp(-ix))^{1/2} = (\exp(0))^{1/2} = 1, \quad x \in \mathbb{R} \end{aligned}$$

$\sin(x)$ and $\cos(x)$ are the projections on the real and imaginary axis, resp.

\sin and \cos are extended on complex arguments by $\exp(iz)$, $z \in \mathbb{C}$. For real arguments \sin and \cos coincide with the known functions.



We consider the polar complex plane.

We have

$$z = r \exp(i\phi) \in \mathbb{C}$$

with $r = |z| \in \mathbb{R}_0^+$ and $\phi \in (-\pi, \pi]$.

Thus for the multiplication of $z_1 = r_1 \exp(i\phi_1)$,
 $z_2 = r_2 \exp(i\phi_2) \in \mathbb{C}$ there holds

$$z_1 z_2 = r_1 r_2 \exp(i(\phi_1 + \phi_2)).$$

Introduction

Basics (sets,
mappings, and
numbers)

Proof techniques

Sequences and
series

Functions

Continuity

Applications for continuous
functions

Inverse functions

Exponentiation and
logarithm

Complex numbers

Trigonometric functions

Differentiation in
1d

Integration in 1d

Summary - outlook
and review

Theorem (Properties of cosine and sine)

For all $x \in \mathbb{R}$ there holds:

- $\cos(x) = \frac{1}{2} (e^{ix} + e^{-ix})$, $\sin(x) = \frac{1}{2i} (e^{ix} - e^{-ix})$
- $\cos(-x) = \cos(x)$, $\sin(-x) = -\sin(x)$
- $\cos^2(x) + \sin^2(x) = 1$
- $\cos : \mathbb{R} \rightarrow \mathbb{R}$ and $\sin : \mathbb{R} \rightarrow \mathbb{R}$ are continuous on \mathbb{R} .



Theorem (Addition theorems of cosine and sine)

For all $x, y \in \mathbb{R}$ there holds:

$$\begin{aligned}\cos(x + y) &= \cos(x) \cos(y) - \sin(x) \sin(y), \\ \sin(x + y) &= \sin(x) \cos(y) + \cos(x) \sin(y).\end{aligned}$$

Theorem (Duplication formulas of cosine and sine)

For all $x \in \mathbb{R}$ there holds:

$$\begin{aligned}\cos(2x) &= \cos^2(x) - \sin^2(x), \\ \sin(2x) &= 2 \sin(x) \cos(x).\end{aligned}$$



Theorem (Sine and cosine as series)

For $x \in \mathbb{R}$ we define

$$\cos(x) := \sum_{k=0}^{\infty} (-1)^k \frac{1}{(2k)!} x^{2k} = 1 - \frac{x^2}{2} + \frac{x^4}{24} + \dots$$

$$\sin(x) := \sum_{k=0}^{\infty} (-1)^k \frac{1}{(2k+1)!} x^{2k+1} = x - \frac{x^3}{6} + \frac{x^5}{120} + \dots$$

Both series converge absolutely.

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Functions

Continuity

Applications for continuous functions

Inverse functions

Exponentiation and logarithm

Complex numbers

Trigonometric functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review



Series representation of (co)sine - proof

Analysis 1

S.-J. Kimmerle

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Functions

Continuity

Applications for continuous functions

Inverse functions

Exponentiation and logarithm

Complex numbers

Trigonometric functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review

Proof.

The exponential series is absolutely convergent. We consider:

$$\exp(ix) = \sum_{k=0}^{\infty} \frac{(ix)^k}{k!}$$

Note that: $i^0 = 1, i^1 = i, i^2 = -1, i^3 = -i, i^4 = 1$

In a cyclic manner: $i^{4k} = 1, i^{4k+1} = i, i^{4k+2} = -1, \dots$

We order the terms (why is it allowed ?) with even $(4k, 4k + 2)$ and odd indices $(4k + 1, 4k + 3)$:

$$\exp(ix) = \sum_{k=0}^{\infty} (-1)^k \frac{1}{(2k)!} x^{2k} + i \sum_{k=0}^{\infty} (-1)^k \frac{1}{(2k+1)!} x^{2k+1}$$

Together with the Euler formula

$$\exp(ix) = \cos(x) + i \sin(x)$$

the statement is verified. □



Since we may evaluate only a finite part of a series, it is important to have estimates for the remainder:

Theorem (Estimates for the remainder for (co)sine)

For all $x \in \mathbb{R}$:

$$\exp(x) = \sum_{k=0}^n \frac{x^k}{k!} + r_n(x),$$

$$\text{with } |r_{n+1}(x)| \leq 2 \frac{|x|^{n+1}}{(n+1)!} \quad \text{for } |x| \leq \frac{n+1}{2}$$

$$\cos(x) = \sum_{k=0}^n (-1)^k \frac{x^{2k}}{(2k)!} + r_{2n+2}(x),$$

$$\text{with } |r_{2n+2}(x)| \leq \frac{|x|^{2n+2}}{(2n+2)!} \quad \text{for } |x| \leq 2n+3$$

$$\sin(x) = \sum_{k=0}^n (-1)^k \frac{x^{2k+1}}{(2k+1)!} + r_{2n+3}(x),$$

$$\text{with } |r_{2n+3}(x)| \leq \frac{|x|^{2n+3}}{(2n+3)!} \quad \text{for } |x| \leq 2n+4$$

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Functions

Continuity

Applications for continuous functions

Inverse functions

Exponentiation and logarithm

Complex numbers

Trigonometric functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review



Theorem (Some limits sin and cos)

For any $x \in \mathbb{R}$ we have

$$\begin{aligned} 1) \quad \lim_{x \rightarrow 0, x \neq 0} \frac{\sin(x)}{x} &= 1, & 2) \quad \lim_{x \rightarrow \infty} \frac{\sin(x)}{x} &= 0, \\ 3) \quad \lim_{x \rightarrow 0, x \neq 0} \frac{\cos x - 1}{x^2} &= \frac{1}{2}, & 4) \quad \lim_{x \rightarrow \infty} \frac{\cos(x)}{x} &= 0 \end{aligned}$$

Lemma (2 estimates for sin and cos)

$$\begin{aligned} \sin(x) &> 0 && \text{for all } x \in (0, 2] \\ \cos(2) &\leq -\frac{1}{3} \end{aligned}$$



Definition (The number π)

The cosine has exactly one zero p in the interval $[0, 2]$.
We define

$$\pi := 2p.$$

The real number π is an infinite, non-periodic decimal fraction, i.e. an irrational number,

$$\pi = 3, 141\,592\,653\,589 \dots$$

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Functions

Continuity

Applications for continuous functions

Inverse functions

Exponentiation and logarithm

Complex numbers

Trigonometric functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review



Periodicities of sine and cosine I

Analysis 1

S.-J. Kimmerle

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Functions

Continuity

Applications for continuous functions

Inverse functions

Exponentiation and logarithm

Complex numbers

Trigonometric functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review

$$\cos(x + 2\pi) = \cos(x)$$

$$\sin(x + 2\pi) = \sin(x)$$

$$\cos(x + \pi) = -\cos(x)$$

$$\sin(x + \pi) = -\sin(x)$$

$$\cos(x) = \sin\left(\frac{\pi}{2} - x\right)$$

$$\sin(x) = \cos\left(\frac{\pi}{2} - x\right)$$



Periodicities of sine and cosine II

Analysis 1

S.-J. Kimmerle

Introduction

Basics (sets,
mappings, and
numbers)

Proof techniques

Sequences and
series

Functions

Continuity

Applications for continuous
functions

Inverse functions

Exponentiation and
logarithm


Complex numbers

Trigonometric functions

Differentiation in
1d

Integration in 1d

Summary - outlook
and review



Sine and cosine with half the periodicity



The quotient of sine and cosine has a certain importance.
It is called the **tangent**

$$\tan(x) = \frac{\sin(x)}{\cos(x)}.$$

Moreover, we define the **cotangent**

$$\cot(x) = \frac{\cos(x)}{\sin(x)}.$$

Note the singularities at zeros of cos or sin.

Remark: There exist further trigonometric functions that we do not consider here:

$$\sec(x) = \frac{1}{\cos(x)}, \quad \csc(x) = \frac{1}{\sin(x)}$$



Special values of trigonometric functions

Analysis 1

S.-J. Kimmerle

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Functions

Continuity

Applications for continuous functions

Inverse functions

Exponentiation and logarithm

Complex numbers

Trigonometric functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review

Radian	degree	sin	cos	tan
0	0°	0	1	0
$\frac{\pi}{6}$	30°	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$
$\frac{\pi}{4}$	45°	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1
$\frac{\pi}{3}$	60°	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$
$\frac{\pi}{2}$	90°	1	0	∞



Special values of co(sine)

Introduction

Basics (sets, mappings, and numbers)

Proof techniques

Sequences and series

Functions

Continuity

Applications for continuous functions

Inverse functions

Exponentiation and logarithm


Complex numbers

Trigonometric functions

Differentiation in 1d

Integration in 1d

Summary - outlook and review



Unit circle with coordinates of important points

Source: [Wikipedia](#)



On the interval $[0, \pi]$ the function \cos is strictly monotone decreasing and maps bijectively to $[-1, 1]$.
The inverse function

$$\arccos(x) : [-1, 1] \rightarrow \mathbb{R}$$

exists and it is called **arccosine**.

On the interval $[-\pi/2, \pi/2]$ the function \sin is strictly monotone increasing and maps bijectively to $[-1, 1]$.
The inverse function

$$\arcsin(x) : [-1, 1] \rightarrow \mathbb{R}$$

exists and it is called **arcsine**.



On the interval $(-\pi/2, \pi/2)$ the function \tan is strictly monotone increasing and maps bijectively to \mathbb{R} .
The inverse function

$$\arctan(x) : \mathbb{R} \rightarrow \mathbb{R}$$

exists and it is called **arctangent**.

Note that the inverse functions are restricted to a “principal branch”.



Let $x \in \mathbb{R}$.

We define the **hyperbolic cosine** as

$$\cosh(x) := \frac{1}{2} (\exp(x) + \exp(-x)),$$

and the **hyperbolic sine** as

$$\sinh(x) := \frac{1}{2} (\exp(x) - \exp(-x)).$$

Applications:

- sagging high voltage lines or chains (e.g. of bridges) have the shape of a cosh
- magnetism
- cosmology
- ...

