

More precisely, we carry out the following computations:

- We calculate the natural homomorphism $O(L_-) \rightarrow O(q_-)$, and obtain an upper bound for the index $[O(L_+, \mathcal{P}_+) : \Gamma]$ by Proposition ???. In fact, we calculate the exact value of this index except for the case $\gamma = (\varepsilon_{12}, 0)$.
- We compute the finite sets $\tilde{\Xi}_\tau$ and Ξ_τ defined in (??) and (??).
- We confirm that $\mathcal{M}(D_0) = \tilde{\mathcal{W}}(D_0)$ holds by showing that Ξ_τ satisfies condition ?? in Corollary ??.
- We calculate a finite generating set of the subgroup $\Sigma_\varepsilon([\tau])$ of $\text{Cen}(\cdot, \varepsilon)$, and compute the orbit decomposition of Ξ_τ under the action of $\Sigma_\varepsilon([\tau])$. Thus we obtain the $\text{Stab}_\Gamma(D_0)$ -orbits in $\mathcal{W}(D_0)$ by Remark ??.
- For each $\text{Stab}_\Gamma(D_0)$ -orbit in $\mathcal{W}(D_0)$, we compute by Lemma ?? the fiber of $\mu: \tilde{\mathcal{W}}(D_0) \rightarrow \mathcal{W}(D_0)$ over a representative of the orbit.

The results are summarized in Table ??.