

Mathematics Stack Exchange is a question and answer site for people studying math at any level and professionals in related fields. It only takes a minute to sign up.

Anybody can ask a question

X

[Sign up to join this community](#)

Anybody can answer

The best answers are voted up and rise to the top

MATHEMATICS

Interesting implicit surfaces in \mathbb{R}^3

Asked 11 years, 6 months ago Modified 8 months ago Viewed 7k times



I have just written a small program in C++ and OpenGL to plot implicit surfaces in \mathbb{R}^3 for a Graphical Computing class and now I'm in need of more interesting surfaces to implement!

28



Some that I've implemented are:



- Basic surfaces like spheres and cylinders;
- Nordstrand's Weird Surface;
- Klein Quartic;
- Goursat's Surface;
- Heart Surface;



So, my question is, what are other interesting implicit surfaces in \mathbb{R}^3 ?

P.S.: I know this is kind of vague, but anything you find interesting will be of use. (:

P.P.S: Turn this into a community wiki, if need be.

[algebraic-geometry](#) [multivariable-calculus](#) [visualization](#) [surfaces](#)

Share Cite Follow

edited Jun 20, 2011 at 21:52

community wiki
8 revs, 5 users 64%
Leonardo Fontoura

2 There is a wealth of documentation on Wolfram's Mathworld: mathworld.wolfram.com/topics/Surfaces.html – [barf](#) Jun 19, 2011 at 1:38

Boy's surface.. – [Cheerful Parsnip](#) Jun 19, 2011 at 1:45

If you type "mathematical surfaces" into Google you get a near endless list, including lots of raytraced models, for example: math.rug.nl/models – [Ryan Budney](#) Jun 19, 2011 at 3:13 ↗

math.rug.nl/models/Clebsch.html is beautiful. Now I want one! Actually: can someone give me the equations of the lines in the Clebsch surface? – [Mariano Suárez-Álvarez](#) Jun 19, 2011 at 3:24 ↗

@Mariano: I can't but [Oliver Lab](#) can (in German) see in particular chapter 3, but the general form looks rather messy. He has some cool pictures on his page algebraicsurface.net – [t.b.](#) Jun 19, 2011 at 15:40

4 Answers

Sorted by: Highest score (default) ↗

▲ A really nice family of implicit surfaces in \mathbb{R}^3 are the Banchoff-Chmutov surfaces

62

$$BC_n := \{(x, y, z) \in \mathbb{R}^3 ; T_n(x) + T_n(y) + T_n(z) = 0\},$$



▼ where T_n denotes the n -th Chebyshev polynomial of the first kind, i.e.





$$T_n(x) := \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} (x^2 - 1)^k x^{n-2k}.$$





• $\leftarrow BC_1 \quad BC_2 \rightarrow$




• $\leftarrow BC_3 \quad BC_4 \rightarrow$



• $\leftarrow BC_5 \quad BC_6 \rightarrow$



• $\leftarrow BC_7 \quad BC_8 \rightarrow$




What is the genus of BC_{2n+2} ? Code for plotting surfaces BC_n in Mathematica 7:

```
BanchoffChmutov[n_] := ContourPlot3D[ChebyshevT[n,x]+ChebyshevT[n,y]+ChebyshevT[n,z], {x,-1.3,1.3}, {y,-1.3,1.3}, {z,-1.3,1.3}, Contours->0.02, AspectRatio->Automatic, Boxed->False, Axes->{False,False,False}, BoxRatios->Automatic, PlotRangePadding->None, PlotPoints->30, ViewPoint->{-2,3,3}]
surfacesBCn = Table[BanchoffChmutov[i], {i, 8}]
Table[ChebyshevT[n,x]+ChebyshevT[n,y]+ChebyshevT[n,z], {n, 8}]
```


Other miscellaneous increasingly pretty examples, that I managed to construct myself, include:

$$\{(x, y, z) \in \mathbb{R}^3 ; (x - 2)^2(x + 2)^2 + (y - 2)^2(y + 2)^2 + (z - 2)^2(z + 2)^2 +$$


$$3(x^2y^2 + x^2z^2 + y^2z^2) + 6 * x * y * z - 10(x^2 + y^2 + z^2) + 22 = 0$$



$$\{(x, y, z) \in \mathbb{R}^3 ; \quad ((x-1)x^2(x+1)+y^2)^2 + \\ ((y-1)y^2(y+1)+z^2)^2 + \\ 0.1y^2 + 0.05(y-1)y^2(y+1) = 0\}$$




$$\{(x, y, z) \in \mathbb{R}^3 ; \quad 15((x-1.2)x^2(x+1.2)+y^2)^2 + 0.8(z-1)z^2(z+1) - 0.1z^2 + \\ 20((y-0.8)y^2(y+0.8)+z^2)^2 + 0.8(x-1)z^2(x+1) - 0.1x^2 = 0\}$$




$$\{(x, y, z) \in \mathbb{R}^3 ; \quad ((x^2 + y^2 - 0.85^2)^2 + (z^2 - 1)^2)*$$


$$\begin{aligned} & ((y^2 + z^2 - 0.85^2)^2 + (x^2 - 1)^2) * \\ & ((z^2 + x^2 - 0.85^2)^2 + (y^2 - 1)^2) - 0.001 = 0 \end{aligned}$$



$$\begin{aligned} & \{(x, y, z) \in \mathbb{R}^3 ; \quad (3(x-1)x^2(x+1) + 2y^2)^2 + (z^2 - 0.85)^2 * \\ & (3(y-1)y^2(y+1) + 2z^2)^2 + (x^2 - 0.85)^2 * \\ & (3(z-1)z^2(z+1) + 2x^2)^2 + (y^2 - 0.85)^2 * -0.12 = 0\} \end{aligned}$$



$$\begin{aligned} & \{(x, y, z) \in \mathbb{R}^3 ; \quad (2.92(x-1)x^2(x+1) + 1.7y^2)^2 * (y^2 - 0.88)^2 + \\ & (2.92(y-1)y^2(y+1) + 1.7z^2)^2 * (z^2 - 0.88)^2 + \\ & (2.92(z-1)z^2(z+1) + 1.7x^2)^2 * (x^2 - 0.88)^2 - 0.02 - 0.1 \} \end{aligned}$$



Hope you enjoy them...

Note also, that all of these examples have the form $\{(x, y, z) \in \mathbb{R}^3; P(x, y, z) = 0\} = P^{-1}(0)$. They are 2-manifolds (surfaces). To see this not only with your eyes but also **in theory**, compute the matrix

$$\left[\frac{\partial P}{\partial x}, \frac{\partial P}{\partial y}, \frac{\partial P}{\partial z} \right]$$

at all the points of $P^{-1}(0)$ and find out (haven't done this myself) that the matrix is non-zero, which by implicit function theorem means that $P^{-1}(0)$ is a $3 - 1 = 2$ manifold.

Also, if you change the definition of $\{(x, y, z) \in \mathbb{R}^3; P(x, y, z) = 0\} = P^{-1}(0)$ to

$$\{(x, y, z) \in \mathbb{R}^3; P(x, y, z) \leq 0\}$$

or

$$\{(x, y, z) \in \mathbb{R}^3; P(x, y, z) \geq 0\},$$

you get a 3-manifold (because 0 is a *regular value* of P) with the boundary $\{(x, y, z) \in \mathbb{R}^3; P(x, y, z) = 0\}$, i.e. the surface $P^{-1}(0)$ with either it's interior or exterior "filled".

ADDITION (how to construct such surfaces): @Soarer, You never try to guess such a complicated polynomial. Here's the key to constructing such surfaces: step by step. You discover on the web, that the torus is

$$\{P(x, y, z) := (x^2 + y^2 - 0.7^2)^2 + z^2 = 0\}.$$

Then you learn that

$$\{P(x - a, y - b, z - c) = 0\}$$

is torus, translated by $(a, b, c) \in \mathbb{R}^3$. Also, you notice, that

$$\{P(y, z, x) = 0\} \text{ and } \{P(z, x, y) = 0\}$$

is a torus with coordinate lines permuted (rotated by 90°) and


$$\{P(ax, by, cz) = 0\}$$

is a torus, stretched in x-direction by factor $1/a$, etc. Next, you see that

$$\{P(x, y, z) \cdot Q(x, y, z) = 0\} = \{P(x, y, z) = 0\} \cup \{Q(x, y, z) = 0\}.$$

Using all these techniques step by step, and being very patient, you manage to discover the following construction

```
implicitSurface[p_String, zoom_, quality_Integer, view_List] := 
ContourPlot3D[ToExpression[p], {x, -zoom, zoom}, {y, -zoom, zoom}, {z, -zoom, zoom}, Contours -> 0.02, Mesh -> Automatic, 
AspectRatio -> Automatic, Boxed -> False, Axes -> {False, False, False}, BoxRatios -> Automatic, PlotRange -> All, 
PlotRangePadding -> None, PlotPoints -> quality, ViewPoint -> view];
surfaces = {
implicitSurface["(x^2+y^2-0.7^2)^2+z^2", 1.2, 10, {0.3, -0.8, 1}],
implicitSurface["(x^2+y^2-0.7^2)^2+(z^2-1)^2-0.05", 1.2, 10, {1.3, -1.0, 2}],
implicitSurface["(1.2y^2-1)^2*(x^2+y^2-1^2)^2+",
(1.2z^2-1)^2*(y^2+z^2-1^2)^2+",
(1.2x^2-1)^2*(z^2+x^2-1^2)^2-0.02", 1.5, 10, {1.3, -1.0, 1.5}],
implicitSurface["((x^2+y^2-0.85^2)^2+(z^2-1)^2)*",
((y^2+z^2-0.85^2)^2+(x^2-1)^2)*",
((z^2+x^2-0.85^2)^2+(y^2-1)^2)-0.001", 1.2, 50, {1.2, -1.0, 1.3}],
implicitSurface["(3(x-1)x^2(x+1)+y^2)^2+2z^2", 1.2, 10, {0.3, -0.8, 1}],
implicitSurface["(3(x-1)x^2(x+1)+y^2)^2+(2z^2-1)^2-0.005", 1.2, 10, {0.3, -0.8, 1}],
implicitSurface["(3(x-1)x^2(x+1)+2y^2)^2+(z^2-0.85)^2*",
(3(y-1)y^2(y+1)+2z^2)^2+(x^2-0.85)^2*",
(3(z-1)z^2(z+1)+2x^2)^2+(y^2-0.85)^2*-0.12", 2, 80, {1, -1.8, 1.7}],
implicitSurface["(2.92(x-1)x^2(x+1)+1.7y^2)^2+(y^2-0.88)^2*",
(2.92(y-1)y^2(y+1)+1.7z^2)^2+(z^2-0.88)^2*",
(2.92(z-1)z^2(z+1)+1.7x^2)^2+(x^2-0.88)^2*-0.02", 2, 20, {0.4, -1.8, 1.3}]]
```



using the code (Mathematica 7):

```
implicitSurface[p_String, zoom_, quality_Integer, view_List] := 
ContourPlot3D[ToExpression[p], {x, -zoom, zoom}, {y, -zoom, zoom}, {z, -zoom, zoom}, Contours -> 0.02, Mesh -> Automatic, 
AspectRatio -> Automatic, Boxed -> False, Axes -> {False, False, False}, BoxRatios -> Automatic, PlotRange -> All, 
PlotRangePadding -> None, PlotPoints -> quality, ViewPoint -> view];
surfaces = {
implicitSurface["(x^2+y^2-0.7^2)^2+z^2", 1.2, 10, {0.3, -0.8, 1}],
implicitSurface["(x^2+y^2-0.7^2)^2+(z^2-1)^2-0.05",
1.2, 10, {1.3, -1.0, 2}],
implicitSurface["(1.2y^2-1)^2*(x^2+y^2-1^2)^2+",
(1.2z^2-1)^2*(y^2+z^2-1^2)^2+",
(1.2x^2-1)^2*(z^2+x^2-1^2)^2-0.02",
1.5, 10, {1.3, -1.0, 1.5}],
implicitSurface["((x^2+y^2-0.85^2)^2+(z^2-1)^2)*",
((y^2+z^2-0.85^2)^2+(x^2-1)^2)*",
((z^2+x^2-0.85^2)^2+(y^2-1)^2)-0.001",
1.2, 50, {1.2, -1.0, 1.3}],
implicitSurface["(3(x-1)x^2(x+1)+y^2)^2+2z^2",
1.2, 10, {0.3, -0.8, 1}],
implicitSurface["(3(x-1)x^2(x+1)+y^2)^2+(2z^2-1)^2-0.005",
1.2, 10, {0.3, -0.8, 1}],
implicitSurface["(3(x-1)x^2(x+1)+2y^2)^2+(z^2-0.85)^2*",
(3(y-1)y^2(y+1)+2z^2)^2+(x^2-0.85)^2*",
(3(z-1)z^2(z+1)+2x^2)^2+(y^2-0.85)^2*-0.12",
```

```

2, 80, {1, -1.8, 1.7}],
implicitSurface[{"(2.92(x-1)x^2(x+1)+1.7y^2)^2*(y^2-0.88)^2+
(2.92(y-1)y^2(y+1)+1.7z^2)^2*(z^2-0.88)^2+
(2.92(z-1)z^2(z+1)+1.7x^2)^2*(x^2-0.88)^2 -0.02",
2, 20, {0.4, -1.8, 1.3}]
}

```

Of course, some of the examples were obtained just with a lot of trying and intelligent guessing.

It is apparent that one can construct surfaces, as complicated as one wishes, via these steps, but each time a new component is added ($P^{-1}(0) \mapsto (P \cdot Q)^{-1}(0)$), the degree of the polynomial increases, which presents considerable numerical problems when plotting the surface.

P.S. I hope I haven't made any mistakes when I copied the code. If so, please inform me, to check with my original Mathematica file.

P.P.S The above surfaces were included, as examples, in my diploma.

Share Cite Follow

edited Jun 20, 2011 at 6:02

community wiki

11 revs

Leon Lampret

10 These are incredibly cool. – Leonardo Fontoura Jun 19, 2011 at 3:45

- 2 How did you write down the equations? Take your "first miscellaneous example" as an example, it's natural to imagine a surface like that, but I have no idea how to write down the equations.. – user325 Jun 19, 2011 at 4:52
- 1 @Soarer: I've added a fairly general "recipe" for construction. – Leo Jun 19, 2011 at 6:49
- 2 beautiful pictures! – Otis Aug 28, 2012 at 6:53
- 2 I can [reproduce](#) your last example but only by increasing the isolevel (e.g. 0.04 instead of 0.02). With 0.02, the result I get is not connex. – Stéphane Laurent Nov 23, 2018 at 8:01

▲ Reposting my comment as requested:

6 There is a wealth of documentation on Wolfram's Mathworld: <http://mathworld.wolfram.com/topics/Surfaces.html>

▼ I've also, not too long ago, seen a pdf file that had pictures and Mathematica code for a lot of these surfaces; if I can dig it out of my history I will post it as well.



Share Cite Follow

answered Jun 19, 2011 at 1:55

community wiki

barf




I like this "tetrahedron":

3




$$x^4 + 2x^2y^2 + 2x^2z^2 + y^4 + 2y^2z^2 + z^4 + 8xyz - 10x^2 - 10y^2 - 10z^2 + 20 = 0.$$

The Entzensberger star:




$$400(x^2y^2 + y^2z^2 + x^2z^2) = (1 - x^2 - y^2 - z^2)^3.$$

On the [higher space forum](#) you can find amazing implicit surfaces, especially those of the member **ICN5D**. These are 3D slices of surfaces in higher dimension. I reproduced some of them with Python, like this one:





Solid Möbius strip, introduced in [this paper](#):



Share Cite Follow

edited Apr 30 at 18:14

community wiki

4 revs

Stéphane Laurent



Complete minimal surfaces, discovered circa 1985. For example, [Meeks' surfaces](#).

2

Share Cite Follow

answered Jun 19, 2011 at 7:13

community wiki

Andrew



Those are really nice, but what are the implicit equations? – [Leo](#) Jun 19, 2011 at 8:17

Maybe this link will give some more explicit information: indiana.edu/~minimal/research/claynotes.pdf – [exchange](#) Nov 1, 2017 at 22:51