

Data Mining and Business Intelligence

Lecture 8: Exponential Smoothing Models

Jing Peng
University of Connecticut

3/12/20

Agenda

- Midterm Survey
- Online class due to coronavirus
- Review of Lecture 7
- Forecasting
- Exponential Smoothing Models
- Applications

Midterm Feedback

- Relate math to real-life cases and explain business insights
- Outline some key points that we should master
- Slow down a bit and spend more time on practices
- Explain concepts more thoroughly and slowly
- Less dig into math
- More concepts than tools. Dive deeper into specific topics
- Provide non-SAS material (e.g., Python)
- More assignments

Stationarity

- What are the three properties of stationary time series?
- Can a stationary time series exhibit trend or seasonality?

Stationarity of AR(1) Model

$$Y_t = \phi_0 + \phi_1 Y_{t-1} + \varepsilon_t$$

- Is the model stationary if $|\phi_1| > 1$?
- Is the model stationary if $\phi_1 = 1$?
- Is the model stationary if $\phi_1 = -1$?
- What if $|\phi_1| < 1$ but $\phi_0 > 1$?

Multiple AR.R

Stationarity of MA(1) Model

$$Y_t = \theta_0 + \theta_1 \varepsilon_{t-1} + \varepsilon_t$$

- Is the model stationary if $|\theta_1| > 1$?

Multiple MA.R



Which of the following statements are true?

- A random walk model has constant mean
- A random walk model has constant variance multiple AR.R
- A random walk with drift model has constant mean
- A random walk with drift model has constant variance

Forecasting

Statistical Forecasting — The Math


- Time Series = Signal + Noise

Systematic Variation Uncertainty

- Forecast = Extrapolated Signal
- Confidence Interval = Extrapolated Signal +/- Uncertainty
- Example: $Y_t = \phi_0 + \phi_1 Y_{t-1} + \varepsilon_t$

Forecast = Extrapolated Signal



Model Evaluation Metrics

- Root Mean Squared Error

$$\text{RMSE} = \sqrt{\frac{1}{n} \sum_{t=1}^n (Y_t - \hat{Y}_t)^2}$$

- Mean Absolute Percent Error:

$$\text{MAPE} = \frac{1}{n} \sum_{t=1}^n |Y_t - \hat{Y}_t| / Y_t$$

- Mean Absolute Error:

$$\text{MAE} = \frac{1}{n} \sum_{t=1}^n |Y_t - \hat{Y}_t|$$

Evaluation Metrics

- Akaike Information Criteria (AIC):

$$\text{AIC} = -2\log(L) + 2k$$

- Schwarz's Bayesian Information Criteria (SBC or BIC):

$$\text{SBC} = -2\log(L) + k\log(n)$$

- L represents the likelihood of the model
- k represents the number of parameters in the model
- n represents number of observations in the data

Evaluation Process

Fit Sample (Fit)

- Used to estimate model parameters for accuracy evaluation

Holdout Sample (Accuracy)

- Used to evaluate model accuracy



Full = Fit + Holdout data is used to fit a **final** deployment model

Choosing the Holdout Sample


- Rules of thumb
 - Cover at least a complete seasonal period in holdout sample
 - Length of fit sample is 4 times more than number of parameters
 - Holdout sample often contains no more than 25% of the series
 - The holdout sample is always at the end of the series
- No holdout sample and base accuracy on the entire series
 - If unique behavior occurs within the holdout sample
 - If there is insufficient data to fit a model without the holdout sample

Exponential Smoothing Models

Exponential Smoothing

- Proposed in the late 1950s (Brown, 1959; Holt, 1957; Winters, 1960)
- Produce forecasts using weighted averages of past observations, with the weights decaying **exponentially** as the observations get older.
- Support data with trend and/or seasonality


Two Special ESM Models



Weights applied to past values to predict Y_9

- Sample mean: equal weights
- Random walk: all weight to the most recent observation

Example: Predicting Y_7




Prediction by sample mean:
Prediction by random walk:

Simple Exponential Smoothing

$$\begin{aligned}\hat{Y}_{t+1} &= \omega Y_t + (1 - \omega) \hat{Y}_t \\ &= \omega Y_t + (1 - \omega) [\omega Y_{t-1} + (1 - \omega) \hat{Y}_{t-1}] \\ &= \omega Y_t + \omega(1 - \omega) Y_{t-1} + (1 - \omega)^2 \hat{Y}_{t-1} \\ &= \omega Y_t + \omega(1 - \omega) Y_{t-1} + (1 - \omega)^2 [\omega Y_{t-2} + (1 - \omega) \hat{Y}_{t-2}] \\ &= \omega Y_t + \omega(1 - \omega) Y_{t-1} + \omega(1 - \omega)^2 Y_{t-2} + \omega(1 - \omega)^3 Y_{t-3} + \dots\end{aligned}$$

Simple Exponential Smoothing



Weights applied to past values to predict Y_9

The larger the parameter, the more that the most recent values are emphasized.

What Happens if ω Approaches Zero

$$\begin{aligned}\hat{Y}_{t+1} &= \omega Y_t + (1 - \omega) \hat{Y}_t \\ &= \omega Y_t + (1 - \omega) [\omega Y_{t-1} + (1 - \omega) \hat{Y}_{t-1}] \\ &= \omega Y_t + \omega(1 - \omega) Y_{t-1} + (1 - \omega)^2 \hat{Y}_{t-1} \\ &= \omega Y_t + \omega(1 - \omega) Y_{t-1} + (1 - \omega)^2 [\omega Y_{t-2} + (1 - \omega) \hat{Y}_{t-2}] \\ &= \omega Y_t + \omega(1 - \omega) Y_{t-1} + \omega(1 - \omega)^2 Y_{t-2} + \omega(1 - \omega)^3 Y_{t-3} + \dots\end{aligned}$$

Since $1 - \omega \approx 1$, the ESM model degenerates to the sample mean model

ESM Parameters

ESM	Parameters	Component
Simple	ω	Level
Double	ω	Level/Trend
Linear (Holt)	ω, γ	Level,Trend
Damped-Trend	ω, γ, ϕ	Level,Trend,Damping

Smoothing Weights

ω Level smoothing weight

γ Trend smoothing weight

ϕ Trend damping weight

The screenshot shows a software window titled "Smoothing Model Specification". It contains the following fields and controls:

- Series:** VALUE
- Model:** Damped Trend Exponential Smoothing
- Smoothing Methods:** A list of checkboxes where "Damped-Trend Smoothing" is selected.
 - ☐ Simple Smoothing
 - ☐ Double (Brown) Smoothing
 - ☐ Seasonal Smoothing
 - ☐ Linear (Holt) Smoothing
 - ☒ Damped-Trend Smoothing
 - ☐ Winters Method - Additive
 - ☐ Winters Method - Multiplicative
- Transformation:** A dropdown menu currently showing a downward arrow.
- Smoothing Weights:** A section with a red border containing:
 - Level:** Optimize
 - Trend:** Optimize
 - Damping:** Optimize
 - Season:** Optimize
 - Bounds:** Zero-One/Additive (with a dropdown arrow)

At the bottom are buttons for OK, Cancel, Reset, Clear, and Help.

Simple Exponential Smoothing Predictions

$$\hat{Y}_1 = Y_0 \text{ (starting value)}$$

$$\hat{Y}_2 = \omega Y_1 + (1 - \omega) \hat{Y}_1$$

...

$$\hat{Y}_{t+1} = \omega Y_t + (1 - \omega) \hat{Y}_t$$

- The starting value Y_0 is often taken to be the mean of the first n observations. SAS TSFS uses $n=6$.
- Does not work well when there is a trend

Double Exponential (Holt) Smoothing

$$L_t = \omega Y_t + (1 - \omega)(L_{t-1} + T_{t-1}) \quad 0 \leq \omega \leq 1$$

Level equation

$$T_t = \gamma(L_t - L_{t-1}) + (1 - \gamma)T_{t-1} \quad 0 \leq \gamma \leq 1$$

Trend equation

$$\hat{Y}_{t+m} = L_t + mT_t \text{ (m-period-ahead forecast)}$$

Prediction equation


- Smoothing for both level and trend
- TSFS uses Double Exponential Smoothing to refer to a simpler model with only one smoothing parameter (see slide 22)
- Damped-trend smoothing: a third weight on T_{t-1}

For more details

<https://onlinecourses.science.psu.edu/stat501/node/363>

<http://www.itl.nist.gov/div898/handbook/pmc/section4/pmc4.htm>

Exponential Smoothing Models in the TSFS



continued...

Exponential Smoothing Models in the TSFS


The screenshot shows a dialog box titled "Smoothing Model Specification". It contains the following fields and options:

- Series:** LEADPROD: Lead Production in Tons
- Model:** (empty text box)
- Smoothing Methods:** A list of seven methods, each with an unchecked checkbox:
 - Simple Smoothing
 - Double (Brown) Smoothing
 - Seasonal Smoothing
 - Linear (Holt) Smoothing
 - Damped-Trend Smoothing
 - Winters Method - Additive
 - Winters Method - Multiplicative
- Transformation:** (empty text box) followed by a dropdown arrow icon.
- Smoothing Weights:** A group box containing five settings, each with a text box:
 - Level: Optimize
 - Trend: Optimize
 - Damping: Optimize
 - Season: Optimize
 - Bounds: Zero-One/Additive (with a dropdown arrow icon)
- Buttons:** OK, Cancel, Reset, Clear, and Help.

continued...

Seasonal Exponential Smoothing

ESM for Seasonal Data



Weights decay with respect to the seasonal factor.

ESM Parameters (Full)

ESM	Parameters
Simple	ω
Double	ω
Linear (Holt)	ω, γ
Damped-Trend	ω, γ, ϕ
Seasonal	ω, δ
Additive Winters	ω, γ, δ
Multiplicative Winters	ω, γ, δ

Smoothing Weights

 ω

Level smoothing weight

 γ

Trend smoothing weight

 δ

Seasonal smoothing weight

 ϕ

Trend damping weight

The choice of Greek letter is arbitrary. The software uses names rather than Greek symbols.

Winters Method — Additive

$$L_t = \omega(Y_t - S_{t-p}) + (1 - \omega)(L_{t-1} + T_{t-1})$$

Level equation

$$T_t = \gamma(L_t - L_{t-1}) + (1 - \gamma)T_{t-1}$$

Trend equation

$$S_t = \delta(Y_t - L_t) + (1 - \delta)S_{t-p}$$

Seasonality equation

$$Y_{t+m} = L_t + mT_t + S_t \text{ (m-period-ahead forecast)}$$

Prediction equation

p is the period of seasonality

Winters Method — Multiplicative

$$L_t = \omega(Y_t/S_{t-p}) + (1 - \omega)(L_{t-1} + T_{t-1})$$

Level equation

$$T_t = \gamma(L_t - L_{t-1}) + (1 - \gamma)T_{t-1}$$

Trend equation

$$S_t = \delta(Y_t/L_t) + (1 - \delta)S_{t-p}$$

Seasonality equation

$$Y_{t+m} = (L_t + mT_t)S_t \text{ (m-period-ahead forecast)}$$

Prediction equation

p is the period of seasonality

Seasonal Exponential Smoothing Models in the TSFS

Smoothing Model Specification

Series:

Model:

Smoothing Methods:

- ☐ Simple Smoothing
- ☐ Double (Brown) Smoothing
- ☒ Seasonal Smoothing
- ☐ Linear (Holt) Smoothing
- ☐ Damped-Trend Smoothing
- ☐ Winters Method - Additive
- ☐ Winters Method - Multiplicative

Transformation: ▼

Smoothing Weights:

Level:

Trend:

Damping:

Season:

Bounds: ▼

OK Cancel Reset Clear Help

continued...

Applications

Fitting ESM for LEADPRD Data

- Models
 - Fit models automatically (use “Diagnose Series” to force SAS to fit all 42 predefined models)
 - Try all non-seasonal ESM models
 - Fit AR(1) and MA(1) models
- Time Series Diagnostics
- Model Residuals Diagnostics

Fitting Seasonal ESM for CONTRACTS Data

- Fit models automatically
- Fit seasonal ESM models
- Setup holdout sample and prediction horizon
- Change evaluation metrics and compare models
- Save predictions

When to Use?

- The prediction of ESM models quickly converges to the mean of time series (plus trend adjustment), not ideal for long-term prediction
 - Simple ESM: short-term prediction
 - Complicated ESM: moderate-term prediction

Readings

- Forecasting Chapter 2
- <http://www.itl.nist.gov/div898/handbook/pmc/section4/pmc4.htm>
- <https://onlinecourses.science.psu.edu/stat501/node/363>
- <https://people.duke.edu/~rnau/411avg.htm>