

Course notes from MITx 14.310x Data Analysis for
Social Scientists (EdX)

James Solomon-Rounce

2018-09-18

Contents

Preface	5
1 Module 1: Introduction to the Course	7
1.1 Introduction to R	7
2 Module 2: Fundamentals of Probability, Random Variables, Joint Distributions + Collecting Data	11
2.1 Fundamentals of Probability	11


Preface

The following notes were taken by me for educational, non-commercial, purposes. If you find the information useful, buy the material/take the course.

Thank you to the original content providers. Additional ramblings are my own.

Core Resources

- Course Schedule
- Grading and Homework Policy
- Honor Code and Collaboration Guide
- Notes - OLS Derivation
- Notes - Matrix Notation
- R Studio Cheatsheets
- R for Data Science Book



Chapter 1

Module 1: Introduction to the Course

Module Sections:

- Welcome to the Course
- Introduction to R
- Introductory Lecture - Data is Beautiful, Insightful, Powerful, Deceitful
- Finger Exercises
- Module 1: Homework

Module Content:

- Module 1 Slides
- Homework 1 Background Paper - The Persistent Effects Of Peru's Mining Mita
- R Instructions
- Intro to R Zip File
- Citations Data for Homework 1

1.1 Introduction to R

First we setup the environment and install the course files

```
library(swirl)
install_course_zip("./files/M1/14_310x_Intro_to_R_.zip",multi=FALSE)
swirl()
```

IF z is a three number vector e.g.

```
z <- c(1.1, 9, 3.14)
```

If we take the square root of z - 1 and assign it to a new variable called my_sqrt e.g.

```
my_sqrt <- sqrt(z - 1)
```

The result is a vector of length three e.g.

```
my_sqrt
```

```
## [1] 0.3162278 2.8284271 1.4628739
```

Next, if we create a new variable called `my_div` that gets the value of `z` divided by `my_sqrt`.

```
my_div <- z / my_sqrt
```

The first element of `my_div` is equal to the first element of `z` divided by the first element of `my_sqrt`, and so on...

```
my_div
```

```
## [1] 3.478505 3.181981 2.146460
```

When given two vectors of the same length, R simply performs the specified arithmetic operation (+, -, *, etc.) element-by-element. If the vectors are of different lengths, R ‘recycles’ the shorter vector until it is the same length as the longer vector.

If the length of the shorter vector does not divide evenly into the length of the longer vector, R will still apply the ‘recycling’ method, but will throw a warning.

```
c(1, 2, 3, 4) + c(0, 10, 100)
```

```
## Warning in c(1, 2, 3, 4) + c(0, 10, 100): longer object length is not a
## multiple of shorter object length
```

```
## [1] 1 12 103 4
```

1.1.1 Module 1 Homework

This is a sample of some of the homework answers. Some questions were observational or required interpretation of maps for example, as such these are not included here.

```
library(tidyverse)
```

```
## -- Attaching packages ----- tidyverse 1.2.1 --
```

```
## v ggplot2 3.0.0      v purrr 0.2.5
## v tibble 1.4.2       v dplyr 0.7.6
## v tidyr 0.8.1       v stringr 1.3.1
## v readr 1.1.1       v forcats 0.3.0
```

```
## -- Conflicts ----- tidyverse_conflicts() --
```

```
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()     masks stats::lag()
```

```
papers <- as_tibble(read_csv("./files/M1/CitesforSara.csv"))
```

```
## Parsed with column specification:
```

```
## cols(
##   .default = col_integer(),
##   journal = col_character(),
##   title = col_character(),
##   au1 = col_character(),
##   au2 = col_character(),
##   au3 = col_character(),
##   past5 = col_double(),
##   aflpn90 = col_double(),
##   aulpn90 = col_double(),
##   aulpn80 = col_double(),
##   aulpn70 = col_double(),
##   lcites = col_double()
```



```
## )
```

```
## See spec(...) for full column specifications.
```

Q. 19 Let's take a look at some of the most popular papers. Using the `filter()` method, how many records exist where there are greater than or equal to 100 citations?

```
#First lets look at our data
```

```
head(papers)
```

```
## # A tibble: 6 x 22
##   journal year cites title au1   au2   au3   female1 female2 female3 page
##   <chr>   <int> <int> <chr> <chr> <chr> <chr>   <int>   <int>   <int> <int>
## 1 Americ~ 1993    31 Jeux~ Kanb~ Keen~ <NA>     0     0    NA    16
## 2 Americ~ 1993     4 Chan~ Jame~ <NA> <NA>     0    NA    NA    22
## 3 Americ~ 1993    28 Fact~ Bert~ <NA> <NA>     0    NA    NA    15
## 4 Americ~ 1993    10 Stra~ Garf~ Oh,-- <NA>     1     0    NA    19
## 5 Americ~ 1993     5 Will~ Coat~ Lour~ <NA>     0     0    NA    21
## 6 Americ~ 1993    21 Merg~ Kim,~ Sing~ <NA>     0     0    NA    21
## # ... with 11 more variables: order <int>, nauthor <int>, past5 <dbl>,
## #   aflpn90 <dbl>, spage <int>, field <int>, subfld <int>, aulpn90 <dbl>,
## #   aulpn80 <dbl>, aulpn70 <dbl>, lcites <dbl>
```

```
arrange(papers,desc(cites), title)
```

```
## # A tibble: 4,182 x 22
##   journal year cites title au1   au2   au3   female1 female2 female3
##   <chr>   <int> <int> <chr> <chr> <chr> <chr>   <int>   <int>   <int>
## 1 Econom~ 1980  2251 A He~ Whit~ <NA> <NA>     0    NA    NA
## 2 Econom~ 1979  2227 Pros~ Kahn~ Tver~ <NA>     0     0    NA
## 3 Econom~ 1987  2164 Co-i~ Engl~ Gran~ <NA>     0     0    NA
## 4 Econom~ 1979  1602 Samp~ Heck~ <NA> <NA>     0    NA    NA
## 5 Econom~ 1978  1217 Spec~ Haus~ <NA> <NA>     0    NA    NA
## 6 Econom~ 1982  1077 Auto~ Engl~ <NA> <NA>     0    NA    NA
## 7 Econom~ 1981  1031 Like~ Dick~ Full~ <NA>     0     0    NA
## 8 Econom~ 1982   983 Larg~ Hans~ <NA> <NA>     0    NA    NA
## 9 Econom~ 1980   864 Macr~ Sims~ <NA> <NA>     0    NA    NA
## 10 Econom~ 1982   563 Time~ Kydl~ Pres~ <NA>     0     0    NA
## # ... with 4,172 more rows, and 12 more variables: page <int>,
## #   order <int>, nauthor <int>, past5 <dbl>, aflpn90 <dbl>, spage <int>,
## #   field <int>, subfld <int>, aulpn90 <dbl>, aulpn80 <dbl>,
## #   aulpn70 <dbl>, lcites <dbl>
```

```
papers %>%
```

```
  filter(cites >= 100)
```

```
## # A tibble: 205 x 22
##   journal year cites title au1   au2   au3   female1 female2 female3
##   <chr>   <int> <int> <chr> <chr> <chr> <chr>   <int>   <int>   <int>
## 1 Americ~ 1994   117 Is I~ Pers~ Tabe~ <NA>     0     0    NA
## 2 Econom~ 1971   149 Furt~ Nerl~ <NA> <NA>     0    NA    NA
## 3 Econom~ 1971   170 The ~ Madd~ <NA> <NA>    NA    NA    NA
## 4 Econom~ 1971   155 Inve~ Luca~ Pres~ <NA>     0     0    NA
## 5 Econom~ 1971   139 Some~ Crag~ <NA> <NA>     0    NA    NA
## 6 Econom~ 1971   108 Iden~ Roth~ <NA> <NA>     0    NA    NA
## 7 Econom~ 1972   164 Meth~ Fair~ Jaff~ <NA>     0     0    NA
## 8 Econom~ 1972   150 Exis~ Radn~ <NA> <NA>     0    NA    NA
```

```
## 9 Econom~ 1973 361 Mani~ Gibb~ <NA> <NA> 0 NA NA
## 10 Econom~ 1973 107 On a~ Kram~ <NA> <NA> 0 NA NA
## # ... with 195 more rows, and 12 more variables: page <int>, order <int>,
## #   nauthor <int>, past5 <dbl>, aflpn90 <dbl>, spage <int>, field <int>,
## #   subfld <int>, aulpn90 <dbl>, aulpn80 <dbl>, aulpn70 <dbl>,
## #   lcites <dbl>
```

Q.20 Use the `group_by()` function to group papers by journal. How many total citations exist for the journal “Econometrica”?

```
papers %>%
  group_by(journal) %>%
  summarise(sum(cites))

## # A tibble: 7 x 2
##   journal                `sum(cites)`
##   <chr>                  <int>
## 1 American-Economic-Review      3738
## 2 Econometrica                75789
## 3 Journal-of-Political-Economy   3398
## 4 Quarterly-Journal-of-Economics 8844
## 5 Review-of-Economic-Studies    21937
## 6 Review-of-Economics-and-Statistics 8473
## 7 <NA>                        14
```

```
#or

summarize(group_by
  (papers, journal),
  SumOfCites = sum(cites))
```

```
## # A tibble: 7 x 2
##   journal                SumOfCites
##   <chr>                  <int>
## 1 American-Economic-Review      3738
## 2 Econometrica                75789
## 3 Journal-of-Political-Economy   3398
## 4 Quarterly-Journal-of-Economics 8844
## 5 Review-of-Economic-Studies    21937
## 6 Review-of-Economics-and-Statistics 8473
## 7 <NA>                        14
```

Q.21 How many distinct primary authors (au1) exist in this dataset?

```
papers %>%
  summarise(n_distinct(au1))
```

```
## # A tibble: 1 x 1
##   `n_distinct(au1)`
##   <int>
## 1          2332
```

```
#or

n_distinct(papers$au1)
```

```
## [1] 2332
```

Chapter 2

Module 2: Fundamentals of Probability, Random Variables, Joint Distributions + Collecting Data

Module Sections:

- Fundamentals of Probability
- Random Variables, Distributions, and Joint Distributions
- Gathering and Collecting Data
- Module 2: Homework

Module Content:

- Module 2 Slides

2.1 Fundamentals of Probability

2.1.1 Set Theory

- A *sample space* is collection of all possible outcomes
- An *event* is any collection of outcomes - could be one, all or none
- If the outcome is a member of an event, the event is said to have *occured*
- Event B is said to be *contained* by A, if all outcomes in B also are in A
- This is the basis of set theory and used widely in probability, although there are some differences between set and probability theory
- If there is no symbol, then this usually means intersection AB in probability - in set theory we would write an inverted U e.g. $A \cap B$
- A and B are mutually exclusive (disjoint in set theory) if they have no outcomes in common
- A and B are exhaustive (complimentary in set theory) if their union is S (the entire sample space)
- A and B are both mutually exclusive and exhaustive, their union is equal to the sample space but they have no events in common - they are a partition of the sample space

2.1.2 Defining Probability

We assign every event a number $P(A)$ which is the prob. the event will occur

1 We require that the probability is greater than one for all events in the sample space - $P(A) \geq 0$ for all $A \subset S$
 2 The entire sample space must be equal to one - $P(S) = 1$
 3 For any sequence of disjoint sets, the prob. of the union of that sequence is equal to the sum of the probabilities of those events - A_1, A_2, \dots , is $P(\cup_i A_i) = \sum_i P(A_i)$

So we have a sample space, and if it satisfies these three properties, then we call it a probability. Sometimes this is referred to as a probability function or a probability distribution, but there is no standard terminology for all probability theory. Set theory helps to prove aspects of probability mathematically, for the purposes on this course, we just need to know what some useful facts are.

- $P(A^c) = 1 - P(A) =$

The probability of A complement, which is the event that contains all of the outcomes that are not in the event A, the probability of A complement is just equal to 1 minus the probability of A. This is useful if the probability of A complement ($P(A^c)$) is difficult to compute, where as the probability of A might be very easy to compute.

- $P(\emptyset) =$

The probability of the empty set is zero.

- If $A \subset B$ then $P(A) \leq P(B) =$

If A is contained in B then the probability of A is less than or equal to the probability of B

- For all A, $0 \leq P(A) \leq 1 =$

For any events, the probability of that event is between 0 and 1.

- $P(A \cup B) = P(A) + P(B) - P(AB) =$

Probability of A union B is just equal to the sum of the probabilities of those two events minus the probability of their intersection.

- $P(AB^c) = P(A) - P(AB) =$

The probability of A times B complement is equal to the probability of A minus the probability of the intersection.