



Course > Unit 7: ... > Lec. 17: ... > 16. Exe...

16. Exercise: Choice of representations

Exercises due Apr 15, 2020 05:29 IST Completed

Exercise: Choice of representations

1/1 point (graded)

We wish to estimate an unknown quantity Θ . Our measuring equipment produces an observation of the form $X = \Theta^3 + W$, where W is a noise term which is small relative to the range of Θ . Which type of linear estimator is preferable in such a situation?

☐ $\hat{\Theta} = aX + b$

☐ $\hat{\Theta} = aX^3 + b$

☒ $\hat{\Theta} = aX^{1/3} + b$



Solution:

If the noise W were completely absent, we would estimate Θ by letting $\hat{\Theta} = X^{1/3}$. In the presence of small noise, our estimator should again have a similar form, which argues in favor of the third option.

Submit

You have used 1 of 1 attempt

Answers are displayed within the problem



Discussion

Hide Discussion

Topic: Unit 7: Bayesian inference:Lec. 17: Linear least mean squares (LLMS) estimation / 16. Exercise: Choice of representations

Show all posts ▼

by recent activity ▼

💬 Don't overthink this question.
And may the force be with you.

15

💬 Would a more generic estimator be better than all the 3 options?
In the previous lecture segment, the professor said that a more general class of estimators usually gives ...

1

💬 hint
"W is a noise term which **is small relative to the range**"

2

© All Rights Reserved

