

# 3. General Random Variables

## Part V: Conditioning

ECE 302 Fall 2009 TR 3-4:15pm

Purdue University, School of ECE

Prof. Ilya Pollak

# Conditioning a continuous random variable on an event

- Conditional PDF of a continuous random variable  $X$ , conditioned on an event  $A$  with  $\mathbf{P}(A)>0$ :

$$\mathbf{P}(X \in B \mid A) = \int_B f_{X|A}(x) dx,$$

for "any"  $B \subset R$

# Computing the conditional PDF of $X$ from the marginal PDF of $X$

$$\mathbf{P}(X \in B \mid A) = \int_B f_{X \mid A}(x) dx,$$

If  $A = \{X \in C\}$  where  $C \subset R$ , then, on the one hand,

$$\mathbf{P}(X \in B \mid X \in C) = \int_B f_{X \mid X \in C}(x) dx$$

# Computing the conditional PDF of $X$ from the marginal PDF of $X$

$$\mathbf{P}(X \in B \mid A) = \int_B f_{X|A}(x) dx,$$

If  $A = \{X \in C\}$  where  $C \subset R$ , then, on the one hand,

$$\mathbf{P}(X \in B \mid X \in C) = \int_B f_{X|X \in C}(x) dx$$

On the other hand,

$$\mathbf{P}(X \in B \mid X \in C) = \frac{\mathbf{P}(\{X \in B\} \cap \{X \in C\})}{\mathbf{P}(\{X \in C\})}$$

# Computing the conditional PDF of $X$ from the marginal PDF of $X$

$$\mathbf{P}(X \in B | A) = \int_B f_{X|A}(x) dx,$$

If  $A = \{X \in C\}$  where  $C \subset R$ , then, on the one hand,

$$\mathbf{P}(X \in B | X \in C) = \int_B f_{X|X \in C}(x) dx$$

On the other hand,

$$\mathbf{P}(X \in B | X \in C) = \frac{\mathbf{P}(\{X \in B\} \cap \{X \in C\})}{\mathbf{P}(\{X \in C\})} = \frac{\mathbf{P}(X \in B \cap C)}{\mathbf{P}(X \in C)}$$

# Computing the conditional PDF of $X$ from the marginal PDF of $X$

$$\mathbf{P}(X \in B \mid A) = \int_B f_{X \mid A}(x) dx,$$

If  $A = \{X \in C\}$  where  $C \subset R$ , then, on the one hand,

$$\mathbf{P}(X \in B \mid X \in C) = \int_B f_{X \mid X \in C}(x) dx$$

On the other hand,

$$\mathbf{P}(X \in B \mid X \in C) = \frac{\mathbf{P}(\{X \in B\} \cap \{X \in C\})}{\mathbf{P}(\{X \in C\})} = \frac{\mathbf{P}(X \in B \cap C)}{\mathbf{P}(X \in C)} = \frac{\int_{B \cap C} f_X(x) dx}{\mathbf{P}(X \in C)}$$

# Computing the conditional PDF of $X$ from the marginal PDF of $X$

$$\mathbf{P}(X \in B \mid A) = \int_B f_{X|A}(x) dx,$$

If  $A = \{X \in C\}$  where  $C \subset R$ , then, on the one hand,

$$\mathbf{P}(X \in B \mid X \in C) = \int_B f_{X|X \in C}(x) dx$$

On the other hand,

$$\mathbf{P}(X \in B \mid X \in C) = \frac{\mathbf{P}(\{X \in B\} \cap \{X \in C\})}{\mathbf{P}(\{X \in C\})} = \frac{\mathbf{P}(X \in B \cap C)}{\mathbf{P}(X \in C)} = \frac{\int_{B \cap C} f_X(x) dx}{\mathbf{P}(X \in C)}$$

$$\text{Therefore, } f_{X|X \in C}(x) = \begin{cases} \frac{f_X(x)}{\mathbf{P}(X \in C)}, & \text{if } x \in C \\ 0, & \text{otherwise} \end{cases}$$

# Computing the conditional PDF of $X$ from the marginal PDF of $X$

$$\mathbf{P}(X \in B \mid A) = \int_B f_{X|A}(x) dx,$$


If  $A = \{X \in C\}$  where  $C \subset \mathcal{R}$ , then, on the one hand,

$$\mathbf{P}(X \in B \mid X \in C) = \int_B f_{X|X \in C}(x) dx$$

On the other hand,

$$\mathbf{P}(X \in B \mid X \in C) = \frac{\mathbf{P}(\{X \in B\} \cap \{X \in C\})}{\mathbf{P}(\{X \in C\})} = \frac{\mathbf{P}(X \in B \cap C)}{\mathbf{P}(X \in C)} = \frac{\int_{B \cap C} f_X(x) dx}{\mathbf{P}(X \in C)}$$

$$\text{Therefore, } f_{X|X \in C}(x) = \begin{cases} \frac{f_X(x)}{\mathbf{P}(X \in C)}, & \text{if } x \in C \\ 0, & \text{otherwise} \end{cases}$$





## Example 3.13: memorylessness of the exponential random variable

- Time  $T$  between two successive buses is an exponential r.v. with mean  $1/\lambda$
- You arrive  $t$  seconds after the last bus. Event  $A = \{T > t\} = \{\text{there has been no buses in the } t \text{ seconds since the last bus}\}$
- $X = T - t$  the time you have to wait until the next bus
- Find  $F_{X|A}(x)$ .

## Example 3.13: memorylessness of the exponential random variable

$$f_T(t) = \begin{cases} \lambda e^{-\lambda t}, & \text{for } t > 0 \\ 0, & \text{for } t \leq 0 \end{cases}$$

$$F_T(t) = \begin{cases} 1 - e^{-\lambda t}, & \text{for } t > 0 \\ 0, & \text{for } t \leq 0 \end{cases}$$

## Example 3.13: memorylessness of the exponential random variable

$$f_T(t) = \begin{cases} \lambda e^{-\lambda t}, & \text{for } t > 0 \\ 0, & \text{for } t \leq 0 \end{cases} \quad F_T(t) = \begin{cases} 1 - e^{-\lambda t}, & \text{for } t > 0 \\ 0, & \text{for } t \leq 0 \end{cases}$$

$$\begin{aligned} \mathbf{P}(X > x \mid A) &= \mathbf{P}(T > t + x \mid T > t) = \frac{\mathbf{P}(T > t + x, T > t)}{\mathbf{P}(T > t)} = \frac{\mathbf{P}(T > t + x)}{\mathbf{P}(T > t)} \\ &= \frac{e^{-\lambda(t+x)}}{e^{-\lambda t}} = e^{-\lambda x} \end{aligned}$$

## Example 3.13: memorylessness of the exponential random variable

$$f_T(t) = \begin{cases} \lambda e^{-\lambda t}, & \text{for } t > 0 \\ 0, & \text{for } t \leq 0 \end{cases} \quad F_T(t) = \begin{cases} 1 - e^{-\lambda t}, & \text{for } t > 0 \\ 0, & \text{for } t \leq 0 \end{cases}$$

$$\begin{aligned} \mathbf{P}(X > x \mid A) &= \mathbf{P}(T > t + x \mid T > t) = \frac{\mathbf{P}(T > t + x, T > t)}{\mathbf{P}(T > t)} = \frac{\mathbf{P}(T > t + x)}{\mathbf{P}(T > t)} \\ &= \frac{e^{-\lambda(t+x)}}{e^{-\lambda t}} = e^{-\lambda x} \end{aligned}$$

$$F_{X|A}(x) = 1 - \mathbf{P}(X > x \mid A) = \begin{cases} 1 - e^{-\lambda x}, & x > 0 \\ 0, & x \leq 0 \end{cases}$$

This distribution is exponential with parameter  $\lambda$ , regardless of the time  $t$  that elapsed between the preceding bus and your arrival.

## Example 3.13: memorylessness of the exponential random variable

$$f_T(t) = \begin{cases} \lambda e^{-\lambda t}, & \text{for } t > 0 \\ 0, & \text{for } t \leq 0 \end{cases} \quad F_T(t) = \begin{cases} 1 - e^{-\lambda t}, & \text{for } t > 0 \\ 0, & \text{for } t \leq 0 \end{cases}$$

$$\begin{aligned} \mathbf{P}(X > x \mid A) &= \mathbf{P}(T > t + x \mid T > t) = \frac{\mathbf{P}(T > t + x, T > t)}{\mathbf{P}(T > t)} = \frac{\mathbf{P}(T > t + x)}{\mathbf{P}(T > t)} \\ &= \frac{e^{-\lambda(t+x)}}{e^{-\lambda t}} = e^{-\lambda x} \end{aligned}$$

$$F_{X|A}(x) = 1 - \mathbf{P}(X > x \mid A) = \begin{cases} 1 - e^{-\lambda x}, & x > 0 \\ 0, & x \leq 0 \end{cases}$$

This distribution is exponential with parameter  $\lambda$ , regardless of the time  $t$  that elapsed between the preceding bus and your arrival.

A process like this where inter-arrival times are exponentially distributed, is called a *Poisson process* (to be studied later in the course).

# Total Probability Theorem for PDFs

Recall that, according to the total probability theorem,  
if  $A_1, \dots, A_n$  partition the sample space, then

$$\mathbf{P}(B) = \sum_{i=1}^n \mathbf{P}(B \mid A_i) \mathbf{P}(A_i)$$

# Total Probability Theorem for PDFs

Recall that, according to the total probability theorem, if  $A_1, \dots, A_n$  partition the sample space, then

$$\mathbf{P}(B) = \sum_{i=1}^n \mathbf{P}(B \mid A_i) \mathbf{P}(A_i)$$

Applying this to the event  $B = \{X \leq x\}$ , we have:

$$\mathbf{P}(X \leq x) = \sum_{i=1}^n \mathbf{P}(X \leq x \mid A_i) \mathbf{P}(A_i)$$

# Total Probability Theorem for PDFs

Recall that, according to the total probability theorem, if  $A_1, \dots, A_n$  partition the sample space, then

$$\mathbf{P}(B) = \sum_{i=1}^n \mathbf{P}(B \mid A_i) \mathbf{P}(A_i)$$

Applying this to the event  $B = \{X \leq x\}$ , we have:

$$\mathbf{P}(X \leq x) = \sum_{i=1}^n \mathbf{P}(X \leq x \mid A_i) \mathbf{P}(A_i)$$

Differentiating both sides with respect to  $x$ , we obtain:

$$f_X(x) = \sum_{i=1}^n f_{X|A_i}(x) \mathbf{P}(A_i)$$



# Conditioning a continuous r.v. on another continuous r.v.

We would like to define  $f_{X|Y}(x|y)$  to be the conditional probability mass of  $X$  per unit length, given that  $Y = y$ .

# Conditioning a continuous r.v. on another continuous r.v.

We would like to define  $f_{X|Y}(x|y)$  to be the conditional probability mass of  $X$  per unit length, given that  $Y = y$ . In other words, we would like:

$$f_{X|Y}(x|y) \cdot \delta \approx \mathbf{P}(x \leq X \leq x + \delta | y \leq Y \leq y + \delta)$$

# Conditioning a continuous r.v. on another continuous r.v.

We would like to define  $f_{X|Y}(x|y)$  to be the conditional probability mass of  $X$  per unit length, given that  $Y = y$ . In other words, we would like:

$$\begin{aligned} f_{X|Y}(x|y) \cdot \delta &\approx \mathbf{P}(x \leq X \leq x + \delta | y \leq Y \leq y + \delta) \\ &= \frac{\mathbf{P}(x \leq X \leq x + \delta, y \leq Y \leq y + \delta)}{\mathbf{P}(y \leq Y \leq y + \delta)} \end{aligned}$$

# Conditioning a continuous r.v. on another continuous r.v.

We would like to define  $f_{X|Y}(x|y)$  to be the conditional probability mass of  $X$  per unit length, given that  $Y = y$ . In other words, we would like:

$$\begin{aligned} f_{X|Y}(x|y) \cdot \delta &\approx \mathbf{P}(x \leq X \leq x + \delta | y \leq Y \leq y + \delta) \\ &= \frac{\mathbf{P}(x \leq X \leq x + \delta, y \leq Y \leq y + \delta)}{\mathbf{P}(y \leq Y \leq y + \delta)} \approx \frac{f_{X,Y}(x,y) \cdot \delta^2}{f_Y(y) \cdot \delta} = \frac{f_{X,Y}(x,y)}{f_Y(y)} \cdot \delta \end{aligned}$$

# Conditioning a continuous r.v. on another continuous r.v.

We would like to define  $f_{X|Y}(x|y)$  to be the conditional probability mass of  $X$  per unit length, given that  $Y = y$ . In other words, we would like:

$$\begin{aligned} f_{X|Y}(x|y) \cdot \delta &\approx \mathbf{P}(x \leq X \leq x + \delta | y \leq Y \leq y + \delta) \\ &= \frac{\mathbf{P}(x \leq X \leq x + \delta, y \leq Y \leq y + \delta)}{\mathbf{P}(y \leq Y \leq y + \delta)} \approx \frac{f_{X,Y}(x,y) \cdot \delta^2}{f_Y(y) \cdot \delta} = \frac{f_{X,Y}(x,y)}{f_Y(y)} \cdot \delta \end{aligned}$$

This motivates the following definition:

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)}, \text{ for all } y \text{ such that } f_Y(y) \neq 0.$$

# Conditioning and independence

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)}, \text{ for all } y \text{ such that } f_Y(y) \neq 0.$$

If  $X$  and  $Y$  are independent, then  $f_{X,Y}(x,y) = f_X(x)f_Y(y)$

# Conditioning and independence

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)}, \text{ for all } y \text{ such that } f_Y(y) \neq 0.$$

If  $X$  and  $Y$  are independent, then  $f_{X,Y}(x,y) = f_X(x)f_Y(y)$ , and therefore  $f_{X|Y}(x|y) = f_X(x)$  whenever  $f_{X|Y}(x|y)$  is defined.

# Conditioning and independence

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)}, \text{ for all } y \text{ such that } f_Y(y) \neq 0.$$

If  $X$  and  $Y$  are independent, then  $f_{X,Y}(x,y) = f_X(x)f_Y(y)$ , and therefore  $f_{X|Y}(x|y) = f_X(x)$  whenever  $f_{X|Y}(x|y)$  is defined.

Similarly, if  $X$  and  $Y$  are independent, then

$$f_{Y|X}(y|x) = f_Y(y) \text{ whenever } f_{Y|X}(y|x) \text{ is defined.}$$



# Conditional Expectation

If  $A$  is an event and  $X$  is a r.v., then

$$E[X | A] = \begin{cases} \int_{-\infty}^{\infty} x f_{X|A}(x) dx & \text{if } X \text{ is a continuous r.v.} \\ \sum_x x p_{X|A}(x) & \text{if } X \text{ is a discrete r.v.} \end{cases}$$

# Conditional Expectation

If  $A$  is an event and  $X$  is a r.v., then

$$E[X | A] = \begin{cases} \int_{-\infty}^{\infty} xf_{X|A}(x)dx & \text{if } X \text{ is a continuous r.v.} \\ \sum_x xp_{X|A}(x) & \text{if } X \text{ is a discrete r.v.} \end{cases}$$

If  $X$  and  $Y$  are r.v.'s, then

$$E[X | Y = y] = \begin{cases} \int_{-\infty}^{\infty} xf_{X|Y}(x | y)dx & \text{if, given } Y=y, X \text{ is a continuous r.v.} \\ \sum_x xp_{X|Y}(x | y) & \text{if, given } Y=y, X \text{ is a discrete r.v.} \end{cases}$$

# Example


$$f_{X,Y}(x,y) = \begin{cases} 3x, & 0 \leq y \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

Find  $f_X(x)$  and  $f_{Y|X}(y|x)$ .

# Example

$$f_{X,Y}(x,y) = \begin{cases} 3x, & 0 \leq y \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases}$$


Find  $f_X(x)$  and  $f_{Y|X}(y|x)$ .



# Example

$$f_{X,Y}(x,y) = \begin{cases} 3x, & 0 \leq y \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

Find  $f_X(x)$  and  $f_{Y|X}(y|x)$ .




To find  $f_X(x_0)$ , integrate along the orange line.

# Example

$$f_{X,Y}(x,y) = \begin{cases} 3x, & 0 \leq y \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

Find  $f_X(x)$  and  $f_{Y|X}(y|x)$ .



To find  $f_X(x_0)$ , integrate along the orange line.

$$f_X(x_0) = \int_{-\infty}^{\infty} f_{X,Y}(x_0, y) dy = \begin{cases} \int_0^{x_0} 3x_0 dy = 3x_0^2, & 0 \leq x_0 \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

# Example

$$f_{X,Y}(x,y) = \begin{cases} 3x, & 0 \leq y \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases} \quad f_X(x) = \begin{cases} 3x^2, & 0 \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

To find  $f_{Y|X}(y|x)$ , use the definition:

$$f_{Y|X}(y|x) = \frac{f_{X,Y}(x,y)}{f_X(x)}$$

# Example

$$f_{X,Y}(x,y) = \begin{cases} 3x, & 0 \leq y \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases} \quad f_X(x) = \begin{cases} 3x^2, & 0 \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

To find  $f_{Y|X}(y|x)$ , use the definition:

$$f_{Y|X}(y|x) = \frac{f_{X,Y}(x,y)}{f_X(x)}$$

This is only defined for the values of  $x$  where  $f_X(x) \neq 0$ , that is, for  $0 < x \leq 1$ .



# Example

$$f_{X,Y}(x,y) = \begin{cases} 3x, & 0 \leq y \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases} \quad f_X(x) = \begin{cases} 3x^2, & 0 \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

To find  $f_{Y|X}(y|x)$ , use the definition:

$$f_{Y|X}(y|x) = \frac{f_{X,Y}(x,y)}{f_X(x)}$$

This is only defined for the values of  $x$  where  $f_X(x) \neq 0$ , that is, for  $0 < x \leq 1$ . For these values of  $x$ , we have:

$$f_{Y|X}(y|x) = \frac{f_{X,Y}(x,y)}{f_X(x)} = \begin{cases} \frac{3x}{3x^2} = \frac{1}{x}, & 0 \leq y \leq x \\ 0, & \text{otherwise} \end{cases}$$

# Example

$$f_{X,Y}(x,y) = \begin{cases} 3x, & 0 \leq y \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases} \quad f_X(x) = \begin{cases} 3x^2, & 0 \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

To find  $f_{Y|X}(y|x)$ , use the definition:

$$f_{Y|X}(y|x) = \frac{f_{X,Y}(x,y)}{f_X(x)}$$

This is only defined for the values of  $x$  where  $f_X(x) \neq 0$ , that is, for  $0 < x \leq 1$ . For these values of  $x$ , we have:

$$f_{Y|X}(y|x) = \frac{f_{X,Y}(x,y)}{f_X(x)} = \begin{cases} \frac{3x}{3x^2} = \frac{1}{x}, & 0 \leq y \leq x \\ 0, & \text{otherwise} \end{cases}$$

For all other values of  $x$ ,  $f_{Y|X}(y|x)$  is undefined.

# Example

$$f_{X,Y}(x,y) = \begin{cases} 3x, & 0 \leq y \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases} \quad f_X(x) = \begin{cases} 3x^2, & 0 \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases}$$


$$f_{Y|X}(y|x) = \begin{cases} \frac{3x}{3x^2} = \frac{1}{x}, & 0 \leq y \leq x \\ 0, & \text{otherwise} \end{cases}$$

# Example

$$f_{X,Y}(x,y) = \begin{cases} 3x, & 0 \leq y \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases} \quad f_X(x) = \begin{cases} 3x^2, & 0 \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

$$f_{Y|X}(y|x) = \begin{cases} \frac{3x}{3x^2} = \frac{1}{x}, & 0 \leq y \leq x \\ 0, & \text{otherwise} \end{cases}$$

Typically, it is most useful to view  $f_{Y|X}(y|x)$ , as a collection of functions of  $y$ , one for each  $x$ .




# Example

$$f_{X,Y}(x,y) = \begin{cases} 3x, & 0 \leq y \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases} \quad f_X(x) = \begin{cases} 3x^2, & 0 \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

$$f_{Y|X}(y|x) = \begin{cases} \frac{3x}{3x^2} = \frac{1}{x}, & 0 \leq y \leq x \\ 0, & \text{otherwise} \end{cases}$$

Typically, it is most useful to view  $f_{Y|X}(y|x)$ , as a collection of functions of  $y$ , one for each  $x$ .

Viewed as a function of  $y$ , with  $x$  fixed,  $f_{Y|X}(y|x)$  is a PDF (i.e., nonnegative and integrates to one).




## Example 3.9

- Romeo & Juliet have a date at a given time, and each will arrive with a delay between 0 and 1 hours.
- $X$  = Romeo's delay.
- $Y$  = Juliet's delay.
- $X$  and  $Y$  are jointly uniform over  $[0,1] \times [0,1]$ .
- The first to arrive will wait for 15 minutes and will leave if the other has not yet arrived.
- $P(\text{they will meet}) = ?$

# Example 3.9: solution


Their joint arrivals  
are uniform over  
the 1x1 square.



# Example 3.9: solution

Their joint arrivals are uniform over the  $1 \times 1$  square.

They meet in the blue region.






# Example 3.9: solution

Their joint arrivals are uniform over the  $1 \times 1$  square.

They meet in the blue region.




The probability of their meeting is the integral of the joint density over the blue region.

# Example 3.9: solution

Their joint arrivals are uniform over the  $1 \times 1$  square.

They meet in the blue region.




The probability of their meeting is the integral of the joint density over the blue region.

The joint density over the square is 1.

# Example 3.9: solution

Their joint arrivals are uniform over the  $1 \times 1$  square.

They meet in the blue region.




The probability of their meeting is the integral of the joint density over the blue region.

The joint density over the square is 1. This integral is therefore the area of the blue region, which is  $1 - (3/4)^2/2 - (3/4)^2/2 = 7/16$ .

# Example

$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .




# Example

$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

The integral of the joint PDF over  $S$  must be equal to 1.

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .




# Example

$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .

The integral of the joint PDF over  $S$  must be equal to 1.

The integral of the joint PDF over  $S$  is equal to  $c$  times the area of  $S$ .



# Example


$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .

The integral of the joint PDF over  $S$  must be equal to 1.

The integral of the joint PDF over  $S$  is equal to  $c$  times the area of  $S$ .


Hence,  $c = 1/\text{area}(S) = 1/4$ .



# Example

$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .



The integral of the joint PDF over  $S$  must be equal to 1.

The integral of the joint PDF over  $S$  is equal to  $c$  times the area of  $S$ .

Hence,  $c = 1/\text{area}(S) = 1/4$ .


To find the marginal PDF of  $X$ , integrate the joint PDF over vertical lines



# Example

$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .



The integral of the joint PDF over  $S$  must be equal to 1.

The integral of the joint PDF over  $S$  is equal to  $c$  times the area of  $S$ .

Hence,  $c = 1/\text{area}(S) = 1/4$ .

To find the marginal PDF of  $X$ , integrate the joint PDF over vertical lines

E.g., the integral of  $f_{X,Y}(x,y)$  along the line  $x=1.5$

# Example

$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .



The integral of the joint PDF over  $S$  must be equal to 1.

The integral of the joint PDF over  $S$  is equal to  $c$  times the area of  $S$ .

Hence,  $c = 1/\text{area}(S) = 1/4$ .

To find the marginal PDF of  $X$ , integrate the joint PDF over vertical lines



E.g., the integral of  $f_{X,Y}(x,y)$  along the line  $x=1.5$  is the marginal PDF  $f_X(1.5)$ .



# Example

$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .



The integral of the joint PDF over  $S$  must be equal to 1.

The integral of the joint PDF over  $S$  is equal to  $c$  times the area of  $S$ .

Hence,  $c = 1/\text{area}(S) = 1/4$ .

To find the marginal PDF of  $X$ , integrate the joint PDF over vertical lines



E.g., the integral of  $f_{X,Y}(x,y)$  along the line  $x=1.5$  is the marginal PDF  $f_X(1.5)$ .

# Example

$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

To find the marginal PDF of  $Y$ ,  
integrate the joint PDF over horizontal lines

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .






# Example

$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .

To find the marginal PDF of  $Y$ ,  
integrate the joint PDF over horizontal lines



E.g., the integral of  $f_{X,Y}(x,y)$  along the line  
 $y=2.5$  is the marginal PDF  $f_Y(2.5)$ .



# Example

$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .





$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)}$$

# Example

$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .





Recall:  $c = 1/4$

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)} = \begin{cases} \text{[plot of } f_{X|Y}(x|y) \text{ for } 1 \leq y \leq 2 \text{ or } 3 \leq y \leq 4 \text{]} & \text{if } 1 \leq y \leq 2 \text{ or } 3 \leq y \leq 4 \\ 0 & \text{otherwise} \end{cases}$$

# Example

$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .



Recall:  $c = 1/4$



$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)} = \begin{cases} \begin{array}{c} \text{[Plot of } f_{X|Y}(x|y) \text{ for } 1 \leq y \leq 2 \text{ or } 3 \leq y \leq 4 \text{, showing a rectangle from } x=1 \text{ to } x=2 \text{ with height } 1 \text{]} \\ \text{if } 1 \leq y \leq 2 \text{ or } 3 \leq y \leq 4 \end{array} \\ \begin{array}{c} \text{[Plot of } f_{X|Y}(x|y) \text{ for } 2 \leq y \leq 3 \text{, showing a rectangle from } x=1 \text{ to } x=3 \text{ with height } 1/2 \text{]} \\ \text{if } 2 \leq y \leq 3 \end{array} \end{cases}$$



# Example

$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .





Recall:  $c = 1/4$

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)} = \begin{cases} \text{[Plot of } f_{X|Y}(x|y) \text{ for } 1 \leq y \leq 2 \text{ or } 3 \leq y \leq 4 \text{]} & \text{if } 1 \leq y \leq 2 \text{ or } 3 \leq y \leq 4 \\ \text{[Plot of } f_{X|Y}(x|y) \text{ for } 2 \leq y \leq 3 \text{]} & \text{if } 2 \leq y \leq 3 \\ \text{undefined} & \text{if } y < 1 \text{ or } y > 4 \end{cases}$$

# Example

$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .





Recall:  $c = 1/4$

$$f_{Y|X}(y|x) = \frac{f_{X,Y}(x,y)}{f_X(x)} = \left[ \begin{array}{l} \text{ } \end{array} \right.$$

# Example


$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .



Recall:  $c = 1/4$

$$f_{Y|X}(y|x) = \frac{f_{X,Y}(x,y)}{f_X(x)} = \begin{cases} \frac{1/4}{3/4} = 1/3, & \text{if } 1 \leq x \leq 2 \text{ and } 1 \leq y \leq 4 \\ \frac{1/4}{1/4} = 1, & \text{if } 2 \leq x \leq 3 \text{ and } 2 \leq y \leq 3 \\ 0, & \text{otherwise} \end{cases}$$





if  $1 \leq x \leq 2$

# Example

$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .





Recall:  $c = 1/4$

$$f_{Y|X}(y|x) = \frac{f_{X,Y}(x,y)}{f_X(x)} = \begin{cases} \begin{array}{c} \text{Graph of } f_{Y|X}(y|x) \text{ for } 1 \leq x \leq 2: \\ \text{A rectangle from } y=1 \text{ to } y=4 \text{ with height } 1/3. \end{array} & \text{if } 1 \leq x \leq 2 \\ \begin{array}{c} \text{Graph of } f_{Y|X}(y|x) \text{ for } 2 \leq x \leq 3: \\ \text{A rectangle from } y=1 \text{ to } y=3 \text{ with height } 1. \end{array} & \text{if } 2 \leq x \leq 3 \end{cases}$$

# Example

$$f_{X,Y}(x,y) = \begin{cases} c, & (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

Find  $c, f_X, f_Y, f_{X|Y}, f_{Y|X}$ .




Recall:  $c = 1/4$

$$f_{Y|X}(y|x) = \frac{f_{X,Y}(x,y)}{f_X(x)} = \begin{cases} \begin{array}{c} \text{[Plot of } f_{Y|X}(y|x) \text{ for } 1 \leq x \leq 2 \text{, showing a uniform distribution from } y=1 \text{ to } y=4 \text{ with height } 1/3 \text{]} \\ \text{if } 1 \leq x \leq 2 \end{array} \\ \begin{array}{c} \text{[Plot of } f_{Y|X}(y|x) \text{ for } 2 \leq x \leq 3 \text{, showing a uniform distribution from } y=2 \text{ to } y=3 \text{ with height } 1 \text{]} \\ \text{if } 2 \leq x \leq 3 \end{array} \\ \text{undefined} & \text{if } x < 1 \text{ or } x > 3 \end{cases}$$



## Problem 3.21

- Stick of length  $\ell$  is broken at a random point, uniformly distributed over the length of the stick.
- $X$  = length of the left piece.
- The left piece is broken at a random point, whose conditional distribution given the left piece is uniform over the left piece.
- $Y$  = length of the resulting left piece.
- Find  $f_{X,Y}$  and  $f_Y$
- Find  $E[Y]$

# Problem 3.21: $f_{X,Y}$





# Problem 3.21: $f_{X,Y}$







# Problem 3.21: $f_{X,Y}$



$$f_{X,Y}(x,y) = f_X(x)f_{Y|X}(y|x) = \begin{cases} \frac{1}{\ell x}, & 0 \leq y \leq x \leq \ell \\ 0, & \text{otherwise} \end{cases}$$




# Problem 3.21: $f_Y$




$$f_{X,Y}(x,y) = f_X(x)f_{Y|X}(y|x) = \begin{cases} \frac{1}{\ell x}, & 0 \leq y \leq x \leq \ell \\ 0, & \text{otherwise} \end{cases}$$

$$f_Y(y_0) = \int_{-\infty}^{\infty} f_{X,Y}(x, y_0) dx$$




# Problem 3.21: $f_Y$




$$f_{X,Y}(x,y) = f_X(x)f_{Y|X}(y|x) = \begin{cases} \frac{1}{\ell x}, & 0 \leq y \leq x \leq \ell \\ 0, & \text{otherwise} \end{cases}$$


$$\begin{aligned} f_Y(y_0) &= \int_{-\infty}^{\infty} f_{X,Y}(x,y_0) dx \\ &= \begin{cases} ? & \text{if } 0 \leq y_0 \leq \ell \\ 0, & \text{if } y_0 < 0 \text{ or } y_0 > \ell \end{cases} \end{aligned}$$



# Problem 3.21: $f_Y$




$$f_{X,Y}(x,y) = f_X(x)f_{Y|X}(y|x) = \begin{cases} \frac{1}{lx}, & 0 \leq y \leq x \leq l \\ 0, & \text{otherwise} \end{cases}$$




$$\begin{aligned} f_Y(y_0) &= \int_{-\infty}^{\infty} f_{X,Y}(x, y_0) dx \\ &= \begin{cases} \int_{y_0}^l \frac{1}{lx} dx & \text{if } 0 \leq y_0 \leq l \\ 0, & \text{if } y_0 < 0 \text{ or } y_0 > l \end{cases} \end{aligned}$$

# Problem 3.21: $f_Y$



$$f_{X,Y}(x,y) = f_X(x)f_{Y|X}(y|x) = \begin{cases} \frac{1}{\ell x}, & 0 \leq y \leq x \leq \ell \\ 0, & \text{otherwise} \end{cases}$$




$$f_Y(y_0) = \int_{-\infty}^{\infty} f_{X,Y}(x, y_0) dx$$

$$= \begin{cases} \int_{y_0}^{\ell} \frac{1}{\ell x} dx = \frac{1}{\ell} \ln x \Big|_{y_0}^{\ell} = \frac{1}{\ell} (\ln \ell - \ln y_0) = \frac{1}{\ell} \ln \frac{\ell}{y_0} & \text{if } 0 \leq y_0 \leq \ell \\ 0, & \text{if } y_0 < 0 \text{ or } y_0 > \ell \end{cases}$$

# Problem 3.21: $E[Y]$ , method 1

$$f_{X,Y}(x,y) = \begin{cases} \frac{1}{\ell x}, & 0 \leq y \leq x \leq \ell \\ 0, & \text{otherwise} \end{cases}$$


$$E[Y] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} y f_{X,Y}(x,y) dx dy$$



# Problem 3.21: $E[Y]$ , method 1

$$f_{X,Y}(x,y) = \begin{cases} \frac{1}{\ell x}, & 0 \leq y \leq x \leq \ell \\ 0, & \text{otherwise} \end{cases}$$


$$E[Y] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} y f_{X,Y}(x,y) dx dy = \int_0^{\ell} \int_0^x y \frac{1}{\ell x} dy dx$$



# Problem 3.21: $E[Y]$ , method 1

$$f_{X,Y}(x,y) = \begin{cases} \frac{1}{\ell x}, & 0 \leq y \leq x \leq \ell \\ 0, & \text{otherwise} \end{cases}$$

$$\begin{aligned} E[Y] &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} y f_{X,Y}(x,y) dx dy = \int_0^{\ell} \int_0^x y \frac{1}{\ell x} dy dx \\ &= \int_0^{\ell} \frac{1}{\ell x} \cdot \frac{y^2}{2} \Big|_0^x dx \end{aligned}$$






# Problem 3.21: $E[Y]$ , method 1

$$f_{X,Y}(x,y) = \begin{cases} \frac{1}{\ell x}, & 0 \leq y \leq x \leq \ell \\ 0, & \text{otherwise} \end{cases}$$

$$\begin{aligned} E[Y] &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} y f_{X,Y}(x,y) dx dy = \int_0^{\ell} \int_0^x y \frac{1}{\ell x} dy dx \\ &= \int_0^{\ell} \frac{1}{\ell x} \cdot \frac{y^2}{2} \Big|_0^x dx = \int_0^{\ell} \frac{x}{2\ell} dx = \frac{x^2}{2\ell} \Big|_0^{\ell} = \frac{\ell}{4} \end{aligned}$$



## Problem 3.21: $E[Y]$ , method 2

$$f_Y(y) = \begin{cases} \frac{1}{\ell} \ln \frac{\ell}{y} & \text{if } 0 \leq y \leq \ell \\ 0, & \text{otherwise} \end{cases}$$

$$E[Y] = \int_{-\infty}^{\infty} y f_Y(y) dy$$

## Problem 3.21: $E[Y]$ , method 2

$$f_Y(y) = \begin{cases} \frac{1}{\ell} \ln \frac{\ell}{y} & \text{if } 0 \leq y \leq \ell \\ 0, & \text{otherwise} \end{cases}$$

$$E[Y] = \int_{-\infty}^{\infty} y f_Y(y) dy = \int_0^{\ell} \frac{y}{\ell} \ln \frac{\ell}{y} dy$$

## Problem 3.21: $E[Y]$ , method 2

$$f_Y(y) = \begin{cases} \frac{1}{\ell} \ln \frac{\ell}{y} & \text{if } 0 \leq y \leq \ell \\ 0, & \text{otherwise} \end{cases}$$

$$E[Y] = \int_{-\infty}^{\infty} y f_Y(y) dy = \int_0^{\ell} \frac{y}{\ell} \ln \frac{\ell}{y} dy$$

Define  $a = \ln \frac{\ell}{y}$ , then  $\frac{y}{\ell} = e^{-a}$ , and  $dy = -\ell e^{-a} da$

## Problem 3.21: $E[Y]$ , method 2

$$f_Y(y) = \begin{cases} \frac{1}{\ell} \ln \frac{\ell}{y} & \text{if } 0 \leq y \leq \ell \\ 0, & \text{otherwise} \end{cases}$$

$$E[Y] = \int_{-\infty}^{\infty} y f_Y(y) dy = \int_0^{\ell} \frac{y}{\ell} \ln \frac{\ell}{y} dy$$

Define  $a = \ln \frac{\ell}{y}$ , then  $\frac{y}{\ell} = e^{-a}$ , and  $dy = -\ell e^{-a} da$

$$E[Y] = -\ell \int_{\infty}^0 a e^{-2a} da =$$

## Problem 3.21: $E[Y]$ , method 2

$$f_Y(y) = \begin{cases} \frac{1}{\ell} \ln \frac{\ell}{y} & \text{if } 0 \leq y \leq \ell \\ 0, & \text{otherwise} \end{cases}$$

$$E[Y] = \int_{-\infty}^{\infty} y f_Y(y) dy = \int_0^{\ell} \frac{y}{\ell} \ln \frac{\ell}{y} dy$$

Define  $a = \ln \frac{\ell}{y}$ , then  $\frac{y}{\ell} = e^{-a}$ , and  $dy = -\ell e^{-a} da$

$$E[Y] = -\ell \int_{\infty}^0 a e^{-2a} da = \frac{\ell}{2} \int_0^{\infty} 2a e^{-2a} da$$

## Problem 3.21: $E[Y]$ , method 2

$$f_Y(y) = \begin{cases} \frac{1}{\ell} \ln \frac{\ell}{y} & \text{if } 0 \leq y \leq \ell \\ 0, & \text{otherwise} \end{cases}$$

$$E[Y] = \int_{-\infty}^{\infty} y f_Y(y) dy = \int_0^{\ell} \frac{y}{\ell} \ln \frac{\ell}{y} dy$$

Define  $a = \ln \frac{\ell}{y}$ , then  $\frac{y}{\ell} = e^{-a}$ , and  $dy = -\ell e^{-a} da$

$$E[Y] = -\ell \int_{\infty}^0 a e^{-2a} da = \frac{\ell}{2} \underbrace{\int_0^{\infty} 2a e^{-2a} da}_{\substack{\text{mean of an exponential} \\ \text{r.v. with parameter} \\ \lambda=2}}$$

## Problem 3.21: $E[Y]$ , method 2

$$f_Y(y) = \begin{cases} \frac{1}{\ell} \ln \frac{\ell}{y} & \text{if } 0 \leq y \leq \ell \\ 0, & \text{otherwise} \end{cases}$$

$$E[Y] = \int_{-\infty}^{\infty} y f_Y(y) dy = \int_0^{\ell} \frac{y}{\ell} \ln \frac{\ell}{y} dy$$

Define  $a = \ln \frac{\ell}{y}$ , then  $\frac{y}{\ell} = e^{-a}$ , and  $dy = -\ell e^{-a} da$

$$E[Y] = -\ell \int_{\infty}^0 a e^{-2a} da = \frac{\ell}{2} \underbrace{\int_0^{\infty} 2a e^{-2a} da}_{\substack{\text{mean of an exponential} \\ \text{r.v. with parameter} \\ \lambda=2}} = \frac{\ell}{2} \cdot \frac{1}{2} = \frac{\ell}{4}$$



## Problem 3.21: $E[Y]$ , method 3

- Proportion of 1<sup>st</sup> stick broken off is  $X/\ell$

## Problem 3.21: $E[Y]$ , method 3

- Proportion of 1<sup>st</sup> stick broken off is  $X/\ell$
- Proportion of 2<sup>nd</sup> stick broken off is  $Y/X$

## Problem 3.21: $E[Y]$ , method 3

- Proportion of 1<sup>st</sup> stick broken off is  $X/\ell$
- Proportion of 2<sup>nd</sup> stick broken off is  $Y/X$
- The two are independent, and each is a continuous r.v., uniform between 0 and 1.

## Problem 3.21: $E[Y]$ , method 3

- Proportion of 1<sup>st</sup> stick broken off is  $X/\ell$
- Proportion of 2<sup>nd</sup> stick broken off is  $Y/X$
- The two are independent, and each is a continuous r.v., uniform between 0 and 1.
- $E[Y] = E[X(Y/X)]$

## Problem 3.21: $E[Y]$ , method 3

- Proportion of 1<sup>st</sup> stick broken off is  $X/\ell$
- Proportion of 2<sup>nd</sup> stick broken off is  $Y/X$
- The two are independent, and each is a continuous r.v., uniform between 0 and 1.
- $E[Y] = E[X(Y/X)] = E[X] E[Y/X]$

## Problem 3.21: $E[Y]$ , method 3

- Proportion of 1<sup>st</sup> stick broken off is  $X/\ell$
- Proportion of 2<sup>nd</sup> stick broken off is  $Y/X$
- The two are independent, and each is a continuous r.v., uniform between 0 and 1.
- $E[Y] = E[X(Y/X)] = E[X] E[Y/X] = (\ell/2)(1/2) = \ell/4$