



8. Exercise: Sample mean bounds

Exercises due May 1, 2020 05:29 IST Completed

Exercise: Sample mean bounds

2/2 points (graded)

By the argument in the last video, if the X_i are i.i.d. with mean μ and variance σ^2 , and if $M_n = (X_1 + \dots + X_n) / n$, then we have an inequality of the form

$$\mathbf{P}(|M_n - \mu| \geq \epsilon) \leq \frac{a\sigma^2}{n},$$

for a suitable value of a .

a) If $\epsilon = 0.1$, then the value of a is:

100

✓ Answer: 100

b) If we change $\epsilon = 0.1$ to $\epsilon = 0.1/k$, for $k \geq 1$ (i.e., if we are interested in k times higher accuracy), how should we change n so that the value of the upper bound does not change from the value calculated in part (a)?

n should

☐ stay the same

☐ increase by a factor of k

☒ increase by a factor of k^2

☐ decrease by a factor of k



☐ none of the above



Solution:

a) Chebyshev's inequality yields

$$\mathbf{P}(|M_n - \mu| \geq \epsilon) \leq \frac{\sigma^2}{n\epsilon^2},$$

so that $a = 1/\epsilon^2 = 1/0.1^2 = 100$.

b) In order to keep the same upper bound, the term $n\epsilon^2$ in the denominator needs to stay constant. If we reduce ϵ by a factor of k , then ϵ^2 gets reduced by a factor of k^2 . Thus, n will have to be increased by a factor of k^2 .

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 8: Limit theorems and classical statistics:Lec. 18:
Inequalities, convergence, and the Weak Law of Large Numbers / 8.
Exercise: Sample mean bounds

Show all posts



by recent activity



Watch the next segment if you're stuck.

The following segment will make this question MUCH clearer if, like me, you didn't find it obvious at first!

4

© All Rights Reserved

