



<u>Course</u> > <u>Unit 6:</u>... > <u>Lec. 11:</u>... > 10. Exe...

10. Exercise: Using the formula for the monotonic case

Exercises due Mar 25, 2020 05:29 IST Completed

Exercise: Using the formula for the monotonic case

5/6 points (graded)

The random variable X is exponential with parameter $\lambda=1$. The random variable Y is defined by $Y=g\left(X\right)=1/\left(1+X\right)$.

a) The inverse function h, for which $h\left(g\left(x
ight)
ight)=x$, is of the form $ay^{b}+c$. Find a, b, and c.

b) For $y\in (0,1]$, the PDF of Y is of the form $f_{Y}\left(y
ight) =y^{a}e^{\left(b/y
ight) +c}$. Find a , b , and c .

$$a=$$
 2 \star Answer: -2 $b=$ -1 \star Answer: -1 $c=$ 1 \star Answer: 1

Solution:

a) If x and y obey the relation $y=g\left(x
ight) =1/\left(1+x
ight)$, then y+yx=1 , so that



$$x=h\left(y
ight) =rac{1-y}{y}=rac{1}{y}-1.$$

Note that we are interested in $x \geq 0$ which restricts y to the range (0,1]. Notice also that the functions g and h are monotonically decreasing on the relevant ranges of values.

b) Note that

$$rac{dh}{dy}(y) = -rac{1}{y^2}.$$

Therefore,

$$f_{Y}\left(y
ight)=f_{X}ig(h\left(y
ight)ig)\Big|rac{dh}{dy}(y)\,\Big|=e^{-(1/y)+1}\cdotrac{1}{y^{2}}.$$

Submit

You have used 3 of 3 attempts

1 Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 6: Further topics on random variables:Lec. 11: Derived distributions / 10. Exercise: Using the formula for the monotonic case

Show all posts

missing minus sign?
what happened to the negative sign in the derivative?

my math has failed me! (hints on what went wrong)
Assuming lambda = 1, f(x) = e^-x, and therefore, y = g(x) = 1 / (1 + e^-x). I proceeded to solve for x in term...

What happens to lambda in fx(h(y))?
If fx(x) = lambda * e^(-lambda *x), do we not just substitute x = h(y)? Please can someone explain step by

hello, could somebody clarify. does the first expression should include a logarithmic expression?

© All Rights Reserved

