



Course > Unit 4: ... > Lec. 6: ... > 17. Exe...

## 17. Exercise: Linearity of expectations drill

Exercises due Feb 28, 2020 05:29 IST Completed

### Exercise: Linearity of expectations drill

1/1 point (graded)

Suppose that  $\mathbf{E}[X_i] = i$  for every  $i$ . Then,

$$\mathbf{E}[X_1 + 2X_2 - 3X_3] = \boxed{-4} \quad \checkmark \text{ Answer: -4}$$

#### Solution:

Using linearity,

$$\begin{aligned}\mathbf{E}[X_1 + 2X_2 - 3X_3] &= \mathbf{E}[X_1] + \mathbf{E}[2X_2] - \mathbf{E}[3X_3] \\ &= \mathbf{E}[X_1] + 2\mathbf{E}[X_2] - 3\mathbf{E}[X_3] \\ &= 1 + 2 \cdot 2 - 3 \cdot 3 \\ &= -4.\end{aligned}$$

Submit

You have used 3 of 3 attempts

**i** Answers are displayed within the problem

## Discussion

Hide Discussion

**Topic:** Unit 4: Discrete random variables:Lec. 6: Variance; Conditioning on an event; Multiple r.v.'s / 17. Exercise: Linearity of expectations drill

Show all posts



by recent



✓ Why is it not 0?

2 new\_ 6

I simply apply the linearity of expectations and the answer reaches 0. What am I missing?

© All Rights Reserved

