

Unified FWA Field Framework: Gravity and Junction Dynamics

Author: Igor Kolesnikov (Mark)

Priority Preprint Framework

— Abstract —

Fractal-Wave Analysis (FWA) is formulated as a single nonlinear field framework governed by a universal operator E_{FWA} . Two independent projections of this operator describe phenomena at radically different scales: (1) galactic dynamics, where FWA yields the observed logarithmic gravitational potential without invoking dark matter; and (2) baryon-junction dynamics, where FWA represents the experimentally observed Y-topology as a coherent three-branch wave state with a calculable transport eigenvalue. The unified structure demonstrates that both macroscopic and microscopic behaviors can arise from the same wave-operator formalism. The framework is falsifiable: its predictions must be computed from the operator itself, not fitted to data.

—

1. Fundamental Equation

FWA is defined by the nonlinear field equation:

$$E_{FWA}[\Psi] = 0$$

where Ψ denotes the relevant wave configuration. Different physical regimes correspond to different projections of the same operator. This paper focuses on two projections:

- **FWA Gravity** — large-scale dynamics of galaxies,
- **FWA Junction** — coherent three-branch baryon structures.

Both projections follow from the same mathematical foundation.

2. FWA Gravity Projection

2.1 Newtonian Limit and Operator Modification

In the Newtonian limit of General Relativity,

$$\nabla^2 \Phi = 4\pi G \rho_b$$

which yields the Keplerian velocity profile:

$$v^2(r) = GM / r$$

Thus GR predicts $v(r) \propto r^{-1/2}$ at large radii. However, spiral galaxies exhibit flat rotation curves:

$$v(r) \rightarrow v_0 \quad (r \rightarrow \infty)$$

implying:

$$d\Phi/dr = v_0^2/r \Rightarrow \Phi(r) \sim v_0^2 \ln r$$

This logarithmic potential is incompatible with GR + baryons alone.

FWA resolves the discrepancy by modifying the operator:

$$\nabla^2 \Phi - 4\pi G \rho_b - \Delta_{FWA}[\Phi] = 0$$

where Δ_{FWA} is the asymptotic wave-operator term. Its large-scale behavior yields:

$$\Phi(r) \sim v_0^2 \ln r$$

without introducing dark matter.

2.2 Effective Density Interpretation

Substituting the logarithmic potential into the Poisson operator gives:

$$\nabla^2 \Phi \sim v_0^2 / r^2$$

corresponding to an effective density:

$$\rho_{\text{eff}}(r) = v_0^2 / (4\pi G r^2)$$

In GR this term is interpreted as a dark-matter halo. In FWA it arises naturally from the operator Δ_{FWA} , not from additional matter.

Thus the gravitational sector illustrates the core FWA principle: modify the operator, not the mass content.

3. FWA Junction Projection

3.1 Coherent Three-Branch Wave State

A baryon junction is represented as a coherent three-branch wave configuration:

$$\Psi_J = \sum_{k=1..3} A_k e^{i\phi_k}, \quad \phi_k = 2\pi k/3 + \delta\phi_k$$

The unperturbed phases satisfy the root-of-unity cancellation:

$$\sum_{k=1..3} e^{i2\pi k/3} = 0$$

interpreted in FWA as a dynamic-zero condition of a perfectly balanced junction. This representation does not replace SU(3) color; it provides an independent wave-based description of the Y-topology.

3.2 Symmetry-Error Functional

Departures from perfect symmetry are quantified by:

$$Q_J^2 = |\sum_{k=1..3} A_k e^{i\phi_k}|^2$$

For equal amplitudes $A_k = 1$ and small distortions $\delta\phi_k$, $e^{i\delta\phi_k} \approx 1 + i\delta\phi_k$, and the zeroth-order term cancels. The leading contribution is therefore quadratic:

$$Q_J^2 \propto |\sum_{k=1..3} \delta\phi_k e^{i2\pi k/3}|^2$$

This is a concrete mathematical prediction: junction symmetry errors scale as $\delta\phi^2$.

3.3 Dynamical Recursion and Transport Law

The junction evolves under the general FWA recursion:

$$\Psi_{n+1} = F(\Psi_n, v_n, d_n, T_n)$$

where T encodes the local medium state. Linearization around a stationary junction yields:

$$\Psi_{n+1} \approx \lambda_J \Psi_n$$

with λ_J the dominant eigenvalue of the linearized operator. Over a rapidity interval Δy :

$$C(\Delta y) = e^{-\lambda_J \Delta y}$$

This leads to the transport law:

$$dN_B/dy = N_0 \exp[-\lambda_J (y - y_0)]$$

The eigenvalue λ_J must be computed from the FWA equations, not fitted to STAR data. Agreement supports the model; disagreement falsifies it.

4. Unified Structure

Both sectors—gravity and junction dynamics—are projections of the same fundamental operator:

$$E_FWA [\Psi] = 0$$

Projections:

- $E_grav [\Phi, \rho_b] = 0$ (galactic scales)
- $E_junction [\Psi_j, T] = 0$ (hadronic scales)

The unification is conceptual and mathematical:

- **Gravity:** FWA modifies the operator to produce the observed logarithmic potential without dark matter.
- **Junctions:** FWA represents Y-topology as a coherent wave state with a calculable transport eigenvalue.

A physical explanation becomes a theory only when its mathematical structure produces consequences that nature can contradict.

5. Final Consolidated Block

[1] Fundamental field equation:

$$E_FWA [\Psi] = 0$$

[2] Gravity projection:

$$\nabla^2 \Phi - 4\pi G \rho_b - \Delta_{FWA}[\Phi] = 0$$
$$\Phi(r) \sim v_0^2 \ln r, \quad v(r) \sim v_0$$

[3] Junction projection:

$$\Psi_j = \sum_k A_k e^{i\phi_k}$$
$$Q_j^2 = |\sum_k A_k e^{i\phi_k}|^2$$
$$\Psi_{n+1} \approx \lambda_j \Psi_n$$
$$dN_B/dy = N_0 e^{-\lambda_j(y-y_0)}$$