

# 1    **Data variability in standardised cell culture experiments**

2    Ian G. Reddin, Tim Fenton, Mark N. Wass\*, Martin Michaelis\*

3    School of Biosciences, University of Kent, Canterbury, UK

4    \* Correspondence to: [m.n.wass@kent.ac.uk](mailto:m.n.wass@kent.ac.uk) (M.N.W.), [m.michaelis@kent.ac.uk](mailto:m.michaelis@kent.ac.uk)

5    (M.M.)

6

7

## 8 Abstract

9 Despite much debate about a perceived 'reproducibility crisis' in the life  
 10 sciences, it remains unclear what level of replicability is technically possible [1,2].  
 11 Here, we analysed the variation among drug response data of the NCI60 project,  
 12 which for decades has tested anti-cancer agents in a 60-cell line panel following a  
 13 standardised protocol [3]. In total, 2.8 million compound/cell line experiments are  
 14 available in the NCI60 resource CellMiner [4]. The largest fold change between the  
 15 lowest and highest GI50 (concentration that reduces cell viability by 50%) in a  
 16 compound/cell line combination was  $3.16 \times 10^{10}$ . All compound/ cell line  
 17 combinations with >100 experiments displayed maximum GI50 fold changes >5,  
 18 99.7% maximum fold changes >10, 87.3% maximum fold changes >100, and 70.5%  
 19 maximum fold changes >1000. FDA-approved drugs and experimental agents  
 20 displayed similar variation. The variability remained very high after removal of  
 21 outliers and among experiments performed in the same month. Hence, our analysis  
 22 shows that high variability is an intrinsic feature of experimentation in biological  
 23 systems, even among highly standardised experiments in a world-leading research  
 24 environment. Thus, a narrow focus on experiment standardisation does not ensure a  
 25 high level of replicability on its own.

26

## 27 Introduction

28 In the life sciences, there is a crisis narrative and a perception of a lack of  
29 data reproducibility ("reproducibility crisis" or "replication crisis") [2,5-10]. However,  
30 the actual scale of the crisis remains unclear and evidence is largely anecdotal [2].  
31 Much of the data are based on researcher views expressed in survey responses [11-  
32 13] or provided as Comments or Correspondence without providing detailed  
33 information [14,15].

34 One project that investigates the replicability of research data is the  
35 'Reproducibility Project: Cancer Biology', which independently repeats influential  
36 preclinical studies ([https://elifesciences.org/collections/9b1e83d1/reproducibility-](https://elifesciences.org/collections/9b1e83d1/reproducibility-project-cancer-biology)  
37 [project-cancer-biology](https://elifesciences.org/collections/9b1e83d1/reproducibility-project-cancer-biology)). So far, 17 replication studies have been completed. Five  
38 studies reported the successful reproduction of the original studies [16-20], while  
39 eight reported a mixed outcome [21-28], and four failed to reproduce the original  
40 findings [29-32]. It is not clear whether these data are representative. It is a small  
41 dataset focused on small, early, and highly cited studies, which are more likely to  
42 overestimate effects [33].

43 There is also a lack of agreement on the expected level of data replicability  
44 [1]. In a dispute about the consistency of two large pharmacogenomic screens in  
45 cancer cell line panels, the Genomics of Drug Sensitivity in Cancer database and the  
46 Cancer Cell Line Encyclopedia [34-37], four analyses by different groups concluded  
47 a reasonable level of consistency [37-40], while six studies, all by the same group,  
48 disagreed [36,41-45].

49 To develop a realistic understanding of the replicability of standardised assays  
50 in a world-leading research environment, we investigated the variation in drug  
51 response data from the NCI60 screen [3]. Since 1985, the NCI60 screen has tested

52 the anti-cancer activity of thousands of compounds multiple times in a 60-cell line  
53 panel following strict standard operating procedures (Figure 1A) [3,4,47-51]. Thus,  
54 the NCI60 database provides an unprecedented wealth of data on the replicability of  
55 findings using highly standardised procedures by highly skilled experts.

56

57

## 58 Results

### 59 NCI60 drug response data are characterised by a high level of variability

60 All drug sensitivity data derived from NCI60 testing are made available via  
 61 Cell Miner [4,47,48,50,51]. In total, 52,585 compounds were tested in the NCI60  
 62 resulting in  $2.8 \times 10^6$  compound/cell line combinations. Two or more (up to 2,286)  
 63 experiments were carried out for 11,841 compounds and 594,450 compound/cell line  
 64 combinations (Figure 1B, Extended Data Table 1; Extended Data Table 2). More  
 65 than 100 experiments in at least one cell line were performed for 18 compounds and  
 66 more than 1,000 experiments for two compounds (Extended Data Table 3).  
 67 Concentration ranges varied from  $10^{1.2}$  to  $10^{12.1}$ . 612 compounds were screened with  
 68 multiple concentration ranges, and the most common concentration range was  $10^4$   
 69 (11,213/ 94.7% of the compounds), representing the standard testing range using  
 70 five 10-fold dilution steps (Extended Data Table 4).

71 The maximum fold change between the lowest and highest GI50  
 72 concentration (reduces cell viability by 50%) was detected for  
 73 cyanomorpholinodoxorubicin in the colorectal cancer cell line COLO 205 ( $3.16 \times$   
 74  $10^{10}$ ) (Extended Data Figure 1A, Extended Data Table 5). 232,315 (39.1%) drug/cell  
 75 line combinations displayed maximum fold changes >2, 108,247 (18.2%) drug/cell  
 76 line combinations fold changes >5, 59,638 (10%) drug/cell line combinations fold  
 77 changes >10, 19,089 (3.2%) drug/cell line combinations >100, and 8320 (1.4%)  
 78 drug/cell line combinations >1,000 (Extended Data Table 5).

79

### 80 Variability increases with the number of experiments


81 The percentage of compound/cell line combinations with high maximum fold  
 82 change strongly increased with the number of experiments (Figure 1C, Extended

83 Data Figure 2A, Extended Data Figure 2B, Extended Data Table 6). The mean and  
84 median GI50 fold changes increased from 41,292 and 1.4 for compound/cell line  
85 combinations with two experiments to 53,343,445 and 1841 for compound/cell line  
86 combinations with >20 experiments (Figure 1C, Figure 1D, Extended Data Table 6).

87 When we considered compound/cell line combinations with a minimum of five  
88 experiments, 25,496 (84.4%) of 30,212 compound/cell line combinations displayed  
89 maximum fold changes >2 and 3,832 (12.7%) compound/cell line combinations  
90 >1000. For compound/ cell line combinations with >100 experiments, 100% of 794  
91 compound/ cell line combinations displayed a maximum fold change > 5 and 70.5%  
92 (560 out of 794) displayed a maximum fold change >1000 (Figure 1E, Extended  
93 Data Table 7).

94 Taken together, maximum GI50 fold changes increase with the number of  
95 experiments. In agreement, a significant correlation was detected between maximum  
96 GI50 fold changes and the number of experiments per compound/cell line  
97 combination (Spearman correlation coefficient = 0.34,  $p < 2.2 \times 10^{-16}$ ) (Extended  
98 Data Figure 3A).

99



**Figure 1 Variability in NCI60 GI50 data.** **A)** Overview of the principle of the NCI 60 screen. **B)** Compound/ cell line combinations with two or more experiments in the NCI60 database. **C)** GI50 fold changes in dependence on the number of experiments per compound/ cell line combination. Numerical data are presented in the adjacent table. **D)** Distribution of maximum GI50 fold changes illustrated by density plots for experimental compound/cell line combination groups with an increasing minimum number of experiments. **E)** Percentage of compound cell line combinations with maximum fold changes above the indicated thresholds in dependence of the number of experiments. Numerical data are presented in the adjacent table. **F)** Distribution of GI50 fold changes in dependence of the

concentration ranges in which compounds were tested. Numerical data are presented in the adjacent table.

## **Variability increases with the concentration range covered**

The observed fold changes also reflected the tested concentration ranges per compound/ cell line combination in addition to the number of experiments, i.e. the broader the range of concentrations that were tested, the larger was the maximum fold change (Figure 1D, Extended Data Table 8). A positive correlation was observed between concentration range and maximum fold change for all compound data (Spearman correlation coefficient = 0.31,  $p < 2.2 \times 10^{-16}$ ) (Extended Data Figure 3B).

The mean and median GI50 fold changes for compound/ cell line combinations for which a maximum concentration range  $<1.0 \times 10^5$  was covered were 146.8 and 1.5, which increased to 68,121,059 and 289, 734 for those with a concentration range of  $\geq 1.0 \times 10^9$  (Figure 1F, Extended Data Table 8).

## **Variability in FDA Approved Drugs**

Since reliable clinical therapy outcomes depend on reproducible drug effects, it may be speculated that FDA-approved drugs are more robust in their drug response data than experimental agents. However, the drug response data observed for FDA-approved drugs displayed a similar variability like that observed across all tested compounds.

The NCI60 database contained data on 181 FDA-approved drugs, which had been tested at least twice, resulting in 399,686 experiments investigating 9,970 individual drug/cell line combinations (Extended Data Table 1). The number of experiments for drug/cell line combinations ranged from 2 to 2,286.



The maximum GI50 fold change was  $1.25 \times 10^{10}$  observed for mithramycin in four cell lines, the colorectal cancer cell line COLO-205 (26 experiments) (Extended Data Figure 1B), the CNS cell lines SF-295 (24 experiments) and U251 (27 experiments), and the ovarian cancer cell line IGROV1 (26 experiments). Mithramycin, a member of the aureolic acid family was approved in 1970 but only temporarily used for testicular carcinoma and other types of cancer due to serious side effects [52]. The second highest GI50 fold change ( $7.28 \times 10^7$ ) was detected for paclitaxel, a stabilising tubulin-binding agent and one of the most commonly used anti-cancer drugs [53], in MDA-MB-435 (Extended Data Figure 1C), which had originally been assumed to be a breast cancer cell line, but was later found to be derived from the melanoma cell line M14 [54].

The maximum GI50 fold changes were higher among the FDA approved drugs than for the non-FDA approved compounds (Figure 2A, Extended Data Table 9), probably because they were tested in more experiments and at bigger concentration ranges (Figure 2A).

When we considered the percentage of FDA-approved drug/ cell line combinations with maximum GI50 fold changes >2, >5, >10, >100, and >1000 for combinations with >5, >10, >20, and >100 experiments (Figure 2B, Figure 2C, Extended Data Table 7, Extended Data Table 10), we obtained similar results to those across all compounds (Figure 1E).


In agreement with the findings across all compound/ cell line combinations, the maximum GI50 fold changes also increased with experiment number when the FDA approved drug/ cell line combinations were grouped into combinations with two experiments, 3 to 5 experiments, 6 to 20 experiments, and >20 experiments (Figure 2D, Suppl. Table 11), and the maximum GI50 fold change was also correlated with

159 the number of experiments performed (Spearman's correlation coefficient = 0.72,  $p <$   
160  $2.2 \times 10^{-16}$ )(Extended Data Figure 3C).

161 Moreover, the maximum GI50 fold change increased with the concentration  
162 range covered (Figure 2E, Extended Data Table 12), and there was a significant  
163 correlation between the concentration range and the maximum GI50 fold change  
164 (Spearman's correlation coefficient = 0.62,  $p < 2.2 \times 10^{-16}$ ) (Extended Data Figure  
165 3D).

166 Taken together, there is no indication that FDA-approved drugs would display  
167 less variability than experimental compounds.

168



169

170 **Figure 2. GI50 variation for FDA-approved drugs. A)** Compound/ cell line combinations with 2 or  
 171 more experiments in the NCI60 database. Numerical data are presented in the adjacent tables. **B)**  
 172 Percentage of FDA-approved drug/ cell line combinations with maximum fold changes above the  
 173 indicated thresholds in dependence of the number of experiments. Numerical data are presented in  
 174 the adjacent table. **C)** Distribution of maximum GI50 fold changes illustrated by density plots for  
 175 experimental compound/cell line combination groups with increasing minimum numbers of  
 176 experiments. **D)** GI50 fold changes in dependence on the number of experiments per compound/ cell  
 177 line combination. Numerical data are presented in the adjacent table. (E) Distribution of GI50 fold

changes in dependence of the concentration ranges in which compounds were tested. Numerical data are presented in the adjacent table.

## **GI50 variability in experiments performed by month**

The reproducibility of results may be affected by parameters such as changes in the reagents, e.g. use of different lots or batches, different experimenters, and using cell lines at different passages [2,55-57]. Hence, experiments performed closely together may be expected to display greater similarity than experiments performed at more distant points in time during the decades of anti-cancer compound testing by the NCI60.


To investigate the effects of the time of testing on data variability, we compared experiments performed in the same month to control samples of the same size that were randomly selected across the whole testing period. For this analysis, we used the 18 FDA-approved drugs that were tested at least 100 times in at least one cell line over periods of 95 to 275 months (Figure 3A, Extended Data Table 13), resulting in 51,872 drug/cell line combinations and in total 321,709 experiments (Figure 3B, Suppl. Table 14).

For every set of experiments performed on the same date, we generated 1,000 random control samples of the same size and compared the value distribution. The variability of GI50 fold changes for same date experiments was indeed lower than for random control samples, but remained very high reaching up to  $1.74 \times 10^8$  (Figure 3C, Extended Data Table 15). Moreover, for 45% of the same date drug/ cell line combinations the GI50 fold change was higher than the mean fold change of the corresponding 1,000 random samples (Figure 3D, Extended Data Table 16).

203           When we looked at the data per cell line, the same date GI50 fold changes  
 204   were higher than the mean random sample fold changes for the majority of drugs in  
 205   ten cell lines, higher for half of the drugs in three cell lines, and lower for the majority  
 206   drugs in the remaining 47 cell lines (Figure 3E, Extended Data Table 17). When we  
 207   looked at the individual drugs, six displayed a majority of drug/cell line combinations  
 208   with higher mean same date GI50 fold changes higher than in the random samples  
 209   and twelve drugs displayed lower ones (Figure 3F, Extended Data Table 18).

210           Taken together, experiments performed in close timely proximity display lower  
 211   variability than experiments performed over a longer time period, but the same date  
 212   variability remains very high.

213



**Figure 3 GI50 variability is high between compound/cell line combination experiments on the same date and is not caused by outliers. A)** Time periods of drug testing for individual drugs. **B)** Testing of individual compound/cell line combinations by date. **C)** Maximum GI50 fold changes in experiments testing compound/ cell line combinations on the same date compared to maximum GI50

fold changes in 1000 random controls of the same sample size. **D)** Percentage of cases in which same date experiments had a higher fold change than control samples randomly picked across the timeline. **E)** Proportion of same date GI50 fold changes in compound/ cell line combinations that are higher or lower than random control samples per cell line. **F)** Proportion of same date GI50 fold changes in compound/ cell line combinations that are higher or lower than random control samples per drug. **G)** GI50 value distribution for maytansine in the prostate cancer cell line DU-145 (outliers indicated in red). **H)** GI50 value distribution for 5-fluorouracil in the lung cancer cell line HOP-62 (outliers indicated in red). **I)** Comparison of maximum GI50 fold changes before and after removal of outliers. Numerical values are presented in the adjacent table. **J)** Maximum GI50 fold changes increase with experiment number after removal of outliers. Numerical values are presented in the adjacent table. **K)** Maximum GI50 fold changes increase with the concentration range covered after removal of outliers. Numerical values are presented in the adjacent table.

## High GI50 variability is not caused by outliers

Finally, we determined GI50 outliers for compound/cell line combinations with 5 or more experiments (738 compounds, 30,212 compound/cell line combinations, 598,243 GI50 values) using the adjusted boxplot method [58]. 5.7% (34,216) of GI50 values were outliers and 43.7% (13,208/30,212) of compound/cell line combinations had at least one GI50 outlier (Extended Data Table 19). This indicates that outliers are not responsible for the observed variability of GI50 in the majority (56.3%) of experiments.

The highest percentage of outliers was 50% (7/14 experiments for maytansine in DU-145 prostate cancer cells) (Figure 3G, Extended Data Table 19). The greatest number of outliers was 291 (16.8%) out of 1731 experiments for 5-fluorouracil in HOP-62 lung cancer cells (Figure 3H, Extended Data Table 19). Outlier number increased with the number of experiments for a compound/cell line combination with a Spearman correlation coefficient of 0.25 ( $p < 2.2 \times 10^{-16}$ ) (Extended Data Figure 4).

The removal of outliers reduced data variability, but the overall variability remained very high with a maximum GI50 fold range of  $2.5 \times 10^9$  detected for maytansine in the ovarian cancer cell line OVCAR-5 over 35 experiments (Figure 3I, Extended Data Table 19).

As detected in the analysis across all experiments, maximum GI50 fold changes increased with the number of experiments and the concentration ranges covered also after the removal of outliers (Figure 3J, Figure 3K, Extended Data Table 19, Extended Data Table 20). A significant correlation was observed between experiment number and maximum GI50 fold change with a Spearman correlation of 0.39 ( $p < 2.2 \times 10^{-16}$ ) (Extended Data Figure 5A) and between concentration range and maximum GI50 fold change with a Spearman correlation of 0.47 ( $p < 2.2 \times 10^{-16}$ ) (Extended Data Figure 5B).

### **No drift in drug sensitivity over time**

Cancer cell lines may display substantial changes in genotype and phenotype over time [55,59]. Hence, part of the variability observed in drug sensitivity may be the consequence of a shift in drug response over time. To investigate this, we established timelines of the GI50 values for the 18 compounds, which had been tested at least 100 times in one or more cell lines. This resulted in time lines for 1080 compound/cell line combinations with time frames ranging from 95 months (vinblastine, floxuridine, cytarabine, daunorubicin, 6-mercaptopurine) to 275 months (Doxorubicin) (Figure 3A, Extended Data Table 13).


Drug/ cell line combinations, in which the fold change between the mean GI50 on the first experimental date and the mean GI50 on the last experimental date was 50% or greater than the maximum GI50 fold change for the data points in between



271 were considered as candidates for a drift in drug sensitivity. Only six (0.56%) out of  
272 1080 drug/cell line combinations fulfilled these criteria (Figure 4, Extended Data  
273 Table 21).

274 The distribution of the individual GI50 values for three of the drug/ cell line  
275 combinations (floxuridine/ SK-OV-3, methotrexate/ BT-549, 6-mercaptopurine/ BT-  
276 549) did not indicate a GI50 shift over time (Figure 4A-C, Extended Data Table 21,  
277 Extended Data Table 22). For the other three drug/ cell line combinations  
278 (vinblastine/ T-47D, bleomycin/ K-562, M-AMSA/ MDA-MB-435) a drift in sensitivity  
279 appears unlikely but cannot be excluded based on the data (Figure 4D-E, Extended  
280 Data Table 21, Extended Data Table 22). However, such observations are very rare.  
281 Moreover, a phenotypic drift in a cell line would be expected to result in changes in  
282 sensitivity to more than one drug over time. Hence, the data provide no evidence  
283 suggesting that the drug sensitivity of individual cell lines may have changed over  
284 time. These findings may also reflect that the NCI60 uses cell lines within a window  
285 of 30 passages [60].

286



287

288 **Figure 4 Experimental time lines for individual compound/cell line combinations.** The six  
 289 experimental timelines for compound/cell line combinations (with more than 100 experiments  
 290 performed in at least one cell line) with a fold change between the first and last mean GI50 that is  
 291 50% greater than the maximum GI50 for the remaining experiments. **A)** floxuridine in the ovarian  
 292 cancer cell line SK-OV-3, **B)** methotrexate in the breast cancer cell line BT-549, **C)** 6-mercaptopurine  
 293 in the breast cancer cell line BT-549, **D)** bleomycin in the leukaemia cell line K-562, **E)** vinblastine in  
 294 the breast cancer cell line T-47D, **F)** M-AMSA in the melanoma cell line MDA-MB-435. Plot lines  
 295 represent mean GI50s while data points represent individual experiments at an experimental date.  
 296

## 297 Discussion

298 To gain real-life insights into the extent of data variability in a standardised  
299 research environment, we here investigated the variation of GI50 values indicating  
300 drug sensitivity for compounds that had been tested multiple times in the NCI60  
301 panel. The variation was large with the highest fold change between the lowest and  
302 highest GI50 in a given compound/cell line combination being  $3.16 \times 10^{10}$ . As might  
303 have been expected, the fold change between the lowest and the highest GI50 in a  
304 specific compound cell line combination increased with the number of experiments  
305 and the concentration range tested.

306 CellMiner contains data on experimental compounds as well as on FDA-  
307 approved drugs that are in clinical use [4,46-51]. Although FDA-approved drugs  
308 might have been expected to result in more robust data, this was not the case and  
309 they displayed a similar data variability as that determined across all compounds.  
310 The variability also remained very high when we only considered experiments that  
311 were performed in the same months or removed outliers.

312 Given that the NCI60 uses highly standardised methods, a high variability of  
313 assay results appears to be an intrinsic feature of lab experiments. The large range  
314 of GI50 values observed for anti-cancer drugs is of potential relevance, given that  
315 cytotoxic anti-cancer drugs are typically used at maximum tolerated doses that  
316 cannot be further increased without unacceptable toxicity [61-63]. Moreover, the  
317 maximum effects of targeted drugs, e.g. antibodies or kinase inhibitors that interfere  
318 with cancer-specific structures or entities, do not further increase beyond the 'optimal  
319 biological dose', i.e. the dose at which the biological target is completely inhibited  
320 [61-64]. Hence, even a two-fold difference in the GI50, which occurred in 25,496  
321 (84.4%) of 30,212 compound/cell line combinations with at least five experiments, is

322 of potential relevance, as a two-fold increase of the clinical dose of an anti-cancer  
323 drug is rarely feasible .

324 Notably, there is awareness of this variability within the NCI60 project as  
325 indicated by strict quality control procedures in the presentation of NCI60 GI50  
326 values in CellMiner, which result in the exclusion of up to 96% of experiments (48 out  
327 of 50) for a given compound/ cell line combination (Extended Data Table 23) [4].

328 However, such knowledge has not penetrated into scientific discourse. Many  
329 authors promote strict standardisation of experimental procedures as a strategy to  
330 improve data quality and reliability [2,5,15,45,65-71].

331 Our analysis of NCI60 data, which is of unprecedented depth and provided by  
332 a world-leading institution with unprecedented transparency, indicates that data  
333 variation remains very high even under ideal conditions that the vast majority of  
334 research groups will not be able to afford. Hence, our data suggest that experiment  
335 heterogenization, the testing of a hypothesis in many different (experimental)  
336 systems and datasets and different laboratories [72-79], is a much better strategy to  
337 generate robust and meaningful data.

338 In conclusion, our analysis demonstrates that the variation of experimental  
339 data is extremely high even under optimal conditions in a world-leading environment  
340 applying the highest standards. This shows that increased standardisation is not a  
341 straightforward way to resolve issues associated with limited replicability. Hence,  
342 increased data robustness will have to include additional strategies such as  
343 independent replication and experiment heterogenization, i.e. multiple testing of the  
344 same hypothesis using different approaches and models. Awareness of the inherent  
345 variability of experimental results, will help researchers to develop a realistic

346 understanding of the meaning of their data and to design more diverse research  
347 strategies that will result in higher data robustness and reliability.

348

349

## Methods

## Data Availability

All data were obtained from CellMiner [80] Version 2.2. Dose concentration range data (June 2018 release) were obtained from the National Cancer Institute DTP NCI bulk data for download pages (<https://wiki.nci.nih.gov/display/NCIDTPdata/NCI-60+Growth+Inhibition+Data>). Of the 52,585 NCI codes given to compounds tested on the NCI-60 cell line panel, 42,794 were given to unnamed compounds and 9,791 were given to 9,027 named compounds. 262 named compounds received two or more individual codes. Some GI50 values represent minimum or maximum drug concentrations where the actual GI50 was not reached [4]. Since such values understate the actual data variation, we did not remove these data. All data generated during this study are included in this published article and its supplementary information files.

## Maximum GI50 fold change calculation

The drug sensitivity data was converted from  $-\log_{10}$  GI50, to the GI50 ( $\mu\text{M}$ ) for all compound, all cell lines and all experiments. Maximum fold changes were calculated for each compound/cell line combination with more than one experiment (594,450) by dividing the maximum GI50 for a cell line by the minimum GI50.

## Number of experiments and experimental groups

The number of experiments for each individual compound/cell line combination was calculated by counting all experiments performed on the same experimental date as well as experiments on different dates. The relationship

between number of experiments and maximum fold change was investigated by using Spearman's correlation coefficient as the distribution of maximum GI50 fold change was not normal.

The compound/cell line combinations were then assigned experimental groupings base on the number of experiments performed: all data, 5 or more experiments, 10 or more experiments, 20 or more experiments, and 100 or more experiments. This allowed comparison of "high" maximum fold changes (>2, >5, >10, >100, and >1000) for combinations with varied number of experiments. Additionally, compound/cell line combinations were assigned to experimental groups: 2 experiments, 3 to 5 experiments, 6 to 20 experiments and over 20 experiments. These experimental groupings enabled comparison of GI50 fold change statistics (mean, median, minimum, maximum, variance) for compound/cell line combinations with number of experiments ranging from lower to higher.

### **Concentration range and experimental groups**

Maximum dose concentration range for a compound/cell line combination was determined by using the minimum and maximum dose concentration used in an experiment for an individual compound on an individual cell line. The minimum concentration range was  $1.0 \times 10^{1.2}$  and the maximum concentration was  $1.0 \times 10^{12.1}$ . The relationship between dose concentration range and maximum fold change was investigated by using Spearman's correlation coefficient as the distribution of maximum GI50 fold change was not normal.

Compound/cell line combinations were assigned to groups based on the dose concentration range for that combinations: maximum concentration range less than  $1.0 \times 10^5$ , maximum concentration range  $1.0 \times 10^5$  to  $1.0 \times 10^9$  exclusive and

400 maximum concentration range  $1.0 \times 10^9$  and above. These experimental groupings  
401 enabled comparison of GI50 fold change statistics (mean, median, minimum,  
402 maximum, variance) for compound/cell line combinations between lower and higher  
403 concentration ranges.

404

#### 405 **FDA-approved compound analysis**

406 All compounds that were classed as FDA-approved drugs by the NCI-60 in  
407 CellMiner Database Version 2.2 and where two or more experiments had been  
408 performed were extracted from the complete dataset. This created an FDA-approved  
409 dataset of 181 drugs for which 399,686 experiments for 9,970 individual drug/cell line  
410 combinations were performed. Analysis of relationship between the number of  
411 experiments/concentration ranges and maximum GI50 fold change for drug/cell line  
412 combinations was performed as for the complete dataset, described above.

413

#### 414 **Experiments on the same date**

415 Month and year of each experiment was available so experimental timelines  
416 were established for compounds by calculating the time between the first and last  
417 experiment date. Multiple experiments were carried out on the same date for many  
418 of the compound/cell line combinations, particularly the 18 compounds with at least  
419 one cell line with 100 total experiments. The data for these 18 compounds, 17 of  
420 which were FDA-approved, was extracted from the complete dataset to create a  
421 subset of data deemed suitable to compare GI50 variability on the same date with  
422 GI50 variability over an experimental timeline.

423 The maximum GI50 fold change on each date where there were multiple  
424 experiments for a compound/cell line combination were calculated by dividing



maximum GI50 by minimum GI50 value. The number of experiments on a specific date for a compound/cell line combination was used to determine the maximum GI50 fold change over the same number of experiments picked randomly from that combination's experimental timeline. This was performed 1000 times so that for every compound/cell line combination and experimental date with a maximum GI50 fold change over multiple experiments there were 1000 corresponding maximum GI50 fold changes calculated from random samples of the same number of experiments on that compound/cell line combination's timeline. The mean maximum GI50 fold change was calculated for the 1000 random samples for each compound/cell line combination and the number of maximum GI50 fold changes for experiments on the same date higher and lower than the random sample mean were counted. For each compound, significance of the difference between same date maximum GI50 fold change and sample mean GI50 fold change was calculated using Wilcoxon Rank Sum Test. This was performed using all cell line data combined for each compound and for each cell line individually for each compound. Where a significant difference between same date and random sample mean maximum GI50 fold changes were observed, the number of times the same date GI50 fold change was higher or lower than the random sample mean maximum GI50 fold change was counted.

#### **Drift in drug sensitivity**

The mean GI50 fold change was calculated for each experimental date (month) for the 18 compounds with 100 or more experiments for at least one cell line. The GI50 fold change between the first experimental date and the last experimental date was calculated using the mean GI50 on those dates. The first/last

450 GI50 fold change was then compared to the maximum GI50 fold change for each  
451 compound/cell line combination and considered a candidate for a drift in sensitivity if  
452 it was 50% or more of the maximum fold change.

453

#### 454 **Removal of outliers**

455 The adjusted boxplot method was used to identify outlier thresholds. This  
456 method was chosen as the data set was highly skewed. To use this method the  
457 medcouple ( $MC$ ), a robust measure of skewness, had to be calculated (where  
458  $X_n = \{x_1, x_2, \dots, x_n\}$  represents data for every compound/cell line combination):

459

$$MC(x_1, \dots, x_n) = \text{med} \frac{(x_j - \text{med}_k) - (\text{med}_k - x_i)}{x_j - x_i}$$

460

461 Where  $\text{med}_k$  is the median of  $X_n$ , and  $i$  and  $j$  have to satisfy  $x_i \leq \text{med}_k \leq x_j$ , and  
462  $x_i \neq x_j$ .

463 Using the  $MC$  the upper ( $U$ ) and lower ( $L$ ) thresholds could be determined. If

464  $MC \geq 0$  :

$$L = Q_1 - 1.5 \times \exp(-3.5MC) \times IQR$$

$$U = Q_3 + 1.5 \times \exp(4MC) \times IQR$$

465

466 If  $MC \leq 0$  :

$$L = Q_1 - 1.5 \times \exp(-4MC) IQR$$

$$U = Q_3 + 1.5 \times \exp(3.5MC) \times IQR$$

467

468 If  $MC = 0$  the adjusted boxplot method was not used but instead the Tukey method  
469 was used:

$$L = Q_1 - 1.5IQR$$

$$U = Q_3 + 1.5IQR$$

470 Where  $Q_1$  is the lower quartile,  $Q_3$  is the upper quartile and  $IQR$  is the interquartile  
471 range.

472 For each compound/cell line combination any GI50 value below  $L$  or  
473 above  $U$  were removed from the dataset. Analyses were performed on this dataset  
474 as previously described for the complete dataset.

475

## 476 **Data processing**

477 Data was carried out using perl version 5.26.0, Microsoft Excel (2011) and R  
478 statistical packages version 3.4.4. Perl modules Statistics:Descriptive and  
479 Statistics::R were used. Packages used in R were robustbase, dplyr, webr,  
480 moonBook, tidyverse, reshape2, scales, gplots, ggpubr, ggExtra, RColorBrewer,  
481 corrplot, ggplot2, and tidyr.

482

483

484

485

486

## 487 **Author contributions**

488 Data acquisition and analysis (I.R.), data interpretation (all authors), study  
489 conception (M.N.W., M.M.), study design (I.R., M.N.W., M.M.), manuscript drafting  
490 (I.R., M.M.), manuscript revision (all authors).

491 All authors have approved the submitted version and agreed both to be  
492 personally accountable for their own contributions and to ensure that questions  
493 related to the accuracy or integrity of any part of the work, even ones in which they  
494 were not personally involved, are appropriately investigated, resolved, and the  
495 resolution documented in the literature.

496

## 497 **Conflicts of interest**

498 The authors declare no competing interests.

499

## 500 **Data availability**

501 All raw data are available from CellMiner [80] Version 2.2. All other data are  
502 provided in the manuscript and its supplements.

503

## 504 **References**

- 505 1. França TF, Monserrat JM. Reproducibility crisis in science or unrealistic  
506 expectations? EMBO Rep. 2018 Jun;19(6). pii: e46008.
- 507 2. Wass MN, Ray L, Michaelis M. Understanding of researcher behavior is required  
508 to improve data reliability. Gigascience. 2019 May 1;8(5). pii: giz017.
- 509 3. Shoemaker RH. The NCI60 human tumour cell line anticancer drug screen. Nat  
510 Rev Cancer. 2006 Oct;6(10):813-23.
- 511 4. Reinhold WC, Sunshine M, Liu H, Varma S, Kohn K, Morris J, Doroshow J and  
512 Pommier Y. CellMiner: a web-based suite of genomic and pharmacologic tools to  
513 explore transcript and drug patterns in the NCI-60 cell line set. Cancer Res. 2012 Jul  
514 15; 72(14): 3499-3511.
- 515 5. Begley CG, Ioannidis JP. Reproducibility in science: improving the standard for  
516 basic and preclinical research. Circ Res. 2015 Jan 2;116(1):116-26.
- 517 6. Frye SV, Arkin MR, Arrowsmith CH, Conn PJ, Glicksman MA, Hull-Ryde EA,  
518 Slusher BS. Tackling reproducibility in academic preclinical drug discovery. Nat Rev  
519 Drug Discov. 2015 Nov;14(11):733-4.
- 520 7. Drucker DJ. Never Waste a Good Crisis: Confronting Reproducibility in  
521 Translational Research. Cell Metab. 2016 Sep 13;24(3):348-360.
- 522 8. Fanelli D. Opinion: Is science really facing a reproducibility crisis, and do we need  
523 it to? Proc Natl Acad Sci U S A. 2018 Mar 13;115(11):2628-2631.
- 524 9. Samsa G, Samsa L. A Guide to Reproducibility in Preclinical Research. Acad  
525 Med. 2019 Jan;94(1):47-52.
- 526 10. Munafò MR, Chambers CD, Collins AM, Fortunato L, Macleod MR. Research  
527 Culture and Reproducibility. Trends Cogn Sci. 2020 Feb;24(2):91-93.

- 528 11. Mobley A, Linder SK, Braeuer R, Ellis LM, Zwelling L. A survey on data  
529 reproducibility in cancer research provides insights into our limited ability to translate  
530 findings from the laboratory to the clinic. PLoS One 2013;8(5):e63221.
- 531 12. Baker M. 1,500 scientists lift the lid on reproducibility. Nature 2016;533:452-4.
- 532 13. Nature Editorial. Checklists work to improve science. Nature 2018;556:273-4.
- 533 14. Prinz F, Schlange T, Asadullah K. Believe it or not: how much can we rely on  
534 published data on potential drug targets? Nat Rev Drug Discov. 2011;10:712.
- 535 15. Begley CG, Ellis LM. Drug development: Raise standards for preclinical cancer  
536 research. Nature 2012;483:531-3.
- 537 16. Aird F, Kandela I, Mantis C; Reproducibility Project: Cancer Biology. Replication  
538 Study: BET bromodomain inhibition as a therapeutic strategy to target c-Myc. Elife  
539 2017;6. pii: e21253.
- 540 17. Kandela I, Aird F; Reproducibility Project: Cancer Biology. Replication Study:  
541 Discovery and preclinical validation of drug indications using compendia of public  
542 gene expression data. Elife 2017;6. pii: e17044.
- 543 18. Shan X, Fung JJ, Kosaka A, Danet-Desnoyers G; Reproducibility Project: Cancer  
544 Biology. Replication Study: Inhibition of BET recruitment to chromatin as an effective  
545 treatment for MLL-fusion leukaemia. Elife 2017;6. pii: e25306.
- 546 19. Showalter MR, Hatakeyama J, Cajka T, VanderVorst K, Carraway KL, Fiehn O;  
547 Reproducibility Project: Cancer Biology. Replication Study: The common feature of  
548 leukemia-associated IDH1 and IDH2 mutations is a neomorphic enzyme activity  
549 converting alpha-ketoglutarate to 2-hydroxyglutarate. Elife 2017;6. pii: e26030.
- 550 20. Essex A, Pineda J, Acharya G, Xin H, Evans J; Reproducibility Project: Cancer  
551 Biology, Iorns E, Tsui R, Denis A, Perfito N, Errington TM, Iorns E, Tsui R, Denis A,  
552 Perfito N, Errington TM. Replication Study: Wnt activity defines colon cancer stem

- 553 cells and is regulated by the microenvironment. *Elife*. 2019 Jun 19;8. pii: e45426.  
554 doi: 10.7554/eLife.45426.
- 555 21. Horrigan SK; Reproducibility Project: Cancer Biology. Replication Study: The  
556 CD47-signal regulatory protein alpha (SIRPa) interaction is a therapeutic target for  
557 human solid tumors. *Elife* 2017;6. pii: e18173.
- 558 22. Horrigan SK, Courville P, Sampey D, Zhou F, Cai S; Reproducibility Project:  
559 Cancer Biology. Replication Study: Melanoma genome sequencing reveals frequent  
560 PREX2 mutations. *Elife* 2017;6. pii: e21634. doi: 10.7554/eLife.21634.
- 561 23. Lewis LM, Edwards MC, Meyers ZR, Talbot CC, Hao H, Blum D, Reproducibility  
562 Project: Cancer Biology, Iorns E, Tsui R, Denis A, Perfito N, Errington TM.  
563 Replication Study: Transcriptional amplification in tumor cells with elevated c-Myc.  
564 *Elife* 2018;7. pii: e30274.
- 565 24. Eaton K, Pirani A, Snitkin ES; Reproducibility Project: Cancer Biology, Iorns E,  
566 Tsui R, Denis A, Perfito N, Errington TM. Replication Study: Intestinal inflammation  
567 targets cancer-inducing activity of the microbiota. *Elife* 2018;7. pii: e34364.
- 568 25. Kim J, Afshari A, Sengupta R, Sebastiano V, Gupta A, Kim YH; Reproducibility  
569 Project: Cancer Biology, Iorns E, Tsui R, Denis A, Perfito N, Errington TM.  
570 Replication study: Melanoma exosomes educate bone marrow progenitor cells  
571 toward a pro-metastatic phenotype through MET. *Elife*. 2018 Dec 11;7. pii: e39944.
- 572 26. Vanden Heuvel JP, Maddox E, Maalouf SW; Reproducibility Project: Cancer  
573 Biology, Iorns E, Tsui R, Denis A, Perfito N, Errington TM. Replication Study:  
574 Systematic identification of genomic markers of drug sensitivity in cancer cells. *Elife*  
575 2018;7. pii: e29747.
- 576 27. Sheen MR, Fields JL, Northan B, Lacoste J, Ang LH, Fiering S; Reproducibility  
577 Project: Cancer Biology, Iorns E, Tsui R, Denis A, Perfito N, Errington TM.

- 578 Replication Study: Biomechanical remodeling of the microenvironment by stromal  
579 caveolin-1 favors tumor invasion and metastasis. *Elife*. 2019 Dec 17;8. pii: e45120.  
580 doi: 10.7554/eLife.45120.
- 581 28. Kerwin J, Khan I; Reproducibility Project: Cancer Biology, Iorns E, Tsui R, Denis  
582 A, Perfito N, Errington TM. Replication Study: A coding-independent function of gene  
583 and pseudogene mRNAs regulates tumour biology. *Elife*. 2020 Apr 21;9:e51019.
- 584 29. Mantis C, Kandela I, Aird F; Reproducibility Project: Cancer Biology. Replication  
585 Study: Coadministration of a tumor-penetrating peptide enhances the efficacy of  
586 cancer drugs. *Elife* 2017;6. pii: e17584. doi: 10.7554/eLife.17584.
- 587 30. Repass J; Reproducibility Project: Cancer Biology, Iorns E, Denis A, Williams  
588 SR, Perfito N, Errington TM. Replication Study: *Fusobacterium nucleatum* infection is  
589 prevalent in human colorectal carcinoma. *Elife* 2018;7. pii: e25801.
- 590 31. Yan X, Tang B, Chen B, Shan Y, Yang H; Reproducibility Project: Cancer  
591 Biology, Iorns E, Tsui R, Denis A, Perfito N, Errington TM. Replication Study: The  
592 microRNA miR-34a inhibits prostate cancer stem cells and metastasis by directly  
593 repressing CD44. *Elife*. 2019 Mar 12;8. pii: e43511.
- 594 32. Wang H, Radomska HS, Phelps MA; Reproducibility Project: Cancer Biology,  
595 Iorns E, Tsui R, Denis A, Perfito N, Errington TM. Replication Study: Coding-  
596 independent regulation of the tumor suppressor PTEN by competing endogenous  
597 mRNAs. *Elife*. 2020 Oct 19;9:e56651.
- 598 33. Fanelli D, Costas R, Ioannidis JP. Meta-assessment of bias in science. *Proc Natl*  
599 *Acad Sci U S A*. 2017;114:3714-9.
- 600 34. Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S,  
601 Wilson CJ, Lehár J, Kryukov GV, Sonkin D, Reddy A, Liu M, Murray L, Berger MF,  
602 Monahan JE, Morais P, Meltzer J, Korejwa A, Jané-Valbuena J, Mapa FA, Thibault



603 J, Bric-Furlong E, Raman P, Shipway A, Engels IH, Cheng J, Yu GK, Yu J, Aspesi P  
604 Jr, de Silva M, Jagtap K, Jones MD, Wang L, Hatton C, Palescandolo E, Gupta S,  
605 Mahan S, Sougnez C, Onofrio RC, Liefeld T, MacConaill L, Winckler W, Reich M, Li  
606 N, Mesirov JP, Gabriel SB, Getz G, Ardlie K, Chan V, Myer VE, Weber BL, Porter J,  
607 Warmuth M, Finan P, Harris JL, Meyerson M, Golub TR, Morrissey MP, Sellers WR,  
608 Schlegel R, Garraway LA. The Cancer Cell Line Encyclopedia enables predictive  
609 modelling of anticancer drug sensitivity. *Nature*. 2012 Mar 28;483(7391):603-7.

610 35. Garnett MJ, Edelman EJ, Heidorn SJ, Greenman CD, Dastur A, Lau KW,  
611 Greninger P, Thompson IR, Luo X, Soares J, Liu Q, Iorio F, Surdez D, Chen L,  
612 Milano RJ, Bignell GR, Tam AT, Davies H, Stevenson JA, Barthorpe S, Lutz SR,  
613 Kogera F, Lawrence K, McLaren-Douglas A, Mitropoulos X, Mironenko T, Thi H,  
614 Richardson L, Zhou W, Jewitt F, Zhang T, O'Brien P, Boisvert JL, Price S, Hur W,  
615 Yang W, Deng X, Butler A, Choi HG, Chang JW, Baselga J, Stamenkovic I,  
616 Engelman JA, Sharma SV, Delattre O, Saez-Rodriguez J, Gray NS, Settleman J,  
617 Futreal PA, Haber DA, Stratton MR, Ramaswamy S, McDermott U, Benes CH.  
618 Systematic identification of genomic markers of drug sensitivity in cancer cells.  
619 *Nature*. 2012 Mar 28;483(7391):570-5.

620 36. Haibe-Kains B, El-Hachem N, Birkbak NJ, Jin AC, Beck AH, Aerts HJ,  
621 Quackenbush J. Inconsistency in large pharmacogenomic studies. *Nature*. 2013 Dec  
622 19;504(7480):389-93.

623 37. Cancer Cell Line Encyclopedia Consortium; Genomics of Drug Sensitivity in  
624 Cancer Consortium. Pharmacogenomic agreement between two cancer cell line data  
625 sets. *Nature*. 2015 Dec 3;528(7580):84-7.

- 626 38. Bouhaddou M, DiStefano MS, Riesel EA, Carrasco E, Holzapfel HY, Jones DC,  
627 Smith GR, Stern AD, Somani SS, Thompson TV, Birtwistle MR. Drug response  
628 consistency in CCLE and CGP. *Nature* 2016;540:E9-E10.
- 629 39. Geeleher P, Gamazon ER, Seoighe C, Cox NJ, Huang RS. Consistency in large  
630 pharmacogenomic studies. *Nature* 2016;540:E1-E2.
- 631 40. Mpindi JP, Yadav B, Östling P, Gautam P, Malani D, Murumägi A, Hirasawa A,  
632 Kangaspeska S, Wennerberg K, Kallioniemi O, Aittokallio T. Consistency in drug  
633 response profiling. *Nature* 2016;540:E5-E6.
- 634 41. Safikhani Z, El-Hachem N, Smirnov P, Freeman M, Goldenberg A, Birkbak NJ,  
635 Beck AH, Aerts HJ, Quackenbush J, Haibe-Kains B. Safikhani et al. reply. *Nature*.  
636 2016 Nov 30;540(7631):E2-E4.
- 637 42. Safikhani Z, El-Hachem N, Smirnov P, Freeman M, Goldenberg A, Birkbak NJ,  
638 Beck AH, Aerts HJ, Quackenbush J, Haibe-Kains B. Safikhani et al. reply. *Nature*.  
639 2016 Nov 30;540(7631):E6-E8.
- 640 43. Safikhani Z, El-Hachem N, Smirnov P, Freeman M, Goldenberg A, Birkbak NJ,  
641 Beck AH, Aerts HJ, Quackenbush J, Haibe-Kains B. Safikhani et al. reply. *Nature*.  
642 2016 Nov 30;540(7631):E11-E12.
- 643 44. Safikhani Z, El-Hachem N, Quevedo R, Smirnov P, Goldenberg A, Juul Birkbak  
644 N, Mason C, Hatzis C, Shi L, Aerts HJ, Quackenbush J, Haibe-Kains B. *F1000Res*.  
645 2016 May 9;5:825.
- 646 45. Safikhani Z, Smirnov P, Freeman M, El-Hachem N, She A, Rene Q, Goldenberg  
647 A, Birkbak NJ, Hatzis C, Shi L, Beck AH, Aerts HJWL, Quackenbush J, Haibe-Kains  
648 B. Revisiting inconsistency in large pharmacogenomic studies. Version 3. *F1000Res*.  
649 2016 Sep 16 [revised 2017 Jan 1];5:2333.

650 46. Monks A, Scudiero DA, Johnson GS, Paull KD, Sausville EA. The NCI anti-  
651 cancer drug screen: a smart screen to identify effectors of novel targets. *Anticancer*  
652 *Drug Des.* 1997 Oct;12(7):533-41.

653 47. Reinhold WC, Varma S, Sousa F, Sunshine M, Abaan OD, Davis SR, Reinhold  
654 SW, Kohn KW, Morris J, Meltzer PS, Doroshow JH, Pommier Y. NCI-60 whole  
655 exome sequencing and pharmacological CellMiner analyses. *PLoS One.* 2014 Jul  
656 17;9(7):e101670.

657 48. Reinhold WC, Sunshine M, Varma S, Doroshow JH, Pommier Y. Using CellMiner  
658 1.6 for Systems Pharmacology and Genomic Analysis of the NCI-60. *Clin Cancer*  
659 *Res.* 2015 Sep 1;21(17):3841-52.

660 49. Chabner BA. NCI-60 Cell Line Screening: A Radical Departure in its Time. *J Natl*  
661 *Cancer Inst.* 2016 Jan 11;108(5). pii: djv388.

662 50. Reinhold WC, Varma S, Sunshine M, Rajapakse V, Luna A, Kohn KW,  
663 Stevenson H, Wang Y, Heyn H, Nogales V, Moran S, Goldstein DJ, Doroshow JH,  
664 Meltzer PS, Esteller M, Pommier Y. The NCI-60 Methylome and Its Integration into  
665 CellMiner. *Cancer Res.* 2017 Feb 1;77(3):601-612.

666 51. Reinhold WC, Varma S, Sunshine M, Elloumi F, Ofori-Atta K, Lee S, Trepel JB,  
667 Meltzer PS, Doroshow JH, Pommier Y. RNA sequencing of the NCI-60: Integration  
668 into CellMiner and CellMiner CDB. *Cancer Res.* 2019 May 21. pii: canres.2047.2018.

669 52. Kormanec J, Novakova R, Csolleiova D, Feckova L, Rezuchova B, Sevcikova B,  
670 Homerova D. The antitumor antibiotic mithramycin: new advanced approaches in  
671 modification and production. *Appl Microbiol Biotechnol.* 2020 Sep;104(18):7701-  
672 7721.

673 53. Gallego-Jara J, Lozano-Terol G, Sola-Martínez RA, Cánovas-Díaz M, de Diego  
674 Puente T. A Compressive Review about Taxol®: History and Future Challenges.  
675 Molecules. 2020 Dec 17;25(24):5986.

676 54. Rae JM, Creighton CJ, Meck JM, Haddad BR, Johnson MD. MDA-MB-435 cells  
677 are derived from M14 melanoma cells--a loss for breast cancer, but a boon for  
678 melanoma research. Breast Cancer Res Treat. 2007 Jul;104(1):13-9.

679 55. Ben-David U, Siranosian B, Ha G, Tang H, Oren Y, Hinohara K, Strathdee CA,  
680 Dempster J, Lyons NJ, Burns R, Nag A, Kugener G, Cimini B, Tsvetkov P, Maruvka  
681 YE, O'Rourke R, Garrity A, Tubelli AA, Bandopadhyay P, Tsherniak A, Vazquez F,  
682 Wong B, Birger C, Ghandi M, Thorner AR, Bittker JA, Meyerson M, Getz G,  
683 Beroukheim R, Golub TR. Genetic and transcriptional evolution alters cancer cell line  
684 drug response. Nature. 2018 Aug;560(7718):325-330.

685 56. Noronha N, Ehx G, Meunier MC, Laverdure JP, Thériault C, Perreault C. Major  
686 multilevel molecular divergence between THP-1 cells from different biorepositories.  
687 Int J Cancer. 2020 Mar 12. doi: 10.1002/ijc.32967.

688 57. Voskuil JLA, Bandrowski A, Begley CG, Bradbury ARM, Chalmers AD, Gomes  
689 AV, Hardcastle T, Lund-Johansen F, Plückthun A, Roncador G, Solache A, Taussig  
690 MJ, Trimmer JS, Williams C, Goodman SL. The Antibody Society's antibody  
691 validation webinar series. MAbs. 2020 Jan-Dec;12(1):1794421.

692 58. Hubert M, Vandervieren E. An adjusted boxplot for skewed distributions.  
693 Computational Statistics and Data Analysis. 2008; 52(12): 5186-5201

694 59. Liu Y, Mi Y, Mueller T, Kriebich S, Williams EG, Van Drogen A, Borel C, Frank M,  
695 Germain PL, Bludau I, Mehnert M, Seifert M, Emmenlauer M, Sorg I, Bezrukov F,  
696 Bena FS, Zhou H, Dehio C, Testa G, Saez-Rodriguez J, Antonarakis SE, Hardt WD,

697 Aebersold R. Multi-omic measurements of heterogeneity in HeLa cells across  
698 laboratories. *Nat Biotechnol.* 2019 Mar;37(3):314-322.

699 60. Lorenzi PL, Reinhold WC, Varma S, Hutchinson AA, Pommier Y, Chanock SJ,  
700 Weinstein JN. DNA fingerprinting of the NCI-60 cell line panel. *Mol Cancer Ther.*  
701 2009 Apr;8(4):713-24.

702 61. Eisenhauer EA, O'Dwyer PJ, Christian M, Humphrey JS. Phase I clinical trial  
703 design in cancer drug development. *J Clin Oncol.* 2000 Feb;18(3):684-92.

704 62. Sachs JR, Mayawala K, Gadamsetty S, Kang SP, de Alwis DP. Optimal Dosing  
705 for Targeted Therapies in Oncology: Drug Development Cases Leading by Example.  
706 *Clin Cancer Res.* 2016 Mar 15;22(6):1318-24.

707 63. Mansinho A, Boni V, Miguel M, Calvo E. New designs in early clinical drug  
708 development. *Ann Oncol.* 2019 Sep 1;30(9):1460-1465.

709 64. Corbaux P, El-Madani M, Tod M, Péron J, Maillet D, Lopez J, Freyer G, You B.  
710 Clinical efficacy of the optimal biological dose in early-phase trials of anti-cancer  
711 targeted therapies. *Eur J Cancer.* 2019 Oct;120:40-46.

712 65. Begley CG. Six red flags for suspect work. *Nature* 2013;497:433-4.

713 66. Ioannidis JP, Greenland S, Hlatky MA, Khoury MJ, Macleod MR, Moher D,  
714 Schulz KF, Tibshirani R. Increasing value and reducing waste in research design,  
715 conduct, and analysis. *Lancet* 2014;383:166-75.

716 67. Hatzis C, Bedard PL, Birkbak NJ, Beck AH, Aerts HJ, Stem DF, Shi L, Clarke R,  
717 Quackenbush J, Haibe-Kains B. Enhancing reproducibility in cancer drug screening:  
718 how do we move forward? *Cancer Res.* 2014 Aug 1;74(15):4016-23.

719 68. Freedman LP, Cockburn IM, Simcoe TS. The Economics of Reproducibility in  
720 Preclinical Research. *PLoS Biol.* 2015;13:e1002165.

721 69. Jarvis MF, Williams M. Irreproducibility in Preclinical Biomedical Research:  
722 Perceptions, Uncertainties, and Knowledge Gaps. Trends Pharmacol Sci.  
723 2016;37:290-302.

724 70. Freedman LP, Venugopalan G, Wisman R. Reproducibility2020: Progress and  
725 priorities. F1000Res. 2017;6:604.

726 71. Kaelin WG Jr. Publish houses of brick, not mansions of straw. Nature  
727 2017;545:387.

728 72. Karp NA, Speak AO, White JK, Adams DJ, Hrabé de Angelis M, Hérault Y, Mott  
729 RF. Impact of temporal variation on design and analysis of mouse knockout  
730 phenotyping studies. PLoS One. 2014 Oct 24;9(10):e111239.

731 73. Ding KF, Finlay D, Yin H, Hendricks WPD, Sereduk C, Kiefer J, Sekulic A,  
732 LoRusso PM, Vuori K, Trent JM, Schork NJ. Analysis of variability in high throughput  
733 screening data: applications to melanoma cell lines and drug responses. Oncotarget.  
734 2017 Apr 25;8(17):27786-27799.

735 74. Kafkafi N, Golani I, Jaljuli I, Morgan H, Sarig T, Würbel H, Yaacoby S, Benjamini  
736 Y. Addressing reproducibility in single-laboratory phenotyping experiments. Nat  
737 Methods. 2017 Apr 27;14(5):462-464.

738 75. Karp NA. Reproducible preclinical research-Is embracing variability the answer?  
739 PLoS Biol. 2018 Mar 5;16(3):e2005413.

740 76. Milcu A, Puga-Freitas R, Ellison AM, Blouin M, Scheu S, Freschet GT, Rose L,  
741 Barot S, Cesarz S, Eisenhauer N, Girin T, Assandri D, Bonkowski M, Buchmann N,  
742 Butenschoen O, Devidal S, Gleixner G, Gessler A, Gigon A, Greiner A, Grignani C,  
743 Hansart A, Kayler Z, Lange M, Lata JC, Le Galliard JF, Lukac M, Mannerheim N,  
744 Müller MEH, Pando A, Rotter P, Scherer-Lorenzen M, Seyhun R, Urban-Mead K,

745 Weigelt A, Zavattaro L, Roy J. Genotypic variability enhances the reproducibility of  
746 an ecological study. *Nat Ecol Evol.* 2018 Feb;2(2):279-287.

747 77. Voelkl B, Vogt L, Sena ES, Würbel H. Reproducibility of preclinical animal  
748 research improves with heterogeneity of study samples. *PLoS Biol.* 2018 Feb  
749 22;16(2):e2003693.

750 78. Bodden C, von Kortzfleisch VT, Karwinkel F, Kaiser S, Sachser N, Richter SH.  
751 Heterogenising study samples across testing time improves reproducibility of  
752 behavioural data. *Sci Rep.* 2019 Jun 3;9(1):8247.

753 79. von Kortzfleisch VT, Karp NA, Palme R, Kaiser S, Sachser N, Richter SH.  
754 Improving reproducibility in animal research by splitting the study population into  
755 several 'mini-experiments'. *Sci Rep.* 2020 Oct 6;10(1):16579.

756 80. Shankavaram UT, Varma S, Kane D, Sunshine M, Chary KK, Reinhold WC,  
757 Pommier Y, Weinstein JN. CellMiner: a relational database and query tool for the  
758 NCI-60 cancer cell lines. *BMC Genomics.* 2009 Jun 23;10:277.

759

760