

SGN - Assignment #1

Student Name, 123456

1 Periodic orbit

Exercise 1

Consider the 3D Earth–Moon Circular Restricted Three-Body Problem with $\mu=0.012150$. Note that the CRTBP has an integral of motion, that is, the Jacobi constant

$$J(x, y, z, v_x, x_y, v_z) := 2\Omega(x, y, z) - v^2 = C$$

where
$$\Omega(x,y,z) = \frac{1}{2}(x^2+y^2) + \frac{1-\mu}{r_1} + \frac{\mu}{r_2} + \frac{1}{2}\mu(1-\mu)$$
 and $v^2 = v_x^2 + v_y^2 + v_z^2$.

1) Find the coordinates of the five Lagrange points L_i in the rotating, adimensional reference frame with at least 10-digit accuracy and report their Jacobi constant C_i .

Solutions to the 3D CRTBP satisfy the symmetry

$$S: (x, y, z, \dot{x}, \dot{y}, \dot{z}, t) \rightarrow (x, -y, z, -\dot{x}, \dot{y}, -\dot{z}, -t).$$

Thus, a trajectory that crosses perpendicularly the y=0 plane twice is a periodic orbit.

2) Given the initial guess $\mathbf{x}_0 = (x_0, y_0, z_0, v_{x0}, v_{y0}, v_{z0})$, with

 $x_0 = 1.068792441776$

 $y_0 = 0$

 $z_0 = 0.071093328515$

 $v_{r0} = 0$

 $v_{v0} = 0.319422926485$

 $v_{\sim 0} = 0$

Find the periodic halo orbit having a Jacobi Constant C=3.09; that is, develop the theoretical framework and implement a differential correction scheme that uses the STM, either approximated through finite differences **or** achieved by integrating the variational equation.

Hint: Consider working on $\varphi(\mathbf{x} + \Delta \mathbf{x}, t + \Delta t)$ and $J(\mathbf{x} + \Delta \mathbf{x})$ and then enforce perpendicular cross of y = 0 and Jacobi energy.

The periodic orbits in the CRTBP exist in families. These can be computed by 'continuing' the orbits along one coordinate or one parameter, e.g., the Jacobi energy C. The numerical continuation is an iterative process in which the desired variable is gradually varied, while the rest of the initial guess is taken from the solution of the previous iteration, thus aiding the convergence process.

3) By gradually decreasing C and using numerical continuation, compute the families of halo orbits until C=3.04.

(8 points)

To prepare the ZIP file for the submission of the Assignment:

- Complete your answers on the Overleaf project you created.
- Red text indicates where answers are needed; be sure there is no red stuff in your report.



- In your answers, <u>be concise</u>: to the point.
- Download the PDF from the Main menu on Overleaf.
- Create a single .zip file containing both the report in PDF and the MATLAB files. The name shall be lastname123456_Assign1.zip.
- Deadline for the submission: Dec 20 2024, 23:59.
- Load the compressed file to the Assignments folder on Webeep.

Write your answer here

- Develop the exercise in the file lastname123456_Assign1_Ex1.m
- Organize the script in sections, one for each point; use local functions if needed.

Fill the table with the required results. Use 10-digits

| | L_1 | L_2 | L_3 | L_4 | L_5 |
|----------------|--------------------|--------------------|---------------------|--------------------|--------------------|
| \overline{x} | ± 0.0000000000 | ± 0.0000000000 | ± 0.0000000000 | ± 0.0000000000 | ± 0.0000000000 |
| \overline{y} | ± 0.0000000000 | ± 0.0000000000 | ± 0.0000000000 | ± 0.0000000000 | ± 0.0000000000 |
| \overline{C} | ± 0.0000000000 | ± 0.0000000000 | ± 0.00000000000 | ± 0.0000000000 | ± 0.0000000000 |

Table 1: Lagrangian points coordinates and Jacobi constants

| \overline{x} | y | z | |
|--------------------|--------------------|--------------------|--|
| ± 0.0000000000 | ± 0.0000000000 | ± 0.0000000000 | |
| | | | |
| v_x | v_x | v_x | |
| | | | |

Table 2: Corrected initial state of the halo orbit with C = 3.09

| x | y | z |
|--------------------|--------------------|--------------------|
| ± 0.0000000000 | ± 0.0000000000 | ± 0.0000000000 |
| | | |
| v_x | v_y | v_x |
| ± 0.0000000000 | ± 0.0000000000 | ± 0.0000000000 |

Table 3: Corrected initial state of the halo orbit with C = 3.04

2 Impulsive guidance

Exercise 2

Consider the two-impulse transfer problem stated in Section 3.1 (Topputo, 2013)*.

^{*}F. Topputo, "On optimal two-impulse Earth–Moon transfers in a four-body model", Celestial Mechanics and Dynamical Astronomy, Vol. 117, pp. 279–313, 2013, DOI: 10.1007/s10569-013-9513-8.



1) Using the procedure in Section 3.2, produce a first guess solution using $\alpha = 0.2\pi$, $\beta = 1.41$, $\delta = 4$, and $t_i = 2$. Plot the solution in both the rotating frame and Earth-centered inertial frame (see Appendix 1 in (Topputo, 2013)). Consider the parameters listed in Table 4 and extract the radius and gravitational parameters of the Earth and Moon from the provided kernels and use the latter to compute the parameter μ .

| Symbol | Value | Units | Meaning |
|------------------|------------------------------|-----------------|------------------------------------|
| $\overline{m_s}$ | 3.28900541×10^5 | - | Scaled mass of the Sun |
| ho | 3.88811143×10^2 | - | Scaled Sun-(Earth+Moon) distance |
| ω_s | $-9.25195985 \times 10^{-1}$ | - | Scaled angular velocity of the Sun |
| ω_{em} | $2.66186135 \times 10^{-1}$ | s^{-1} | Earth-Moon angular velocity |
| l_{em} | 3.84405×10^8 | \mathbf{m} | Earth-Moon distance |
| h_i | 167 | km | Altitude of departure orbit |
| h_i | 100 | km | Altitude of arrival orbit |
| \overline{DU} | 3.84405000×10^5 | km | Distance Unit |
| TU | 4.34256461 | days | Time Unit |
| VU | 1.02454018 | $\mathrm{km/s}$ | Velocity Unit |

Table 4: Constants to be considered to solve the PCRTBP. The units of distance, time, and velocity are used to map scaled quantities into physical units.

- 2) Considering the first guess in 1) and using $\{\mathbf{x}_i, t_i, t_f\}$ as variables, solve the problem in Section 3.1 with simple shooting in the following cases
 - a) without providing any derivative to the solver, and
 - b) by providing the derivatives and by estimating the state transition matrix with variational equations.
- 3) Considering the first guess solution in 1) and the procedure in Section 3.3, solve the problem with multiple shooting taking N=4 and using the variational equation to compute the Jacobian of the nonlinear equality constraints.
- 4) Perform an n-body propagation using the solution $\{\mathbf{x}_i, t_i, t_f\}$ obtained in point 2), transformed in Earth-centered inertial frame and into physical units. To move from 2-D to 3-D, assume that the position and velocity components in inertial frame are $r_z(t_i) = 0$ and $v_z(t_i) = 0$. To perform the propagation it is necessary to identify the epoch t_i . This can be done by mapping the relative position of the Earth, Moon and Sun in the PCRTBP to a similar condition in the real world:
 - a) Consider the definition of $\theta(t)$ provided in Section 2.2 to compute the angle $\theta_i = \theta(t_i)$. Note that this angle corresponds to the angle between the rotating frame x-axis, aligned to the position vector from the Earth-Moon System Barycenter (EMB) to the Moon, and the Sun direction.
 - b) The angle θ ranges between $[0, 2\pi]$ and it covers this domain in approximately the revolution period of the Moon around the Earth.
 - c) Solve a zero-finding problem to determine the epoch at which the angle Moon-EMB-Sun is equal to θ_i , considering as starting epoch 2024 Sep 28 00:00:00.000 TDB. **Hints**: Exploit the SPK kernels to define the orientation of the rotating frame axes in the inertial frame for an epoch t. Consider only the projection of the EMB-Sun position vector onto the so-defined x-y plane to compute the angle (planar motion).

Plot the propagated orbit and compare it to the previously found solutions.

(11 points)

Write your answer here

- Develop the exercise in the file lastname123456_Assign1_Ex2.m
- Organize the script in sections, one for each point; use local functions if needed.

$$r_{x,0} [DU]$$
 $r_{y,0} [DU]$ $v_{x,0} [VU]$ $v_{y,0} [VU]$ ± 0.000000 ± 0.000000 ± 0.000000

Table 5: Initial guess in Earth-Moon rotating frame.

| ${f Gradients}$ | $r_{x,0}$ [DU] | $r_{y,0} [DU]$ | $v_{x,0} [VU]$ | $v_{y,0} [VU]$ | $t_i [TU]$ | $t_f [TU]$ |
|-----------------|----------------|----------------|----------------|----------------|------------|------------|
| False | ± 0.000000 | ± 0.000000 | ± 0.000000 | ± 0.000000 | 0.000 | 0.000 |
| True | ± 0.000000 | ± 0.000000 | ± 0.000000 | ± 0.000000 | 0.000 | 0.000 |

Table 6: Simple shooting solutions in the Earth-Moon rotating frame.

Table 7: Multiple shooting solution in the Earth-Moon rotating frame.



| Symbol | Calendar epoch (UTC) | | | |
|------------------|----------------------|------------------|------------------|--|
| $\overline{t_i}$ | YYY | Y-MM-DDTH | H:MM:SS.sss | |
| t_f | YYY | Y-MM-DDTH | H:MM:SS.sss | |
| - | | | | |
| $r_{x,0}$ [A | km] | $r_{y,0} [km]$ | $r_{z,0} [km]$ | |
| ± 0.0000 | 00000 | ± 0.00000000 | 0.0 | |
| | | | | |
| $v_{x,0}$ [km | n/s] | $v_{y,0} [km/s]$ | $v_{z,0} [km/s]$ | |
| ± 0.00000 | 0000 | ± 0.00000000 | 0.0 | |

Table 8: Initial epoch, final epoch, and initial state in Earth-centered inertial frame.



3 Continuous guidance

Exercise 3

To be disclosed on Oct 30 2024.

(11 points)

Write your answer here

- Develop the exercise in the file lastname123456_Assign1_Ex3.m
- Organize the script in sections, one for each point; use local functions if needed.