

Clique is hard on average for regular resolution

Ilario Bonacina, UPC Barcelona Tech

July 20, 2018

RaTLoCC, Bertinoro

How hard is to certify that a graph is Ramsey?

Ilario Bonacina, UPC Barcelona Tech

July 20, 2018

RaTLoCC, Bertinoro

Talk based on a joint work with:



A. Atserias



S. de Rezende



M. Lauria



J. Nordström



A. Razborov

How hard is to certify that a graph is Ramsey?

A graph G with n vertices we say that is k -Ramsey if it has no set of k vertices forming a clique or an independent set.

If $k = \lceil 2 \log_2 n \rceil$ we just say that G is Ramsey.

How hard is to certify that a graph is **Ramsey**?

A graph G with n vertices we say that is **k -Ramsey** if it has no set of k vertices forming a clique or an independent set.

If $k = \lceil 2 \log_2 n \rceil$ we just say that G is **Ramsey**.

Erdős-Rényi random graphs

A graph $G = (V, E) \sim \mathcal{G}(n, p)$ is such that $|V| = n$ and each edge $\{u, v\} \in E$ independently with prob. $p \in [0, 1]$

How hard is to certify that a graph is **Ramsey**?

A graph G with n vertices we say that is **k -Ramsey** if it has no set of k vertices forming a clique or an independent set.

If $k = \lceil 2 \log_2 n \rceil$ we just say that G is **Ramsey**.

Erdős-Rényi random graphs

A graph $G = (V, E) \sim \mathcal{G}(n, p)$ is such that $|V| = n$ and each edge $\{u, v\} \in E$ independently with prob. $p \in [0, 1]$

- if $p \ll n^{-2/(k-1)}$ then a.a.s. $G \sim \mathcal{G}(n, p)$ has no k -cliques
- A.a.s. $G \sim \mathcal{G}(n, \frac{1}{2})$ is Ramsey

How hard is to certify that a graph is Ramsey?

Construct a propositional formula $\Psi_{G,k}$ unsatisfiable if and only if
“ G is k -Ramsey”

How hard is to certify that a graph is Ramsey?

Construct a propositional formula $\Psi_{G,k}$ unsatisfiable if and only if
“ G is k -Ramsey”

$x_{v,j} \equiv$ “ v is the j -th vertex of a k -clique in G
or the j -th vertex of a k -independent set”.

How hard is to certify that a graph is Ramsey?

Construct a propositional formula $\Psi_{G,k}$ unsatisfiable if and only if “ G is k -Ramsey”

$x_{v,j} \equiv$ “ v is the j -th vertex of a k -clique in G
or the j -th vertex of a k -independent set”.

$$\bigvee_{v \in V} x_{v,i} \quad \text{for } i \in [k]$$

and

$$y \vee \neg x_{u,i} \vee \neg x_{v,j} \quad \text{for } i \neq j \in [k], u \neq v \in V, (u, v) \notin E$$

and

$$\neg y \vee \neg x_{u,i} \vee \neg x_{v,j} \quad \text{for } i \neq j \in [k], u \neq v \in V, (u, v) \in E$$

How hard is **to certify** that a graph is Ramsey?

Resolution

$$\neg y \vee \neg z$$

$$\neg x$$

$$clause_1 \vee var$$

$$clause_2 \vee \neg var$$

$$clause_1 \vee clause_2$$

$$y \vee \neg c$$


$$x \vee c$$

$$\neg x \vee z$$

$$\neg y$$


How hard is **to certify** that a graph is Ramsey?

Resolution




How hard is **to certify** that a graph is Ramsey?

Resolution




How hard is **to certify** that a graph is Ramsey?

Resolution




How hard is **to certify** that a graph is Ramsey?

Resolution




How hard is **to certify** that a graph is Ramsey?

Resolution




How hard is **to certify** that a graph is Ramsey?

Resolution



How hard is **to certify** that a graph is Ramsey?


Resolution




Tree-like = the proof DAG is a tree

Regular = no variable resolved twice in any source-to-sink path


Size = # of nodes in the proof DAG



Regular?



Regular? No.



Regular? **No**. And none of the shortest proofs is regular [HY87].

[HY87] Huang and Yu, 1987. *A DNF without regular shortest consensus path.*

What is Resolution good for?

- algorithms routinely used to solve NP-complete problems (hardware verification, ...) are *somewhat* formalizable in resolution
- the state-of-the-art algorithms to solve k-clique (Bron-Kerbosch, Östergård, Russian dolls algorithms, ...) are formalizable in *regular* resolution

What is Resolution good for?

- algorithms routinely used to solve NP-complete problems (hardware verification, ...) are *somewhat* formalizable in resolution
- the state-of-the-art algorithms to solve k-clique (Bron-Kerbosch, Östergård, Russian dolls algorithms, ...) are formalizable in *regular* resolution

[HKM16] All possible 2-colorings of $\{1, \dots, 7825\}$ have a monochromatic Pythagorean triple.

[HKM16] M. Heule, O. Kullmann and V. Marek, 2016. *Solving and Verifying the Boolean Pythagorean Triples problem via Cube-and-Conquer*

What is Resolution good for?

- algorithms routinely used to solve NP-complete problems (hardware verification, ...) are *somewhat* formalizable in resolution
- the state-of-the-art algorithms to solve k-clique (Bron-Kerbosch, Östergård, Russian dolls algorithms, ...) are formalizable in *regular* resolution

[HKM16] All possible 2-colorings of $\{1, \dots, 7825\}$ have a monochromatic Pythagorean triple.

This slide is too small to contain the 200Terabyte resolution proof...

[HKM16] M. Heule, O. Kullmann and V. Marek, 2016. *Solving and Verifying the Boolean Pythagorean Triples problem via Cube-and-Conquer*

Resolution size

Let ϕ be a conjunction of clauses in N variables with $|\phi| = N^{\mathcal{O}(1)}$

$S(\phi)$ = minimum size of a resolution refutation of ϕ

$S_{\text{tree}}(\phi)$ = minimum size of a **tree-like** resolution refutation of ϕ

$S_{\text{reg}}(\phi)$ = minimum size of a **regular** resolution refutation of ϕ

Resolution size

Let ϕ be a conjunction of clauses in N variables with $|\phi| = N^{\mathcal{O}(1)}$

$S(\phi)$ = minimum size of a resolution refutation of ϕ

$S_{tree}(\phi)$ = minimum size of a **tree-like** resolution refutation of ϕ

$S_{reg}(\phi)$ = minimum size of a **regular** resolution refutation of ϕ

- for every ϕ , $S(\phi) \leq S_{reg}(\phi) \leq S_{tree}(\phi)$
(and there are examples of exponential separations)
- for every ϕ , $S_{tree}(\phi) = 2^{\mathcal{O}(N)}$

How **hard** is to certify that a graph is Ramsey?

Theorem? (folklore)

$\Psi_{G,k}$, whenever unsatisfiable, has $S_{tree}(\Psi_{G,k}) = n^{\mathcal{O}(k)}$

How **hard** is to certify that a graph is Ramsey?

Theorem? (folklore)

$\Psi_{G,k}$, whenever unsatisfiable, has $S_{tree}(\Psi_{G,k}) = n^{\mathcal{O}(k)}$

Theorem [LPRT17]

If G is a Ramsey graph in n vertices and $k = \lceil 2 \log n \rceil$ then $S_{tree}(\Psi_{G,k}) = n^{\Omega(\log n)}$.

[LPRT17] Lauria, Pudlák, Rödl, and Thapen, 2017. *The complexity of proving that a graph is Ramsey.*

How **hard** is to certify that a graph is Ramsey?

Theorem? (folklore)

$\Psi_{G,k}$, whenever unsatisfiable, has $S_{tree}(\Psi_{G,k}) = n^{\mathcal{O}(k)}$

Theorem [LPRT17]

If G is a Ramsey graph in n vertices and $k = \lceil 2 \log n \rceil$ then $S_{tree}(\Psi_{G,k}) = n^{\Omega(\log n)}$.

Theorem

If $G \sim \mathcal{G}(n, \frac{1}{2})$ (hence in particular a.a.s. G is Ramsey) and $k = \lceil 2 \log n \rceil$ then $S_{reg}(\Psi_{G,k}) \stackrel{\text{a.a.s.}}{=} n^{\Omega(\log n)}$.

[LPRT17] Lauria, Pudlák, Rödl, and Thapen, 2017. *The complexity of proving that a graph is Ramsey.*

How **hard** is to certify that a graph is Ramsey?

Theorem? (folklore)

$\Psi_{G,k}$, whenever unsatisfiable, has $S_{tree}(\Psi_{G,k}) = n^{\mathcal{O}(k)}$

Theorem [LPRT17]

If G is a Ramsey graph in n vertices and $k = \lceil 2 \log n \rceil$ then $S_{tree}(\Psi_{G,k}) = n^{\Omega(\log n)}$.

Theorem

If $G \sim \mathcal{G}(n, \frac{1}{2})$ (hence in particular a.a.s. G is Ramsey) and $k = \lceil 2 \log n \rceil$ then $S_{reg}(\Psi_{G,k}) \stackrel{\text{a.a.s.}}{=} n^{\Omega(\log n)}$.

Open Problem

Let G be a Ramsey graph in n vertices and let $k = \lceil 2 \log n \rceil$. Is it true that $S(\Psi_{G,k}) = n^{\Omega(\log n)}$?

[LPRT17] Lauria, Pudlák, Rödl, and Thapen, 2017. *The complexity of proving that a graph is Ramsey.*

How hard is to certify that **a graph does not contain a k -clique?**

Construct a propositional formula $\Phi_{G,k}$ unsatisfiable if and only if
“ G does not contain a k -clique”

We already have it: $\Phi_{G,k} = \Psi_{G,k} \upharpoonright_{y=0}$

How hard is to certify that **a graph does not contain a k -clique?**

Construct a propositional formula $\Phi_{G,k}$ unsatisfiable if and only if
“ G does not contain a k -clique”

We already have it: $\Phi_{G,k} = \Psi_{G,k} \upharpoonright_{y=0}$

$x_{v,j} \equiv$ “ v is the j -th vertex of a k -clique in G ”.

How hard is to certify that **a graph does not contain a k -clique?**

Construct a propositional formula $\Phi_{G,k}$ unsatisfiable if and only if
“ G does not contain a k -clique”

We already have it: $\Phi_{G,k} = \Psi_{G,k} \upharpoonright_{y=0}$

$x_{v,j} \equiv$ “ v is the j -th vertex of a k -clique in G ”.

$$\bigvee_{v \in V} x_{v,i} \quad \text{for } i \in [k]$$

and

$$\neg x_{u,i} \vee \neg x_{v,j} \quad \text{for } i \neq j \in [k], u \neq v \in V, (u, v) \notin E$$

How hard is to certify that **a graph does not contain a k -clique?**

Construct a propositional formula $\Phi_{G,k}$ unsatisfiable if and only if
“ G does not contain a k -clique”

We already have it: $\Phi_{G,k} = \Psi_{G,k} \upharpoonright_{y=0}$

$x_{v,j} \equiv$ “ v is the j -th vertex of a k -clique in G ”.

$$\bigvee_{v \in V} x_{v,i} \quad \text{for } i \in [k]$$

and

$$\neg x_{u,i} \vee \neg x_{v,j} \quad \text{for } i \neq j \in [k], u \neq v \in V, (u, v) \notin E$$

How hard is to certify that **a graph does not contain a k -clique?**

Construct a propositional formula $\Phi_{G,k}$ unsatisfiable if and only if
“ G does not contain a k -clique”

We already have it: $\Phi_{G,k} = \Psi_{G,k} \upharpoonright_{y=0}$

$x_{v,j} \equiv$ “ v is the j -th vertex of a k -clique in G ”.

$$\bigvee_{v \in V} x_{v,i} \quad \text{for } i \in [k]$$

and

$$\neg x_{u,i} \vee \neg x_{v,j} \quad \text{for } i \neq j \in [k], u \neq v \in V, (u, v) \notin E$$

lower bounds on $S(\Phi_{G,k})$ imply lower bounds on $S(\Psi_{G,k})$

Overview of the literature: Upper Bounds

[~BGL13] if G is $(k - 1)$ -colorable then

$$S_{reg}(\Phi_{G,k}) \leq 2^k k^2 n^2$$

[folklore] $\Phi_{G,k}$, whenever unsatisfiable, has

$$S_{tree}(\Phi_{G,k}) = n^{\mathcal{O}(k)}$$

[BGL13] Beyersdorff, Galesi and Lauria 2013. *Parameterized complexity of DPLL search procedures*.

Overview of the literature: Lower Bounds

[BGL13] If G is the complete $(k - 1)$ -partite graph, then $S_{tree}(\Phi_{G,k}) = n^{\Omega(k)}$.

The same holds for $G \sim \mathcal{G}(n, p)$ with suitable edge density p .

[BIS07] for $n^{5/6} \ll k < \frac{n}{3}$ and $G \sim \mathcal{G}(n, p)$ (with suitable edge density p), then $S(\Phi_{G,k}) \stackrel{\text{a.a.s.}}{=} 2^{n^{\Omega(1)}}$

[LPRT17] if we encode k -clique using some other propositional encodings (e.g. in binary) we get $n^{\Omega(k)}$ size lower bounds for resolution

[BIS07] Beame, Impagliazzo and Sabharwal, 2007. *The resolution complexity of independent sets and vertex covers in random graphs.*

[LPRT17] Lauria, Pudlák, Rödl, and Thapen, 2017. *The complexity of proving that a graph is Ramsey.*

Main Result (simplified versions)

Main Theorem (version 1)

Let $G \sim \mathcal{G}(n, p)$ be an Erdős-Rényi random graph with, for simplicity, $p = n^{-4/(k-1)}$ and let $k \leq n^{1/2-\epsilon}$ for some arbitrary small ϵ . Then, $S_{\text{reg}}(\Phi_{G,k}) \stackrel{\text{a.a.s.}}{=} n^{\Omega(k)}$.

Main Result (simplified versions)

Main Theorem (version 1)

Let $G \sim \mathcal{G}(n, p)$ be an Erdős-Rényi random graph with, for simplicity, $p = n^{-4/(k-1)}$ and let $k \leq n^{1/2-\epsilon}$ for some arbitrary small ϵ . Then, $S_{\text{reg}}(\Phi_{G,k}) \stackrel{\text{a.a.s.}}{=} n^{\Omega(k)}$.

the actual lower bound decreases smoothly w.r.t. p

Main Result (simplified versions)

Main Theorem (version 1)

Let $G \sim \mathcal{G}(n, p)$ be an Erdős-Rényi random graph with, for simplicity, $p = n^{-4/(k-1)}$ and let $k \leq n^{1/2-\epsilon}$ for some arbitrary small ϵ . Then, $S_{\text{reg}}(\Phi_{G,k}) \stackrel{\text{a.a.s.}}{=} n^{\Omega(k)}$.

the actual lower bound decreases smoothly w.r.t. p

Main Theorem (version 2)

Let $G \sim \mathcal{G}(n, \frac{1}{2})$, then

$$S_{\text{reg}}(\Phi_{G,k}) \stackrel{\text{a.a.s.}}{=} n^{\Omega(\log n)} \text{ for } k = \mathcal{O}(\log n)$$

and

$$S_{\text{reg}}(\Phi_{G,k}) \stackrel{\text{a.a.s.}}{=} n^{\omega(1)} \text{ for } k = o(\log^2 n).$$

Focus on proving the following.

Theorem

Let $k = \lceil 2 \log n \rceil$ and $G \sim \mathcal{G}(n, \frac{1}{2})$, then $S_{reg}(\Phi_{G,k}) \stackrel{\text{a.a.s.}}{=} n^{\Omega(\log n)}$

$\hat{N}_W(R)$ is the set of common neighbors of R in W

$\hat{N}_W(R)$ is the set of common neighbors of R in W

W is (r, q) -dense if for every subset $R \subseteq V$ of size $\leq r$, it holds
 $|\hat{N}_W(R)| \geq q$

$\hat{N}_W(R)$ is the set of common neighbors of R in W

W is (r, q) -dense if for every subset $R \subseteq V$ of size $\leq r$, it holds $|\hat{N}_W(R)| \geq q$

Theorem 1

Let $k = \lceil 2 \log n \rceil$. A.a.s. $G = (V, E) \sim \mathcal{G}(n, \frac{1}{2})$ satisfies the following:

- (\star) V is $(\frac{k}{50}, \Theta(n^{0.9}))$ -dense; and
- ($\star\star$) For every $(\frac{k}{10000}, \Theta(n^{0.9}))$ -dense $W \subseteq V$ there exists $S \subseteq V$, $|S| \leq \sqrt{n}$ s.t. for every $R \subseteq V$, with $|R| \leq \frac{k}{50}$ and $|\hat{N}_W(R)| < \tilde{\Theta}(n^{0.6})$ it holds that $|R \cap S| \geq \frac{k}{10000}$.

$\hat{N}_W(R)$ is the set of common neighbors of R in W

W is (r, q) -dense if for every subset $R \subseteq V$ of size $\leq r$, it holds $|\hat{N}_W(R)| \geq q$

Theorem 1


Let $k = \lceil 2 \log n \rceil$. A.a.s. $G = (V, E) \sim \mathcal{G}(n, \frac{1}{2})$ satisfies the following:

- (\star) V is $(\frac{k}{50}, \Theta(n^{0.9}))$ -dense; and
- ($\star\star$) For every $(\frac{k}{10000}, \Theta(n^{0.9}))$ -dense $W \subseteq V$ there exists $S \subseteq V$, $|S| \leq \sqrt{n}$ s.t. for every $R \subseteq V$, with $|R| \leq \frac{k}{50}$ and $|\hat{N}_W(R)| < \tilde{\Theta}(n^{0.6})$ it holds that $|R \cap S| \geq \frac{k}{10000}$.


Theorem 2

Let $k = \lceil 2 \log n \rceil$. For every G satisfying properties (\star) and ($\star\star$), $S_{\text{reg}}(\Phi_{G,k}) = n^{\Omega(\log n)}$

Regular resolution \equiv Read-Once Branching Programs



Regular resolution \equiv Read-Once Branching Programs



Haken bottleneck counting idea

Haken bottleneck counting idea

“Lemma 1”

Every random path $\gamma \sim \mathcal{D}$ in the ROBP passes through a **bottleneck** node.

Haken bottleneck counting idea

“Lemma 1”

Every random path $\gamma \sim \mathcal{D}$ in the ROBP passes through a **bottleneck** node.

“Lemma 2”

Given any **bottleneck** node b in the ROBP,

$$\Pr_{\gamma \sim \mathcal{D}} [b \in \gamma] \leq n^{-\Theta(k)}.$$

Haken bottleneck counting idea

“Lemma 1”

Every random path $\gamma \sim \mathcal{D}$ in the ROBP passes through a **bottleneck** node.

“Lemma 2”

Given any **bottleneck** node b in the ROBP,

$$\Pr_{\gamma \sim \mathcal{D}} [b \in \gamma] \leq n^{-\Theta(k)}.$$

Then, it is trivial to conclude:

$$\begin{aligned} 1 &= \Pr_{\gamma \sim \mathcal{D}} [\exists b \in \text{ROBP } b \text{ bottleneck and } b \in \gamma] \\ &\leq |\text{ROBP}| \cdot \max_{\substack{b \text{ bottleneck} \\ \text{in the ROBP}}} \Pr_{\gamma \sim \mathcal{D}} [b \in \gamma] \\ &\leq |\text{ROBP}| \cdot n^{-\Theta(k)} \end{aligned}$$

The real bottleneck counting

$\beta(c) = \max$ (partial) assignment contained in all paths from the source to c

$\beta(c) = \max$ (partial) assignment contained in all paths from the source to c

$j \in [k]$ is **forgotten** at c if no sink reachable from c has label $\bigvee_{v \in V} x_{v,j}$

$\beta(c) = \max$ (partial) assignment contained in all paths from the source to c

$j \in [k]$ is **forgotten** at c if no sink reachable from c has label $\bigvee_{v \in V} x_{v,j}$

The random path γ

- if j forgotten at c or $\beta(c) \cup \{x_{v,j} = 1\}$ falsifies a short clause of $\Phi_{G,k}$ then continue with $x_{v,j} = 0$
- otherwise toss a coin and with prob. $\Theta(n^{-0.6})$ continue with $x_{v,j} = 1$

$$V_j^0(a) = \{v \in V : \beta(a)(x_{v,j}) = 0\}$$

$$V_j^0(a) = \{v \in V : \beta(a)(x_{v,j}) = 0\}$$

Lemma 1

For every random path γ , there exists two nodes a, b in the ROBP s.t.

$$V_j^0(a) = \{v \in V : \beta(a)(x_{v,j}) = 0\}$$

Lemma 1

For every random path γ , there exists two nodes a, b in the ROBP s.t.

1. γ touches a , sets $\leq \lceil \frac{k}{200} \rceil$ variables to 1 and then touches b ;

$$V_j^0(a) = \{v \in V : \beta(a)(x_{v,j}) = 0\}$$

Lemma 1

For every random path γ , there exists two nodes a, b in the ROBP s.t.

1. γ touches a , sets $\leq \lceil \frac{k}{200} \rceil$ variables to 1 and then touches b ;
2. there exists a $j^* \in [k]$ not-forgotten at b and such that $V_{j^*}^0(b) \setminus V_{j^*}^0(a)$ is $(\frac{k}{10000}, \Theta(n^{0.9}))$ -dense.

$$V_j^0(a) = \{v \in V : \beta(a)(x_{v,j}) = 0\}$$

Lemma 1

For every random path γ , there exists two nodes a, b in the ROBP s.t.

1. γ touches a , sets $\leq \lceil \frac{k}{200} \rceil$ variables to 1 and then touches b ;
2. there exists a $j^* \in [k]$ not-forgotten at b and such that $V_{j^*}^0(b) \setminus V_{j^*}^0(a)$ is $(\frac{k}{10000}, \Theta(n^{0.9}))$ -dense.

Lemma 2

For every pair of nodes (a, b) in the ROBP satisfying point (2) of Lemma 1,

$$\Pr_{\gamma}[\gamma \text{ touches } a, \text{ sets } \leq \lceil \frac{k}{200} \rceil \text{ vars to 1 and then touches } b] \leq n^{-\Theta(k)}$$

Proof sketch of Lemma 2

Let $E = \{\gamma \text{ touches } a, \text{ sets } \leq \lceil k/200 \rceil \text{ vars to } 1 \text{ and then touches } b\}$
and let $W = V_{j^*}^0(b) \setminus V_{j^*}^0(a)$

Proof sketch of Lemma 2

Let $E = \text{"}\gamma \text{ touches } a, \text{ sets } \leq \lceil k/200 \rceil \text{ vars to 1 and then touches } b\text{"}$
and let $W = V_{j^*}^0(b) \setminus V_{j^*}^0(a)$

Case 1: $V^1(a) = \{v \in V : \exists i \in [k] \beta(a)(x_{v,i}) = 1\}$ has large size ($\geq k/20000$). Then $\Pr[E] \leq n^{-\Theta(k)}$ because of the prob. of 1s in the random path γ and a Markov chain argument.

Proof sketch of Lemma 2

Let $E = \text{"}\gamma \text{ touches } a, \text{ sets } \leq \lceil k/200 \rceil \text{ vars to 1 and then touches } b\text{"}$
and let $W = V_{j^*}^0(b) \setminus V_{j^*}^0(a)$

Case 1: $V^1(a) = \{v \in V : \exists i \in [k] \beta(a)(x_{v,i}) = 1\}$ has large size ($\geq k/20000$). Then $\Pr[E] \leq n^{-\Theta(k)}$ because of the prob. of 1s in the random path γ and a Markov chain argument.

Case 2.1: $V^1(a)$ is not large but many ($\geq \tilde{\Theta}(n^{0.6})$) vertices in W are set to 0 by coin tosses.

So $\Pr[E \wedge W \text{ has many coin tosses}] \leq n^{-\Theta(k)}$ again by a Markov chain argument as in **Case 1**.

Proof sketch of Lemma 2

Let $E = \text{“}\gamma \text{ touches } a, \text{ sets } \leq \lceil k/200 \rceil \text{ vars to 1 and then touches } b\text{”}$
and let $W = V_{j^*}^0(b) \setminus V_{j^*}^0(a)$

Case 1: $V^1(a) = \{v \in V : \exists i \in [k] \beta(a)(x_{v,i}) = 1\}$ has large size ($\geq k/20000$). Then $\Pr[E] \leq n^{-\Theta(k)}$ because of the prob. of 1s in the random path γ and a Markov chain argument.

Case 2.1: $V^1(a)$ is not large but many ($\geq \tilde{\Theta}(n^{0.6})$) vertices in W are set to 0 by coin tosses.

So $\Pr[E \wedge W \text{ has many coin tosses}] \leq n^{-\Theta(k)}$ again by a Markov chain argument as in **Case 1**.

Case 2.2: $V^1(a)$ is not large and not many vertices in W are set to 0 by coin tosses. Then many of the 1s set by the random path γ between a and b must belong to a set of size at most \sqrt{n} , by the new combinatorial property ($\star\star$).

So $\Pr[E \wedge W \text{ has not many coin tosses}] \leq n^{-\Theta(k)}$.

Conclusions

Open Problem: How hard is to prove that a graph is Ramsey?

Let G be a Ramsey graph in n vertices and let $k = \lceil 2 \log n \rceil$. Is it true that $S(\Psi_{G,k}) = n^{\Omega(\log n)}$?

([LPRT17] proved this but for a binary encoding of “ G is Ramsey”)



full paper

Thanks!