

# Research summary

My work addresses the question of mass and angular momentum transfer in High Mass X-ray Binaries (HMXB), systems where a compact object orbits a high mass donor star. More specifically, I am interested in the behavior of the stellar material being accreted by the compact object on time scales within the scope of observing campaigns. The aim of my investigations has been twofold : to understand the accretion process onto wind-fed compact objects, since the star rarely fills its Roche lobe in SgXB, and to constrain the properties of the line-driven wind from the O/B supergiant donor star.

I use state-of-the-art magneto-hydrodynamics codes to follow the flow from the stellar surface down to the vicinity of the accretor and identify features susceptible to explain observations. I have been project leader for 4 successful computing time proposals on national supercomputers (115,000 CPUdays over the last 3 years) and I am co-PI of 2 XMM-Newton proposals to observe a classic and an obscured Supergiant X-ray binary (SgXB).

Constrain the

## Time-variability of the mass accretion rate in Supergiant X-ray binaries

- [1] *Hydrodynamical Bondi-Hoyle accretion onto a compact object*, El Mellah & Casse, MNRAS 2015
- [2] *Accretion from a clumpy massive-star wind in SgXB*, El Mellah, Sundqvist & Keppens, MNRAS 2017
- [3] *The clumpy absorber in the HMXB Vela X-1*, Grinberg, Hell, El Mellah et al., A&A 2017

Probe the clumpiness of the wind, using the orbiting X-ray source as a probe => connection to massive stars community. Clumpiness important since alters mass loss rates observed and mass loss rates important since (i) significantly alter stellar evolution and the mass of the stellar remnant left in the end and (ii) to know angular momentum evolution (spins and shrinking of the orbit).

Supergiant Fast X-ray transients

## Enhanced accretion and wind-captured discs in High Mass X-ray binaries

- [4] *A numerical investigation of wind accretion in persistent SgXB*, El Mellah & Casse, MNRAS 2016
- [5] *Formation of wind-captured discs in SgXB : the case of Vela X-1 and Cygnus X-1*  
El Mellah, Sanders, Sundqvist & Keppens, submitted
- [6] *Wind Roche lobe overflow in HMXB : a mass transfer mechanism for Ultraluminous X-ray sources*  
El Mellah, Sundqvist & Keppens, submitted

A better understanding of transfer of angular momentum in HMXBs is essential to predict the number of compact binaries with periods short enough to merge within a Hubble time.

## Code development & Kepler data analysis

- [7] *MPI-AMRVAC 2.0 for Solar and Astrophysical Applications*, Xia, Teunissen, El Mellah et al., ApJS 2018
- [8] *A study of the shortest-period planets found with Kepler*, Sanchis-Ojeda et al., ApJ 2014
- [9] *Triple-star candidates among the Kepler binaries*, Rappaport et al., ApJ 2013
- [10] *Possible disintegrating short-period super-Mercury orbiting KIC 12557548*  
Rappaport, Levine, Chiang, El Mellah et al., ApJ 2012

During my 3 years of PhD, I concentrated on mass transfer in binaries via wind accretion, the low angular momentum counterpart of the more comprehensively understood Roche lobe overflow (RLOF) mechanism. Supergiant X-ray binaries (SgXB), where a compact object (generally a NS) orbits an evolved O/B supergiant, are the ideal stage for wind accretion to occur. Indeed, the latter displays intense outflows, a fraction of which being captured by the NS. The rapid increase since the late 2000's in the number of SgXB (Walter et al. 2015) and the ambiguous status of the newly discovered SFXTs (Negueruela et al. 2006) only increased the appeal of this burning topic.

In a first attempt to better understand the wind accretion process, I confronted the analytical prescriptions given by Bondi, Hoyle and Lyttleton (BHL, Hoyle & Lyttleton 1939, Bondi & Hoyle 1944) to a hydrodynamical (HD) representation of the flow. To do so, I used and developed the explicitly flux-conserving finite volume transport code MPI-AMRVAC, a code whose origins trace back to the mid 90's when Gábor Tóth and Rony Keppens first tackled the question (Tóth 1996, Tóth et al. 1998). The new version I contributed to now addresses hydrodynamical or magneto-hydrodynamical problems, in Cartesian, cylindrical or spherical geometry, with or without polytropic prescriptions, source terms, etc (Xia et al. 2017). For wind speeds similar to the ones observed in SgXB ( $\sim 1,000 \text{ km} \cdot \text{s}^{-1}$ ), the main challenge is the contrast between the scale at which the gravitational beaming of the fast inflow by the accretor becomes significant (the accretion radius) and the size of the compact accretor, typically 4 to 5 orders of magnitude smaller. Since most of the emitted light comes from the immediate vicinity of the accretor, it is important to follow the flow through these scales. To uniformly resolve the incoming planar flow, I implemented a radially stretched grid in a 2D spherical geometry. With suitable boundary conditions, I reached a numerically relaxed state and spanned the 5 required orders of magnitude thanks to the computing time I was granted on the CINES Tier-1 cluster (see Figure 1). In El Mellah & Casse (2015), I characterized the structure of the flow, which forms a stable detached bow shocked as it is beamed towards the wake of the accretor, and the dependence of the mass accretion rate on the Mach number of the inflow. For the first time, we monitored the flow deep enough to also confirm the




Figure 1: Successive zoom in on the innermost parts of a planar flow (coming from the top) being deflected by a central accretor for different Mach numbers at infinity. In white are represented the streamlines while the dotted black lines represent the Mach=1 surfaces.

analytical prediction by Foglizzo & Ruffert (1997) concerning the topology of the inner sonic surface (where the shocked flow becomes supersonic again) which has to be anchored into the accretor.

In a realistic SgXB though, the incoming wind is not planar due to the orbital effects. It carries a non-zero angular momentum which could, in some cases, lead to the formation of a wind-capture disc. To identify the favorable configurations, I designed a model of supersonic line-driven wind propagation in SgXB, coupling the stellar, orbital, wind and accretion parameters (El Mellah & Casse 2016). I identified the minimal set of dimensionless degrees of freedom of the problem to optimally explore the space of parameters. This investigation showed how sheared and beamed the wind is when it enters the region around the accretor where the shock is expected to develop - i.e. where the ballistic assumption breaks up and where HD simulations similar to the ones above are required. The need to connect the orbital scale motion, essentially ballistic, and the accretion region, centered on the compact object, became apparent.

### Postdoctoral research activity

Since the beginning of my postdoctoral activity one year ago, I started to consider more realistic internal structure for the incoming wind in SgXB than the uniform flow I had worked with during my PhD. Indeed, the line-driven winds of massive stars are notoriously inhomogeneous, due to the line-deshadowing instability (Owocki & Rybicki 1984) which leads to the formation of internal shocks. The serendipitous accretion of these overdense regions, or clumps, has been suggested as a possible explanation to the time variability of the X-ray luminosity in SgXB, of the order of 100 peak-to-peak. Using a two-dimensional pseudo-planar grid sampling a restricted angular




Figure 2: *(left)* Principle of the clumpy wind accretion simulations : we inject into the simulation space (upper right insert) a wind whose micro-structure has been computed out of radiative HD simulations by Sundqvist et al. (2017). *(right)* Two-slices representation of the upstream hemisphere of the simulation space, with the wind coming from the upper left. We overlaid a logarithmic density map to show the typical size of the inhomogeneities to resolve. The accretor lies in the bottom right corner.

region, Sundqvist et al. (2017) recently managed to compute the micro-structure of the wind and by then, the dimensions of the clumps, for an isolated massive stars. To evaluate the impact of clumps on the accretion process, I plunged a compact object in the wind ("CO" in the left panel in Figure 2), at different orbital separations, and injected the corresponding wind computed by Sundqvist et al. (2017) within the simulation space (right panel in Figure 2). By coupling the stretching of the mesh to the Adaptive Mesh Refinement (AMR) of MPI-AMRVAC, I could design 3D spherical setups spanning several orders of magnitude at an affordable computational cost and resolve small scale off-centered features like clumps injected from the upstream hemisphere. In El Mellah et al. (2017), we were able to follow the clumps as they cross the shock and to monitor the time variability at the inner border of the simulation space, corresponding approximately to the dimensions of the NS magnetosphere. With this work, we discovered how tempering the shock could be, which led to variations of the inner mass accretion rate an order of magnitude smaller than the observed variations of the X-ray luminosity in these systems. Thus, if the stochastic variations at low X-ray luminosity seem to match the variability induced by the clumps alone, the high luminosity levels can only be reached due to other underlying mechanisms, possibly within the NS magnetosphere (eg the propeller effect, Bozzo et al. 2016). Concerning the column density levels, we retrieve average values compatible with what has been observed recently in Vela X-1 by Grinberg et al. (2017). However, in the latter paper, I evaluated the time variability associated to unaccreted clumps passing by the line-of-sight and concluded that this type of micro-structure within the wind can not explain by itself the variations in column density observed by Grinberg et al. (2017).


In Xia et al. (2017), I carried out a numerical validation of the stretched grid implementation I had made during my PhD by confronting quantitative simulation results of Bondi spherical accretion to the analytical expectations on the mass accretion rate and the location of the sonic point for different adiabatic indexes. We also studied the propagation of a trans-Alfvénic wind from the solar surface to the Earth orbit to validate the compatibility of the stretched grid with the magneto-HD solver and Powell's method for the cleaning of the divergence of the magnetic field.

## Project outline

In X-ray binaries, the challenge of scales can be alleviate thanks to the different dominant physical effects at stake at different scales. For instance, the magnetic field has generally little influence at the orbital scale but is decisive to fully appreciate the conditions in which the accreted flow emits most of the light we observe. Besides, in SgXB, the magnetic field dominates the last phase of accretion possibly through magnetic gating. It is also a key-ingredient of the ejection mechanisms responsible for self-collimated jets we often observe in accreting systems.

## RLOF discs

To guarantee that we work with physically consistent accretion discs, I am now performing numerical simula-



tions of RLOF configurations where the expanded atmosphere of the donor star is channeled into the Roche lobe of the accretor through the first Lagrangian point. This model includes a proxy on viscosity, similar to the one introduced by Shakura & Sunyaev (1973), and energy losses through radiative cooling. The former stands for the turbulent viscosity associated to the magnetic rotational instability (Balbus & Hawley 1991). Together with spiral shocks, they participate to the evacuation of angular momentum which makes the accretion possible. The ballistic solution is then superseded by the actual formation of a disc around the accretor. Since the plasma largely exceeds 10,000K during the accretion process, a significant fraction of the elements is ionized. We make use of the SPEX cooling tables for solar abundances to compute the cooling function (van Marle & Keppens 2011), which yields cooling rates large enough to impact the thickness of the flow. However, the numerically convenient optically thin assumption we currently make breaks up in the disc and must be complemented with a flux-limited diffusion method I plan to implement in MPI-AMRVAC. These improvements could first lead to insights concerning the origin of negative superhumps in cataclysmic variables (CV, Murray & Armitage 1998). Replacing the inner accretor with a BH or a lowly magnetized NS, the wrapping of the disc could also be studied and numerical results obtained in the context of Smooth Particle Hydrodynamics simulations could be confronted (Foulkes et al. 2006). If the disc turns out to be misaligned with the orbital plane, it could have a serious impact on jet-launching conditions since a misalignment with the BH spin is likely to induce a precession of the jets Liska et al. (2017).

## Diving into the magnetosphere

Once I obtain a satisfying self-generated RLOF disc, I will use it as a fruitful landscape to explore two kinds of systems. First, with Zakaria Meliani, we wish to make this setup a scaled up version of a NS accretor in a Low Mass X-ray binary. A magnetized white dwarf (WD) in a CV intermediate polar alleviates the contrast between the orbital separation and the size of the compact object, while still retaining most of the geometry of the system: a RLOF donor star feeding a disc truncated in its innermost regions by the magnetic field of the accretor (Ghosh et al. 1977). Until now, the studies which have been carried on make an adiabatic assumption to bypass the question of cooling (Ju et al. 2017). Working with a physically-motivated vertical profile of the disc, we think we can shed light on the question of the spin-down efficiency of the accretion process on the WD. We would also be in possession of a reliable tool to elucidate the disc reformation during the afterglow phase of novae (Ness et al. 2012). With U Scorpii expected to go off in a couple of years, we started to collaborate with Jan-Uwe Ness to provide the modeling support for observer proposals.

This twofold setup also serves another purpose : diving into the magnetosphere of an ac-

creting NS. Numerically, we recently implemented and validated a method of magnetic splitting generalized to non-potential fields (Xia et al. 2017). It enables us to handle more accurately the magnetic field evolution, in particular in low- $\beta$  shock-dominated plasmas, and to clean more easily the non-zero divergence. With this new feature available, we want to study how the innermost parts of the wind accreted flow derived in El Mellah et al. (2017) behave. Is the angular momentum carried by the clumps large enough to form a transient disc-like structure or does the NS magnetic field come into play before they can do so? The possible magnetic gating undergone by the innermost parts of the flow has never been explored with physical inputs accounting for the upper scales. Hence, it has produced fruitful results but ambiguous since the prescriptions employed have been independent from the large scale parameters (such as the stellar ones). We also want to address the case where the NS magnetospheric radius is much larger, of the order of the accretion radius. How does it alter the shock?

To conclude, the numerical expertise I have developed in Computational Astrophysics enables me to make the most of the high performance computing technologies available. They have ushered in a particularly exciting period to address a wide range of questions pertaining to accretion in X-ray binaries. To progress on these questions in the following 5 years a tenure track at the University of Amsterdam would offer me, I need to keep connecting with observers and modelers in this field. The research activity of the Anton Pannekoek Institute for Astronomy shows a shared interest in the key-questions of accretion-ejection processes and the neutron star equation-of-state. The former intersects the question of the disc structure we try to address, while the latter would take advantage of more constraining predictions on the properties of the accreted flow. To raise the funds necessary to such a project, I intend to apply to ambitious funding such as the ERC grants. Beyond their autonomous requirements and satisfactions, I also expect my teaching duties to be opportunities to arouse the interest of the students in pursuing in this research domain. Should it be the case, I would gladly mentor them, as I have been mentored on my way to the endlessly mesmerizing field of Astrophysics.

## References

- Balbus, S. A. & Hawley, J. F. (1991), *Astrophys. J.* **376**, 214.
- Bondi, H. & Hoyle, F. (1944), *Mon. Not. R. Astron. Soc.* **104**, 273.
- Bozzo, E., Oskinova, L., Feldmeier, A. & Falanga, M. (2016), *Astron. Astrophys.* **589**.
- El Mellah, I. & Casse, F. (2015), *Mon. Not. R. Astron. Soc.* **454**(3), 2657–2667.
- El Mellah, I. & Casse, F. (2016), *Mon. Not. R. Astron. Soc.* **467**(3), 2585–2593.
- El Mellah, I., Sundqvist, J. O. & Keppens, R. (2017), *MNRAS* .
- Foglizzo, & Ruffert, (1997), *Astron. Astrophys.* .
- Foulkes, S. B., Haswell, C. A. & Murray, J. R. (2006), *Mon. Not. R. Astron. Soc.* **366**(4), 1399–1409.
- Ghosh, P., Pethick, C. J. & Lamb, F. K. (1977), *Astrophys. J.* **217**, 578–596.
- Giacconi, R., Gursky, H., Paolini, F. R. & Rossi, B. B. (1962), *Phys. Rev. Lett.* **9**(11), 439–443.
- Grinberg, V., Hell, N., El Mellah, I., Neilsen, J., Sander, A. A. C., Leutenegger, M. A., Fürst, F., Huenemoerder, D. P., Kretschmar, P., Kühnel, M., Martínez-Núñez, S., Niu, S., Pottschmidt, K., Schulz, N. S., Wilms, J. & Nowak, M. A. (2017), *MNRAS* .
- Hoyle, F. & Lyttleton, R. A. (1939), *Math. Proc. Cambridge Philos. Soc.* **35**(03), 405–415.
- Ju, W., Stone, J. M. & Zhu, Z. (2017), *Astrophys. J.* **841**(1), 29.
- Liska, M., Hesp, C., Tchekhovskoy, A., Ingram, A., van der Klis, M. & Markoff, S. (2017), *eprint arXiv:1707.06619* 5(November), 1–5.
- Murray, J. R. & Armitage, P. J. (1998), *Mon. Not. R. Astron. Soc.* **300**(2), 561–566.
- Negueruela, I., Smith, D. M., Reig, P., Chaty, S. & Torrejón, J. M. (2006), *Proc. X-ray Universe 2005 (ESA SP-604). 26–30 Sept. 2005* .
- Ness, J. U. U., Schaefer, B. E., Dobrotka, A., Sadowski, A., Drake, J. J., Barnard, R., Talavera, A., Gonzalez-Riestra, R., Page, K. L., Hernanz, M., Sala, G. & Starrfield, S. (2012), *Astrophys. J.* **745**(1).
- Owocki, S. P. & Rybicki, G. B. (1984), *Astrophys. J.* **284**, 337.
- Shakura, N. I. & Sunyaev, R. a. (1973), *Astron. Astrophys.* **24**, 337–355.
- Sundqvist, J. O., Owocki, S. P. & Puls, J. (2017), (2).
- Tóth, G. (1996), *Magnetodynamic Phenom. Sol. Atmos. prototypes stellar Magn. Act. Proc. 153rd Colloq. Int. Astron. Union, held Makuhari, near Tokyo, May 22–27, 1995* **471**, 471.
- Tóth, G., Keppens, R. & Botchev, M. A. (1998), *Astron. Astrophys.* **332**, 1159–1170.
- van Marle, A. J. & Keppens, R. (2011), *Comput. Fluids* **42**(1), 44–53.
- Walter, R., Lutovinov, A. A., Bozzo, E. & Tsygankov, S. S. (2015), *Astron. Astrophys. Rev.* **23**(1), 2.
- Xia, C., Teunissen, J., El Mellah, I., Chané, E. & Keppens, R. (2017), *submitted* .

