

Computational Methods for Astrophysical Applications

Ileyk El Mellah & Jon Sundqvist



Centre for mathematical Plasma Astrophysics
Instituut voor Sterrenkunde
KU Leuven

Lesson 2 :

Finite Difference Approximation for linear hyperbolic PDE

- **Spatial discretization**

Mesh and data collocation

Discrete equations

- **Time advance**

Von Neumann stability analysis

Explicit linear schemes


- 1 Preliminary concepts
 - Mesh and data collocation
 - Consistency
 - Stability

- 2 Spatial discretization
 - Discretized function
 - Discretized first order derivatives
 - Discretized second order derivatives

- 3 Temporal discretization
 - Explicit methods
 - Implicit methods
 - Accuracy and positivity

Mesh

- Numerical schemes require spatial and temporal discretization



Data collocation

- Finite difference (FD): values are located at points, not necessarily regularly distributed. There is no cell.
 - ⇒ Intuitive & computationally fast but not conservative and no straightforward refinement strategy
- Finite volume (FV): values are associated to the whole cell : the value of the function is constant over the cell. Conservative by Gauss' law (a.k.a Green-Ostrogradsky).
 - ⇒ Reconstruction at interfaces required => slower

Consistency of a method

- A FD approximation of a PDE is *consistent* if and only if

$$\text{FD} \xrightarrow{\Delta t, \Delta x \rightarrow 0} \text{PDE}$$

\Rightarrow Balsara 2.4

- Empirical order of convergence (EOC) : u analytic solution and \tilde{u}_i numerical solution on 1D uniform grid x_i with $i \in [1, N_\nu = 2^\nu]$. Compute the maximal error $\epsilon_\nu = \|u - \tilde{u}\|_\infty$ and deduce EOC :

$$\text{EOC}_\nu = \frac{\log(\epsilon_\nu / \epsilon_{\nu-1})}{\log(N_{\nu-1} / N_\nu)}$$

\Rightarrow quantify the overall quality of a method, on analytic test-cases

\Rightarrow see [Project# 1](#) of Köln school

\Rightarrow see Balsara chapter 2, slides# 27-32

Von Neumann stability analysis

- Balsara 2.5 and intro of 2.7
- round-off errors should not grow during time progression
 - \Rightarrow numerical solution = exact + round-off error $\epsilon(x, t)$
 - \Rightarrow represent $\epsilon(x, t)$ in Fourier series, analyse Fourier term

$$\epsilon_i^n(k) = \hat{\epsilon}_k e^{\lambda t^n} e^{ikx_i}$$

- numerically stable scheme \forall spatial wavenumbers k if


$$\left| \frac{\epsilon_i^{n+1}(k)}{\epsilon_i^n(k)} \right| = |e^{\lambda \Delta t}| \leq 1$$

- Lax equivalence theorem :
consistent + stable \Leftrightarrow convergent

- 1 Preliminary concepts
 - Mesh and data collocation
 - Consistency
 - Stability
- 2 Spatial discretization
 - Discretized function
 - Discretized first order derivatives
 - Discretized second order derivatives
- 3 Temporal discretization
 - Explicit methods
 - Implicit methods
 - Accuracy and positivity

Discretized function

- Balsara 2.3
- We now want to obtain a FD discretized form of the equations to solve



⇒ variables evaluated at the points

Discretized spatial derivatives

- Assume $N + 1$ points regularly spaced by Δx and u_i the associated values
- Key-question: how to get a FD approximation of the spatial derivatives $u'_i, u''_i \dots$?
 - \Rightarrow Use Taylor series expansions

Backward, forward & central FD

- For instance, to 1st order accuracy in Δx , we have the *1st order forward difference* :

$$u'_i \sim \frac{u_{i+1} - u_i}{\Delta x}$$

\Rightarrow but we also have the *1st order backward difference* :

$$u'_i \sim \frac{u_i - u_{i-1}}{\Delta x}$$

\Rightarrow or the *2nd order central difference* :

$$u'_i \sim \frac{u_{i+1} - u_{i-1}}{2\Delta x}$$

- Which one to choose? Properties (e.g. stencil, accuracy)?

Truncation error and order of accuracy

- Taylor expansions linking u_i to the continuous function u :

$$u_{i+1} = u(0) + u'(0) \Delta x + u''(0) \frac{(\Delta x)^2}{2!} + u^{(3)}(0) \frac{(\Delta x)^3}{3!} + \dots \quad (1)$$

$$u_i = u(0) \quad (2)$$

$$u_{i-1} = u(0) - u'(0) \Delta x + u''(0) \frac{(\Delta x)^2}{2!} - u^{(3)}(0) \frac{(\Delta x)^3}{3!} + \dots \quad (3)$$

$$\Rightarrow (1)-(3) : u'(0) = \frac{u_{i+1} - u_{i-1}}{2\Delta x} + \underbrace{O((\Delta x)^2)}_{\text{truncation error}}$$

- Order of accuracy (o.a.) = order of first truncated term
 $\Rightarrow (\Delta x \rightarrow \Delta x/2) \Rightarrow$ accuracy improved by $2^{\text{o.a.}}$
- Stencil : points required, increases with o.a.

2nd order derivative

- Same principle

$$(1)+(3)-2\times(2) : u''(0) = \frac{u_{i+1}-2u_i+u_{i-1}}{(\Delta x)^2} + O((\Delta x)^2)$$

⇒ O.a.? Stencil?

- Exo

⇒ 3rd, 4th & 5th order central difference representations of u'_i ?

⇒ 2nd order forward/backward differences?

- 1 Preliminary concepts
 - Mesh and data collocation
 - Consistency
 - Stability
- 2 Spatial discretization
 - Discretized function
 - Discretized first order derivatives
 - Discretized second order derivatives
- 3 Temporal discretization
 - Explicit methods
 - Implicit methods
 - Accuracy and positivity

Principle

- Consider the 1D linear advection equation with constant speed v :

$$\partial_t \rho + v \partial_x \rho = 0 \quad (4)$$

- Explicit time integration: values at t^{n+1} computed from available information on time level $t^n = t^{n+1} - \Delta t$

Principle

- Example: explicit forward 2nd order central difference (a.k.a. Euler's Forward Time Central Space, FTCS)

$$\frac{u^{n+1} - u^n}{\Delta t} = -v \frac{u_{i+1}^n - u_{i-1}^n}{2\Delta x}$$

⇒ rearrange to

$$u_i^{n+1} = u_i^n - v \frac{\Delta t}{\Delta x} \frac{u_{i+1}^n - u_{i-1}^n}{2} \quad (5)$$

⇒ 1st order accurate in time, 2nd order in space

⇒ Is this FD approximation consistent?

- Sandbox: Write your 1st solver using the Euler's FTCS above (and periodic boundary conditions, ~ infinite simulation space). Apply it to the Riemann problem (advection of a step function). Is the numerical solution stable?

Von Neumann stability analysis

- Balsara 2.7.1
- Von Neumann stability analysis

$$e^{\lambda \Delta t} = 1 - \frac{v \Delta t}{\Delta x} \frac{e^{ik\Delta x} - e^{-ik\Delta x}}{2} = 1 - i \frac{v \Delta t}{\Delta x} \sin(k\Delta x)$$

\Rightarrow scheme is unconditionally unstable since for all k

$$\left| \frac{\epsilon_k^{n+1}}{\epsilon_k^n} \right| > 1$$

- 3 possible solutions to stabilize
 - \Rightarrow add numerical diffusivity to damp nonphysical instability
 - \Rightarrow impose same space-time symmetry as original PDE
 - \Rightarrow use implicit scheme (next section)

Numerical diffusivity : the Lax-Friedrichs scheme

- adding diffusion: advection-diffusion equation has form

$$\frac{\partial u}{\partial t} = -v \frac{\partial u}{\partial x} + \mathcal{D} \frac{\partial^2 u}{\partial x^2} \quad \text{with } \mathcal{D} \text{ a diffusion coefficient}$$

- Lax-Friedrichs scheme : in (5), if we replace u_i^n by spatial average between $i-1$ and $i+1$, we get :

$$u_i^{n+1} = \frac{u_{i+1}^n + u_{i-1}^n}{2} - v \frac{\Delta t}{\Delta x} \frac{u_{i+1}^n - u_{i-1}^n}{2}$$

which can be rearranged to form

$$\frac{u_i^{n+1} - u_i^n}{\Delta t} = -v \frac{u_{i+1}^n - u_{i-1}^n}{2\Delta x} + \frac{(\Delta x)^2}{2\Delta t} \frac{u_{i+1}^n - 2u_i^n + u_{i-1}^n}{(\Delta x)^2}$$

\Rightarrow numerical dissipation with $\mathcal{D} \equiv (\Delta x)^2 / 2\Delta t$

\Rightarrow 1st order accurate in time, 2nd order in space

CFL condition

- perform von Neumann stability analysis for Lax-Friedrichs

$$e^{\lambda \Delta t} = \cos(k \Delta x) - i \frac{v \Delta t}{\Delta x} \sin(k \Delta x)$$

⇒ conditional stability requiring Courant number C

$$C \equiv \frac{|v| \Delta t}{\Delta x} \leq 1$$

⇒ limitation of the time step Δt for a given resolution Δx

⇒ Courant-Friedrichs-Lewy condition (1928)

⇒ necessary condition for stability

- Rk: see stencil of Lax-Friedrichs scheme in Table on slide 26
- Exo : retrieve the figures on slide# 62 of Balsara 2.7.2
⇒ very diffusive

The Lax-Wendroff scheme

- Lax-Friedrichs has different o.a. in space (2nd) and time (1st)
 \Rightarrow 2nd order Taylor expansion in time of $u_i(t)$, yields :

$$u_i(t + \Delta t) = u_i(t) + u'_i(t)\Delta t + u''_i(t)\frac{(\Delta t)^2}{2} \quad (6)$$

$$\Rightarrow (4) \Rightarrow u'_i \sim -v\partial_x u$$

\Rightarrow to find u''_i , $\partial_t(4)$ and (illegally) switch ∂_t and ∂_x in mixed term :

$$u''_i \sim v^2\partial_{xx}u$$

\Rightarrow finally, reinject in (6) and use a central difference for spatial derivatives :

$$u_i^{n+1} = u_i^n - \frac{1}{2}\frac{\Delta t}{\Delta x} v (u_{i+1}^n - u_{i-1}^n) + \frac{1}{2}\frac{(\Delta t)^2}{(\Delta x)^2} v^2 (u_{i+1}^n - 2u_i^n + u_{i-1}^n)$$

The Lax-Wendroff scheme

- 2nd order in space and time
- Conditionally stable
 - \Rightarrow for $\Delta t = \Delta x/v$, we retrieve the Lax-Friedrichs scheme
- Exo : retrieve the figures on slide# 67 of Balsara 2.7.3 and compare the empirical order of convergence with respect to Lax-Friedrichs
 - \Rightarrow non positivity-preserving (\leq monotonicity-preserving)

Space-time symmetry

- How to maintain the space-time symmetry of the PDE in the FD approximation?
 - ⇒ use central discretisation for both time and space
 - ⇒ leapfrog scheme

$$u_i^{n+1} = u_i^{n-1} - v \frac{\Delta t}{\Delta x} \frac{u_{i+1}^n - u_{i-1}^n}{2}$$

- ⇒ conditionally stable and second-order accurate
 - ⇒ multiple time levels involved: $n-1$, n , $n+1$
 - ⇒ potential problem: even/odd time levels may 'decouple'
- stencil of leapfrog on slide 26

First order upwind scheme

- Balsara 2.7.5
- Physically-speaking, isn't it silly that, for $v > 0$ for instance, u_i^{n+1} depends on u_{i+1}^n ?
⇒ Information should propagate wave speeds (see previous lesson)

$$\frac{u_i^{n+1} - u_i^n}{\Delta t} = -v \frac{u_i^n - u_{i-1}^n}{\Delta x}$$

⇒ also called the donor cell scheme

First order upwind scheme

- Positivity preserving (" > 0 "), but 1st order accurate




Figure 3.2: (*upper panel*) A central differencing numerical solving of the uni-dimensional linear advection equation of a step function which went wrong. The dotted line stands for the analytical solution while the markers and solid line indicate the numerical answer. (*lower panel*) The same numerical problem solved with an upwind differencing scheme. From [Dullemond \(2009\)](#).

Overview of explicit methods








Name	Difference Equations	Stencil
Backward Euler	$U_j^{n+1} = U_j^n - \frac{k}{2k} A(U_{j+1}^n - U_{j-1}^n)$	
One-sided	$U_j^{n+1} = U_j^n - \frac{k}{k} A(U_j^n - U_{j-1}^n)$	
One-sided	$U_j^{n+1} = U_j^n - \frac{k}{k} A(U_{j+1}^n - U_j^n)$	
Lax-Friedrichs	$U_j^{n+1} = \frac{1}{2}(U_{j-1}^n + U_{j+1}^n) - \frac{k}{2k} A(U_{j+1}^n - U_{j-1}^n)$	
Leapfrog	$U_j^{n+1} = U_j^{n-1} - \frac{k}{2k} A(U_{j+1}^n - U_{j-1}^n)$	
Lax-Wendroff	$U_j^{n+1} = U_j^n - \frac{k}{2k} A(U_{j+1}^n - U_{j-1}^n) + \frac{k^2}{2k^2} A^2(U_{j+1}^n - 2U_j^n + U_{j-1}^n)$	
Beam-Warming	$U_j^{n+1} = U_j^n - \frac{k}{2k} A(3U_j^n - 4U_{j-1}^n + U_{j-2}^n) + \frac{k^2}{2k^2} A^2(U_j^n - 2U_{j-1}^n + U_{j-2}^n)$	

TABLE 10.1
Finite difference methods for the linear problem $u_t + Au_x = 0$.

$h=dx$
 $k=dt$

1st order schemes safe, can be > 0 BUT very diffusive
 2nd order schemes accurate \Rightarrow great for smooth profiles (e.g. Gaussian) but terrible for discontinuous ones (e.g. top hat) because non > 0

Principle

- third cure to instability of Euler FTCS scheme: evaluate spatial derivative at t^{n+1}

$$u_i^{n+1} = u_i^n - v \frac{\Delta t}{\Delta x} \frac{u_{i+1}^{n+1} - u_{i-1}^{n+1}}{2}$$

⇒ Backwards in Time, Central in Space Euler scheme (BTCS)

⇒ u_i^{n+1} not expressed in terms of values at time t^n : implicit

- Stencil?

Von Neumann stability analysis

- von Neumann stability analysis for BTCS scheme

$$|e^{\lambda \Delta t}| = \frac{1}{|1 + i \frac{v \Delta t}{\Delta x} \sin(k \Delta x)|} < 1 \quad \text{for all } k$$

\Rightarrow unconditionally stable, any time step Δt allowed

- But stability \neq accuracy
 \Rightarrow the larger Δt , the lower the accuracy
- BTCS is 1st order in time
- Implicit methods more used for elliptic PDE \Rightarrow not covered by the current course

Godunov's theorem

- Explicit schemes

Scheme o.a.	Can be stable? > 0 ?
FTCS 1 st	noyes
Lax-Friedrichs 1 st	yesyes
Lax-Wendroff 2 nd	yesno
Upwind 1 st	yesno

"All linear positivity-preserving schemes for linear advection are condemned to be 1st order accurate"

- We want :

- \Rightarrow accuracy where the variables are smooth (i.e. \sim Gaussian)
- \Rightarrow positivity where they are discontinuous (i.e. \sim top hat)