$$P(N) = \begin{cases} \frac{\theta^{-1}}{x^{\theta}} & N \ge 1 \\ 0 & K \le 1 \end{cases}$$

$$L(X, \theta) = \frac{(\theta^{-1})^n}{\int_{-1}^{\infty} x^{\theta}} L_n L_{2n} h(\theta^{-1}) - \theta \frac{2}{x^{\theta}} h_{x},$$

$$\left( h_{1} L_{2}^{-\frac{1}{\theta}} - \frac{2}{x^{\theta}} h_{2}^{-\frac{1}{\theta}} \right)$$

Del une:
$$(\int_{K^{-}}^{\Phi-1} dx)_{0}^{1} = x^{1-0}h_{1} x \Rightarrow mageh caumo recuplingum.$$

$$\int_{K^{-}}^{\Phi-1} dx)_{0}^{1} = x^{1-0}h_{1} x \Rightarrow mageh caumo recuplingum.$$

$$\int_{K^{-}}^{\Phi-1} dx = -\frac{1}{x^{2}} + 1 = \frac{1}{2} \hat{x} = 0 \quad \frac{1}{6}$$

$$\int_{K^{-}}^{\Phi-1} dx = -\frac{1}{x^{2}} + 1 = \frac{1}{2} \hat{x} = 0 \quad \frac{1}{6}$$

$$\int_{K^{-}}^{\Phi-1} dx = -\frac{1}{x^{2}} + 1 = \frac{1}{2} \hat{x} = 0 \quad \frac{1}{6}$$

$$\int_{K^{-}}^{\Phi-1} dx = -\frac{1}{x^{2}} + 1 = \frac{1}{2} \hat{x} = 0 \quad \frac{1}{6}$$

$$\int_{K^{-}}^{\Phi-1} dx = -\frac{1}{x^{2}} + 1 = \frac{1}{2} \hat{x} = 0 \quad \frac{1}{6} \hat{x} = 0$$

$$\int_{K^{-}}^{\Phi-1} dx = -\frac{1}{x^{2}} + 1 = \frac{1}{2} \hat{x} = 0 \quad \frac{1}{6} \hat{x} = 0$$

$$\int_{K^{-}}^{\Phi-1} dx = -\frac{1}{x^{2}} \hat{x} = 0 \quad \frac{1}{6} \hat{x} = 0 \quad \frac{1}{6} \hat{x} = 0$$

$$\int_{K^{-}}^{\Phi-1} dx = -\frac{1}{2} \hat{x} = 0 \quad \frac{1}{6} \hat{x} = 0 \quad$$