

Comparing tree diversity between two forest plots, BCI and Paracou

Ilke Geladi^{1*}

Abstract

BCI is one of the best-studied forest plots on earth, however we must consider the value of other similar forest plots such as Paracou in French Guiana. Here, we compare the diversity between the two through diversity profiles and rarefaction of the data. It was found that Paracou is more diverse than BCI. An increased diversity in a forest plot holds potential for novel and unique studies that BCI does not offer.

Keywords

Diversity, BCI, Paracou, Diversity Profiles, Hill numbers

¹UMR EcoFoG, AgroParistech, CNRS, Cirad, INRA, Université de Guyane.
Campus Agronomique, 97310 Kourou, France.

*Contact: ilke.geladi@ecofog.gf, <http://www.ecofog.gf/spip.php?article47>

Contents

1	Introduction	1
2	Materials and Methods	1
2.1	Study Sites	1
	BCI • Paracou	
2.2	Diversity Comparison	2
3	Results	2
4	Discussion	2

1. Introduction

Tropical rainforests are the most biodiverse ecosystems in the world, housing two to three thirds of the species in the world (Laurance, 2007; Noss, 1999; Bierregaard *et al.*, 1992). Reliable information on the status and condition in each forest studied, including the diversity of trees, is crucial to expanding our knowledge in fields such as forest dynamics, community ecology, and biotic interactions to ensure the best possible management and conservation strategies (Condit, 1995; Noss, 1999). However, to collect such knowledge, long-term studies are necessary as they provide a more wholistic view through continuous monitoring and data on patterns and rates of change in forest ecosystems (Comiskey & Dallmeier, 1998). Permanent forest plots are widely used for these sorts of studies (Comiskey & Dallmeier, 1998; Condit, 1995), and many have been established in various ecosystems worldwide (Condit *et al.*, 2014).

In the tropics, arguably the most famous long-term forest plot is Barro Colorado Island (BCI), a 1500-ha island in Panama (Leigh, 1999). Scientists have been conducting studies at BCI since 1916 and it has since become one of the best studied areas of tropical rainforest in the world (Leigh, 1999). Despite this, BCI is not a very diverse or pristine habitat (Leigh, 1999), and there are many advantages to conducting studies at other less-famous forest plots, such as Paracou in French Guiana (Gourlet-Fleury *et al.*, 2004). Offering more than 100 hectares of plots, Paracou

was established in 1982 and has also been home to many studies on forest functioning and forestry research programs (Charles-Dominique, 2001; Gourlet-Fleury *et al.*, 2004).

In this study we will compare the neutral forest diversity, focusing solely on trees, of BCI and Paracou. With Paracou being connected to mainland South America thus having higher colonization rates compared to BCI which is an island (MacArthur, 1965), we hypothesize that Paracou is more diverse than BCI. Habitats with increased diversity, such as Paracou, might increase potential research questions and provide different opportunities to further increase our knowledge of tropical rainforests compared to BCI.

2. Materials and Methods

2.1 Study Sites

2.1.1 BCI

For the first study site, we used tree diversity data from Barro Colorado Island (BCI) (Croat, 1978), which can be found in the “vegan” package (Oksanen *et al.*, 2010). BCI became an island with the damming of the Chagres River in Panama between 1911 and 1914 and was subsequently named a biological reserve in 1923, which is currently supervised by the Smithsonian Tropical Research Institute (STRI) (Croat, 1978; Leigh, 1999). It is a 1,560-hectare island that welcomes hundreds scientists from around the world each year (Croat, 1978).

2.1.2 Paracou

For the second study site, we used tree diversity data from Paracou (Degen *et al.*, 2006), which can be found in the “entropart” package (Marcon & Hérault, 2015).

Paracou is located in French Guiana, and has been used since 1982 for basic ecological research on lowland rainforest (Gourlet-Fleury *et al.*, 2004). The site is more than 1000 hectares large and trees have been fully mapped as Paracou was used as an experimental design to study the impact of disturbances on forest stands and tree populations (Gourlet-Fleury *et al.*, 2004).




Figure 1. Diversity Profiles for Paracou (top line) and BCI (bottom line). Confidence intervals are obtained by bootstrap from 1000 simulations.

2.2 Diversity Comparison

To compare the two study sites, we first compared their asymptotic diversity profiles. To do this we used the “CommunityProfile” function in the “entropart” package (Marcon & Hérault, 2015) to create the diversity profile for both sites followed by the “CEnvelope” function (Marcon & Hérault, 2015) to add the BCI diversity profile to that of Paracou. Hill Numbers were then calculated when q is equal to 0, 1 and 2 using the “Diversity” function (Marcon & Hérault, 2015).

Next, we rarefied the data to the same coverage as an alternative method to compare the two communities whilst accounting for the difference in study site size. Through this method, we simulate the decrease in the size of the large community, in this case Paracou, to reduce it to the size of the small community, BCI (Marcon, 2017). In the process, we lose some data, however it is better to lose some data than have a lot of uncertainty, which is the case when estimating the asymptotic diversity. To do so, we used the function “estimatedD” of the package iNEXT (Hsieh *et al.*, 2016), and compared the data for values of diversity of $q=0$, 1 and 2, rarefied to the same coverage of 80%. All above analyses and calculations were performed in R (R Core Team, 2017).

3. Results

Asymptotic diversity profiles reveal Paracou to be more diverse than BCI (See 1). As can be seen in Figure 1, the Paracou diversity profile curve is above the BCI curve for all values of q . This is confirmed by calculating the Hill numbers for both BCI and Paracou. It was found that Paracou was more diverse at all order’s of q where for when q equals 0, 1 and 2, the corresponding Hill numbers for Paracou were 315, 133 and 73, and for BCI were 239, 72 and 38.

BCI and Paracou data rarefied to the same coverage (80%) also reveals Paracou to have a greater diversity at order of diversities 0, 1 and 2 compared to BCI. Species richness ($q=0$) of Paracou ($S=136$) is almost double that of BCI ($S=69$). The exponential of Shannon’s entropy index

($q=1$) of Paracou is 89, whereas that of BCI is only 47. Finally, when $q=2$ (the inverse of Simpson’s concentration index), diversity of Paracou is once again almost double that of BCI where Paracou’s is at 61 and BCI’s at 31.

4. Discussion

Diversity Profiles show that tree diversity is more neutrally diverse in Paracou than in BCI, as the Paracou curve is above the BCI curve. It can further be said that BCI and Paracou are separable as the communities diversity profiles do not intersect (Marcon, 2017). There has been much debate as to which diversity indices to use as there exist many. However, diversity profiles play an important role in converging these indices by portraying them graphically as the value of diversity is plotted in function to the order q , which represent different indices (Tothmeresz, 1995; Marcon, 2017). The raw result of the Hill numbers calculated support the above conclusion that Paracou is more diverse than BCI. Hill numbers provide many advantages over other frequent used measures of diversity, such as species richness, through incorporating both relative abundance and species richness, amongst other reasons (Chao *et al.*, 2014). Overall, diversity profiles and Hill numbers, which are two more reliable methods of calculating diversity, reveal that Paracou has a higher tree diversity than BCI.

Rarefaction results support the conclusions from the diversity profiles, taking it a step further and revealing the tree diversity in Paracou is almost double that of BCI (when $q=0$ and $q=2$). Rarefaction of the data is important as it standardizes the data on the basis of sample size or sample completeness thus facilitating and providing a more accurate view of the comparison between both forest plots (Chao *et al.*, 2014). Having both diversity profiles and rarefied data support that Paracou is more diverse than BCI, we can be certain of this conclusion.

A logical explanation as to why Paracou is more diverse than BCI is related to island biogeography theory where it predicts that with increased colonization, there is more diversity (MacArthur, 1965). Seeing as Paracou is on continental land, it is very connected to other communities where colonization is facilitated compared to BCI which is an island. Furthermore, island populations have a much higher risk of extinction than mainland populations (Frankham, 1997). However, it must also be recognized that often islands have higher speciation rates and thus more endemic species, being valuable in a different way than mainland ecosystems, which is not taken into account by these calculations (Kier *et al.*, 2009).

Studying diversity is very important in ecological theory and practice (Tothmeresz, 1995). Diversity indices are often used as indicators of the ‘well-being’ of an ecosystem and are thus implicated in conservation management and environmental monitoring (Tothmeresz, 1995; Chao *et al.*, 2014).

In conclusion, Paracou is more diverse than BCI. Studying diversity is important as it informs us about the state of the ecosystem, and has implications in management strategies. Eventhough BCI has the advantage of being very well-studied and potentially having more endemic species than

Paracou, Paracou has a higher diversity than BCI which can provide very interesting research opportunities compared to BCI.

References

- Bierregaard, R.O., Lovejoy, T.E., Kapos, V., dos Santos, A.A. & Hutchings, R.W. (1992) The biological dynamics of tropical rainforest fragments : a prospective comparison of fragments and continuous forest. *BioScience*, **42**, 859–866.
- Chao, A., Gotelli, N.J., Hsieh, T.C., Sander, E.L., Ma, K.H., Colwell, R.K. & Ellison, A.M. (2014) Rarefaction and extrapolation with Hill numbers: A framework for sampling and estimation in species diversity studies. *Ecological Monographs*, **84**, 45–67.
- Charles-Dominique, P. (2001) The Field Station BT. F. Bongers, P. Charles-Dominique, P.M. Forget & M. Théry, eds., *Nouragues: Dynamics and Plant-Animal Interactions in a Neotropical Rainforest*, pp. 1–8. Springer Netherlands, Dordrecht.
- Comiskey, J.A. & Dallmeier, F. (1998) *Forest Biodiversity in North, Central and South America, and the Caribbean: Research and Monitoring*. Taylor & Francis.
- Condit, R. (1995) Research in large, long-term tropical forest plots. *Trends in Ecology & Evolution*, **10**, 18–22.
- Condit, R., Lao, S., Singh, A., Esufali, S. & Dolins, S. (2014) Data and database standards for permanent forest plots in a global network. *Forest Ecology and Management*, **316**, 21–31.
- Croat, T.B. (1978) *Flora of Barro Colorado Island*. Stanford University Press.
- Degen, B., Blanc, L., Caron, H., Maggia, L., Kremer, A. & Gourlet-Fleury, S. (2006) Impact of selective logging on genetic composition and demographic structure of four tropical tree species. *Biological Conservation*, **131**, 386–401.
- Frankham, R. (1997) Do island populations have less genetic variation than mainland populations? *Heredity*, **78**, 311–327.
- Gourlet-Fleury, S., Guehl, J.M. & Laroussinie, O. (2004) *Ecology & management of a neotropical rainforest. Lessons drawn from Paracou, a long-term experimental research site in French Guiana*. Elsevier, Paris.
- Hsieh, T.C., Ma, K.H. & Chao, A. (2016) iNEXT: an R package for rarefaction and extrapolation of species diversity (Hill numbers). *Methods in Ecology and Evolution*, **7**, 1451–1456.
- Kier, G., Kreft, H., Lee, T.M., Jetz, W., Ibsch, P.L., Nowicki, C., Mutke, J. & Barthlott, W. (2009) A global assessment of endemism and species richness across island and mainland regions. *Proceedings of the National Academy of Sciences*, **106**, 9322–9327.
- Laurance, W.F. (2007) Have we overstated the tropical biodiversity crisis? *Trends in Ecology & Evolution*, **22**, 65–70.
- Leigh, E.G. (1999) *Tropical Forest Ecology: A View from Barro Colorado Island*. Oxford University Press.
- MacArthur, R.H. (1965) Patterns of Species Diversity. *Biological Reviews*, **40**, 510–533.
- Marcon, E. (2017) *Mesures de la Biodiversité*. UMR EcoFoG, Kourou.
- Marcon, E. & Hérault, B. (2015) entropart, an R Package to Measure and Partition Diversity. *Journal of Statistical Software*, **67**, 1–26.
- Noss, R.F. (1999) Assessing and monitoring forest biodiversity: A suggested framework and indicators. *Forest Ecology and Management*, **115**, 135–146.
- Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H. & Wagner, H. (2010) vegan: Community Ecology Package.
- R Core Team (2017) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Tothmeresz, B. (1995) Comparison of different methods for diversity ordering. *Journal of Vegetation Science*, **6**, 283–290.