

Femobiome SecundaScreen

Patient name: ...
Date of birth: ...
Sampling date: ...
Sample type:
Sample ID:
Physician: ...
Medical record: ...
Referring organization: ...

Control parameters	Ig GE/mL	Reference
Human genomic DNA	5,2	≥ 3,5
Total bacterial load	6,8	≥ 3,5

Test result interpretation



- Relative amount of *Lactobacillus* spp. within the reference range: 85-100%.
- Detected: CMV.

Microbiota status

	Ig GE/mL	%	Reference	
NORMAL MICROBIOTA, total proportion		85-100%	≥ 80%	
<i>Lactobacillus</i> spp. ¹	6,8	85-100%		
AEROBES, total proportion		8-10%		
<i>Staphylococcus</i> spp.	3,1	< 1%		
<i>Streptococcus</i> spp.	5,7	7-9%		
<i>Enterobacteriaceae</i>	—	—		
<i>Enterococcus</i> spp.	—	—		
<i>Haemophilus</i> spp.	4,6	< 1%		
ANAEROBES, total proportion		< 1%		
<i>Gardnerella vaginalis</i>	4,7	< 1%		
<i>Fannyhessea (Atopobium) vaginae</i>	3,1	< 1%		
<i>Mobiluncus</i> spp.	—	—		
MYCOPLASMAS, total proportion		< 1%		
<i>Ureaplasma urealyticum</i>	—	—	< 4,0 lg	
<i>Ureaplasma parvum</i>	< 4,0	< 1%	< 4,0 lg	
<i>Mycoplasma hominis</i>	—	—	< 4,0 lg	

¹ <i>Lactobacillus</i> spp.	Ig GE/mL	Species proportion, %
<i>L.non-iners</i> *	6,8	100%
<i>L.iners</i>	—	—

*Species representatives:

L.crispatus, *L.jensenii*, *L.mulieris*, *L.gasseri*, *L.paragasseri*

Yeast fungi	Ig GE/mL	Reference
<i>Candida</i> spp.	< 4,0	< 4,0
<i>Candida albicans</i>	—	< 4,0

Date of report 2025-08-12

Performed by

Laboratory director

Full name

Signature

Full name

Signature

Pathogens

	Result	Ig GE/mL	Reference
OBLIGATE PATHOGENS (STI)			
<i>Chlamydia trachomatis</i>	—	—	—
<i>Mycoplasma genitalium</i>	—	—	—
<i>Neisseria gonorrhoeae</i>	—	—	—
<i>Trichomonas vaginalis</i>	—	—	—
HERPESVIRUSES			
HSV-1	—	—	—
HSV-2	—	—	—
CMV	DETECTED	< 3,0	—

Terminology and symbols

Aerobes — facultative-anaerobic opportunists. Anaerobes — obligate anaerobic opportunists. Mycoplasmas — opportunistic genital mycoplasmas.

Dash (—) = negative result; "I" = combined detection.

Femobiome SecundaScreen

Patient name: ...
Date of birth: ...
Sampling date: ...
Sample type:
Sample ID:
Physician: ...
Medical record: ...
Referring organization: ...

Control parameters	Ig GE/mL	Reference
Human genomic DNA	4,8	≥ 3,5
Total bacterial load	6,5	≥ 3,5

Test result interpretation



- Relative amount of *Lactobacillus* spp. within the reference range: 85-100%, predominance of *L.iners*.
- No pathogens detected.

Microbiota status

	Ig GE/mL	%	Reference	
NORMAL MICROBIOTA, total proportion		85-100%	≥ 80%	
<i>Lactobacillus</i> spp. ¹	6,5	85-100%		
AEROBES, total proportion		0%		
<i>Staphylococcus</i> spp.	—	—		
<i>Streptococcus</i> spp.	—	—		
<i>Enterobacteriaceae</i>	—	—		
<i>Enterococcus</i> spp.	—	—		
<i>Haemophilus</i> spp.	—	—		
ANAEROBES, total proportion		0%		
<i>Gardnerella vaginalis</i>	—	—		
<i>Fannyhessea (Atopobium) vaginae</i>	—	—		
<i>Mobiluncus</i> spp.	—	—		
MYCOPLASMAS, total proportion		0%		
<i>Ureaplasma urealyticum</i>	—	—	< 4,0 lg	
<i>Ureaplasma parvum</i>	—	—	< 4,0 lg	
<i>Mycoplasma hominis</i>	—	—	< 4,0 lg	

¹ <i>Lactobacillus</i> spp.	Ig GE/mL	Species proportion, %
<i>L.non-iners</i> *	—	—
<i>L.iners</i>	6,5	100%

*Species representatives:
L.crispatus, *L.jensenii*, *L.mulieris*, *L.gasseri*, *L.paragasseri*

Yeast fungi	Ig GE/mL	Reference
<i>Candida</i> spp.	< 4,0	< 4,0
<i>Candida albicans</i>	—	< 4,0

Date of report 2025-08-12

Performed by

Laboratory director

Full name

Signature

Full name

Signature

Pathogens

	Result	Ig GE/mL	Reference
OBLIGATE PATHOGENS (STI)			
<i>Chlamydia trachomatis</i>	—	—	—
<i>Mycoplasma genitalium</i>	—	—	—
<i>Neisseria gonorrhoeae</i>	—	—	—
<i>Trichomonas vaginalis</i>	—	—	—
HERPESVIRUSES			
HSV-1	—	—	—
HSV-2	—	—	—
CMV	—	—	—

Terminology and symbols

Aerobes — facultative-anaerobic opportunists. Anaerobes — obligate anaerobic opportunists. Mycoplasmas — opportunistic genital mycoplasmas.

Dash (—) = negative result; "I" = combined detection.