

Enterobiome Kids Research result

FULL NAME:

SEX:

DATE OF BIRTH:

DATE OF SAMPLING:

CONTAINER ID:

PHYSICIAN:

STATE OF INTESTINAL MICROBIOTA

Proportion of normal microbiota is within the normal range – **99.9%**. Total amount of bifidobacteria is within the normal range – **8.5 GE/g**. Taxonomic diversity of normal microbiota is within the normal range - **6**. The diversity of metabolically active “child” bifidobacteria is within the normal range – **4**. The proportion of metabolically active “child” bifidobacteria is within the normal range** – **100.0%**. The proportion of metabolically active bifidobacteria is within the normal range and amounts at **> 10%**. *Lactobacillaceae* agents are absent. Proportion of opportunistic microbiota is within normal limits – **0.1%**. The order of Enterobacterales is represented by *E.coli*. Yeast fungi are not detected.

IMPORTANT: the interpretation of the results must be made strictly by the attending physician

NAME OF RESEARCH	RESULT	REFERENCE INTERVAL	UNIT MEASUREMENTS
Total bacterial mass (TBM)	8.9*	> 6.0	Lg (GE/g)
Normal microbiota			
proportion	99.9	≥ 80	%
diversity, number of taxa	6	≥ 4	PCS
Bifidobacterium spp.			
total amount	8.5*	> 5.3	Lg (GE/g)
metabolically active species, proportion	> 10	≥ 10	%
metabolically active “child” species, proportion**	100.0	> 5	%
metabolically active “child” species, diversity	4	≥ 2	PCS
Lactobacillaceae, quantity	-	> 0.0	Lg (GE/g)
Bacteroidetes, presence		Not assessed at this age	
Firmicutes/Bacteroidetes, ratio:		Not assessed at this age	
Yeast fungi, quantity	0.0*	< 4.8	Lg (GE/g)
Opportunistic pathogens, proportion	0.1	< 20	%
Markers of pathogenicity and resistance, presence			
pathogenic agents <i>Enterobacterales</i>	not detected	not detected	
<i>Clostridioides difficile</i>		Not assessed at this age	
<i>tcdA, tcdB</i>		Not assessed at this age	
<i>Staphylococcus aureus</i>	3.8*	< 5.8	Lg (GE/g)
<i>mecA</i>	not detected	not detected	
<i>Streptococcus agalactiae</i> with <i>srr2</i>		Not assessed at this age	
<i>Candida albicans</i>	not detected	not detected	

Assay date: 25.09.2023

The study was carried out by _____ (Full name)

* Lg X values are given – means 10^x

** the value of the proportion of the number of metabolically active bifidobacteria is indicated

FULL NAME:

CONTAINER ID:

DATE OF SAMPLING:

NAME OF RESEARCH	ABSOLUTE, Lg (GE/g feces)*		RELATIVE, %	
	Value*	Reference interval	Value	Reference interval
Total bacterial mass	8.9	> 6.0		
NORMAL MICROBIOTA	8.6	7.0 - 10.2	99.9	80 - 100
Actinobacteria				
Bifidobacterium spp	8.5	5.3 - 10.1	70.9	25.7 - 98.3
Metabolically active bifidobacteria species, proportion			> 10	≥ 10
Metabolically active "infant" bifidobacteria **	7.6	5.0 - 8.6	100.0	> 5
Bifidobacterium longum subsp. infantis	4.7		0.1	
Bifidobacterium longum subsp. longum	7.6		99.2	
Bifidobacterium bifidum	5.4		0.6	
Bifidobacterium breve	3.9		< 0.1	
Metabolically active "adult" bifidobacteria **	-	0.0 - 6.6	-	0 - 40
Bifidobacterium adolescentis	-		-	
Bifidobacterium catenulatum ssp	-		-	
Bifidobacterium animalis subsp. lactis	-		-	
Bifidobacterium dentium	-		-	
Coriobacteriia	5.4	0.0 - 8.8	0.1	0.0 - 5.0
Firmicutes				
Clostridium leptum gr	-	0.0 - 8.4	-	0.0 - 5.7
Dialister+Allisonella+Megasphaera+Veillonella	6.4	0.0 - 8.7	0.6	0.0 - 10.0
Faecalibacterium prausnitzii	-	0.0 - 5.7	-	0.0 - 0.1
Lachnospiraceae	4.8	0.0 - 9.6	< 0.1	0.0 - 47.1
Lactobacillaceae	-	0.0 - 7.8	-	0.0 - 5.0
Streptococcus spp	5.8	5.1 - 8.6	0.1	0.0 - 20.0
Lactococcus lactis	-	0.0 - 6.0	-	0.0 - 1.0
Bacteroidetes				
Alistipes spp	-	0.0 - 5.8	-	0.0 - 0.1
Bacteroides spp	8.1	0.0 - 9.4	28.2	0.0 - 26.2
Butyricimonas spp	-	0.0 - 4.2	-	0.0 - 0.1
Parabacteroides spp	-	0.0 - 8.4	-	0.0 - 4.7
Prevotella spp	-	0.0 - 6.5	-	0.0 - 1.0
Other bacteria				
Akkermansia muciniphila	-	0.0 - 8.5	-	0.0 - 8.0
Desulfovibrio spp	-	0.0 - 4.5	-	0.0 - 0.1
Methanobrevibacter spp	-	0.0	-	0.0 - 1.0
Diversity, number of taxa:	6	≥ 4		
Firmicutes/Bacteroidetes ratio:	-	≥ 1.5		
OPPORTUNISTIC PATHOGENS	5.8	6.0 - 9.3	0.1	0 - 20
Clostridium difficile gr	-	0.0 - 7.8	-	0.0 - 5.0
Clostridium perfringens gr	-	0.0 - 8.1	-	0.0 - 6.0
Enterobacterales	5.8	6.9 - 9.2	0.1	0.0 - 20.0
E.coli	5.8	4.1 - 8.1	0.1	0.0 - 10.0
Enterococcus spp	-	5.6 - 8.5	-	0.0 - 11.0
Erysipelotrichaceae	-	0.0 - 8.5	-	0.0 - 6.9
Fusobacteriaceae	-	0.0 - 5.9	-	0.0 - 0.1
Peptoniphilaceae	-	0.0 - 6.7	-	0.0 - 1.0
Pseudomonas spp	-	0.0 - 5.0	-	0.0 - 0.1
Staphylococcus spp	4.1	0.0 - 6.3	< 0.1	0.0 - 1.0
MARKERS OF PATHOGENICITY AND RESISTANCE				
Clostridioides difficile	-	0.0 - 6.0	-	0.0 - 0.1
tcdA tcdB	-			
Staphylococcus aureus	3.8	0.0 - 5.8	< 0.1	0.0 - 0.1
mecA	-			
Streptococcus agalactiae	-	0.0 - 5.0	-	0.0 - 0.1
srr2	-			
YEAST FUNGI				
Candida spp	-	0.0 - 4.8		
C.albicans	-	0.0		

* Lg X values are given – means 10^x

** the value of the proportion of the number of metabolically active bifidobacteria is indicated