

Female microbiota assessment

Name:

Age:

Date of collection:

Locus of biomaterial collection: V

Container ID:

Doctor:

Control biomaterial parameters


	Ig GE/ml	Reference
Human genomic DNA	3,8	≥ 3,5
Total Bacterial Load	7,3	≥ 4,0

Conclusion



■ Microbiota state – **severe anaerobic dysbiosis**: normobiota is decreased to 6%, relative amount of *Lactobacillus* spp. 4% (*L.iners*, *L.crispatus*, *L.jensenii*/*L.mulieris*, *L.gasseri*/*L.paragasseri*), *Bifidobacterium* spp. 2%, relative amount of anaerobes is increased

■ **Detected**: HPV 16, HPV 31-68 (see the note)



Microbiota state

	Ig GE/ml	%	Reference	
NORMOBIOTA, total proportion		5-7%	≥80%	
<i>Lactobacillus</i> spp. ¹	5,9	3-5%		
<i>Bifidobacterium</i> spp.	5,6	1-3%		
AEROBES, total proportion		< 1%		
<i>Staphylococcus</i> spp.	—	—		
<i>Streptococcus</i> spp. ²	3,3	<1%		
<i>Enterobacteriaceae</i>	—	—		
<i>Enterococcus</i> spp.	—	—		
<i>Haemophilus</i> spp.	3,0	<1%		
ANAEROBES, total proportion		80-100%		
<i>Gardnerella vaginalis</i>	6,9	32-44%		
<i>Fannyhessea (Atopobium) vaginae</i>	4,2	< 1%		
<i>Mobiluncus</i> spp.	—	—		
<i>Anaerococcus</i> spp.	—	—		
<i>Peptostreptococcus</i> spp.	4,3	< 1%		
<i>Bacteroides</i> spp. / <i>Porphyromonas</i> spp. / <i>Prevotella</i> spp.	6,1	5-7%		
<i>Sneathia</i> spp. / <i>Leptotrichia</i> spp. / <i>Fusobacterium</i> spp.	6,8	25-35%		
<i>Megasphaera</i> spp. / <i>Veillonella</i> spp. / <i>Dialister</i> spp.	5,5	1-3%		
BVAB1 / BVAB2 / BVAB3	6,6	16-22%		
MYCOPLASMAS, total proportion		< 1%		
<i>Ureaplasma urealyticum</i>	—	—	< 4,0 Ig	
<i>Ureaplasma parvum</i>	< 4,0	< 1%	< 4,0 Ig	
<i>Mycoplasma hominis</i>	—	—	< 4,0 Ig	

¹ <i>Lactobacillus</i> spp.	Ig GE/ml	Proportion of species in <i>Lactobacillus</i> spp.,%
<i>L.crispatus</i>	4,4	2%
<i>L.gasseri</i> / <i>L. paragasseri</i>	3,4	<1%
<i>L.jensenii</i> / <i>L. mulieris</i>	4,2	1%
<i>L.iners</i>	6,1	97%

Yeast fungi	Ig GE/ml	Reference
<i>Candida</i> spp.	—	< 4,0
<i>Candida albicans</i>	—	< 4,0
² Group B streptococcus		
<i>Streptococcus agalactiae</i>	—	—

Female microbiota assessment

Name: _____

Container ID: _____

PATHOGENS

	Result	Ig GE/ml	Reference
SEXUALLY TRANSMITTED INFECTIONS (STI)			
<i>Chlamydia trachomatis</i>	—	—	—
<i>Mycoplasma genitalium</i>	—	—	—
<i>Neisseria gonorrhoeae</i>	—	—	—
<i>Trichomonas vaginalis</i>	—	—	—
HERPESVIRUSES			
HSV-1	—	—	—
HSV-2	—	—	—
CMV	—	—	—
HUMAN PAPILLOMAVIRUSES (HPV)			
HPV 16	DETECTED	< 3,0	—
HPV 18	—	—	—
HPV 45	—	—	—
HPV 31/33/35/39/51/52/56/58/59/66/68	DETECTED	5,9	—

NOTE:

Automated conclusion is only formed if the locus of biomaterial collection (V or C) is specified. Result interpretation is carried out by the physician.

Date of report 2025-08-13

Assay performed by

Name

Signature

Laboratory Manager

Name

Signature

Note

Reference for *Streptococcus agalactiae*:

Gestational age over 37-38 weeks – negative result, all other cases – <4,0 Ig GE/ml.

Types of Infographics in the test report

- Color indicator – overall assessment of the microbiota state (eubiosis / moderate or severe dysbiosis);
- Linear histogram – display of the proportions of normobiota, aerobes, anaerobes, and mycoplasmas in the microbiota;
- Column histogram – individual profile of microorganisms.

Terminology and Designations

Aerobes – facultative anaerobic opportunistic microorganisms. Anaerobes – obligate anaerobic opportunistic microorganisms. Mycoplasmas – opportunistic genital mycoplasmas.

HPV 31-68 (11 types) – total detection of HPV 31/33/35/39/51/52/56/58/59/66/68 without differentiation.

A dash (—) means a negative result; a listing separated by a "/" indicates pooled detection.