

An open multimodal benchmark for LLM molecular structure elucidation reveals a recall-bound bottleneck that forward verification exploits

Ilkham Yabbarov¹ (*corresponding: ilkhamfy@gmail.com*), Rodrigo A. Vargas-Hernández^{1 1}

Department of Chemistry and Chemical Biology, McMaster University, Hamilton, Ontario, Canada

Manuscript draft prepared for Digital Discovery (RSC).

Abstract

Large language models (LLMs) have been reported to predict NMR spectra and propose molecular structures from them, but the supporting evaluation is small, curated, NMR-only, hinted, and drawn from a single model family — leaving the real-world question open: given an unfiltered experimental spectrum from the literature, how often does a frontier LLM recover the correct structure? We address this with three contributions. First, **IRexp**, to our knowledge the largest *permissively-licensed, redistributable* dataset of *experimental* infrared spectra — 121,233 records mined from open-access full text, of which **42,842 are linked to a resolved 2D structure** and **40,491 are full IR + ¹H + ¹³C + structure quadruples** — assembled by a browser-free literature-mining agent and released under permissive licences (larger view-only databases exist but are not bulk-redistributable; §2.1). Second, **IRSpectra-Bench**, an open, blind, mechanically scored benchmark of **194 compounds** (134 spectrally-validated plus a 60-compound controlled set) for spectrum→structure elucidation, stratified across molecular complexity. We find that a frontier LLM (Claude Opus), given molecular formula + IR + ¹H + ¹³C, recovers the exact constitution of **28.4%** of real compounds (95% CI 22–35; 33.5% within three candidates), with a sharp, tight gradient — 48% on simple molecules versus 8% on complex ones, and 60% for small (≤15 heavy atoms) versus 7% for large (>25). This sits far below the ~100% implied by curated evaluations, a gap we attribute largely to compound difficulty, candidate ranking, and hints; a within-compound control shows that **solving each compound in an independent context with tool access roughly triples accuracy over a single fatigued pass (5%→15%)**, i.e. methodology, not raw capability, dominates reported numbers. Third, we use a training-free **forward-verification** probe (we fine-tune nothing; the LLM is pretrained) to *localise* this gap rather than close it: candidates from the (hard) inverse direction are re-ranked by forward-predicting each one’s ¹³C spectrum and matching it to the observed one. The decomposition is our central diagnostic finding — **in our measurements the binding constraint is candidate recall, not verification**: when the true structure is among the candidates, forward verification selects it 84% of the time versus 73% for the model’s own ranking — a real but bounded gain — while the true structure is generated for only 31% of compounds. Acting on it (generating six regiochemistry-aware candidates and forward-verifying) lifts measured exact top-1 from 23% to **30%** (recall 41%), a genuine no-training improvement that nonetheless stays *below its own ceiling* (recall plateaus; verification precision falls 84%→72% as near-degenerate regioisomers accumulate), and the match distance does **not** double as a calibrated confidence signal (selective prediction is null). The honest summary is a *diagnosis plus a bounded improvement*, not a solution. We confirm the same recall-bound regime in a **battery-electrolyte domain** (46 scored compounds, **26% top-1**, the same broad regime, overlapping the headline CI) along a chemically interpretable gradient (sp³-C-F easiest at 50%; sulfonyl and nitrile hardest at 12%), and cast forward-verification as a training-free *analog* —

not an equivalent — of the computational-NMR / NMR-crystallography validation chemists trust, sharing its forward-prediction principle but forgoing DFT’s calibrated error model. All experiments run with no model training and no paid API, using LLM agents under a standard subscription, and all data, predictions, and code are released as an open resource.

1. Introduction

Determining a molecule’s structure from its spectra is a central, time-consuming task in synthetic and analytical chemistry. The dominant machine-learning approach trains specialised encoder–decoder models to map spectra to structures; the recent *Spectro* model, for example, learns $^1\text{H}/^{13}\text{C}/\text{IR} \rightarrow \text{SELFIES}$ from a corpus of 6,833 molecules.¹ In parallel, general-purpose LLMs have been reported to perform the same task off-the-shelf: a 2026 industrial white paper (a non-peer-reviewed company report) found that Claude Opus matched or beat commercial NMR-prediction software in the forward direction (structure→spectrum, ± 0.08 ppm ^1H) and “recovered all eight simpler structures on every attempt” in the inverse direction (spectrum→structure).² We treat its numbers as a motivating claim to be tested against peer-reviewed benchmarks, not as an established baseline.

These results are striking, but the evaluation that supports them is narrow: 15 inverse problems on curated single-ring or two-fragment molecules, NMR only, with the seven harder targets additionally given the *starting-material structure* as a hint, and “recovery” scored leniently over three runs and three ranked candidates. The question a practising chemist actually faces — *take an arbitrary experimental spectrum from a paper and recover the structure* — is left open. Answering it requires (i) a large, diverse, real benchmark; (ii) a blind, reproducible scoring protocol; and (iii) honest accounting for the methodological choices that inflate or deflate the apparent score.

We provide all three. We build IRexp (§2), a literature-mined dataset that supplies the IR modality absent from prior LLM evaluations alongside paired NMR and resolved structures. We define a blind benchmark on it (§3) and measure inverse-elucidation performance under matched and mismatched methodologies (§4), isolating a large solver-methodology effect with a within-compound control. We then introduce forward-verification elucidation (§5), a training-free method that turns the model’s *strong* direction (forward prediction) against its *weak* one (inverse regiochemistry), and use it to localise the bottleneck. **Figure 1** summarises the end-to-end design. Throughout, the solver and verifier are LLM agents run under a consumer subscription — no fine-tuning, no API spend — making every experiment cheaply reproducible. This is, by construction, a data-science and open-resource contribution in the remit of *Digital Discovery*: an openly licensed, structure-linked spectral dataset; a pre-registered, mechanically scored benchmark with released ground truth and scorer; and an end-to-end reproducible, training-free, zero-paid-API methodology in which every figure regenerates from the released code and predictions. We prioritise an honest, reusable measurement of where current models stand over a leaderboard-topping number.

Contributions. 1. **IRexp** — the largest openly *redistributable* experimental-IR dataset (121,233 records; 42,842 structure-linked; 40,491 IR+ ^1H + ^{13}C +structure): larger by raw record count than any prior collection and uniquely released under permissive licences for bulk model development (view-only resources such as SDBS are larger in structure-linked spectra but are not redistributable; §2.1), with a reproducible mining pipeline. 2. An **open multimodal benchmark** and the first blind, mechanically scored, complexity-stratified evaluation of frontier-LLM structure elucidation on real data — measured in depth on one model family (Claude) — reconciling the gap to optimistic

prior reports. 3. A **diagnostic decomposition** of LLM elucidation into recall and verification, showing that **recall — not verification — bounds current performance**, obtained via a training-free forward-verification probe that yields a bounded improvement (top-1 23%→30%) but, by our own measurements, cannot exceed a recall/precision ceiling without sharper verification or 2D-NMR data.

1.1 Related work

Trained spectra→structure models. The dominant line trains sequence or graph decoders to emit structures from spectra: *Spectro* ($^1\text{H}/^{13}\text{C}/\text{IR}\rightarrow\text{SELFIES}$)¹, the multitask CNN+transformer of routine 1D-NMR¹¹, and set/graph transformers such as NMRTrans¹². These reach high accuracy *in-distribution* but require a labelled spectra→structure corpus, which is exactly the scarce resource our IReXP pipeline targets, and they are retrained per modality. Closest in spirit to our multimodal setting is **NMIRacle**¹³, a generative model conditioned jointly on $\text{IR} + ^1\text{H} + ^{13}\text{C}$; it is a strong trained baseline, whereas our contribution is the *open experimental-IR data it (and others) can train on*, plus a *training-free* protocol and a blind benchmark to measure it.

LLMs as elucidators, and how this work differs. General-purpose LLMs have been applied off-the-shelf — Anthropic’s forward/inverse white paper² (a non-peer-reviewed company report), SpectraLLM and MolSpectLLM (multimodal LLMs over multi-spectral input)¹⁴, and knowledge-enhanced tree-search reasoning¹⁵ — and dedicated multimodal *benchmarks* now exist, most prominently **MolPuzzle**²⁰ ($\text{IR}+\text{MS}+^1\text{H}+^{13}\text{C}$ elucidation puzzles with the molecular formula supplied). These establish that the task is worth benchmarking and that LLMs can be scored on multi-spectral input; we claim priority on neither. What is new here is three things that, to our knowledge, no prior benchmark combines. (i) **Real, literature-mined experimental spectra at scale:** prior multimodal benchmarks and trained baselines^{1 13} rely on simulated, software-predicted, or hand-curated puzzle spectra, whereas IReXP-Bench is drawn from IReXP’s experimental $\text{IR} + ^1\text{H} + ^{13}\text{C}$ of out-of-distribution compounds, so it measures the simulated-to-real gap rather than in-distribution skill. (ii) **Blind, fully specified, mechanical scoring:** reported MolPuzzle exact-match accuracy for one model (GPT-4o) ranges from 1.4%²⁰ to 27.8%¹⁵ depending only on the scoring harness — a $\sim 20\times$ swing — which is itself the argument for the single, pre-registered, RDKit-InChIKey protocol with bootstrap CIs that we adopt and release. (iii) **A recall/verification decomposition:** existing benchmarks report a single aggregate score, whereas we factor performance into generation recall and verification precision and show *where* the task is lost — the result that transfers across models, domains, and verifier choices.

Computational NMR for structure validation. Our forward-verification method is the LLM analog of a workflow chemists already trust: assign a structure by computing the spectrum each candidate *would* give and matching it to experiment. In solution, this is the DP4 / DP4+ probabilistic framework over GIAO-DFT shifts^{16 17}; in the solid state it is **NMR crystallography**, where GIPAW-computed shifts adjudicate between candidate structures^{18 19}. We replace the quantum-chemical predictor with a forward LLM, trading accuracy for zero setup cost, and inherit the same core principle — *verification by forward prediction is easier than inverse generation*.

2. The IRExp dataset

2.1 Motivation

Open experimental IR is scarce *in a redistributable, ML-ready form*. The largest freely *downloadable* collections are the NIST WebBook⁸ (~16k IR spectra) and the Chemotion electronic-lab-notebook⁷ deposit (~2k). The AIST SDBS²² is larger (~54k FT-IR, all structure-linked) but is *view-only*: it caps downloads at 50 spectra/day, forbids commercial use, and ships no permissive licence or bulk export, so it cannot be used to train or redistribute open models at scale. Commercial libraries (e.g. Wiley KnowItAll, ~10⁵ spectra) are larger still but closed. IR — which directly encodes functional groups complementary to NMR — has thus had no large, *permissively-licensed*, structure-linked, bulk-reusable resource. IRExp fills this gap, and is the largest experimental-IR collection by raw record count.

2.2 Construction

A per-compound IR band list, together with co-reported ¹H/¹³C NMR, follows a remarkably stable textual convention in the experimental sections of organic chemistry papers. We exploit this with a browser-free harvesting agent:

- **Discovery.** Open-access primary literature is enumerated through the NCBI E-utilities and harvested in bulk from the PMC Open-Access Subset⁶ on AWS S3 (plain HTTPS, no anti-bot, fully redistributable CC-BY content), supplemented by the Chemotion FT-IR deposit (RADAR4Chem, CC-BY-SA-4.0).
- **Extraction.** A deterministic parser segments experimental text into per-compound records and extracts IR wavenumbers and ¹H/¹³C shift lists, with quality gates that reject instrument scan-range artefacts and prose false-positives (band-list density, ≥ 4 bands, plausible 400–4000 cm⁻¹ window).
- **Structure resolution.** In-text IUPAC names are converted to SMILES with OPSIN³, canonicalised with RDKit⁴ (InChIKey, SELFIES⁵), with a PubChem fallback for trivial/natural-product names. A key engineering finding was that the dominant open-access main-text convention labels compounds with *letter-prefixed* series labels (e.g. “...carbothioamide (**B1**):”) rather than the digit-first “(3a)” of supporting-information sections; capturing these and cleaning narrative/PDF artefacts before resolution raised structure coverage from 24% to **35%**.

The pipeline is browser-free by design. This is not an anti-bot bypass — the bulk corpus is fully open — but a throughput choice: plain HTTP from a bulk corpus parses at hundreds of papers per second, which is what makes a 10⁵-scale harvest feasible where a per-page browser driver (as used to build prior 10³-scale sets) is not.

2.3 Contents and licensing

Table 1. IRExp dataset contents and provenance.

field	value
experimental IR records	121,233
...co-reporting ¹ H and/or ¹³ C NMR	87,075 (72%)
...with a resolved 2D structure	42,842 (35%)
full IR + ¹ H + ¹³ C + structure quadruples	40,491

A structure-complete split, **irexp_resolved** (42,842 records, 100% structure-linked), is the training-/benchmark-ready subset and is $\sim 6\times$ the 6,833-molecule set used to train Spectro¹ (Fig. 2). Provenance is 119,345 PMC-OA records (CC-BY) plus 1,888 Chemotion records (CC-BY-SA); the two licences are kept as separable pools. Each record is DOI-/accession-traceable. Re-resolution is additive and content-keyed, so the dataset can be re-enriched without re-mining.

3. Benchmark design (IRSpectra-Bench)

From **irexp_resolved** we draw **IRSpectra-Bench**, 194 blind elucidation problems. Each problem presents the **molecular formula** (as from HRMS), the **IR band list**, and the **¹H and ¹³C shift lists** (with multiplicities and J-couplings where reported), and asks for the structure. No name, SMILES, or hint is given. Every main-round ground-truth structure is **spectrally validated** by an automated RDKit consistency check (¹³C peak count vs symmetry-unique carbons, molecular-formula match, SELFIES round-trip), excluding records with merged or incomplete spectra (6/140 in the main round, leaving 134). The 60 compounds of the controlled rounds are retained as fixed, pre-registered sets for the difficulty and within-compound controls, with the same self-consistency audit reported separately rather than used to exclude (57/60 pass; §7). Problems are stratified by RDKit ring analysis: a compound is **simple** iff it has at most two rings (single ring or two separate ring fragments), no fused/spiro/bridgehead system, and ≤ 22 heavy atoms; every other compound — any fused/spiro/bridged or ≥ 3 -ring system, **or** more than 22 heavy atoms — is **complex** (98 simple / 96 complex). The criterion is therefore exhaustive (the 23–24-atom band, 13 compounds, is classed complex by size). InChIKey de-duplication is applied across all rounds to prevent leakage.

Scoring is mechanical. A prediction is *correct* if its RDKit InChIKey connectivity layer (first 14 characters) matches the reference — i.e. we score *constitution* (atom-and-bond connectivity); stereochemistry is reported separately. This is a deliberate floor: a candidate with the correct constitution but wrong stereochemistry counts as correct. The exposure is bounded — only 10.3% (20/194) of benchmark answers carry a *defined* (assigned R/S) stereocentre (29.9% any potential) — and 1D ¹H/¹³C/IR rarely fixes absolute configuration, so a stereochemistry-aware score is neither well-posed from the given data nor materially different on this set.

Metrics. We report five quantities. **Top-1 (exact constitution)** is the fraction of compounds whose single best-ranked candidate matches the reference at the InChIKey connectivity layer. **Recovered (top-3)** is the fraction for which the reference appears among the up-to-three ranked candidates returned (matching the lenient “recovery” protocol of ref. 2). **Generation recall** is the fraction for which the reference is present in the candidate pool *before* re-ranking — the ceiling any verifier can reach. **Conditional-on-recall precision** is the verifier’s hit rate over recall-positive compounds only, isolating verification quality from generation. The forward verifier ranks candidates by a symmetric **chamfer distance** between predicted and observed ¹³C peak sets (for each predicted peak, the distance to its nearest observed peak, summed, and symmetrically for each observed peak; lower is better, with no equal-count requirement). We also report Morgan(2, 2048)⁹ Tanimoto as a graded “right scaffold/family” signal. CIs are bootstrap 95% over compounds; model-vs-model differences use McNemar’s exact test with Holm correction.

Solvers are LLM agents run under a consumer subscription. A frontier LLM (Claude Opus) is invoked as an independent sub-agent per batch of problems; agents are closed-book — configured with no web/tool access beyond an RDKit formula check and no access to ground truth, verified by grep-auditing their task transcripts at run time (audit logs available on request; the

committed artefacts are the parsed per-compound predictions). Adherence to the supplied formula is high but imperfect: 91.3% (115/126) of forward-verification candidates match the given molecular formula exactly (100% of true structures, 89.6% of decoys), and 76.6% across the full top-3 pool — a residual generator error orthogonal to regiochemistry. This makes the benchmark free to run and reproducible without API credits.

4. How well do LLMs elucidate real structures?

4.1 Headline performance

Decoupled per-compound agents solve each problem in an independent context (formula + IR + ^1H + ^{13}C , blind, up to three ranked candidates). Over the full **194-compound benchmark** (134 spectrally-validated compounds + the 60 from the controlled rounds), with bootstrap 95% confidence intervals:

Table 2. Headline elucidation performance on IRSpectra-Bench (n=194), overall and by difficulty stratum, with bootstrap 95% CIs.

metric	overall (n=194)	simple (n=98)	complex (n=96)
top-1 exact	28.4% [22–35]	48.0% [39–57]	8.3% [3–15]
constitution			
recovered (within	33.5% [27–40]	54.1% [44–63]	12.5% [6–20]
top-3)			
scaffold-level (best	56%	73%	39%
Tanimoto ≥ 0.45)			
mean best Tanimoto	0.59	0.73	0.45

As a cohort-robustness check, restricting the controlled rounds to their spectrally-clean subsets as well (134 main-clean + 57 controlled-clean = 191 strictly-validated compounds) leaves the headline unchanged: top-1 **28.8%** (95% CI 23–36), recall 34.0%, with simple 49.0% / complex 8.4% — within 1.7 points of the n=194 figures and with fully overlapping intervals, so the asymmetric inclusion of the pre-registered controlled sets does not drive the result.

The gradient is sharp and the intervals are tight. The 48%→8% simple→complex separation (Fig. 3) confirms the benchmark is discriminating across a realistic difficulty range, and accuracy falls monotonically with molecular size in step with it — top-1 **60.5%** for ≤ 15 heavy atoms, 28.3% for 16–25, and **7.0%** above 25 (Fig. 4). The model reliably recovers molecular formula, functional groups, and scaffold, but the exact constitution far less often, failing predominantly on **regiochemistry** — *which* position a substituent occupies. This is consistent with an information limit of 1D data: many regioisomers have similar $^1\text{H}/^{13}\text{C}$ shifts, which is precisely why 2D experiments (HMBC, NOESY) exist.

4.2 Reconciling with prior reports

Our 28% top-1 sits far below the ~100% on “simple” molecules reported, in a non-peer-reviewed company white paper, for the same model class.² Because that figure has not been independently

scored, we reconcile it against the *peer-reviewed* record (the MolPuzzle benchmark²⁰ and its scorings¹⁵, and the trained baselines¹⁻¹³) rather than treating it as settled. The gap is largely attributable to methodology and scoring choices, not raw capability:

- **Difficulty.** “Single ring” by ring-count is not “easy”: our simple stratum includes, e.g., a hexasubstituted benzene whose regiochemistry has many realisations. Ring count is a poor proxy for elucidation difficulty.
- **Scoring.** Prior work counts a recovery if the reference appears among three ranked candidates over three independent runs; we report single-run top-1 and top-3.
- **Hints.** Prior hard targets received the starting-material SMILES, which fixes most of the scaffold; we give none.
- **Curation.** Prior compounds were hand-selected; ours are scraped and unfiltered for solvability.

On a like-for-like easy/hinted/lenient setup the numbers rise; on the realistic, hint-free, scraped regime, ~28% is the honest figure.

Versus trained models — a bound, not a leaderboard. A like-for-like comparison against specialised trained models is not available: no system has been scored on an identical test set, and published numbers differ in the three respects that most move the score — spectrum realism (simulated/curated vs. real), hints, and how “exact match” is defined. The strongest trained baselines report their accuracy *in-distribution on simulated spectra*. Spectro (¹H/¹³C/IR→SELFIES, 6,833 training molecules) reaches ~90% top-1 exact recovery — but on a 1,366-molecule held-out split whose IR is plotted from reference data and whose NMR is software-*predicted*, not experimental¹; and NMIracle, which like us conditions jointly on IR+¹H+¹³C with *no* hints, reports 48% top-1 / 66% top-15 exact-SMILES recovery — again on held-out molecules from a *simulated* corpus drawn from the training distribution¹³. We measure 28.4% top-1 (33.5% top-3; 30% with forward verification) on **blind, real, literature-mined experimental** spectra of out-of-distribution compounds. These are not comparable as a leaderboard; read as a *bound on the simulated-to-real gap*, the contrast suggests that high in-distribution accuracies substantially overstate real-world performance — the same gap we document for the LLM above. The instability of the metric itself reinforces the caution: as noted in §1.1, the *same* model (GPT-4o) on MolPuzzle scores anywhere from 1.4%²⁰ to 27.8%¹⁵ depending only on the harness — a ~20× swing, in the peer-reviewed literature, that bounds how much weight any single unaudited near-100% claim (ref 2) can bear.

4.3 Methodology dominates: a within-compound control

The same 20 molecules were solved two ways: (a) by a single LLM context handling all of them sequentially with no tools, and (b) by independent per-compound agents with RDKit formula-checking. On the *identical* compounds, recovery rose from **5% to 15%** (1/20 → 3/20) and top-1 from 0% to 15% (0/20 → 3/20) — a 3× methodology effect with zero sample confound, though on a small set (the 15% point estimate carries a Wilson 95% CI of roughly 5–36%), so it is read as a directional within-compound demonstration rather than a precise multiplier. Small rounds also swing widely (15–40% across n=20–40 draws), which is why the headline is the full **194-compound** figure (28.4% top-1, 95% CI 22–35) rather than any single round. The practical lesson — fresh per-compound context plus tool access roughly triples apparent performance — also explains part of the gap to optimistic prior reports, whose per-problem API calls implicitly used method (b).

4.4 Model comparison: the benchmark ranks capability (and separates the extremes)

A benchmark is only useful if it separates models. On a fixed 24-compound subset solved blind by four Claude models — spanning a wide capability range, including the newest (Table 5) — under the identical protocol (Fig. 5):

Table 3. Four-model comparison on a fixed 24-compound subset under the identical blind protocol.

model	top-1	recovered	top-1 95% CI
Claude Fable 5	45%	54%	[25, 67]
Claude Opus	25%	29%	[8, 42]
Claude Sonnet	20%	25%	[8, 38]
Claude Haiku	0%	4%	[0, 0]

Three signals matter here. First, the four models rank in **monotonic capability order**, and the outcomes are **strictly nested** (Haiku Sonnet Opus Fable — each stronger model solves a superset of the compounds the weaker one does), with the smallest **floored at 0% exact** (4% recovered) on the same problems — so the benchmark is **capability-sensitive** and not a coin-flip. The nesting makes the *ranking* robust, but at $n=24$ the subset is **underpowered to separate adjacent models**: by McNemar’s exact test only the Fable-vs-Haiku gap survives multiple-comparison correction (Holm-adjusted $p=0.006$), Fable-vs-Opus is not significant ($p=0.063$), and Opus vs Sonnet are statistically indistinguishable — the bootstrap CIs above overlap accordingly. Second, two mid-tier frontier models agree closely (Opus 25%, Sonnet 20%), so the recall-bound $\sim 25\%$ regime of §4.1 is **not an artefact of a single model**. Third, the newest model nearly **doubles** the next-best top-1 (45% vs 25% on identical compounds; a large but, at this n , not-yet-significant gap) yet still misses the majority — IRSpectra-Bench is **hard and far from saturated even for the strongest model available**, with clear headroom. We ran four Claude-family models because they are callable for free under one subscription; a true cross-vendor sweep (GPT-, Gemini-class, open models) needs API access we deliberately did without, but the monotonic, large capability spread makes it unlikely the recall-bound pattern is specific to one model lineage. This is the behaviour a benchmark needs to remain informative as models improve.

4.5 Domain case study: battery-electrolyte chemistry

Structure elucidation from spectra is the daily inverse problem of the electrolyte-and-interphase community: the soluble decomposition products of carbonate and sulfone electrolytes, the additives that build the solid-electrolyte interphase, and the fluorophosphate/fluorosulfonyl species that NMR is routinely used to assign are exactly the molecules whose constitution must be read back out of IR and multinuclear NMR. To test whether the recall-bound regime above holds for *this* chemistry — rather than for organic molecules at large — we curated **IRSpectra-Bench-Electrolyte**, a 48-compound subset (eight per class as curated; two later yielded no parseable candidate, leaving **46 scored**) of structure-resolved IRExp records selected by substructure for the six functional families that dominate lithium- and sodium-battery electrolytes and their breakdown products: **carbonate** (linear/cyclic, the EC/DMC backbone), **sulfonyl/sulfonate** (sulfones, sulfonamides, the $-\text{SO}_2\text{CF}_3$ motif of imide salts), **nitrile** (acetonitrile- and adiponitrile-type high-voltage additives), **$\text{sp}^3\text{-C-F}$** (fluorinated solvents and additives), **phosphoryl** (phosphates/phosphonates, the LiPF_6 -derived OPF chemistry), and **glyme/oligoether** (the ether solvents and PEG linkers). These are literature

compounds bearing the *functional chemistry* of electrolytes, drawn from the open corpus; they are **not** operando or in-cell degradation spectra, and we make no claim that they are. They are excluded from every other split in this paper. Each compound was solved blind under the identical decoupled-agent protocol (Opus, closed-book, up to three ranked candidates, RDKit only for formula/parse checks).

Performance lands in the **same broad regime as the headline benchmark** — overall **top-1 26%, recovered 28%** (12/46 and 13/46; two compounds received no parseable candidate; at n=46 the 95% CI is wide and overlaps the headline) — consistent with the bottleneck being a property of the elucidation task rather than of any one chemical neighbourhood. The per-class breakdown (Fig. 6) is itself informative:

Table 4. Per-class performance on IRSpectra-Bench-Electrolyte (n=46).

electrolyte class	n	top-1	recovered (top-3)
sp ³ -C-F	8	50%	50%
carbonate	7	29%	43%
phosphoryl	8	25%	25%
glyme / oligoether	7	29%	29%
sulfonyl / sulfonate	8	12%	12%
nitrile	8	12%	12%

The gradient is chemically legible. **sp³-C-F** centres are the easiest: a C-F coupling fingerprint (the large $^1\text{J}/^2\text{J}(\text{C},\text{F})$ splittings) localises the fluorine and pins regiochemistry, removing exactly the degeneracy that defeats elucidation elsewhere. **Sulfonyl** and **nitrile** are the hardest, for two distinct reasons that recur in real electrolyte analysis: sulfur/phosphorus **oxidation-state ambiguity** — sulfide vs. sulfoxide vs. sulfone, $-\text{SCF}_3$ vs. $-\text{SO}_2\text{CF}_3$, the very distinctions an interphase study must make — is poorly constrained by $^1\text{H}/^{13}\text{C}$ shifts alone (it lives in the IR and in heteronuclei the benchmark does not score); and nitrile-bearing targets in the corpus sit on heavily substituted heteroaromatic cores whose regiochemistry the 1D spectra underdetermine. This is the recall-bound failure of §4.1 reappearing in a domain-specific guise, and it is precisely where the **forward-verification** recipe of §5 — compute the spectrum each candidate *would* give and match it to the one observed — plays a role *analogous* to (not a replacement for) computational-NMR / NMR-crystallography structure validation, sharing its forward-prediction logic while trading DFT’s calibrated accuracy for zero setup cost. The subset, its per-class answers, and the scorer are released with the benchmark.

5. Forward-verification elucidation

5.1 Method

The inverse direction is the model’s hard, isomer-blind direction; the **forward** direction (structure→spectrum) is its easy, accurate one.² Regioisomers, crucially, have *different* forward-predicted ^{13}C shifts. We therefore close a generator–verifier loop:

generate candidate structures (inverse) → **forward-predict** each candidate’s ^{13}C spectrum, blind to the observed spectrum → **re-rank** candidates by the distance between predicted and observed ^{13}C → return the best match.

This mirrors how a chemist confirms a structure (“if it were that isomer, C-3 would be at ~120 ppm; we observe 135, so it is the other”), exploits the standard principle that verification is easier than generation, and requires no training. **Figure 7** shows the mechanism on a real benchmark pair — the picolinamide / nicotinamide regioisomers, which the inverse task cannot separate but whose forward-predicted ^{13}C spectra match the observed one at 0.42 vs 1.30 ppm. We implemented the verifier as independent LLM agents that predict ^{13}C shift lists from SMILES alone (candidates shuffled and anonymised so isomers of one target never co-occur; pure reasoning, no tools), and matched predicted to observed ^{13}C with a symmetric chamfer distance over peak sets.

For balance — and because conditional precision is 72–84%, not 100% — the verifier also produces clear **false positives** on near-degenerate pairs. On the 2-(nitrophenyl)-2,3-dihydroquinazolin-4(1*H*)-one targets ($\text{C}_8\text{H}_7\text{N}_3\text{O}_3$), the forward predictor cannot separate the *ortho*- and *meta*-nitrophenyl regioisomers: their predicted ^{13}C lie within its ~2 ppm error, and it ranks the wrong (2-nitrophenyl) isomer first by a chamfer of 1.35 ppm versus 1.36 ppm for the true (3-nitrophenyl) one — a 0.01 ppm margin, i.e. effectively a coin flip. This is the precision-loss mechanism of §5.3 made concrete: when candidates are near-degenerate the cheap predictor cannot resolve them, and it is exactly these cases — not generation failures — that a sharper (DFT-level) or 2D-NMR-grounded verifier would need to fix. We report it alongside the success so the gain is read against its failure mode, not in isolation.

Conceptually this is **analogous to NMR-crystallography logic, with an LLM in place of the quantum chemistry** — an analogy, not an equivalence: where DP4/DP4+ rank candidates by GIAO-DFT shifts^{16 17} and NMR crystallography adjudicates polymorphs and connectivity by GIPAW-computed shifts^{18 19}, we rank by the shifts a forward LLM predicts. The shared principle is *verification by forward prediction*; what we do **not** inherit is DFT’s calibrated error model. The trade is deliberate — we forgo the ~1–2 ppm accuracy and the rigorous, probabilistic error model of DFT (the very thing that lets DP4 emit a posterior probability) for a predictor that runs in seconds at zero setup, and we quantify below exactly how far that cheaper, uncalibrated verifier carries the inverse problem.

5.2 Result

On the 60 benchmark compounds (126 candidate structures from the solver agents):

Table 5. Forward-verification decomposition on the 60 forward-verify compounds.

	value
generation recall (true structure among candidates)	31%
top-1, solver self-ranking	23%
top-1, forward-verified re-ranking	26%
conditional on recall — self-ranking	14/19 (73%)
conditional on recall — forward-verification	16/19 (84%)

The decomposition is the finding. **When the true structure is among the candidates, forward verification selects it 84% of the time, versus 73% for the model’s own ranking (+11 points)** — a real, exploitable generator–verifier gap. The overall top-1 moves only 23%→26% because the binding constraint is **generation recall**: the true structure was never proposed for 41 of 60 compounds, which no re-ranking can repair.

LLM elucidation therefore factorises into two near-independent levers: the **verifier is already strong (84%)**; the **generator (31% recall) is the wall**.

5.3 Generate-wide: testing the recipe

The decomposition implies a recipe — *generate wide, verify by forward prediction* — a chemistry analog of self-consistency sampling, where many independent generations are pooled and the answer is chosen by agreement rather than from a single pass.¹⁰ We tested it directly: ten independent solver agents proposed up to **six regiochemistry-aware candidates per compound**, pooled with the originals, and the 65 new candidates were forward-predicted and re-ranked as before. The measured result:

Table 6. Generate-wide vs original on the same 60 compounds.

	original	generate-wide
recall (true structure among candidates)	31%	41%
forward-verified top-1	26%	30%
verification precision (conditional on recall)	84%	72%

Wide generation lifts recall +10 points and exact top-1 +7 points (23%→30% over the self-ranking baseline on the same 60 compounds; Fig. 8) — a real, measured gain with no training. But it does **not** reach the ~50% a naïve extrapolation would predict, for two instructive reasons: **(i) recall plateaus at 41%** — exotic and large targets (selenium heterocycles, poly-aryl polyketones, eleven-nitrogen polyamines) resist even six regiochemical variants; and **(ii) verification precision falls 84%→72%** as more near-degenerate regioisomers enter the pool and the ~2-ppm forward predictor can no longer separate them. This recall/precision tension is the honest ceiling of the training-free approach: the recall-bound diagnosis stands, and closing the gap further requires either sharper verification or 2D-NMR constraints, not merely more candidates.

5.4 Does a deterministic verifier help? A HOSE-code ablation

§5.3 and a natural reading of the verifier-precision story suggest an obvious fix: swap the LLM forward-predictor for a *deterministic* ¹³C predictor with a rigorous error model. We tested the canonical choice — a **HOSE-code-style lookup trained on nmrshiftdb2** (RDKit radial-environment bins, spheres r=4→1 with a hybridisation prior fallback; 31,000 molecules, 332,595 assigned carbons; held-out **MAE 3.23 ppm, median 1.73 ppm**) — as a drop-in replacement for the verifier, re-ranking the same §5.2 candidates.

It does **not** help. The HOSE verifier recovers **14/19 (73%)** of the in-set compounds — identical to the solver’s own ranking and *below* the LLM verifier’s **16/19 (84%)** on the same set. The reason is coverage, not the method: of the 2,244 candidate carbons, only **2%** match a training environment at the most specific sphere (r=4) and **67%** resolve only at coarse spheres (r≤2), because the benchmark’s exotic chemistry (selenium heterocycles, polyaryl ketones, large polyamines) is under-represented in nmrshiftdb2. A coarse environment cannot separate the regioisomers the verifier exists to separate, so the deterministic predictor degrades to the self-rank baseline exactly where discrimination is needed — whereas the LLM forward-predictor generalises across novel scaffolds. The practical lesson is the opposite of the naïve one: on hard, real chemistry the LLM’s *breadth* is an asset for verification, and the genuine fix is sharper still — **compound-specific (DFT-level) shift accuracy or orthogonal 2D-NMR constraints**, not a generic lookup table.

5.5 Negative control

Permutation negative control (Y-randomisation analog). A re-ranker earns the interpretation that it exploits genuine predicted-vs-observed ^{13}C agreement only if that agreement is destroyed when the pairing is broken. Borrowing the Y-randomisation discipline of QSAR validation, we re-paired — as a *derangement*, so no compound keeps its own spectrum — which observed ^{13}C spectrum each candidate set is scored against, and re-ran the verifier 1,000 times. Conditional-on-recall precision falls from the true **84.2% (16/19)** to a permuted mean of **66.4%** (95% range 52.6–78.9%; one-sided empirical $p = 0.019$, two-sided $p = 0.038$) — the verifier acts on real spectral agreement, not a candidate-list artefact. The honest caveat is the height of the chance floor: because recall-positive compounds carry few and near-identical (regioisomeric) candidates, even a random pairing lands on the correct structure $\sim 66\%$ of the time, so the verifier’s genuine margin over chance is **~ 18 points**, not the full 84% — real and significant, but to be read against this high baseline.

Confidence calibration (a negative result). We also asked whether the chamfer match-distance doubles as a confidence signal that would support selective prediction. It does not. Ranking the 48 multi-candidate compounds by their chamfer margin (the gap between the best and second-best predicted-vs-observed distance) and answering only the most-confident fraction leaves top-1 essentially flat and non-monotonic with coverage ($\approx 21\%$ overall, neither rising nor falling reliably as the threshold tightens). Compounds with a single proposed candidate have no margin at all and must be excluded; an earlier analysis that retained them produced a spurious “improvement” that was entirely an artefact of those trivial cases. We therefore do **not** claim the verifier distance as a calibrated abstention gauge — a sharper, DFT/2D-NMR-grounded confidence estimate (§5.4) remains the route to reliable selective prediction.

6. Discussion

The frontier LLMs we tested (the Claude family) are not “solving structure elucidation” on realistic 1D data, but neither are they failing: they are reliable **scaffold-level** elucidators (scaffold-correct on 56% of compounds — 73% of simple ones; mean best Tanimoto 0.59) and good **verifiers** (84% conditional), with exact recovery throttled by candidate recall and by the regiochemical underdetermination intrinsic to 1D NMR. The contribution of this paper is therefore best read as a **diagnosis with a bounded, training-free improvement attached**, not as a method that solves the task. The diagnostic result is the durable take-away: across models, domains, and verifiers, the wall is **generation recall**, not verification. The accompanying improvement is real but deliberately modest — forward verification with wide generation lifts top-1 from 23% to 30%, and we show by direct measurement (§5.3) that this stays *below its own ceiling* (recall plateaus at 41%; verification precision falls to 72% as near-degenerate regioisomers accumulate), while the match distance fails as a confidence gauge for selective prediction (§5.5, a null result we report rather than bury). This reframes the engineering problem. Rather than training a bespoke spectra $\rightarrow$ structure model — a target the frontier already meets at the scaffold level and that ages out each model generation — the durable levers are (i) **open, hard, honestly scored benchmarks** that expose specific failure modes (here, regiochemistry and recall), and (ii) **inference-time scaffolding** (decoupled solving, generate-and-verify) that needs no training and improves with each model. IRExp’s IR modality and 2D-NMR-ready provenance position it for the obvious next step: supplying the HMBC/COSY/NOESY constraints that would attack regiochemistry at the source.

Protocol over fashionable levers. That *how* the problem is posed and verified — decoupled per-compound solving, generate-and-verify — moves accuracy more than raw model capability echoes a recurring lesson across cheminformatics: careful pipeline and protocol design dominates the fashionable component. The same pattern appears in molecular property prediction, where a recent systematic benchmark reports that learned, pretrained molecular embeddings rarely outperform classical ECFP fingerprints once evaluation is controlled²¹ — the modelling fashion underperforms the well-engineered baseline. We read our within-compound control (§4.3) and forward-verification recipe (§5) in the same light: the durable gains in LLM elucidation come from honest benchmarking and inference-time scaffolding rather than from waiting on the next, larger model. We draw this parallel narrowly and claim no formal equivalence between the two settings.

That the bottleneck reproduces inside a single application domain (§4.5: 26% top-1 on battery-electrolyte chemistry, n=46, against 28% overall) is consistent with it being structural rather than incidental — and it makes the forward-verification recipe directly relevant to the magnetic-resonance workflows (electrolyte-decomposition assignment, NMR crystallography) where computing a candidate’s spectrum to confirm it is already standard practice.

For practitioners, two operational findings transfer immediately: solve each problem in a fresh context with tool access ($\approx 3\times$ over a single long context), and trust a proposed structure in proportion to how well its forward-predicted spectrum matches the observed one — while remembering the abstention caveat below.

7. Limitations

Several limitations of earlier drafts are now resolved with controlled experiments; we state what remains plainly.

Resolved. **Extraction noise:** an RDKit self-consistency audit (¹³C peak count vs symmetry-unique carbons, formula, SELFIES round-trip) finds **57/60 ground truths spectrally clean**, and all headline metrics are unchanged on that clean subset (top-1 24% vs 23%, recall 33% vs 31%) — the conclusions are not driven by scraping artefacts. **Verifier precision / abstention:** the generate-wide experiment (§5.3) quantifies the recall/precision tension directly — verification precision is 72–84% conditional on recall and degrades as near-degenerate regioisomers accumulate, so forward-match distance is a strong *re-ranker* but a soft *confidence* gauge. We further tested the obvious deterministic fix (§5.4): a nmrshiftdb2-trained HOSE-code ¹³C predictor does **not** beat the LLM verifier (73% vs 84% conditional), because its specific-environment coverage collapses on the benchmark’s exotic chemistry — so the identified fix is compound-specific DFT-level accuracy or 2D-NMR constraints, not a generic lookup. **Projection:** §5.2’s extrapolation is replaced by the **measured** §5.3 result (top-1 30%, recall 41%) — and is honestly below the optimistic estimate.

Independence checks. Scoring throughout is mechanical RDKit (not LLM-judged), and all solver/verifier runs were transcript-audited *at generation time* for zero web and zero ground-truth access; the committed artefacts are the parsed per-compound predictions, and the underlying transcripts are available on request. To confirm that the forward-verifier’s measured advantage reflects real predicted-vs-observed spectral agreement rather than leakage or a candidate-list artefact, we ran a permutation negative control (Y-randomisation analog, §5.5): re-pairing — as a derangement — which observed ¹³C spectrum each candidate set is scored against, over 1,000 permutations, collapses conditional-on-recall precision from 84.2% to a chance mean of 66.4%

(two-sided $p=0.038$) — the signal weakens to the chance floor under label-shuffle, as a correctly isolated blind evaluation requires. A second model family (Sonnet) re-solved a 12-compound subset under the identical protocol and is comparably recall-bound (recall 33% vs Opus 41%, with cross-family ensembling adding no recall), indicating the recall-bound result is a property of the task, not an artefact of one model.

Remaining. (i) **Human audit — prepared but not yet run.** Solver and verifier are both LLMs, so the one validation we cannot perform ourselves is an expert-chemist review of a sample of elucidations and forward predictions. We have therefore **built and frozen** a blinded, pre-registered audit package — a difficulty-stratified 30-compound sample (9 recall-positive), rendered candidate structures, a separate withheld answer key, and a mechanical scoring sheet — released at `data/audit/` with its protocol at `docs/EXPERT_AUDIT_PROTOCOL.md` (regenerable via `scripts/make_audit_sample.py`), designed to test the two load-bearing claims (misses are predominantly consistent regioisomers, §4; forward verification is a trustworthy re-ranker, §5). **We have not yet run the panel**; until expert results are in, those claims should be read as machine-validated (RDKit InChIKey) but not yet human-validated.

(ii) **Single vendor and an underpowered cross-model comparison.** Every number comes from one vendor’s models. The headline (28.4% top-1) is a single frontier model (Claude Opus), and the cross-model evidence (§4.4) is four **Claude-family** models on a shared **24-compound** subset. That comparison is statistically underpowered: only the extreme Fable-vs-Haiku contrast survives multiple-comparison correction (Holm-adjusted $p=0.006$); Fable-vs-Opus is not significant ($p=0.063$) and Opus-vs-Sonnet are indistinguishable, with overlapping bootstrap CIs. What is robust is the **ranking** — outcomes are strictly nested (Haiku < Sonnet < Opus < Fable) and the weakest floors at 0% — so the benchmark is capability-sensitive; what is *not* established at $n=24$ is the size of any adjacent gap. Quantitative claims should therefore be read as holding **for the Claude family**, and a true cross-vendor sweep (GPT-, Gemini-class, open-weight models) — the test of whether the pattern is a property of the task rather than one lineage — was forgone because it needs paid API access incompatible with our zero-cost protocol; we flag it as the most important external-validity experiment left open.

(iii) **Domain-subset scope:** the battery-electrolyte case study (§4.5) comprises *literature compounds bearing electrolyte-relevant functional chemistry* drawn from the open corpus — not operando or in-cell decomposition spectra — so it demonstrates functional-class transfer of the elucidation bottleneck, not direct assignment of authentic interphase/degradation products, which would require a dedicated operando set.

(iv) **Constitution-only scoring:** correctness is judged on InChIKey connectivity, so a correct-constitution / wrong-stereochemistry prediction is counted correct. This affects at most the 10.3% (20/194) of targets with a defined stereocentre and is intrinsic to what 1D NMR/IR can determine; a stereochemistry-sensitive benchmark would need 2D/chiroptical data.

(v) **Single-sample scoring:** each headline compound is scored from one solver prediction set (one decoupled run), so reported top-1/recall carry no run-to-run (LLM-sampling) variance estimate — the bootstrap CIs reflect compound sampling only. The generate-wide experiment (§5.3) pools ten independent generation passes (recall 31%→41%), indicating generator stochasticity that single-pass scoring understates.

(vi) **Chemical-space coverage:** the benchmark is drawn from open-access organic methodology and total-synthesis literature, covering small-to-medium drug-like and synthetic-organic space; it is **not** representative of organometallic/coordination compounds, large biomolecules (peptides,

oligonucleotides, oligosaccharides), or stereochemistry-heavy targets, and our results should not be extrapolated to them.

8. Methods

Mining and resolution. PMC-OA full text was fetched from `s3://pmc-oa-opendata`; records were extracted with a deterministic parser and resolved with OPSIN, RDKit, and SELFIES, with an optional cached PubChem fallback. Durable content-keyed caches make resolution additive and restart-safe.

Benchmark and agents. Problems were sampled from `irexp_resolved`, stratified by RDKit ring analysis, with cross-round de-duplication by InChIKey. The battery-electrolyte subset (§4.5) was drawn from the same corpus by SMARTS substructure filters for six electrolyte functional classes (carbonate, sulfonyl/sulfonate, nitrile, $\text{sp}^3\text{-C-F}$, phosphoryl, glyme/oligoether), balanced to eight compounds per class (48 curated; 46 scored after two yielded no parseable candidate), excluding every compound used elsewhere, and J-enriched and spectrally validated identically to the main rounds. Solver and forward-prediction agents were independent Claude-Opus sub-agents invoked under a consumer subscription; agents were instructed closed-book and audited via transcript grep for zero web/answer access. Scoring used RDKit InChIKey-connectivity matching and Morgan(2, 2048) Tanimoto; forward-verification used a symmetric chamfer distance over ^{13}C peak sets. No model was trained or fine-tuned; no paid API was used.

Models and versions. All experiments used Anthropic Claude models via the consumer subscription (Claude Code / `claude.ai`). The §4.4 comparison spanned, in capability order, Claude Haiku, Claude Sonnet, Claude Opus, and Claude Fable 5; the headline benchmark (§4.1) and forward-verification (§5) used Claude Opus. Exact model snapshot identifiers and the access window are recorded in `docs/MODELS.md` so the benchmark can be re-run against the same checkpoints.

Reproducibility. Every round is frozen: questions, ground-truth answers, per-agent raw outputs, predictions, and scorer outputs are released, and the sampler, scorer, and forward-verification harness are scripted end-to-end.

Figures

- **Fig. 1** (`docs/figures/fig0_overview.png`) — study design: open multimodal data (IRexp) → blind, complexity-stratified benchmark → decoupled blind solving → forward-verification re-ranking; training-free throughout.
- **Fig. 2** (`docs/figures/fig4_dataset.png`) — IRexp composition: IR records → NMR-paired → structure-linked → full $\text{IR}+^1\text{H}+^{13}\text{C}$ +structure quadruples.
- **Fig. 3** (`docs/figures/fig1_difficulty.png`) — top-1 and recovered accuracy on IRSpectra-Bench by difficulty (all / simple / complex, $n=194$) with bootstrap 95% CIs.
- **Fig. 4** (`docs/figures/fig2_size.png`) — accuracy vs molecular size (heavy-atom bucket); the monotonic 60%→7% top-1 gradient.
- **Fig. 5** (`docs/figures/fig5_models.png`) — four-model comparison on a 24-compound subset: Fable 5 45% > Opus 25% > Sonnet 20% > Haiku 0% top-1 (strictly nested); the benchmark discriminates capability monotonically and is far from saturated even for the newest model.

- **Fig. 6** (docs/figures/fig6_electrolyte.png) — top-1 and recovered accuracy on IRspectra-Bench-Electrolyte by battery-electrolyte chemical class (n=46): sp³-C-F easiest (50%), sulfonyl and nitrile hardest (12%); overall 26%/28%, the same regime as the headline benchmark.
- **Fig. 7** (docs/figures/fig_mechanism.png) — forward-verification on a real benchmark regioisomer pair (picolinamide vs nicotinamide): forward-predicted ¹³C matches the true isomer (chamfer 0.42 vs 1.30 ppm), the LLM analog of NMR-crystallography.
- **Fig. 8** (docs/figures/fig3_method.png) — forward-verification inference ladder on the same 60 compounds: solver self-ranking → + forward-verify → + generate-wide (23%/26%/30% top-1).

Data and code availability

All data and code are released in the project repository: IRexp and the `irexp_resolved` split (`data/irexp/`, `data/irexp_resolved/`); the benchmark rounds and within-compound control (`data/benchmark*`), scored by `scripts/benchmark_v2.py`; the battery-electrolyte case-study subset (`data/benchmark_electrolyte/`, built by `scripts/build_electrolyte_bench.py`, scored by `scripts/score_electrolyte.py`); the ground-truth integrity audit (`scripts/validate_benchmark.py`, `data/benchmark*/clean_qids.json`); the forward-verification and generate-wide experiments (`data/fverify/`, `data/gw/`, `data/fverify2/`, `scripts/forward_verify.py`); the deterministic HOSE-code verifier ablation (`scripts/hose_predict.py`, `data/fverify/hose_results.txt`); and the mining/resolution pipeline (`scripts/`). Companion technical notes: `docs/BENCHMARK.md`, `docs/BENCHMARK_v2.md`, `docs/BENCHMARK_v3.md`, `docs/FORWARD_VERIFY.md`, and `data/SPECTRO_TRAINING_DATA`. The verifier negative-control and selective-prediction analyses (`scripts/verifier_diagnostics.py`), the recall-headroom and scaffold-enumeration study (`scripts/analyze_recall_headroom.py`, `scripts/enumerate_isomers.py`, `scripts/closing_the_gap.py`), the modality-ablation harness (`scripts/modality_ablation.py`, `docs/MODALITY_ABLATION.md`), and the blinded expert-audit package (`data/audit/`, `docs/EXPERT_AUDIT_PROTOCOL.md`) are released likewise. **Archival deposit:** a complete frozen snapshot (dataset, benchmark, answer keys, predictions, scripts, figure-regeneration, and the expert-audit package) will be archived on Zenodo under DOI [TODO: 10.5281/zenodo.XXXXXXX — mint on submission]; the GitHub repository is the development mirror and the Zenodo record the citable version of record. (The held-out answer keys `data/audit/key.jsonl` and `data/modality/key.json` are deposited but flagged “withhold from blinded reviewers”.)

Licensing and attribution

IRexp is derived from two open-access source pools and redistributed under terms compatible with each (see `data/NOTICE`). **(a) Redistribution:** we release only *extracted numeric data* — IR band lists, ¹H/¹³C shift lists, and resolved structures (SMILES/SELFIES/InChIKey) — plus each record’s source DOI/accession, not source full text, figures, or PDFs. **(b) Two separable pools:** 119,345 records derive from the PMC Open-Access Subset (**CC-BY-4.0**) and 1,888 from the Chemotion RADAR4Chem FT-IR deposit (**CC-BY-SA-4.0**); the pools are kept separate and individually labelled, so users may take the CC-BY pool alone, while any combined or Chemotion-derived release carries CC-BY-SA-4.0 to honour the ShareAlike term. Code is released under [TODO: MIT / Apache-2.0 — authors to confirm]. **(c) Attribution:** re-users must cite this dataset (Zenodo DOI above) and attribute the original publications via each record’s `source_doi`.

Author Contributions

(*CRedit* — authors to confirm/adjust.) **I.Y.:** conceptualization, methodology, software, formal analysis, investigation, data curation, visualization, writing — original draft. **R.A.V.-H.:** conceptualization, methodology, supervision, writing — review and editing.

Conflicts of Interest

The authors declare no competing interests.

Acknowledgements

(To be completed: funding sources and compute/infrastructure acknowledgements.)

References

1. *Spectro: transformer-based molecular structure elucidation from spectra*. ChemRxiv, 2024, doi:10.26434/chemrxiv-2024-37v2j.
2. Kamber, D. (Anthropic). *Making Claude a Chemist: how Claude performs on NMR prediction and structure elucidation*. Non-peer-reviewed company white paper, 5 June 2026. anthropic.com/research/making-claude-a-chemist.
3. Lowe, D. M. et al. *Name to Structure (OPSIN)*. J. Chem. Inf. Model. 2011, 51, 739.
4. Landrum, G. *RDKit: Open-source cheminformatics*. rdkit.org.
5. Krenn, M. et al. *SELFIES*. Mach. Learn.: Sci. Technol. 2020, 1, 045024.
6. *NCBI PMC Open Access Subset*. National Library of Medicine.
7. *Chemotion repository / RADAR4Chem*, doi:10.22000/OGoeEQGlSZGEIrgst.
8. *NIST Chemistry WebBook*, SRD 69.
9. Rogers, D.; Hahn, M. *Extended-connectivity fingerprints*. J. Chem. Inf. Model. 2010, 50, 742.
10. Wang, X. et al. *Self-consistency improves chain-of-thought reasoning*. ICLR 2023.
11. *Accurate and efficient structure elucidation from routine one-dimensional NMR spectra using multitask machine learning*. arXiv:2408.08284, 2024.
12. *NMRTrans: structure elucidation from experimental NMR spectra via set transformers*. arXiv:2602.10158, 2026.
13. Ottomano, F.; Li, Y.; Ganose, A. M. *NMIRacle: multi-modal generative molecular elucidation from IR and NMR spectra*. arXiv:2512.19733, 2025.
14. *SpectraLLM / MolSpectLLM: multimodal language models for molecular structure elucidation from multi-spectral input*. arXiv:2508.08441; arXiv:2509.21861, 2025.
15. *Boosting LLM molecular structure elucidation with knowledge-enhanced tree-search reasoning*. arXiv:2506.23056, 2025.
16. Smith, S. G.; Goodman, J. M. *Assigning stereochemistry to single diastereoisomers by GIAO NMR calculation: the DP₄ probability*. J. Am. Chem. Soc. 2010, 132, 12946.
17. Grimblat, N.; Zanardi, M. M.; Sarotti, A. M. *Beyond DP₄: an improved probability for the stereochemical assignment of isomeric compounds (DP₄+)*. J. Org. Chem. 2015, 80, 12526.
18. Pickard, C. J.; Mauri, F. *All-electron magnetic response with pseudopotentials: NMR chemical shifts (GIPAW)*. Phys. Rev. B 2001, 63, 245101.
19. Ashbrook, S. E.; McKay, D. *Combining solid-state NMR spectroscopy with first-principles calculations — a guide to NMR crystallography*. Chem. Commun. 2016, 52, 7186.

20. Guo, K. et al. *MolPuzzle: a benchmark for molecular structure elucidation with multimodal large language models*. NeurIPS 2024 (Datasets and Benchmarks Track).
21. *Benchmarking pretrained molecular embedding models for molecular representation learning*. arXiv:2508.06199, 2025.
22. *Spectral Database for Organic Compounds (SDBS)*. National Institute of Advanced Industrial Science and Technology (AIST), Japan. <https://sdb.sdb.aist.go.jp> (accessed 2026; view-only, not bulk-redistributable).

Figure plates




Fig. 1: Study design: open multimodal data (IReXP) → blind, complexity-stratified benchmark → decoupled blind solving → forward-verification re-ranking; training-free throughout.




Fig. 2: IRexp composition: IR records, NMR-paired, structure-linked, and full IR+¹H+¹³C+structure quadruples.




Fig. 3: Top-1 and recovered accuracy on IRSpectra-Bench by difficulty (all / simple / complex, n=194) with bootstrap 95% CIs.




Fig. 4: Accuracy versus molecular size; the monotonic 60% $\rightarrow$ 7% top-1 gradient with heavy-atom count.




Fig. 5: Four-model comparison on a 24-compound subset: Fable 5 45% > Opus 25% > Sonnet 20% > Haiku 0% top-1 (strictly nested; underpowered to separate adjacent models at n=24).




Fig. 6: IRSpectra-Bench-Electrolyte by battery-electrolyte class (n=46): sp³-C-F easiest (50%), sulfonyl and nitrile hardest (12%).




Fig. 7: Forward-verification on a real benchmark regioisomer pair (picolinamide vs nicotinamide): forward-predicted ¹³C matches the true isomer (chamfer 0.42 vs 1.30 ppm) — the LLM analog of NMR-crystallography.




Fig. 8: Forward-verification inference ladder on the same 60 compounds: solver self-ranking $\rightarrow$ + forward-verify $\rightarrow$ + generate-wide (23%/26%/30% top-1).