

# Locally Minimax Optimal and Dimension-Agnostic Discrete Argmin Inference

Ilmun Kim

Department of Statistics & Data Science  
Yonsei University



# Joint work with



Aaditya Ramdas  
(Carnegie Mellon University)

# Problem Setup: Discrete Argmin Inference

- Suppose  $X_1, \dots, X_{2n} \stackrel{\text{i.i.d.}}{\sim} \mathbf{P}$  in  $\mathbb{R}^d$  with mean  $\mu = (\mu_1, \dots, \mu_d)^\top$  and let

$$\Theta = \Theta(\mathbf{P}) := \arg \min_{k \in [d]} \mu_k$$

# Problem Setup: Discrete Argmin Inference

- Suppose  $X_1, \dots, X_{2n} \stackrel{\text{i.i.d.}}{\sim} \mathbf{P}$  in  $\mathbb{R}^d$  with mean  $\mu = (\mu_1, \dots, \mu_d)^\top$  and let

$$\Theta = \Theta(\mathbf{P}) := \arg \min_{k \in [d]} \mu_k$$

i.e., the set of all coordinates whose mean equals the *smallest* in  $\mu$

# Problem Setup: Discrete Argmin Inference

- Suppose  $X_1, \dots, X_{2n} \stackrel{\text{i.i.d.}}{\sim} \mathbf{P}$  in  $\mathbb{R}^d$  with mean  $\mu = (\mu_1, \dots, \mu_d)^\top$  and let

$$\Theta = \Theta(\mathbf{P}) := \arg \min_{k \in [d]} \mu_k$$

i.e., the set of all coordinates whose mean equals the *smallest* in  $\mu$

- We would like to form a **confidence set**  $\widehat{\Theta}$  for  $\Theta$

# Problem Setup: Discrete Argmin Inference

- Suppose  $X_1, \dots, X_{2n} \stackrel{\text{i.i.d.}}{\sim} \mathbf{P}$  in  $\mathbb{R}^d$  with mean  $\mu = (\mu_1, \dots, \mu_d)^\top$  and let

$$\Theta = \Theta(\mathbf{P}) := \arg \min_{k \in [d]} \mu_k$$

i.e., the set of all coordinates whose mean equals the *smallest* in  $\mu$

- We would like to form a **confidence set**  $\widehat{\Theta}$  for  $\Theta$

Each  $r \in \Theta$  is included in  $\widehat{\Theta}$  with high probability:


$$\inf_{\mathbf{P} \in \mathcal{P}} \inf_{r \in \Theta(\mathbf{P})} P(r \in \widehat{\Theta}) \geq 1 - \alpha$$



$\mathcal{P}$ : class of distributions  
 $\alpha$ : target error level


# Modern Application

**Prompt:** “Generate cat images”



# Modern Application

**Prompt:** “Generate cat images”




• • •



- The best model minimizes the **population risk**

# Modern Application

**Prompt:** “Generate cat images”




• • •



- The best model minimizes the **population risk**
- But **the population risk is unknown** → **the empirical risk**

# Modern Application

**Prompt:** “Generate cat images”



- The best model minimizes the **population risk**
- But **the population risk is unknown** → **the empirical risk**
- We must **account for statistical uncertainty** to determine which models are plausibly **optimal with statistical confidence**

# We aim to develop a method for argmin inference with

I. Dimension-agnostic validity: works in both low- and high-dimensional settings

# We aim to develop a method for argmin inference with

- I. Dimension-agnostic validity: works in both **low-** and **high-dimensional** settings
2. Powerful inference: attains **local minimax rates** across diverse regimes

# We aim to develop a method for argmin inference with

- I. **Dimension-agnostic validity:** works in both **low-** and **high-dimensional** settings
2. **Powerful inference:** attains **local minimax rates** across diverse regimes
3. **Robustness to complex data:** handles **ties** and strong **dependence**

# We aim to develop a method for argmin inference with


- I. **Dimension-agnostic validity:** works in both **low-** and **high-dimensional** settings
2. **Powerful inference:** attains **local minimax rates** across diverse regimes
3. **Robustness to complex data:** handles **ties** and strong **dependence**
4. **Model-agnostic applicability:** requires **no parametric** assumptions

# We aim to develop a method for argmin inference with

- I. **Dimension-agnostic validity:** works in both **low-** and **high-dimensional** settings
2. **Powerful inference:** attains **local minimax rates** across diverse regimes
3. **Robustness to complex data:** handles **ties** and strong **dependence**
4. **Model-agnostic applicability:** requires **no parametric** assumptions
5. **Tuning-free implementation:** requires **no** (non-trivial) **tuning** parameters

# We aim to develop a method for argmin inference with

1. **Dimension-agnostic validity:** works in both **low-** and **high-dimensional** settings
2. **Powerful inference:** attains **local minimax rates** across diverse regimes
3. **Robustness to complex data:** handles **ties** and strong **dependence**
4. **Model-agnostic applicability:** requires **no parametric** assumptions
5. **Tuning-free implementation:** requires **no (non-trivial) tuning** parameters



**Sample Splitting + Studentization (Kim and Ramdas, 2024)**

# Dimension-Agnostic Argmin Inference

## Formal (Primal) Goal.

- We seek a **dimension-agnostic confidence set**  $\widehat{\Theta}$  for  $\Theta$  that such that

$$\liminf_{n \rightarrow \infty} \inf_{P \in \mathcal{P}} \inf_{r \in \Theta(P)} P(r \in \widehat{\Theta}) \geq 1 - \alpha, \text{ regardless of the sequence } (d_n)_{n=1}^{\infty}$$

## Formal (Primal) Goal.

- We seek a **dimension-agnostic confidence set**  $\widehat{\Theta}$  for  $\Theta$  that such that

$$\liminf_{n \rightarrow \infty} \inf_{P \in \mathcal{P}} \inf_{r \in \Theta(P)} P(r \in \widehat{\Theta}) \geq 1 - \alpha, \text{ regardless of the sequence } (d_n)_{n=1}^{\infty}$$

- We also seek a **confidence set**  $\widehat{\Theta}$  for  $\Theta$  such that its **expected cardinality** is small and **ideally optimal**

# Formal (Dual) Goal.

- Given some fixed  $r \in [d]$ , consider the **null** and **alternative** hypotheses:

$$H_0 : r \in \Theta \quad \text{versus} \quad H_1 : r \notin \Theta$$

## Formal (Dual) Goal.

- Given some fixed  $\textcolor{red}{r} \in [d]$ , consider the **null** and **alternative** hypotheses:

$$H_0 : \textcolor{red}{r} \in \Theta \quad \text{versus} \quad H_1 : \textcolor{red}{r} \notin \Theta$$

- Let  $\psi_{\textcolor{red}{r}} : \{X_1, \dots, X_{2n}\} \rightarrow \{0,1\}$  denote a test function that rejects  $H_0$  if  $\psi_{\textcolor{red}{r}} = 1$

# Formal (Dual) Goal.

- Given some fixed  $\mathbf{r} \in [d]$ , consider the **null** and **alternative** hypotheses:

$$H_0 : \mathbf{r} \in \Theta \quad \text{versus} \quad H_1 : \mathbf{r} \notin \Theta$$

- Let  $\psi_{\mathbf{r}} : \{X_1, \dots, X_{2n}\} \rightarrow \{0,1\}$  denote a test function that rejects  $H_0$  if  $\psi_{\mathbf{r}} = 1$
- We seek a **dimension-agnostic test**  $\psi_{\mathbf{r}}$  such that

$$\limsup_{n \rightarrow \infty} \sup_{P \in \mathcal{P}_{0,\mathbf{r}}} P(\psi_{\mathbf{r}} = 1) \leq \alpha, \quad \text{regardless of the sequence } (d_n)_{n=1}^{\infty}$$



# Formal (Dual) Goal.

- Given some fixed  $\mathbf{r} \in [d]$ , consider the **null** and **alternative** hypotheses:

$$H_0 : \mathbf{r} \in \Theta \quad \text{versus} \quad H_1 : \mathbf{r} \notin \Theta$$

- Let  $\psi_{\mathbf{r}} : \{X_1, \dots, X_{2n}\} \rightarrow \{0,1\}$  denote a test function that rejects  $H_0$  if  $\psi_{\mathbf{r}} = 1$
- We seek a **dimension-agnostic test**  $\psi_{\mathbf{r}}$  such that

$$\limsup_{n \rightarrow \infty} \sup_{P \in \mathcal{P}_{0,\mathbf{r}}} P(\psi_{\mathbf{r}} = 1) \leq \alpha, \quad \text{regardless of the sequence } (d_n)_{n=1}^{\infty}$$

- We can construct a **DA confidence set** using **DA tests**



# DA tests yield a DA confidence set via duality

- Suppose that each  $\psi_k$  satisfies the **dimension-agnostic (DA)** property:

$$\limsup_{n \rightarrow \infty} \sup_{P \in \mathcal{P}_{0,k}} P(\psi_k = 1) \leq \alpha, \text{ regardless of the sequence } (d_n)_{n=1}^{\infty}$$

# DA tests yield a DA confidence set via duality

- Suppose that each  $\psi_k$  satisfies the **dimension-agnostic (DA)** property:

$$\limsup_{n \rightarrow \infty} \sup_{P \in \mathcal{P}_{0,k}} P(\psi_k = 1) \leq \alpha, \text{ regardless of the sequence } (d_n)_{n=1}^{\infty}$$

- We run  $d$  such **DA** tests  $\psi_1, \dots, \psi_d$  and let

$$\widehat{\Theta} = \{k \in [d] : \psi_k = 0\}$$

# DA tests yield a DA confidence set via duality


- Suppose that each  $\psi_k$  satisfies the **dimension-agnostic (DA)** property:

$$\limsup_{n \rightarrow \infty} \sup_{P \in \mathcal{P}_{0,k}} P(\psi_k = 1) \leq \alpha, \text{ regardless of the sequence } (d_n)_{n=1}^{\infty}$$

- We run  $d$  such **DA** tests  $\psi_1, \dots, \psi_d$  and let

$$\widehat{\Theta} = \{k \in [d] : \psi_k = 0\}$$

For example,



# DA tests yield a DA confidence set via duality

- Suppose that each  $\psi_k$  satisfies the **dimension-agnostic (DA)** property:

$$\limsup_{n \rightarrow \infty} \sup_{P \in \mathcal{P}_{0,k}} P(\psi_k = 1) \leq \alpha, \text{ regardless of the sequence } (d_n)_{n=1}^{\infty}$$

- We run  $d$  such **DA** tests  $\psi_1, \dots, \psi_d$  and let

$$\widehat{\Theta} = \{k \in [d] : \psi_k = 0\}$$

which results in the **DA confidence set**

$$\liminf_{n \rightarrow \infty} \inf_{P \in \mathcal{P}} \inf_{r \in \Theta(P)} P(r \in \widehat{\Theta}) \geq 1 - \alpha, \text{ regardless of the sequence } (d_n)_{n=1}^{\infty}$$

# Procedures

# Reformulation

- Recall our (dual) goal is to test

$$H_0 : \mathbf{r} \in \Theta \quad \text{vs.} \quad H_1 : \mathbf{r} \notin \Theta$$

# Reformulation

- Recall our (dual) goal is to test

$$H_0 : \mathbf{r} \in \Theta \quad \text{vs.} \quad H_1 : \mathbf{r} \notin \Theta$$

- We can reformulate this as

$$H_0 : \mu_{\mathbf{r}} - \mu_{\mathbf{s}} \leq 0 \quad \text{vs.} \quad H_1 : \mu_{\mathbf{r}} - \mu_{\mathbf{s}} > 0$$

where  $\mathbf{s} = \arg \min_{k \in [d] \setminus \{\mathbf{r}\}} \mu_k$

# Reformulation

- Recall our (dual) goal is to test

$$H_0 : \mathbf{r} \in \Theta \quad \text{vs.} \quad H_1 : \mathbf{r} \notin \Theta$$

- We can reformulate this as

$$H_0 : \mu_{\mathbf{r}} - \mu_{\mathbf{s}} \leq 0 \quad \text{vs.} \quad H_1 : \mu_{\mathbf{r}} - \mu_{\mathbf{s}} > 0$$

where  $\mathbf{s} = \arg \min_{k \in [d] \setminus \{\mathbf{r}\}} \mu_k$

i.e., Target  $\mu_{\mathbf{r}}$  is greater than the smallest mean  $\mu_{\mathbf{s}}$ ?

# Reformulation

- Recall our (dual) goal is to test

$$H_0 : \mathbf{r} \in \Theta \quad \text{vs.} \quad H_1 : \mathbf{r} \notin \Theta$$

- When  $\mathbf{s}$  is known, use a one-sided t-test

$$\frac{\bar{X}_{\mathbf{r}} - \bar{X}_{\mathbf{s}}}{\hat{\sigma}_{\mathbf{r}, \mathbf{s}}} > z_{1-\alpha}$$

- We can reformulate this as

$$H_0 : \mu_{\mathbf{r}} - \mu_{\mathbf{s}} \leq 0 \quad \text{vs.} \quad H_1 : \mu_{\mathbf{r}} - \mu_{\mathbf{s}} > 0$$

where  $\mathbf{s} = \arg \min_{k \in [d] \setminus \{\mathbf{r}\}} \mu_k$

i.e., Target  $\mu_{\mathbf{r}}$  is greater than the smallest mean  $\mu_{\mathbf{s}}$ ?

# Reformulation

- Recall our (dual) goal is to test

$$H_0 : \mathbf{r} \in \Theta \quad \text{vs.} \quad H_1 : \mathbf{r} \notin \Theta$$

- We can reformulate this as

$$H_0 : \mu_{\mathbf{r}} - \mu_{\mathbf{s}} \leq 0 \quad \text{vs.} \quad H_1 : \mu_{\mathbf{r}} - \mu_{\mathbf{s}} > 0$$

where  $\mathbf{s} = \arg \min_{k \in [d] \setminus \{\mathbf{r}\}} \mu_k$

- When  $\mathbf{s}$  is known, use a one-sided t-test

$$\frac{\bar{X}_{\mathbf{r}} - \bar{X}_{\mathbf{s}}}{\hat{\sigma}_{\mathbf{r}, \mathbf{s}}} > z_{1-\alpha}$$

- But  $\mathbf{s}$  is unknown: one idea (Mogstad et al. 2024)

$$\max_{k \in [d] \setminus \{\mathbf{r}\}} \frac{\bar{X}_{\mathbf{r}} - \bar{X}_k}{\hat{\sigma}_{\mathbf{r}, k}} > c_{1-\alpha}$$

i.e., Target  $\mu_{\mathbf{r}}$  is greater than the smallest mean  $\mu_{\mathbf{s}}$ ?

# Reformulation

- Recall our (dual) goal is to test

$$H_0 : \mathbf{r} \in \Theta \quad \text{vs.} \quad H_1 : \mathbf{r} \notin \Theta$$

- We can reformulate this as

$$H_0 : \mu_{\mathbf{r}} - \mu_{\mathbf{s}} \leq 0 \quad \text{vs.} \quad H_1 : \mu_{\mathbf{r}} - \mu_{\mathbf{s}} > 0$$

where  $\mathbf{s} = \arg \min_{k \in [d] \setminus \{\mathbf{r}\}} \mu_k$

- When  $\mathbf{s}$  is known, use a one-sided t-test

$$\frac{\bar{X}_{\mathbf{r}} - \bar{X}_{\mathbf{s}}}{\hat{\sigma}_{\mathbf{r}, \mathbf{s}}} > z_{1-\alpha}$$

- But  $\mathbf{s}$  is unknown: one idea (Mogstad et al. 2024)

$$\max_{k \in [d] \setminus \{\mathbf{r}\}} \frac{\bar{X}_{\mathbf{r}} - \bar{X}_k}{\hat{\sigma}_{\mathbf{r}, k}} > c_{1-\alpha}$$

- Hard to calibrate  $c_{1-\alpha}$  in high-dimensions

i.e., Target  $\mu_{\mathbf{r}}$  is greater than the smallest mean  $\mu_{\mathbf{s}}$ ?

# Dimension-Agnostic Argmin Test


💡 Idea: use  $\mathcal{D}_1$  to estimate  $s$  and use  $\mathcal{D}_2$  for inference

## Step I (Sample Splitting)

- Split the data into two

$$\begin{array}{c} \mathcal{D}_1 \\ \vdots \\ \mathcal{D}_2 \end{array} \quad \left[ \begin{array}{c} X_1 \\ X_2 \\ \vdots \\ X_{n-1} \\ X_n \end{array} \right] \quad \left[ \begin{array}{c} X_{n+1} \\ X_{n+2} \\ \vdots \\ X_{2n-1} \\ X_{2n} \end{array} \right]$$

# Dimension-Agnostic Argmin Test




$$s = \arg \min_{k \in [d] \setminus \{r\}} \mu_k$$

**Idea:** use  $\mathcal{D}_1$  to estimate  $s$  and use  $\mathcal{D}_2$  for inference

## Step I (Sample Splitting)

- Split the data into two

$$\mathcal{D}_1 \quad \mathcal{D}_2$$
$$\begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_{n-1} \\ X_n \end{bmatrix} \quad \begin{bmatrix} X_{n+1} \\ X_{n+2} \\ \vdots \\ X_{2n-1} \\ X_{2n} \end{bmatrix}$$



## Step II (Model Selection)

- Based on  $\mathcal{D}_1$ , compute

### I. Plug-in version

$$\hat{s} = \arg \max_{k \in [d] \setminus \{r\}} \bar{X}_{\mathbf{r}}^{(2)} - \bar{X}_k^{(2)}$$

### 2. Noise-adjusted version


$$\hat{s} = \arg \max_{k \in [d] \setminus \{r\}} \frac{\bar{X}_{\mathbf{r}}^{(2)} - \bar{X}_k^{(2)}}{\hat{\sigma}_{\mathbf{r}, k}^{(2)} \vee \kappa}$$

# Dimension-Agnostic Argmin Test

💡 Idea: use  $\mathcal{D}_1$  to estimate  $s$  and use  $\mathcal{D}_2$  for inference

## Step I (Sample Splitting)

- Split the data into two

$$\mathcal{D}_1 \quad \mathcal{D}_2$$
$$\begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_{n-1} \\ X_n \end{bmatrix} \quad \begin{bmatrix} X_{n+1} \\ X_{n+2} \\ \vdots \\ X_{2n-1} \\ X_{2n} \end{bmatrix}$$


## Step II (Model Selection)


- Based on  $\mathcal{D}_1$ , compute

### I. Plug-in version

$$\hat{s} = \arg \max_{k \in [d] \setminus \{\mathbf{r}\}} \bar{X}_{\mathbf{r}}^{(2)} - \bar{X}_k^{(2)}$$

### 2. Noise-adjusted version

$$\hat{s} = \arg \max_{k \in [d] \setminus \{\mathbf{r}\}} \frac{\bar{X}_{\mathbf{r}}^{(2)} - \bar{X}_k^{(2)}}{\hat{\sigma}_{\mathbf{r}, k}^{(2)} \vee \kappa}$$



# Dimension-Agnostic Argmin Test


 **Idea:** use  $\mathcal{D}_1$  to estimate  $s$  and use  $\mathcal{D}_2$  for inference

## Step III (Student's t-statistic)

- Given  $\hat{s} \in [d] \setminus \{r\}$ , compute a one-sided t-statistic

$$T = \frac{\bar{X}_{\color{red}r}^{(1)} - \bar{X}_{\color{blue}\hat{s}}^{(1)}}{\hat{\sigma}_{\color{red}r, \color{blue}\hat{s}}^{(1)}}$$

based on  $\mathcal{D}_2$



# Dimension-Agnostic Argmin Test



 **Idea:** use  $\mathcal{D}_1$  to estimate  $s$  and use  $\mathcal{D}_2$  for inference

## Step III (Student's t-statistic)

- Given  $\hat{s} \in [d] \setminus \{r\}$ , compute a one-sided t-statistic


$$T = \frac{\bar{X}_{\hat{s}}^{(1)} - \bar{X}_r^{(1)}}{\hat{\sigma}_{r,\hat{s}}^{(1)}}$$

based on  $\mathcal{D}_2$



## Step IV (Decision)


- Reject** the null if  $T > z_{1-\alpha}$
- Accept** the null o.w.



$$\Phi(z_{1-\alpha}) = 1 - \alpha$$

# Theoretical Properties

- ▶ I. Asymptotic Validity
- 2. Power Analysis




$X = (X^{(1)}, \dots, X^{(d)})$

# Asymptotic Validity

## Assumption (Truncated 2nd Moment Condition)

Let  $W_k := (X^{(\textcolor{red}{r})} - \mu_{\textcolor{red}{r}}) - (X^{(k)} - \mu_k)$  and




$X = (X^{(1)}, \dots, X^{(d)})$

# Asymptotic Validity

## Assumption (Truncated 2nd Moment Condition)

Let  $W_k := (X^{(\textcolor{red}{r})} - \mu_{\textcolor{red}{r}}) - (X^{(k)} - \mu_k)$  and

$$M_k := \sup_{P \in \mathcal{P}_{0,\textcolor{red}{r}}} \mathbb{E}_P \left[ \frac{W_k^2}{\mathbb{E}_P[W_k^2]} \min \left\{ 1, \frac{|W_k|}{n^{1/2}(\mathbb{E}_P[W_k^2])^{1/2}} \right\} \right]$$



$X = (X^{(1)}, \dots, X^{(d)})$


# Asymptotic Validity

## Assumption (Truncated 2nd Moment Condition)

Let  $W_k := (X^{(\textcolor{red}{r})} - \mu_{\textcolor{red}{r}}) - (X^{(k)} - \mu_k)$  and

$$M_k := \sup_{P \in \mathcal{P}_{0,\textcolor{red}{r}}} \mathbb{E}_P \left[ \frac{W_k^2}{\mathbb{E}_P[W_k^2]} \min \left\{ 1, \frac{|W_k|}{n^{1/2}(\mathbb{E}_P[W_k^2])^{1/2}} \right\} \right]$$

Assume that  $\max_{k \in [d] \setminus \{\textcolor{red}{r}\}} M_k = o(1)$



# Asymptotic Validity


## Assumption (Truncated 2nd Moment Condition)

Let  $W_k := (X^{(\textcolor{red}{r})} - \mu_{\textcolor{red}{r}}) - (X^{(k)} - \mu_k)$  and

$$M_k := \sup_{P \in \mathcal{P}_{0,\textcolor{red}{r}}} \mathbb{E}_P \left[ \frac{W_k^2}{\mathbb{E}_P[W_k^2]} \min \left\{ 1, \frac{|W_k|}{n^{1/2}(\mathbb{E}_P[W_k^2])^{1/2}} \right\} \right]$$

Assume that  $\max_{k \in [d] \setminus \{\textcolor{red}{r}\}} M_k = o(1)$

- **Equivalent** to Lindeberg's condition (weaker than Lyapunov)



# Asymptotic Validity


## Assumption (Truncated 2nd Moment Condition)

Let  $W_k := (X^{(\textcolor{red}{r})} - \mu_{\textcolor{red}{r}}) - (X^{(k)} - \mu_k)$  and

$$M_k := \sup_{P \in \mathcal{P}_{0,\textcolor{red}{r}}} \mathbb{E}_P \left[ \frac{W_k^2}{\mathbb{E}_P[W_k^2]} \min \left\{ 1, \frac{|W_k|}{n^{1/2}(\mathbb{E}_P[W_k^2])^{1/2}} \right\} \right]$$

Assume that  $\max_{k \in [d] \setminus \{\textcolor{red}{r}\}} M_k = o(1)$

- **Equivalent** to Lindeberg's condition (weaker than Lyapunov)
- If  $X \sim N(\mu, \Sigma)$ , it only requires  $\text{Var}[W_k] > 0$



$X = (X^{(1)}, \dots, X^{(d)})$

# Asymptotic Validity

## Assumption (Truncated 2nd Moment Condition)

Let  $W_k := (X^{(\textcolor{red}{r})} - \mu_{\textcolor{red}{r}}) - (X^{(k)} - \mu_k)$  and

$$M_k := \sup_{P \in \mathcal{P}_{0,\textcolor{red}{r}}} \mathbb{E}_P \left[ \frac{W_k^2}{\mathbb{E}_P[W_k^2]} \min \left\{ 1, \frac{|W_k|}{n^{1/2}(\mathbb{E}_P[W_k^2])^{1/2}} \right\} \right]$$

Assume that  $\max_{k \in [d] \setminus \{\textcolor{red}{r}\}} M_k = o(1)$

- **Equivalent** to Lindeberg's condition (weaker than Lyapunov)
- If  $X \sim N(\mu, \Sigma)$ , it only requires  $\text{Var}[W_k] > 0$
- **Arbitrary dependence** among the components except  $X^{(\textcolor{red}{r})}$

# Asymptotic Validity

$$M_k := \sup_{P \in \mathcal{P}_{0,r}} \mathbb{E}_P \left[ \frac{W_k^2}{\mathbb{E}_P[W_k^2]} \min \left\{ 1, \frac{|W_k|}{n^{1/2}(\mathbb{E}_P[W_k^2])^{1/2}} \right\} \right]$$

Assume that  $\max_{k \in [d] \setminus \{r\}} M_k = o(1)$

**Theorem** Kim and Ramdas (2025)

Let  $\mathcal{P}_{0,r}$  be the class of null distributions with  $H_0 : r \in \Theta$  that satisfy the **truncated 2nd moment condition**. Then the DA test is asymptotically valid as

$$\limsup_{n \rightarrow \infty} \sup_{P \in \mathcal{P}_{0,r}} P(\psi_r = 1) \leq \alpha, \text{ regardless of the sequence } (d_n)_{n=1}^\infty$$

# Theoretical Properties

- I. Asymptotic Validity
- ▶ 2. Power Analysis

# Confusion Set

- The problem **difficulty** depends on the cardinality of a **confusion set**  $\mathbb{C}_r$

$$\mathbb{C}_r := \left\{ k \in [d] \setminus \{\textcolor{red}{r}\} : \frac{\mu_{\textcolor{red}{r}} - \mu_\star}{2} \leq \mu_k - \mu_\star \leq C_n \sqrt{\frac{\log(d)}{n}} \right\}$$

# Confusion Set

- The problem **difficulty** depends on the cardinality of a **confusion set**  $\mathbb{C}_r$

$$\mathbb{C}_r := \left\{ k \in [d] \setminus \{\textcolor{red}{r}\} : \frac{\mu_{\textcolor{red}{r}} - \mu_\star}{2} \leq \mu_k - \mu_\star \leq \textcolor{red}{C}_n \sqrt{\frac{\log(d)}{n}} \right\}$$

$C_n$  : any increasing sequence

# Confusion Set

$$\mu_{\star} = \min_{i \in [d]} \mu_i$$

- The problem **difficulty** depends on the cardinality of a **confusion set**  $\mathbb{C}_r$

$$\mathbb{C}_r := \left\{ k \in [d] \setminus \{\textcolor{red}{r}\} : \frac{\mu_{\textcolor{red}{r}} - \mu_{\star}}{2} \leq \mu_k - \mu_{\star} \leq C_n \sqrt{\frac{\log(d)}{n}} \right\}$$

$\mathcal{A}$

- Intuition for  $\mathcal{A}^c$

$C_n$  : any increasing sequence

When  $\mu_k$  is far from  $\mu_{\star}$ , it is unlikely that  $\hat{s} = k$   
 $\therefore$  Well-separated set

# Confusion Set

$$\mu_{\star} = \min_{i \in [d]} \mu_i$$

- The problem **difficulty** depends on the cardinality of a **confusion set**  $\mathbb{C}_r$

$$\mathbb{C}_r := \left\{ k \in [d] \setminus \{\textcolor{red}{r}\} : \frac{\mu_{\textcolor{red}{r}} - \mu_{\star}}{2} \leq \mu_k - \mu_{\star} \leq C_n \sqrt{\frac{\log(d)}{n}} \right\}$$

- **Intuition for  $\mathcal{A}^c$**

$C_n$  : any increasing sequence

When  $\mu_k$  is far from  $\mu_{\star}$ , it is unlikely that  $\hat{s} = k$   
 $\therefore$  Well-separated set

- **Intuition for  $\mathcal{B}^c$**

$$\frac{\mu_{\textcolor{red}{r}} - \mu_{\star}}{2} > \mu_k - \mu_{\star} \iff \mu_{\textcolor{red}{r}} - \mu_k > \frac{\mu_{\textcolor{red}{r}} - \mu_{\star}}{2}$$

$\therefore$  Comparable signal

# Power Analysis

- Let  $\mathcal{P}$  be a collection of **sub-Gaussian** distributions with  $\sigma^2$

# Power Analysis

- Let  $\mathcal{P}$  be a collection of **sub-Gaussian** distributions with  $\sigma^2$
- Define a class of **alternative** distributions

$$\mathcal{P}_{1,\textcolor{red}{r}}(\varepsilon; \tau) := \left\{ P \in \mathcal{P} : \mu_{\textcolor{red}{r}} - \mu_{\star} \geq \varepsilon \text{ and } |\mathbb{C}_{\textcolor{red}{r}}| = \tau \right\}$$

# Power Analysis

- Let  $\mathcal{P}$  be a collection of **sub-Gaussian** distributions with  $\sigma^2$
- Define a class of **alternative** distributions

$$\mathcal{P}_{1,\textcolor{red}{r}}(\varepsilon; \tau) := \left\{ P \in \mathcal{P} : \mu_{\textcolor{red}{r}} - \mu_{\star} \geq \varepsilon \text{ and } |\mathbb{C}_{\textcolor{red}{r}}| = \tau \right\}$$

- The **critical radius**  $\varepsilon^{\star}$  is defined as

$$\varepsilon^{\star} = \varepsilon^{\star}(\tau) = \sqrt{\frac{1 \vee \log(\tau)}{n}}$$

# Power Analysis

**Theorem** Kim and Ramdas (2025)

For any  $\tau$ , suppose that  $\varepsilon \gg \varepsilon^*$ . Then the asymptotic uniform power of the DA test is equal to one:

$$\lim_{n \rightarrow \infty} \inf_{P \in \mathcal{P}_{1,r}(\varepsilon; \tau)} P(\psi_r = 1) = 1$$



$$\varepsilon^* = \sqrt{\frac{1 \vee \log(\tau)}{n}}$$

# Power Analysis

**Theorem** Kim and Ramdas (2025)

For any  $\tau$ , suppose that  $\varepsilon \gg \varepsilon^*$ . Then the asymptotic uniform power of the DA test is equal to one:

$$\lim_{n \rightarrow \infty} \inf_{P \in \mathcal{P}_{1,r}(\varepsilon; \tau)} P(\psi_r = 1) = 1$$

- Adaptivity to unknown  $|\mathbb{C}_r|$
- The rate changes from  $1/\sqrt{n}$ -rate to  $\sqrt{\log(d)/n}$



$$\varepsilon^* = \sqrt{\frac{1 \vee \log(\tau)}{n}}$$

# Power Analysis

**Theorem** Kim and Ramdas (2025)

For any  $\tau$ , suppose that  $\varepsilon \gg \varepsilon^*$ . Then the asymptotic uniform power of the DA test is equal to one:

$$\lim_{n \rightarrow \infty} \inf_{P \in \mathcal{P}_{1,r}(\varepsilon; \tau)} P(\psi_r = 1) = 1$$

- Adaptivity to unknown  $|\mathbb{C}_r|$
- The rate changes from  $1/\sqrt{n}$ -rate to  $\sqrt{\log(d)/n}$

**Question.** Can we further **improve** the **separation rate**?



$$\varepsilon^* = \sqrt{\frac{1 \vee \log(\tau)}{n}}$$

# Local Minimax Optimality

**Theorem** Kim and Ramdas (2025)

Let  $\Psi_\alpha$  be the set of all asymptotic level- $\alpha$  tests over  $\mathcal{P}_{0,\textcolor{red}{r}}$ ,

$$\Psi_{\textcolor{teal}{\alpha}} := \left\{ \psi : \limsup_{n \rightarrow \infty} \sup_{\textcolor{blue}{P} \in \mathcal{P}_{0,\textcolor{red}{r}}} \textcolor{blue}{P}(\psi = 1) \leq \alpha \right\}$$

# Local Minimax Optimality

**Theorem** Kim and Ramdas (2025)

Let  $\Psi_\alpha$  be the set of all asymptotic level- $\alpha$  tests over  $\mathcal{P}_{0,\textcolor{red}{r}}$ ,

$$\Psi_\alpha := \left\{ \psi : \limsup_{n \rightarrow \infty} \sup_{P \in \mathcal{P}_{0,\textcolor{red}{r}}} P(\psi = 1) \leq \alpha \right\}$$

If  $\varepsilon \ll \varepsilon^*$ , then the **asymptotic type II error** is at least  $\beta$ :

$$\liminf_{n \rightarrow \infty} \inf_{\psi \in \Psi_\alpha} \sup_{P \in \mathcal{P}_{1,\textcolor{red}{r}}(\varepsilon; \tau)} P(\psi = 0) \geq \beta$$



$$\varepsilon^* = \sqrt{\frac{1 \vee \log(\tau)}{n}}$$

# Summary

- We have introduced a DA method for **high-dimensional argmin inference** problem based on **sample splitting** and **studentization**
- The proposed method achieves the **locally minimax separation rate** and adapts to the intrinsic difficulty of the problem characterized by the **confusion set**
- We have demonstrated its **strong empirical performance** under various settings

Thank you!



# Coverage Guarantees

## I. Pointwise coverage

Each  $r \in \Theta$  is included in  $\widehat{\Theta}$  with high probability:

$$\inf_{P \in \mathcal{P}} \inf_{r \in \Theta(P)} P(r \in \widehat{\Theta}) \geq 1 - \alpha$$

- **Less** demanding → **smaller** expected set size
- **Higher** power but **weaker** protection

## 2. Uniform coverage

The entire set  $\Theta$  is contained in  $\widehat{\Theta}$  with high probability:

$$\inf_{P \in \mathcal{P}} P(\Theta \subseteq \widehat{\Theta}) \geq 1 - \alpha$$

- **More** demanding → **larger** expected set size
- **Stronger** protection but **more** conservative



$\mathcal{P}$ : class of distributions  
 $\alpha$ : target error level

# Related Work

- **Classical work:** Bechhofer (1954), Gupta (1956, 1965), Futschik and Pflug (1995) etc  
→ rely on parametric models, independence between coordinates, absence of ties

# Related Work

- **Classical work:** Bechhofer (1954), Gupta (1956, 1965), Futschik and Pflug (1995) etc  
→ rely on parametric models, independence between coordinates, absence of ties
- **Hansen et al. (2011):** propose a sequential procedure for MCS with uniform coverage  
→ widely cited (2550+), produce wide sets; extremely slow to run

# Related Work



- **Classical work:** Bechhofer (1954), Gupta (1956, 1965), Futschik and Pflug (1995) etc  
→ rely on parametric models, independence between coordinates, absence of ties
- **Hansen et al. (2011):** propose a sequential procedure for MCS with uniform coverage  
→ widely cited (2550+), produce wide sets; extremely slow to run
- **Mogstad et al. (2024):** introduce a bootstrap approach for rank inference (can be tweaked)  
→ efficiency not analyzed; results limited to fixed dimensional settings

# Related Work



- **Classical work:** Bechhofer (1954), Gupta (1956, 1965), Futschik and Pflug (1995) etc  
→ rely on parametric models, independence between coordinates, absence of ties
- **Hansen et al. (2011):** propose a sequential procedure for MCS with uniform coverage  
→ widely cited (2550+), produce wide sets; extremely slow to run
- **Mogstad et al. (2024):** introduce a bootstrap approach for rank inference (can be tweaked)  
→ efficiency not analyzed; results limited to fixed dimensional settings
- **Zhang et al. (2024):** introduce a cross-validation + privacy approach for argmin inference  
→ require careful tuning and fall short of minimax optimality

# Confusion Set

- Set  $\mu_r = \mu_1$  and  $\mu_\star = \mu_2$





# Local Minimax Optimality



# Empirical Results

# Power and Validity in High-Dimensional Settings



- **L0O:** Zhang et al. (2024)
- **Bonferroni:** multiple correction
- **csranks:** Mogstad et al. (2024)
- **DA-plug:** plug-in  $\hat{s}$
- **DA-adj:** noise-adjusted  $\hat{s}$

## • Null Setting

$$\mu = (0, 0, 1, \dots, 1)^\top$$
$$\Sigma_{11} = \Sigma_{22} = 1 \text{ & } \Sigma_{33} = \dots = \Sigma_{dd} = 20$$

## • Alternative Setting

$$\mu = (0.15, 0, 1, \dots, 1)^\top$$
$$\Sigma_{11} = \Sigma_{22} = 1 \text{ & } \Sigma_{33} = \dots = \Sigma_{dd} = 20$$


# Power under Various Settings

| Method                   | $\mu^{(a)} + \text{unequal variance}$ |              |              | $\mu^{(b)} + \text{unequal variance}$ |              |              | $\mu^{(c)} + \text{unequal variance}$ |              |              |
|--------------------------|---------------------------------------|--------------|--------------|---------------------------------------|--------------|--------------|---------------------------------------|--------------|--------------|
|                          | $\rho = 0$                            | $\rho = 0.4$ | $\rho = 0.8$ | $\rho = 0$                            | $\rho = 0.4$ | $\rho = 0.8$ | $\rho = 0$                            | $\rho = 0.4$ | $\rho = 0.8$ |
| L00                      | 0.084                                 | 0.115        | 0.380        | 0.000                                 | 0.001        | 0.181        | <b>0.258</b>                          | <b>0.351</b> | <b>0.703</b> |
| Bonferroni               | 0.171                                 | 0.130        | 0.055        | 0.166                                 | 0.103        | 0.030        | 0.017                                 | 0.006        | 0.003        |
| csranks                  | <b>0.184</b>                          | <b>0.381</b> | <b>0.962</b> | 0.162                                 | 0.363        | 0.961        | 0.019                                 | 0.041        | 0.223        |
| MCS                      | 0.004                                 | 0.002        | 0.004        | 0.000                                 | 0.000        | 0.000        | 0.140                                 | 0.156        | 0.166        |
| DA-plug                  | 0.049                                 | 0.052        | 0.042        | 0.062                                 | 0.067        | 0.059        | 0.098                                 | 0.128        | 0.202        |
| DA-plug $^{ \times 10 }$ | 0.050                                 | 0.052        | 0.050        | 0.080                                 | 0.080        | 0.073        | 0.125                                 | 0.145        | 0.240        |
| DA-adj                   | 0.122                                 | 0.259        | 0.841        | 0.217                                 | 0.384        | 0.916        | 0.135                                 | 0.188        | 0.462        |
| DA-adj $^{ \times 10 }$  | 0.160                                 | 0.343        | 0.946        | <b>0.294</b>                          | <b>0.517</b> | <b>0.982</b> | 0.164                                 | 0.251        | 0.605        |

# Power under Various Settings

| Method                 | $\mu^{(a)}$ + unequal variance |              |              | $\mu^{(b)}$ + unequal variance |              |              | $\mu^{(c)}$ + unequal variance |              |              |
|------------------------|--------------------------------|--------------|--------------|--------------------------------|--------------|--------------|--------------------------------|--------------|--------------|
|                        | $\rho = 0$                     | $\rho = 0.4$ | $\rho = 0.8$ | $\rho = 0$                     | $\rho = 0.4$ | $\rho = 0.8$ | $\rho = 0$                     | $\rho = 0.4$ | $\rho = 0.8$ |
| L0O                    | 0.084                          | 0.115        | 0.380        | 0.000                          | 0.001        | 0.181        | <b>0.258</b>                   | <b>0.351</b> | <b>0.703</b> |
| Bonferroni             | 0.171                          | 0.130        | 0.055        | 0.166                          | 0.103        | 0.030        | 0.017                          | 0.006        | 0.003        |
| csranks                | <b>0.184</b>                   | <b>0.381</b> | <b>0.962</b> | 0.162                          | 0.363        | <b>0.961</b> | 0.019                          | 0.041        | 0.223        |
| MCS                    | 0.004                          | 0.002        | 0.004        | 0.000                          | 0.000        | 0.000        | 0.140                          | 0.156        | 0.166        |
| DA-plug                | 0.049                          | 0.052        | 0.042        | 0.062                          | 0.067        | 0.059        | 0.098                          | 0.128        | 0.202        |
| DA-plug $^{\times 10}$ | 0.050                          | 0.052        | 0.050        | 0.080                          | 0.080        | 0.073        | 0.125                          | 0.145        | 0.240        |
| DA-adj                 | 0.122                          | 0.259        | 0.841        | 0.217                          | 0.384        | 0.916        | 0.135                          | 0.188        | 0.462        |
| DA-adj $^{\times 10}$  | 0.160                          | 0.343        | 0.946        | <b>0.294</b>                   | <b>0.517</b> | <b>0.982</b> | 0.164                          | 0.251        | 0.605        |

- DA-plug: plug-in  $\hat{s}$
- DA-adj: noise-adjusted  $\hat{s}$
- DA-plug $^{\times 10}$ : 10 splits + average
- DA-adj $^{\times 10}$ : 10 splits + average



1. **Multiple splits** improve the power
2. **Noise-adjusted** version performs better

# Power under Various Settings

| Method                  | $\mu^{(a)}$ + unequal variance |              |              | $\mu^{(b)}$ + unequal variance |              |              | $\mu^{(c)}$ + unequal variance |              |              |
|-------------------------|--------------------------------|--------------|--------------|--------------------------------|--------------|--------------|--------------------------------|--------------|--------------|
|                         | $\rho = 0$                     | $\rho = 0.4$ | $\rho = 0.8$ | $\rho = 0$                     | $\rho = 0.4$ | $\rho = 0.8$ | $\rho = 0$                     | $\rho = 0.4$ | $\rho = 0.8$ |
| L00                     | 0.084                          | 0.115        | 0.380        | 0.000                          | 0.001        | 0.181        | <b>0.258</b>                   | <b>0.351</b> | <b>0.703</b> |
| Bonferroni              | 0.171                          | 0.130        | 0.055        | 0.166                          | 0.103        | 0.030        | 0.017                          | 0.006        | 0.003        |
| csranks                 | <b>0.184</b>                   | <b>0.381</b> | <b>0.962</b> | 0.162                          | 0.363        | 0.961        | 0.019                          | 0.041        | 0.223        |
| MCS                     | 0.004                          | 0.002        | 0.004        | 0.000                          | 0.000        | 0.000        | 0.140                          | 0.156        | 0.166        |
| DA-plug                 | 0.049                          | 0.052        | 0.042        | 0.062                          | 0.067        | 0.059        | 0.098                          | 0.128        | 0.202        |
| DA-plug $^{ \times 10}$ | 0.050                          | 0.052        | 0.050        | 0.080                          | 0.080        | 0.073        | 0.125                          | 0.145        | 0.240        |
| DA-adj                  | 0.122                          | 0.259        | 0.841        | 0.217                          | 0.384        | 0.916        | 0.135                          | 0.188        | 0.462        |
| DA-adj $^{ \times 10}$  | 0.160                          | 0.343        | 0.946        | <b>0.294</b>                   | <b>0.517</b> | <b>0.982</b> | 0.164                          | 0.251        | 0.605        |

DA-adj $^{ \times 10}$  achieves the **best or second-best** performance  
 (with only a **small** margin)

## Mean Structures

$$\mu^{(a)} = (0.1, 0, 0.1, \dots, 0.1)^\top \quad \mu^{(b)} = (0.1, 0.019, \dots, 0.99, 1)^\top$$

# Power under Various Settings

| Method                   | $\mu^{(a)} + \text{unequal variance}$ |              |              | $\mu^{(b)} + \text{unequal variance}$ |              |              | $\mu^{(c)} + \text{unequal variance}$ |              |              |
|--------------------------|---------------------------------------|--------------|--------------|---------------------------------------|--------------|--------------|---------------------------------------|--------------|--------------|
|                          | $\rho = 0$                            | $\rho = 0.4$ | $\rho = 0.8$ | $\rho = 0$                            | $\rho = 0.4$ | $\rho = 0.8$ | $\rho = 0$                            | $\rho = 0.4$ | $\rho = 0.8$ |
| LOO                      | 0.084                                 | 0.115        | 0.380        | 0.000                                 | 0.001        | 0.181        | <b>0.258</b>                          | <b>0.351</b> | <b>0.703</b> |
| Bonferroni               | 0.171                                 | 0.130        | 0.055        | 0.166                                 | 0.103        | 0.030        | 0.017                                 | 0.006        | 0.003        |
| csranks                  | <b>0.184</b>                          | <b>0.381</b> | <b>0.962</b> | 0.162                                 | 0.363        | 0.961        | 0.019                                 | 0.041        | 0.223        |
| MCS                      | 0.004                                 | 0.002        | 0.004        | 0.000                                 | 0.000        | 0.000        | 0.140                                 | 0.156        | 0.166        |
| DA-plug                  | 0.049                                 | 0.052        | 0.042        | 0.062                                 | 0.067        | 0.059        | 0.098                                 | 0.128        | 0.202        |
| DA-plug $^{ \times 10 }$ | 0.050                                 | 0.052        | 0.050        | 0.080                                 | 0.080        | 0.073        | 0.125                                 | 0.145        | 0.240        |
| DA-adj                   | 0.122                                 | 0.259        | 0.841        | 0.217                                 | 0.384        | 0.916        | 0.135                                 | 0.188        | 0.462        |
| DA-adj $^{ \times 10 }$  | 0.160                                 | 0.343        | 0.946        | <b>0.294</b>                          | <b>0.517</b> | <b>0.982</b> | 0.164                                 | 0.251        | 0.605        |

LOO performs **best** and DA-adj $^{ \times 10 }$  performs **second best**  
 But LOO **does not** control the type I error rate

## Mean Structure

$$\boldsymbol{\mu}^{(c)} = (0.05, 0, 0, 0, 10, \dots, 10)^\top$$

# Power under Various Settings

| Method                  | $\mu^{(a)}$ + unequal variance |              |              | $\mu^{(b)}$ + unequal variance |              |              | $\mu^{(c,0)}$ + unequal variance |              |              |
|-------------------------|--------------------------------|--------------|--------------|--------------------------------|--------------|--------------|----------------------------------|--------------|--------------|
|                         | $\rho = 0$                     | $\rho = 0.4$ | $\rho = 0.8$ | $\rho = 0$                     | $\rho = 0.4$ | $\rho = 0.8$ | $\rho = 0$                       | $\rho = 0.4$ | $\rho = 0.8$ |
| LOO                     | 0.084                          | 0.115        | 0.380        | 0.000                          | 0.001        | 0.181        | 0.070                            | 0.064        | 0.065        |
| Bonferroni              | 0.171                          | 0.130        | 0.055        | 0.166                          | 0.103        | 0.030        | 0.002                            | 0.001        | 0.001        |
| csranks                 | 0.184                          | 0.381        | 0.962        | 0.162                          | 0.363        | 0.961        | 0.002                            | 0.001        | 0.002        |
| MCS                     | 0.004                          | 0.002        | 0.004        | 0.000                          | 0.000        | 0.000        | 0.042                            | 0.042        | 0.034        |
| DA-plug                 | 0.049                          | 0.052        | 0.042        | 0.062                          | 0.067        | 0.059        | 0.048                            | 0.052        | 0.048        |
| DA-plug $^{ \times 10}$ | 0.050                          | 0.052        | 0.050        | 0.080                          | 0.080        | 0.073        | 0.053                            | 0.046        | 0.047        |
| DA-adj                  | 0.122                          | 0.259        | 0.841        | 0.217                          | 0.384        | 0.916        | 0.054                            | 0.052        | 0.050        |
| DA-adj $^{ \times 10}$  | 0.160                          | 0.343        | 0.946        | 0.294                          | 0.517        | 0.982        | 0.053                            | 0.049        | 0.050        |

LOO performs **best** and DA-adj $^{ \times 10}$  performs **second best**  
 But LOO **does not** control the type I error rate


## Mean Structure

$$\boldsymbol{\mu}^{(c)} = (0.05, 0, 0, 0, 10, \dots, 10)^\top$$

# Real-World Data

- We revisit **CMU 36-462 prediction competition** analyzed by Zhang et al. (2024)
  - Spring 2023:  $n = 183, p = 44$
  - Spring 2024:  $n = 1246, p = 39$
- $X_{i,k} \in \{0,1\}$ : correct/incorrect prediction by team  $k$  at test point  $i$
- $\mu_k$ : the population prediction risk of team  $k$  (unknown)
- Methods to compare
  - Bonferroni correction (**Bonf**)
  - Hansen et al. (2011) (**MCS**)
  - Mogstad et al. (2024) (**CSR**)
  - Zhang et al. (2024) (**LOO**)

# Real-World Data



- **Colored points** indicate the indices included in **each confidence set**
- Our method (**DA-adj $\times 10$** ) produces confidence sets with **smallest cardinality**

# Future directions

- General **rank-k** inference
- Confidence sets for  $\mu_{(k)}$
- **Computationally efficient** multiple splitting approach