

# Local Permutation Tests for Conditional Independence

Ilmun Kim


Department of Statistics & Data Science  
Yonsei University




# Joint work with



Matey Neykov  
(Northwestern)



Sivaraman  
Balakrishnan  
(CMU)



Larry Wasserman  
(CMU)

# Conditional Independence (CI) Testing

- Suppose that we observe  $\{(X_i, Y_i, Z_i)\}_{i=1}^n \stackrel{i.i.d.}{\sim} P_{X,Y,Z}$
- We would like to **determine** whether

$$H_0 : X \perp\!\!\!\perp Y | Z \quad \text{versus} \quad H_1 : X \not\perp\!\!\!\perp Y | Z$$

$$(P_{X,Y,Z} = P_{X|Z}P_{Y|Z}P_Z) \quad (P_{X,Y,Z} \neq P_{X|Z}P_{Y|Z}P_Z)$$

# Applications: Variable Importance

Is the predictor  $X$  important to predict the response variable  $Y$ ?

# Applications: Variable Importance

Is the predictor  $X$  important to predict the response variable  $Y$ ?

## I. Parametric methods

$$Y = \beta X + \gamma Z + \varepsilon$$

- e.g., F-test
- $H_0 : \beta = 0$  vs.  $H_1 : \beta \neq 0$
- Not meaningful when applied beyond linear models

# Applications: Variable Importance

Is the predictor  $X$  important to predict the response variable  $Y$ ?

## I. Parametric methods

$$Y = \beta X + \gamma Z + \epsilon$$

- e.g., F-test

$$H_0 : \beta = 0 \text{ vs. } H_1 : \beta \neq 0$$

- Not meaningful when applied beyond linear models

## 2. Blackbox methods

- Feature importance from e.g., random forests and XGboost
- Model dependent & does not present uncertainty

# Applications: Variable Importance

Is the predictor  $X$  important to predict the response variable  $Y$ ?

## I. Parametric methods

$$Y = \beta X + \gamma Z + \epsilon$$

- e.g., F-test

$$H_0 : \beta = 0 \text{ vs. } H_1 : \beta \neq 0$$

- Not meaningful when applied beyond linear models

## 2. Blackbox methods

- Feature importance from e.g., random forests and XGboost
- Model dependent & does not present uncertainty

We can formulate it as a **hypothesis testing problem!**

$$H_0 : Y \perp\!\!\!\perp X | Z \quad \text{versus} \quad H_1 : Y \not\perp\!\!\!\perp X | Z$$

# Prior methods

## Asymptotic method

Su and White (2008)

Zhang et al. (2012)

Huang (2010)

Wang et al. (2015)

Strobl et al. (2019)

Zhou et al. (2020)

Shah and Peters (2020)

Huang et al. (2022)

Lundborg et al. (2024)


⋮

# Prior methods

## Asymptotic method

Su and White (2008)  
Zhang et al. (2012)  
Huang (2010)  
Wang et al. (2015)  
Strobl et al. (2019)  
Zhou et al. (2020)  
Shah and Peters (2020)  
Huang et al. (2022)  
Lundborg et al. (2024)

⋮

- Pivotal Statistic  $\xrightarrow{d}$    $N(0,1)$
- Determine the **critical value** based on the **limiting distribution**

# Prior methods

## Asymptotic method

- Su and White (2008)
- Zhang et al. (2012)
- Huang (2010)
- Wang et al. (2015)
- Strobl et al. (2019)
- Zhou et al. (2020)
- Shah and Peters (2020)
- Huang et al. (2022)
- Lundborg et al. (2024)
- ⋮

## Model-X framework

- Candés et al. (2018)
- Berrett et al. (2020)
- Bellot and van der Schaar (2019)
- Katsevish and Ramdas (2022)
- Watson and Wright (2021)
- Liu et al. (2022)
- ⋮

# Prior methods

## Asymptotic method

Su and White (2008)  
Zhang et al. (2012)  
Huang (2010)  
Wang et al. (2015)  
Strobl et al. (2019)  
Zhou et al. (2020)  
Shah and Peters (2020)  
Huang et al. (2022)  
Lundborg et al. (2024)  
⋮

## Model-X framework

Candés et al. (2018)  
Berrett et al. (2020)  
Bellot and van der  
Schaar (2019)  
Katsevish and Ramdas  
(2022)  
Watson and Wright  
(2021)  
Liu et al. (2022)  
⋮

- Assume  $F_{X|Z}$  is known
- Generate  $\tilde{X}$  from  $F_{X|Z}$
- Construct an exchangeable sequence under the null

$$T, T_1, \dots, T_B$$

# Prior methods

## Asymptotic method

- Su and White (2008)
- Zhang et al. (2012)
- Huang (2010)
- Wang et al. (2015)
- Strobl et al. (2019)
- Zhou et al. (2020)
- Shah and Peters (2020)
- Huang et al. (2022)
- Lundborg et al. (2024)
- ⋮

## Model-X framework

- Candés et al. (2018)
- Berrett et al. (2020)
- Bellot and van der Schaar (2019)
- Katsevish and Ramdas (2022)
- Watson and Wright (2021)
- Liu et al. (2022)
- ⋮

## Local permutation

- Margaritis (2005)
- Fukumizu et al. (2008)
- Doran et al. (2014)
- Sen et al. (2017)
- Neykov et al. (2021)
- ⋮

# Prior methods

## Asymptotic method

- Su and White (2008)
- Zhang et al. (2012)
- Huang (2010)
- Wang et al. (2015)
- Strobl et al. (2019)
- Zhou et al. (2020)
- Shah and Peters (2020)
- Huang et al. (2022)
- Lundborg et al. (2024)
- ⋮

## Model-X framework

- Candés et al. (2018)
- Berrett et al. (2020)
- Bellot and van der Schaar (2019)
- Katsevish and Ramdas (2022)
- Watson and Wright (2021)
- Liu et al. (2022)
- ⋮

## Local permutation

- Margaritis (2005)
- Fukumizu et al. (2008)
- Doran et al. (2014)
- Sen et al. (2017)
- Neykov et al. (2021)
- ⋮

We aim to enhance our understanding of  
**local permutation tests:**

*“when does it work and when does it fail?”*

# Local Permutation Tests


- **Partition** the domain of  $Z$  into  $B_1, \dots, B_M$  bins

# Local Permutation Tests

- **Partition** the domain of  $Z$  into  $B_1, \dots, B_M$  bins
- **Transform**  $\{(X_i, Y_i, Z_i)\}_{i=1}^n \mapsto \{(X_i, Y_i, \tilde{Z}_i)\}_{i=1}^n$  where  $\tilde{Z}_i = k$  if  $Z_i \in B_k$


# Local Permutation Tests

- **Partition** the domain of  $Z$  into  $B_1, \dots, B_M$  bins
- **Transform**  $\{(X_i, Y_i, Z_i)\}_{i=1}^n \mapsto \{(X_i, Y_i, \tilde{Z}_i)\}_{i=1}^n$  where  $\tilde{Z}_i = k$  if  $Z_i \in B_k$




# Local Permutation Tests

- **Partition** the domain of  $Z$  into  $B_1, \dots, B_M$  bins
- **Transform**  $\{(X_i, Y_i, Z_i)\}_{i=1}^n \mapsto \{(X_i, Y_i, \tilde{Z}_i)\}_{i=1}^n$  where  $\tilde{Z}_i = k$  if  $Z_i \in B_k$




# Local Permutation Tests

- **Partition** the domain of  $Z$  into  $B_1, \dots, B_M$  bins
- **Transform**  $\{(X_i, Y_i, Z_i)\}_{i=1}^n \mapsto \{(X_i, Y_i, \tilde{Z}_i)\}_{i=1}^n$  where  $\tilde{Z}_i = k$  if  $Z_i \in B_k$




# Local Permutation Tests

- Partition the domain of  $Z$  into  $B_1, \dots, B_M$  bins
- Transform  $\{(X_i, Y_i, Z_i)\}_{i=1}^n \mapsto \{(X_i, Y_i, \tilde{Z}_i)\}_{i=1}^n$  where  $\tilde{Z}_i = k$  if  $Z_i \in B_k$




# Local Permutation Tests

- Partition the domain of  $Z$  into  $B_1, \dots, B_M$  bins
- Transform  $\{(X_i, Y_i, Z_i)\}_{i=1}^n \mapsto \{(X_i, Y_i, \tilde{Z}_i)\}_{i=1}^n$  where  $\tilde{Z}_i = k$  if  $Z_i \in B_k$




# Local Permutation Tests

- Partition the domain of  $Z$  into  $B_1, \dots, B_M$  bins
- Transform  $\{(X_i, Y_i, Z_i)\}_{i=1}^n \mapsto \{(X_i, Y_i, \tilde{Z}_i)\}_{i=1}^n$  where  $\tilde{Z}_i = k$  if  $Z_i \in B_k$



# Local Permutation Tests

- Partition the domain of  $Z$  into  $B_1, \dots, B_M$  bins
- Transform  $\{(X_i, Y_i, Z_i)\}_{i=1}^n \mapsto \{(X_i, Y_i, \tilde{Z}_i)\}_{i=1}^n$  where  $\tilde{Z}_i = k$  if  $Z_i \in B_k$



# Local Permutation Tests

- **Partition** the domain of  $Z$  into  $B_1, \dots, B_M$  bins
- **Transform**  $\{(X_i, Y_i, Z_i)\}_{i=1}^n \mapsto \{(X_i, Y_i, \tilde{Z}_i)\}_{i=1}^n$  where  $\tilde{Z}_i = k$  if  $Z_i \in B_k$

$\tilde{Z} = 1$

$$\begin{aligned}(X_{1,1}, Y_{1,1}) \\ (X_{2,1}, Y_{2,1}) \\ \vdots \\ (X_{\sigma_1,1}, Y_{\sigma_1,1})\end{aligned}$$

$\tilde{Z} = 2$

$$\begin{aligned}(X_{1,2}, Y_{1,2}) \\ (X_{2,2}, Y_{2,2}) \\ \vdots \\ (X_{\sigma_2,2}, Y_{\sigma_2,2})\end{aligned}$$

$\tilde{Z} = 3$

$$\begin{aligned}(X_{1,3}, Y_{1,3}) \\ (X_{2,3}, Y_{2,3}) \\ \vdots \\ (X_{\sigma_3,3}, Y_{\sigma_3,3})\end{aligned}$$


⋮ ⋮ ⋮

$\tilde{Z} = M$

$$\begin{aligned}(X_{1,M}, Y_{1,M}) \\ (X_{2,M}, Y_{2,M}) \\ \vdots \\ (X_{\sigma_M,M}, Y_{\sigma_M,M})\end{aligned}$$


# Local Permutation Tests

- **Partition** the domain of  $Z$  into  $B_1, \dots, B_M$  bins
- **Transform**  $\{(X_i, Y_i, Z_i)\}_{i=1}^n \mapsto \{(X_i, Y_i, \tilde{Z}_i)\}_{i=1}^n$  where  $\tilde{Z}_i = k$  if  $Z_i \in B_k$



# Local Permutation Tests


- Compute a binning-based test statistic  $T := T(\mathcal{D}_1, \mathcal{D}_2, \dots, \mathcal{D}_M)$



# Local Permutation Tests

- Compute a binning-based test statistic  $T := T(\mathcal{D}_1, \mathcal{D}_2, \dots, \mathcal{D}_M)$

e.g.,  $T = \sum_{i=1}^M \chi^2(\mathcal{D}_i)$  or  $T = \max_{1 \leq i \leq M} \chi^2(\mathcal{D}_i)$



# Local Permutation Tests

$\tilde{Z} = 1$

$$\begin{aligned}(X_{1,1}, Y_{1,1}) \\ (X_{2,1}, Y_{2,1}) \\ \vdots \\ (X_{\sigma_1,1}, Y_{\sigma_1,1})\end{aligned}$$

$\tilde{Z} = 2$

$$\begin{aligned}(X_{1,2}, Y_{1,2}) \\ (X_{2,2}, Y_{2,2}) \\ \vdots \\ (X_{\sigma_2,2}, Y_{\sigma_2,2})\end{aligned}$$

$\tilde{Z} = 3$

$$\begin{aligned}(X_{1,3}, Y_{1,3}) \\ (X_{2,3}, Y_{2,3}) \\ \vdots \\ (X_{\sigma_3,3}, Y_{\sigma_3,3})\end{aligned}$$


• • •

$\tilde{Z} = M$

$$\begin{aligned}(X_{1,M}, Y_{1,M}) \\ (X_{2,M}, Y_{2,M}) \\ \vdots \\ (X_{\sigma_M,M}, Y_{\sigma_M,M})\end{aligned}$$


# Local Permutation Tests

$\tilde{Z} = 1$




Permute within  
 $Y$  values

$\tilde{Z} = 2$



Permute within  
 $Y$  values


$\tilde{Z} = 3$



Permute within  
 $Y$  values




⋮ ⋮ ⋮ ⋮

$\tilde{Z} = M$




Permute within  
 $Y$  values

# Local Permutation Tests




• • •



# Local Permutation Tests

- Compute a binning-based test statistic  $T^\pi := T(\mathcal{D}_1^{\pi_1}, \mathcal{D}_2^{\pi_2}, \dots, \mathcal{D}_M^{\pi_M})$

e.g.,  $T^\pi = \sum_{i=1}^M \chi^2(\mathcal{D}_i^{\pi_1})$  or  $T^\pi = \max_{1 \leq i \leq M} \chi^2(\mathcal{D}_i^{\pi_1})$



# Algorithm: Local Permutation Tests

- Compute a binning-based test statistic  $T := T(\mathcal{D}_1, \mathcal{D}_2, \dots, \mathcal{D}_M)$

*For*  $i = 1, \dots, K$

    Permute  $Y$  values within bins

$T_i^\pi := T(\mathcal{D}_1^{\pi_1}, \mathcal{D}_2^{\pi_2}, \dots, \mathcal{D}_M^{\pi_M})$

*End*

- Reject the null if  $T > q_{1-\alpha}$

The  $1 - \alpha$  quantile of  $\{T, T_1^\pi, \dots, T_K^\pi\}$

# Algorithm: Local Permutation Tests

- Compute a binning-based test statistic  $T := T(\mathcal{D}_1, \mathcal{D}_2, \dots, \mathcal{D}_M)$

**For**  $i = 1, \dots, K$

    Permute  $Y$  values within bins

$$T_i^\pi := T(\mathcal{D}_1^{\pi_1}, \mathcal{D}_2^{\pi_2}, \dots, \mathcal{D}_M^{\pi_M})$$

**End**

- Reject the null if  $T > q_{1-\alpha}$

The  $1 - \alpha$  quantile of  $\{T, T_1^\pi, \dots, T_K^\pi\}$

# Algorithm: Local Permutation Tests

- Compute a binning-based test statistic  $T := T(\mathcal{D}_1, \mathcal{D}_2, \dots, \mathcal{D}_M)$


*For*  $i = 1, \dots, K$

    Permute  $Y$  values within bins

$$T_i^\pi := T(\mathcal{D}_1^{\pi_1}, \mathcal{D}_2^{\pi_2}, \dots, \mathcal{D}_M^{\pi_M})$$

*End*

- Reject the null if  $T > q_{1-\alpha}$



The  $1 - \alpha$  quantile of  $\{T, T_1^\pi, \dots, T_K^\pi\}$

# Local Permutation Tests

## Remark

- Unlike the two-sample and independence testing problem, there is **no guarantee** that the **local permutation test** is **valid** under the null
- In fact, **any valid CI test** has power *at most* its size  $\alpha$  against **any alternative!** (*Shah & Peters 2020*)

# Local Permutation Tests

## Remark

- Unlike the two-sample and independence testing problem, there is **no guarantee** that the **local permutation test** is **valid** under the null
- In fact, **any valid CI test** has power *at most* its size  $\alpha$  against **any alternative!** (*Shah & Peters 2020*)

# Hardness Result for CI Testing

## Theorem [Shah & Peters 2020]

- Let  $\mathcal{P}_{\text{conti}}$  be the **class of continuous distributions** on  $\mathbb{R}^d$ ,  $\mathcal{P}_{0,\text{conti}}$  be the subset of  $\mathcal{P}_{\text{conti}}$  such that  $X \perp\!\!\!\perp Y | Z$ , and  $\mathcal{P}_{1,\text{conti}} = \mathcal{P}_{\text{conti}} \setminus \mathcal{P}_{0,\text{conti}}$ .

# Hardness Result for CI Testing

Distributions under  $H_0$

## Theorem [Shah & Peters 2020]

- Let  $\mathcal{P}_{\text{conti}}$  be the **class of continuous distributions** on  $\mathbb{R}^d$ ,  $\mathcal{P}_{0,\text{conti}}$  be the subset of  $\mathcal{P}_{\text{conti}}$  such that  $X \perp\!\!\!\perp Y | Z$ , and  $\mathcal{P}_{1,\text{conti}} = \mathcal{P}_{\text{conti}} \setminus \mathcal{P}_{0,\text{conti}}$ .



Distributions under  $H_1$

# Hardness Result for CI Testing

## Theorem [Shah & Peters 2020]

- Let  $\mathcal{P}_{\text{conti}}$  be the **class of continuous distributions** on  $\mathbb{R}^d$ ,  $\mathcal{P}_{0,\text{conti}}$  be the subset of  $\mathcal{P}_{\text{conti}}$  such that  $X \perp\!\!\!\perp Y | Z$ , and  $\mathcal{P}_{1,\text{conti}} = \mathcal{P}_{\text{conti}} \setminus \mathcal{P}_{0,\text{conti}}$ .
- Suppose that a test  $\phi$  satisfies

$$\sup_{P \in \mathcal{P}_{0,\text{conti}}} \mathbb{E}_P[\phi] \leq \alpha.$$

# Hardness Result for CI Testing

## Theorem [Shah & Peters 2020]

- Let  $\mathcal{P}_{\text{conti}}$  be the **class of continuous distributions** on  $\mathbb{R}^d$ ,  $\mathcal{P}_{0,\text{conti}}$  be the subset of  $\mathcal{P}_{\text{conti}}$  such that  $X \perp\!\!\!\perp Y | Z$ , and  $\mathcal{P}_{1,\text{conti}} = \mathcal{P}_{\text{conti}} \setminus \mathcal{P}_{0,\text{conti}}$ .
- Suppose that a test  $\phi$  satisfies

$$\sup_{P \in \mathcal{P}_{0,\text{conti}}} \mathbb{E}_P[\phi] \leq \alpha.$$


Then for **any**  $P \in \mathcal{P}_{1,\text{conti}}$ , the power of  $\phi$  is **bounded above** by

$$\mathbb{E}_P[\phi] \leq \alpha.$$

# Intuition for the Hardness Result


- When  $Z$  is **continuous**, we **never** observe the same value of  $Z$

→ *The effective sample size is one!*



# Intuition for the Hardness Result

- When  $Z$  is continuous, we **never** observe the same value of  $Z$   
→ *The effective sample size is one!*



## Question.

What about scenarios where  $Z$  follows a **discrete** or a **mixture** distribution? Is CI testing still **hard**?

# New Hardness Result for CI Testing

## Theorem [KNBW 2022]

- For an arbitrary  $J \geq n(n - 1)$ , let  $\rho_{J,P} = \mathbb{P}_P(Z_1, \dots, Z_J \text{ are distinct})$

Random sample from  $P_Z$

No-collision probability

# New Hardness Result for CI Testing

## Theorem [KNBW 2022]

- For an arbitrary  $J \geq n(n - 1)$ , let  $\rho_{J,P} = \mathbb{P}_P(Z_1, \dots, Z_J \text{ are distinct})$
- Suppose that a test  $\phi$  satisfies

$$\sup_{P \in \mathcal{P}_{0,\text{disc}}} \mathbb{E}_P[\phi] \leq \alpha$$

Class of null distributions including discrete  $P_Z$

# New Hardness Result for CI Testing

## Theorem [KNBW 2022]

- For an arbitrary  $J \geq n(n - 1)$ , let  $\rho_{J,P} = \mathbb{P}_P(Z_1, \dots, Z_J \text{ are distinct})$
- Suppose that a test  $\phi$  satisfies

$$\sup_{P \in \mathcal{P}_{0,\text{disc}}} \mathbb{E}_P[\phi] \leq \alpha$$

Then for **any**  $P \in \mathcal{P}_1$ , the power of  $\phi$  is **bounded above** by

$$\mathbb{E}_P[\phi] \leq \alpha \times \rho_{J,P} + (1 - \rho_{J,P}) + \frac{n(n - 1)}{J}$$

# New Hardness Result for CI Testing

## Theorem [KNBW 2022]

- For an arbitrary  $J \geq n(n - 1)$ , let  $\rho_{J,P} = \mathbb{P}_P(Z_1, \dots, Z_J \text{ are distinct})$
- Suppose that a test  $\phi$  satisfies

$$\sup_{P \in \mathcal{P}_{0,\text{disc}}} \mathbb{E}_P[\phi] \leq \alpha$$

When  $\rho_{J,P}$  is close to one, then the power becomes close to the size

Then for **any**  $P \in \mathcal{P}_1$ , the power of  $\phi$  is **bounded above** by

$$\mathbb{E}_P[\phi] \leq \alpha \times \rho_{J,P} + (1 - \rho_{J,P}) + \frac{n(n - 1)}{J}$$

# New Hardness Result for CI Testing

## Examples

- **Continuous** distributions ( $\rho_{J,P} = 1$  for any  $J \geq 2$ )

Any **valid** CI test has the power **upper bounded** by

$$\mathbb{E}_P[\phi] \leq \alpha$$

# New Hardness Result for CI Testing

## Examples

- Continuous distributions ( $\rho_{J,P} = 1$  for any  $J \geq 2$ )

Any **valid** CI test has the power **upper bounded** by

$$\mathbb{E}_P[\phi] \leq \alpha$$

- Discrete uniform distribution on  $\{1, 2, \dots, M\}$  with  $n^4/M \rightarrow 0$

Any **valid** CI test has the power **upper bounded** by

$$\mathbb{E}_P[\phi] \leq \alpha + o(1)$$


We need **assumptions** to make CI testing **feasible**  
especially in continuous settings

# Validity of Local Permutation Tests

Joint distribution of the original data

$$\{(X_i, Y_i, Z_i)\}_{i=1}^n$$

$$P_{X,Y,Z}^n$$



# Validity of Local Permutation Tests



Joint distribution of the original data

$$\{(X_i, Y_i, Z_i)\}_{i=1}^n$$

Joint distribution of the binned data

$$\{(X_i, Y_i, \tilde{Z}_i)\}_{i=1}^n$$


Discretization of  $Z$



# Validity of Local Permutation Tests

Joint distribution of the original data  
 $\{(X_i, Y_i, Z_i)\}_{i=1}^n$

Joint distribution of the binned data  
 $\{(X_i, Y_i, \tilde{Z}_i)\}_{i=1}^n$



CI projection of  $Q_{X,Y,\tilde{Z}}^n$

# Validity of Local Permutation Tests

Joint distribution of the original data  
 $\{(X_i, Y_i, Z_i)\}_{i=1}^n$

Joint distribution of the binned data  
 $\{(X_i, Y_i, \tilde{Z}_i)\}_{i=1}^n$

Discretization of  $Z$

$$P_{X,Y,Z}^n$$

$$Q_{X,Y,\tilde{Z}}^n$$

$$\tilde{Q}_{X,Y,\tilde{Z}}^n$$

$$X \perp\!\!\!\perp Y | \tilde{Z}$$

Need to be **small** to ensure validity

CI projection of  $Q_{X,Y,\tilde{Z}}^n$

# Validity of Local Permutation Tests

## Lemma [KNBW 2022]

- Suppose that the distribution  $P_{X,Y,Z}$  satisfies  $X \perp\!\!\!\perp Y | Z$ .

# Validity of Local Permutation Tests

## Lemma [KNBW 2022]

- Suppose that the distribution  $P_{X,Y,Z}$  satisfies  $X \perp\!\!\!\perp Y | Z$ .
- Then for any  $\alpha \in (0,1)$ , the **Type I error** of the local permutation test is bounded above by

$$\mathbb{P}_{P_{X,Y,Z}^n}(T > q_{1-\alpha}) \leq \alpha + d_{\text{TV}}(Q_{X,Y,\tilde{Z}}^n, \widetilde{Q}_{X,Y,\tilde{Z}}^n)$$

\* **Total variation distance**

$$d_{\text{TV}}(P, Q) = \sup_{A \in \mathcal{F}} |P(A) - Q(A)| = \frac{1}{2} \|P - Q\|_1$$

# Validity of Local Permutation Tests

## Lemma [KNBW 2022]

- Suppose that the distribution  $P_{X,Y,Z}$  satisfies  $X \perp\!\!\!\perp Y | Z$ .
- Then for any  $\alpha \in (0,1)$ , the **Type I error** of the local permutation test is bounded above by

$$\mathbb{P}_{P_{X,Y,Z}^n}(T > q_{1-\alpha}) \leq \alpha + d_{\text{TV}}(Q_{X,Y,\tilde{Z}}^n, \widetilde{Q}_{X,Y,\tilde{Z}}^n)$$

CI projection of  $Q_{X,Y,\tilde{Z}}^n$

Discretized Distribution

# Validity of Local Permutation Tests

## Lemma [KNBW 2022]

- Suppose that the distribution  $P_{X,Y,Z}$  satisfies  $X \perp\!\!\!\perp Y | Z$ .
- Then for any  $\alpha \in (0,1)$ , the **Type I error** of the local permutation test is bounded above by

$$\mathbb{P}_{P_{X,Y,Z}^n}(T > q_{1-\alpha}) \leq \alpha + d_{\text{TV}}(Q_{X,Y,\tilde{Z}}^n, \widetilde{Q}_{X,Y,\tilde{Z}}^n)$$

The bound is **universal** but **abstract**

**Next goal:** make this bound more **explicit** under smoothness conditions

# Validity of Local Permutation Tests

## *Definition* [Generalized Hellinger distance]

Given  $\gamma \geq 1$ , the **generalized Hellinger distance** with parameter  $\gamma$  between  $P$  and  $Q$  is defined as

$$d_\gamma(P, Q) = \left( \frac{1}{2} \int |P^{1/\gamma} - Q^{1/\gamma}|^\gamma d\mu \right)^{1/\gamma}$$

# Validity of Local Permutation Tests

## *Definition* [Generalized Hellinger distance]

Given  $\gamma \geq 1$ , the **generalized Hellinger distance** with parameter  $\gamma$  between  $P$  and  $Q$  is defined as

$$d_\gamma(P, Q) = \left( \frac{1}{2} \int |p^{1/\gamma} - q^{1/\gamma}|^\gamma d\mu \right)^{1/\gamma}$$

**For example**

$$d_\gamma(P, Q) = \text{TV}(P, Q) \quad \text{when } \gamma = 1$$

$$d_\gamma(P, Q) = \text{Hellinger}(P, Q) \quad \text{when } \gamma = 2$$

# Validity of Local Permutation Tests

## *Definition* [ $\gamma$ -Hellinger Lipschitzness]

Let  $\mathcal{P}_{0,\gamma}(L) \subset \mathcal{P}_0$  be the collection of null distributions such that for all  $\textcolor{red}{z}, \textcolor{blue}{z}' \in \mathcal{Z}$

$$d_\gamma(P_{X|Z=\textcolor{red}{z}}, P_{X|Z=\textcolor{blue}{z}'}) \leq L\delta(\textcolor{red}{z}, \textcolor{blue}{z}') \quad \text{and} \quad d_\gamma(P_{Y|Z=\textcolor{red}{z}}, P_{Y|Z=\textcolor{blue}{z}'}) \leq L\delta(\textcolor{red}{z}, \textcolor{blue}{z}')$$

where  $\delta(z, z')$  is a distance between  $z$  and  $z'$  (e.g., Euclidean distance)

# Validity of Local Permutation Tests

## Definition [ $\gamma$ -Hellinger Lipschitzness]

Let  $\mathcal{P}_{0,\gamma}(L) \subset \mathcal{P}_0$  be the collection of null distributions such that for all  $\textcolor{red}{z}, \textcolor{blue}{z}' \in \mathcal{Z}$

$$d_\gamma(P_{X|Z=\textcolor{red}{z}}, P_{X|Z=\textcolor{blue}{z}'}) \leq L\delta(\textcolor{red}{z}, \textcolor{blue}{z}') \quad \text{and} \quad d_\gamma(P_{Y|Z=\textcolor{red}{z}}, P_{Y|Z=\textcolor{blue}{z}'}) \leq L\delta(\textcolor{red}{z}, \textcolor{blue}{z}')$$

where  $\delta(z, z')$  is a distance between  $z$  and  $z'$  (e.g., Euclidean distance)

## In a nutshell

Both  $P_{X|Z=z}$  and  $P_{Y|Z=z}$  are **smooth** functions with respect to  $z$

# Validity of Local Permutation Tests

## Theorem [KNBW 2022]

- Let  $h$  be the maximum diameter of bins  $B_1, \dots, B_M$

# Validity of Local Permutation Tests

## Theorem [KNBW 2022]

- Let  $h$  be the maximum diameter of bins  $B_1, \dots, B_M$
- For any  $\alpha \in (0,1)$ , the **Type I error** of the local permutation test over  $\mathcal{P}_{0,\gamma}(L)$  is bounded above by

$$\sup_{P_{X,Y,Z} \in \mathcal{P}_{0,\gamma}(L)} \mathbb{P}_{P_{X,Y,Z}^n}(T > q_{1-\alpha}) \leq \begin{cases} \alpha + O(\sqrt{n}h^\gamma), & \text{if } \gamma \in [1,2] \\ \alpha + O(\sqrt{n}h^2), & \text{if } \gamma > 2 \end{cases}$$

# Validity of Local Permutation Tests

$$\sup_{P_{X,Y,Z} \in \mathcal{P}_{0,\gamma}(L)} \mathbb{P}_{P_{X,Y,Z}^n}(T > q_{1-\alpha}) \leq \begin{cases} \alpha + O(\sqrt{n} h^\gamma), & \text{if } \gamma \in [1,2] \\ \alpha + O(\sqrt{n} h^2), & \text{if } \gamma > 2 \end{cases}$$

## Remark

- This result holds for **any binning-based test statistic  $T$**

# Validity of Local Permutation Tests

$$\sup_{P_{X,Y,Z} \in \mathcal{P}_{0,\gamma}(L)} \mathbb{P}_{P_{X,Y,Z}^n}(T > q_{1-\alpha}) \leq \begin{cases} \alpha + O(\sqrt{n} h^\gamma), & \text{if } \gamma \in [1,2] \\ \alpha + O(\sqrt{n} h^2), & \text{if } \gamma > 2 \end{cases}$$

## Remark

- This result holds for **any binning-based test statistic  $T$**
- If  $\sqrt{n} h^\gamma \rightarrow 0$ , then the local permutation test is **asymptotically valid**

# Validity of Local Permutation Tests

$$\sup_{P_{X,Y,Z} \in \mathcal{P}_{0,\gamma}(L)} \mathbb{P}_{P_{X,Y,Z}^n}(T > q_{1-\alpha}) \leq \begin{cases} \alpha + O(\sqrt{n}h^\gamma), & \text{if } \gamma \in [1,2] \\ \alpha + O(\sqrt{n}h^2), & \text{if } \gamma > 2 \end{cases}$$

## Remark

- This result holds for **any binning-based test statistic  $T$**
- If  $\sqrt{nh^\gamma} \rightarrow 0$ , then the local permutation test is **asymptotically valid**
- If  $\sqrt{nh^\gamma} \rightarrow \infty$ , then we can construct  $T$  such that **the type I error blows up**

# Validity of Local Permutation Tests

$$\sup_{P_{X,Y,Z} \in \mathcal{P}_{0,\gamma}(L)} \mathbb{P}_{P_{X,Y,Z}^n}(T > q_{1-\alpha}) \leq \begin{cases} \alpha + O(\sqrt{n}h^\gamma), & \text{if } \gamma \in [1,2] \\ \alpha + O(\sqrt{n}h^2), & \text{if } \gamma > 2 \end{cases}$$


## Remark

- This result holds for **any binning-based test statistic  $T$**
- If  $\sqrt{nh^\gamma} \rightarrow 0$ , then the local permutation test is **asymptotically valid**
- If  $\sqrt{nh^\gamma} \rightarrow \infty$ , then we can construct  $T$  such that **the type I error blows up**
- We need to take  $h$  to be small to ensure **type I error control**

# Trade-off between Type I error and Type II error




Small  $h$




Large  $h$


# Trade-off between Type I error and Type II error



# Trade-off between Type I error and Type II error



# Trade-off between Type I error and Type II error




We need to **balance** this **trade-off**

**Local permutation tests** can achieve a certain  
**minimax optimality** by choosing  $h$  carefully

# Detour: Minimax Testing Framework

- $\text{Risk}(\epsilon) = \inf_{\phi \in \Phi} \left\{ \sup_{P \in \mathcal{P}_0} \mathbb{E}_P[\phi] + \sup_{P \in \mathcal{P}_1(\epsilon)} \mathbb{E}_P[1 - \phi] \right\}$
- Worst-case type I error      Worst-case type II error
- A set of all possible tests




# Detour: Minimax Testing Framework

- $\text{Risk}(\epsilon) = \inf_{\phi \in \Phi} \left\{ \sup_{P \in \mathcal{P}_0} \mathbb{E}_P[\phi] + \sup_{P \in \mathcal{P}_1(\epsilon)} \mathbb{E}_P[1 - \phi] \right\}$ 

Worst-case type I error      Worst-case type II error

A set of all possible tests

- $\epsilon^* = \inf \{ \epsilon : \text{Risk}(\epsilon) \leq 0.1 \}$




# Detour: Minimax Testing Framework

$$\bullet \text{ Risk}(\epsilon) = \inf_{\phi \in \Phi} \left\{ \sup_{P \in \mathcal{P}_0} \mathbb{E}_P[\phi] + \sup_{P \in \mathcal{P}_1(\epsilon)} \mathbb{E}_P[1 - \phi] \right\}$$

Worst-case type I error      Worst-case type II error

A set of all possible tests

- $\epsilon^* = \inf \{ \epsilon : \text{Risk}(\epsilon) \leq 0.1 \}$
- We say that  $\phi$  is **minimax optimal** if
$$\sup_{P \in \mathcal{P}_0} \mathbb{E}_P[\phi] + \sup_{P \in \mathcal{P}_1(\epsilon^*)} \mathbb{E}_P[1 - \phi] \leq 0.1$$




# Detour: Minimax Testing Framework

- $$\text{Risk}(\epsilon) = \inf_{\phi \in \Phi} \left\{ \sup_{P \in \mathcal{P}_0} \mathbb{E}_P[\phi] + \sup_{P \in \mathcal{P}_1(\epsilon)} \mathbb{E}_P[1 - \phi] \right\}$$

Worst-case type I error      Worst-case type II error

A set of all possible tests

- $$\epsilon^* = \inf \{ \epsilon : \text{Risk}(\epsilon) \leq 0.1 \}$$
  - We say that  $\phi$  is **minimax “rate” optimal** if
 
$$\sup_{P \in \mathcal{P}_0} \mathbb{E}_P[\phi] + \sup_{P \in \mathcal{P}_1(\epsilon)} \mathbb{E}_P[1 - \phi] \leq 0.1$$
- whenever  $\epsilon \asymp \epsilon^*$



# Optimality results of Neykov et al. (2021)

Let  $(X, Y, Z) \in [0,1]^3$

# Optimality results of Neykov et al. (2021)

Let  $(X, Y, Z) \in [0,1]^3$

## Class of null distributions

$\mathcal{P}_{0,\text{NBL}}$ : TV or  $\chi^2$  smooth ( $Z$ ) +  $X \perp\!\!\!\perp Y | Z$

# Optimality results of Neykov et al. (2021)

Let  $(X, Y, Z) \in [0,1]^3$

## Class of null distributions

$\mathcal{P}_{0,\text{NBL}}$ : TV or  $\chi^2$  smooth ( $Z$ ) +  $X \perp\!\!\!\perp Y | Z$

## Class of alternative distributions

$\mathcal{P}_{1,\text{NBL}}(\epsilon)$ : TV smooth ( $Z$ ) +  $s$ -Hölder smooth ( $X, Y$ ) +  $\inf_{P \in \mathcal{P}_0} d_{TV}(P, Q) \geq \epsilon$

# Optimality results of Neykov et al. (2021)

Let  $(X, Y, Z) \in [0,1]^3$

**Class of null distributions**

$\mathcal{P}_{0,\text{NBL}}$ : TV or  $\chi^2$  smooth ( $Z$ ) +  $X \perp\!\!\!\perp Y | Z$

**Class of alternative distributions**

$\mathcal{P}_{1,\text{NBL}}(\epsilon)$ : TV smooth ( $Z$ ) +  $s$ -Hölder smooth ( $X, Y$ ) +  $\inf_{P \in \mathcal{P}_0} d_{TV}(P, Q) \geq \epsilon$

Neykov et al. (2021) prove that the **minimax testing rate** for this problem is

$$\epsilon_n^\star \asymp n^{-\frac{2s}{5s+2}}$$

## In our work [KNBW 2022]

- We show that the same test statistic calibrated by the **local permutation method** achieves the **same minimax rate**  $\epsilon_n^* \asymp n^{-\frac{2s}{5s+2}}$
- **However**, the optimality can be obtained over a slightly smaller class of null distributions (**Hellinger smoothness** instead of **TV smoothness**)

⇒ Double-binning method

## In our work [KNBW 2022]

- We show that the same test statistic calibrated by the **local permutation method** achieves the **same minimax rate**  $\epsilon_n^* \asymp n^{-\frac{2s}{5s+2}}$
- **However**, the optimality can be obtained over a slightly smaller class of null distributions (**Hellinger smoothness** instead of **TV smoothness**)



**Double-binning method**

## In our work [KNBW 2022]

- We show that the same test statistic calibrated by the **local permutation method** achieves the **same minimax rate**  $\epsilon_n^* \asymp n^{-\frac{2s}{5s+2}}$
- **However**, the optimality can be obtained over a slightly smaller class of null distributions (**Hellinger smoothness** instead of **TV smoothness**)

⇒ **Double-binning method**

# Summary

- We presented a **theoretical foundation** of local permutation tests
- We revisited the **hardness of CI testing** and showed that the hardness is determined by “**no-collision**” probability
- We illustrated that the local permutation test with a careful binning scheme can lead to **minimax optimality**

# Summary


- We presented a **theoretical foundation** of local permutation tests
- We revisited the **hardness of CI testing** and showed that the hardness is determined by “**no-collision**” probability
- We illustrated that the local permutation test with a careful binning scheme can lead to **minimax optimality**

# Summary

- We presented a **theoretical foundation** of local permutation tests
- We revisited the **hardness of CI testing** and showed that the hardness is determined by “**no-collision**” probability
- We illustrated that the local permutation test with a careful binning scheme can lead to **minimax optimality**


Thank you!

# Double-binning method




- Previously, we **permuted** a sample and **computed** a test statistic within **the same bin**

# Double-binning method



- In **double-binning**, we consider bins of **two** distinct resolutions
  - **Permute** the observations within the **finer bins**
  - **Compute** test statistics over the **coarser bins**

# Double-binning method




- In double-binning, we consider bins of **two** distinct resolutions
    - Permute the observations within the **finer bins**
    - Compute test statistics over the **coarser bins**
- Decrease the type I error**
- Increase the power**

# Optimality results of Neykov et al. (2021)

## Procedure


- Discretize  $(X, Y, Z)$  into  $d_X \times d_Y \times d_Z$  bins



# Optimality results of Neykov et al. (2021)

## Procedure


- Discretize  $(X, Y, Z)$  into  $d_X \times d_Y \times d_Z$  bins
- Set  $d_X = d_Y = \lceil n^{\frac{2}{5s+2}} \rceil$  and  $d_Z = \lceil n^{\frac{2s}{5s+2}} \rceil$



# Optimality results of Neykov et al. (2021)

## Procedure

- Discretize  $(X, Y, Z)$  into  $d_X \times d_Y \times d_Z$  bins
- Set  $d_X = d_Y = \lceil n^{\frac{2}{5s+2}} \rceil$  and  $d_Z = \lceil n^{\frac{2s}{5s+2}} \rceil$
- Compute the test statistic  $T$  proposed by  
**Canonne et al. (2018)** for discrete CI testing




$$T = \sum_{i=1}^{d_Z} w_i U_i(\mathcal{D}_i)$$

Weighted sum of U-statistics

# Optimality results of Neykov et al. (2021)

## Procedure


- Discretize  $(X, Y, Z)$  into  $d_X \times d_Y \times d_Z$  bins
- Set  $d_X = d_Y = \lceil n^{\frac{2}{5s+2}} \rceil$  and  $d_Z = \lceil n^{\frac{2s}{5s+2}} \rceil$
- Compute the test statistic  $T$  proposed by  
**Canonne et al. (2018)** for discrete CI testing
- **Reject** the null when  $T > Cd_Z^{1/2}$  for a sufficiently large  $C$



# Optimality results of Neykov et al. (2021)

## Procedure

- Discretize  $(X, Y, Z)$  into  $d_X \times d_Y \times d_Z$  bins
- Set  $d_X = d_Y = \lceil n^{\frac{2}{5s+2}} \rceil$  and  $d_Z = \lceil n^{\frac{2s}{5s+2}} \rceil$
- Compute the test statistic  $T$  proposed by  
**Canonne et al. (2018)** for discrete CI testing
- **Reject** the null when  $T > Cd_Z^{1/2}$  for a sufficiently large  $C$




- It achieves the **minimax optimal rate**

# Optimality results of Neykov et al. (2021)

## Procedure

- Discretize  $(X, Y, Z)$  into  $d_X \times d_Y \times d_Z$  bins
- Set  $d_X = d_Y = \lceil n^{\frac{2}{5s+2}} \rceil$  and  $d_Z = \lceil n^{\frac{2s}{5s+2}} \rceil$
- Compute the test statistic  $T$  proposed by  
**Canonne et al. (2018)** for discrete CI testing
- **Reject** the null when  $T > Cd_Z^{1/2}$  for a sufficiently large  $C$




- It achieves the **minimax optimal rate**
- **However**, the unspecified number  $C$  makes this procedure **impractical**

# Simulation Results

# Simulation settings

## ■ Under the null

- $Z \sim \text{Uniform}[0,1]$
- $X = \sin\{2\pi\Phi^{-1}(Z)\}e^{-\Phi^{-1}(Z)^2/2} + 0.3\varepsilon_1$
- $Y = \cos\{2\pi\Phi^{-1}(Z)\}e^{-\Phi^{-1}(Z)^2/2} + 0.3\varepsilon_1$
- $\varepsilon_1, \varepsilon_2 \stackrel{i.i.d.}{\sim} N(0,1)$



# Simulation settings

## ■ Under the null

- $Z \sim \text{Uniform}[0,1]$
- $X = \sin\{2\pi\Phi^{-1}(Z)\}e^{-\Phi^{-1}(Z)^2/2} + 0.3\varepsilon_1$
- $Y = \cos\{2\pi\Phi^{-1}(Z)\}e^{-\Phi^{-1}(Z)^2/2} + 0.3\varepsilon_1$
- $\varepsilon_1, \varepsilon_2 \stackrel{i.i.d.}{\sim} N(0,1)$

## ■ Under the alternative

- $Z \sim \text{Uniform}[0,1]$
- $X = \sin\{2\pi\Phi^{-1}(Z)\}e^{-\Phi^{-1}(Z)^2/2} + 0.3\varepsilon_1$
- $Y = \textcolor{blue}{0.15X} + \cos\{2\pi\Phi^{-1}(Z)\}e^{-\Phi^{-1}(Z)^2/2} + 0.3\varepsilon_1$
- $\varepsilon_1, \varepsilon_2 \stackrel{i.i.d.}{\sim} N(0,1)$

# Simulation settings

## ■ Under the null

- $Z \sim \text{Uniform}[0,1]$
- $X = \sin\{2\pi\Phi^{-1}(Z)\}e^{-\Phi^{-1}(Z)^2/2} + 0.3\varepsilon_1$
- $Y = \cos\{2\pi\Phi^{-1}(Z)\}e^{-\Phi^{-1}(Z)^2/2} + 0.3\varepsilon_1$
- $\varepsilon_1, \varepsilon_2 \stackrel{i.i.d.}{\sim} N(0,1)$

## ■ Under the alternative

- $Z \sim \text{Uniform}[0,1]$
- $X = \sin\{2\pi\Phi^{-1}(Z)\}e^{-\Phi^{-1}(Z)^2/2} + 0.3\varepsilon_1$
- $Y = 0.15X + \cos\{2\pi\Phi^{-1}(Z)\}e^{-\Phi^{-1}(Z)^2/2} + 0.3\varepsilon_1$
- $\varepsilon_1, \varepsilon_2 \stackrel{i.i.d.}{\sim} N(0,1)$

■ Test statistic:  $T = \sum_{i=1}^M U_i(W_i)$

$U_i(W_i)$ : a test statistic for independence between  $X$  and  $Y$

# Simulation settings

## ■ Under the null

- $Z \sim \text{Uniform}[0,1]$
- $X = \sin\{2\pi\Phi^{-1}(Z)\}e^{-\Phi^{-1}(Z)^2/2} + 0.3\varepsilon_1$
- $Y = \cos\{2\pi\Phi^{-1}(Z)\}e^{-\Phi^{-1}(Z)^2/2} + 0.3\varepsilon_1$
- $\varepsilon_1, \varepsilon_2 \stackrel{i.i.d.}{\sim} N(0,1)$

## ■ Under the alternative

- $Z \sim \text{Uniform}[0,1]$
- $X = \sin\{2\pi\Phi^{-1}(Z)\}e^{-\Phi^{-1}(Z)^2/2} + 0.3\varepsilon_1$
- $Y = 0.15X + \cos\{2\pi\Phi^{-1}(Z)\}e^{-\Phi^{-1}(Z)^2/2} + 0.3\varepsilon_1$
- $\varepsilon_1, \varepsilon_2 \stackrel{i.i.d.}{\sim} N(0,1)$

- Test statistic:  $T = \sum_{i=1}^M U_i(W_i)$
- Sample size:  $n = 1000$

$U_i(W_i)$ : a test statistic for independence between  $X$  and  $Y$

# Type I error vs. Number of bins

Significance level  
 $\alpha = 0.05$

The number of bins

	5	10	15	20	25	30	35
HSIC	1.000	0.443	0.105	0.055	0.057	0.048	0.040
Dcov	1.000	0.668	0.159	0.074	0.050	0.043	0.036
Cov	1.000	0.755	0.185	0.078	0.070	0.049	0.051
XiCov	0.986	0.103	0.058	0.043	0.065	0.058	0.053
Tau	1.000	0.679	0.158	0.073	0.066	0.054	0.048
TauStar	1.000	0.623	0.144	0.065	0.061	0.051	0.049
MIT	0.991	0.121	0.066	0.057	0.062	0.056	0.049

As # of bins **increases** (i.e., maximum diameter decreases),  
the type I error **decreases** consistently

# Type I error vs. Number of bins

Significance level  
 $\alpha = 0.05$

The number of bins

	5	10	15	20	25	30	35
HSIC	1.000	0.443	0.105	0.055	0.057	0.048	0.040
Dcov	1.000	0.668	0.159	0.074	0.050	0.043	0.036
Cov	1.000	0.755	0.185	0.078	0.070	0.049	0.051
XiCov	0.986	0.103	0.058	0.043	0.065	0.058	0.053
Tau	1.000	0.679	0.158	0.073	0.066	0.054	0.048
TauStar	1.000	0.623	0.144	0.065	0.061	0.051	0.049
MIT	0.991	0.121	0.066	0.057	0.062	0.056	0.049

As # of bins **increases** (i.e., maximum diameter decreases),  
the type I error **decreases** consistently

# Power vs. Number of bins

Significance level  
 $\alpha = 0.05$

The number of bins

	5	10	15	20	25	30	35
HSIC	1.000	0.996	0.870	0.653	0.501	0.410	0.369
Dcov	1.000	1.000	0.982	0.900	0.782	0.696	0.612
Cov	1.000	1.000	0.998	0.955	0.878	0.801	0.734
XiCov	1.000	0.558	0.270	0.211	0.196	0.160	0.154
Tau	1.000	1.000	0.989	0.908	0.805	0.714	0.659
TauStar	1.000	1.000	0.974	0.856	0.722	0.604	0.537
MIT	1.000	0.627	0.356	0.294	0.226	0.192	0.196

As # of bins **increases** (i.e., maximum diameter decreases),  
the power **decreases** consistently

# Power vs. Number of bins

Significance level  
 $\alpha = 0.05$

The number of bins

	5	10	15	20	25	30	35
HSIC	1.000	0.996	0.870	0.653	0.501	0.410	0.369
Dcov	1.000	1.000	0.982	0.900	0.782	0.696	0.612
Cov	1.000	1.000	0.998	0.955	0.878	0.801	0.734
XiCov	1.000	0.558	0.270	0.211	0.196	0.160	0.154
Tau	1.000	1.000	0.989	0.908	0.805	0.714	0.659
TauStar	1.000	1.000	0.974	0.856	0.722	0.604	0.537
MIT	1.000	0.627	0.356	0.294	0.226	0.192	0.196

As # of bins **increases** (i.e., maximum diameter decreases),  
the power **decreases** consistently

# Double-binning approach: Type I error

Significance level  
 $\alpha = 0.05$

		The number of bins							
		5	10	15	20	25	30	35	Double-Binning
Test Statistics	HSIC	1.000	0.443	0.105	0.055	0.057	0.048	0.040	0.055
	Dcov	1.000	0.668	0.159	0.074	0.050	0.043	0.036	0.065
	Cov	1.000	0.755	0.185	0.078	0.070	0.049	0.051	0.077
	XiCov	0.986	0.103	0.058	0.043	0.065	0.058	0.053	0.047
	Tau	1.000	0.679	0.158	0.073	0.066	0.054	0.048	0.057
	TauStar	1.000	0.623	0.144	0.065	0.061	0.051	0.049	0.062
	MIT	0.991	0.121	0.066	0.057	0.062	0.056	0.049	0.066

# of coarser bin:  $\lceil n^{0.3} \rceil$   
# of finer bin:  $\lceil n^{0.3} \rceil \times \lceil n^{0.3} \rceil$

# Double-binning approach: Power

Significance level  
 $\alpha = 0.05$

The number of bins

	5	10	15	20	25	30	35	Double-Binning
Test Statistics	X	X	X	0.653	0.501	0.410	0.369	0.928
Dcov	X	X	X	0.900	0.782	0.696	0.612	0.987
Cov	X	X	X	0.955	0.878	0.801	0.734	0.988
XiCov	X	X	X	0.211	0.196	0.160	0.154	0.411
Tau	X	X	X	0.908	0.805	0.714	0.659	0.987
TauStar	X	X	X	0.856	0.722	0.604	0.537	0.976
MIT	X	X	X	0.294	0.226	0.192	0.196	0.480

(X) They do not control the type I error;  
Hence ignored in power comparisons

# of coarser bin:  $\lceil n^{0.3} \rceil$   
# of finer bin:  $\lceil n^{0.3} \rceil \times \lceil n^{0.3} \rceil$