

Figure 2. Composite overdose-vulnerability index, Dallas County

Equal-weighted index of overdose mortality, poverty, uninsurance, naloxone distance, and transit access; Jenks natural-break tiers



Vulnerability tiers (Jenks natural breaks)

Tier 1 - highest, ≥ 0.434 (170 tracts, heavy outline)

Tier cuts marked on colour bar:
 Tier 2 = 0.338-0.434 (303 tracts)
 Tier 3 = < 0.338 (170 tracts)

Death count NOT suppressed (52 of 643 ranked tracts; all others CDC-suppressed)

Excluded - zero population (2 tracts)



Sources: CDC NCHS drug-overdose death rates, model-based (4day-mt2f), 12-month period 2025-01/2025-12, data as of 2026-07-22 | ACS 2020-2024 5-year (S1701 poverty; S2701 uninsured), released Jan 2026 | CDC/ATSDR SVI 2022 | TX DSHS Narcansites (Apr 2026), SAMHSA findtreatment.gov & ArcGIS naloxone inventory (verified 2024-2026) | DART GTFS (via context module, 2026-07-23) | County & tract boundaries: US Census TIGER/Line 2024 | Projection: NAD83 / Texas North Central (EPSG:32138). CDC suppresses tract-level death counts of 1-9; 592 of 645 tracts (91.8%) carry suppressed counts, so model-based rates are smoothed estimates, not exact counts.