

Physics-Informed Neural Quantum States for Quantum Dots:
Tangent-Space Geometry, Message Passing, and Wigner
Molecules

Aleksander Sekkelsten

September 7, 2026

Abstract

A neural network used as a variational wavefunction is judged by its energy. This thesis asks what the energy cannot see: two networks landing on the same number need not be the same object, a preconditioner decisive in one training regime does nothing in another, and an objective that reproduces a Markov chain’s answer in one corner of the phase diagram departs from it in the next. The system is the two-dimensional harmonic quantum dot, from the kinetic-energy-dominated droplet to the dilute Wigner molecule, for $N \in \{2, 6, 12, 20\}$ electrons and confinements ω from 1 to 10^{-3} .

The ansatz is a Slater determinant dressed by a neural Jastrow correlator, which reshapes the amplitude, and a coordinate backflow, which reshapes the nodes. Both are built to be safe under a Laplacian objective: trap-unit coordinates, soft-core pair features, an analytic electron–electron cusp. Each exists in a separable form and a message-passing one. Training is by residual pretraining followed by stochastic reconfiguration, and separately by importance-sampled collocation that places no Markov chain in the gradient. The trained states are read through the geometry of their log-derivatives—the quantum geometric tensor and the neural tangent kernel—and through structural diagnostics of the electron arrangement. Energies reproduce published fixed-node diffusion Monte Carlo wherever it exists, to 0.0008–0.04%; below $\omega = 0.1$ for $N \geq 6$ none exists, and those states are predictions.

Message passing changes the variational geometry before it changes the energy. At strong confinement a message-passing correlator reaches the same reference-quality solution along about half as many effective tangent directions as a separable network of equal or greater capacity, and at every confinement far more of its tangent weight lies inside a compact dictionary of collective operators. Its local-energy variance is lower throughout, by factors of 1.2 to 3.4, while the two agree in energy to four figures. In the nodes the same channel shows as a collapse averted: toward the Wigner regime a conventional backflow’s displacement field collapses spectrally to rank one and loses half a percent of the energy, where the message-passing field keeps its rank. Holding the backflow fixed and swapping only the correlator moves the collapse boundary, which points to one relational bottleneck rather than two deficits.

Natural-gradient preconditioning earns its cost where the tangent metric is anisotropic and still *estimable*. Conditioning alone is not the criterion: under $|\Psi|^2$ sampling the empirical metric is badly conditioned on exactly the models where stochastic reconfiguration and Adam agree to four decimals. Under a fixed collocation proposal it is what makes the optimisation converge, and once the sampling that builds the metric collapses, inverting it does harm.

The trained states then serve as instruments, and the crossover they resolve is *staged*: the cloud expands and its shells stiffen radially well before the electrons acquire angular order within them. A polygon shell model fitted on held-out frames reproduces the measured pair distribution to cosine similarity 0.987–0.999 against 0.936–0.996 for an angle-randomised null. At $\omega = 10^{-3}$ the six-electron dot shows persistent mixing between the (1, 5) and (0, 6) sectors, the twelve-electron dot forms a two-shell molecule with a modal (3, 9) split, and the twenty-electron dot a three-shell molecule that collapses to a single ring within one decade in ω .

Those measurements return to the solver. A generic collocation proposal loses the Wigner state entirely; a proposal rebuilt from the measured shell geometry lifts the effective sample size by 15–32 $\times$ with no training at all and carries Markov-chain-free training to $\omega = 0.0035$, where a different limit takes over—a transfer instability in the Slater–Jastrow cancellation that no change of proposal repairs.

Two limits bound all of it. Every state is the lowest found within a fixed spin-projection sector, and total spin is not scanned. The architecture comparison is seeded at $N=6$, rests on fewer models at $N=12$ and on a single production state at $N=20$, and the two backflow variants are not capacity-matched.

*To go wrong in one's own way is better than to go right in
someone else's.*

FYODOR DOSTOEVSKY
Crime and Punishment

*Life must be understood backwards. But they forget the
other proposition, that it must be lived forwards.*

SØREN KIERKEGAARD
Journals

Preface

When I came to UiO, Morten read my background and said: “You know PINNs, you know many-body quantum systems. I want you to do PINNs on quantum dots.” I said “sure” and then asked the only sensible follow-up question I had: “Where do I start?”

I did not realise then that this was the question I had been assigned to answer. This thesis is the long answer. The short answer is: start by arranging to be wrong in a way that can be measured.

That is what variational methods make possible. The variational principle does not say how to improve a trial wavefunction, but it does guarantee that its energy lies above the ground-state energy. Improvement therefore has a direction: lower energy is better. For a project with few fixed landmarks, that was enough to begin. The harder task was learning what a disappointing result meant.

It was a solitary problem, not for want of literature—neural quantum states, FermiNet, PauliNet, and neural backflow offered much to learn from—but because the particular route I was trying to take had little precedent. When an experiment failed, it was rarely clear whether the method was unsound, the architecture inadequate, the optimiser poorly matched to the problem, or the code quietly broken. For long stretches, I did not know. The work became an attempt to construct tests that could tell these explanations apart.

The most important lessons came from failures. An error in the importance-sampling code went undetected for months because its signature—loss of accuracy at larger particle numbers and weaker confinement—was exactly what an inadequate architecture might have produced. A small per-particle error in a log-density accumulates over the N particles and is then exponentiated into the importance weights; the bug was invisible where experiments were cheap and ruinous where they mattered. Discovering it damaged my confidence in everything else. Re-establishing that confidence, rather than assuming it, became a substantial part of the later work.

Other failures taught the same lesson from different directions. A Langevin refinement step introduced bias into the gradient. A second-order optimiser became least reliable when its estimated metric was too noisy to resolve. Neither result was useful merely because it was negative. Each exposed an assumption that had been invisible until the computation violated it. Negative results of this kind are not a concession or an embarrassment; they are part of the method, and I have written them up as such.

That diagnostic stance also changed what I thought the network was for. I began by asking whether a machine-learning construction could solve the many-body Schrödinger equation. I ended up using the trained state as an apparatus: something to examine for the structure it had captured and the directions of variation that mattered. That shift, from treating a network as an answer to treating it as an object of study, is the thread running through this thesis.

The work also made a connection harder to ignore. Its obstacles have a double character: each is at once a physics problem and an optimisation problem. Stochastic reconfiguration, developed for variational wavefunctions, is natural gradient descent on the variational manifold. The REINFORCE estimator from reinforcement learning is the score-function gradient of the variational energy. The Fisher information matrix of statistics is, for a real amplitude and up to a factor of four, the quantum geometric tensor of physics. The same mathematics appears under different names in different communities. Recognising that felt less like discovering a bridge between fields than like realising that the two fields had been approaching the same problem from opposite sides.

Looking back, it is easy to make a catalogue of what I would now do differently. Most of that

knowledge, however, was purchased by the failures and could not have been read off in advance. The regret and the result are the same object seen from two sides.

The most consistent source of inspiration, motivation, and steadiness through all of it has been my girlfriend, Elida. Her support made it possible to begin this work and to stay with it when it became difficult. I am more grateful than I can put down here.

I owe a warm and sincere thanks to my family, who have supported me in every undertaking I have had and have simply always been there; and to my friends, inside CCSE and out, who supplied the joy, the laughter, and a good deal of the sanity this project required.

To Morten, again, and properly: I thank you for the assignment, for the two years we've had together, and above all for a genuine and undisguised interest in what I was finding at every stage—including the stages where what I was finding was that I had been wrong. That interest is most of what encouragement actually consists of.

Which is, at last, the answer to the question I asked on the first day. You start by being willing to be wrong on the record, and then you keep the record.

Contents

Abstract	1
Preface	3
I Introduction	11
II Theory	16
1 The many-body problem	17
1.1 The many-body problem	17
1.2 Correlation, closed shells, and the cost of exactness	17
1.3 Quantum Dots	18
1.3.1 Model and units	18
1.3.2 One-particle structure and shell filling	18
1.3.3 Spin structure and Slater form	18
1.3.4 Kato’s Cusp Conditions (2D Coulomb systems)	19
1.4 Structural diagnostics and Wigner markers	20
1.5 Quantum Monte Carlo benchmarks	20
1.5.1 Variational Monte Carlo	21
1.5.2 Diffusion Monte Carlo, and the fixed-node constraint	21
1.5.3 Requirements on the trial state	22
2 Neural quantum states and variational geometry	23
2.1 The variational learning objective	23
2.2 Optimisation	23
2.2.1 First-order methods	24
2.2.2 Second-order methods and natural gradient	24
2.2.3 The variational tangent space: geometric tensor, tangent kernel, and effective dimension	24
2.2.4 Tangent-space diagnostics	25
2.3 Smooth networks for Laplacian objectives	25
2.4 Architectures for many-body wavefunctions	26
2.4.1 Permutation symmetry and equivariance	27
2.4.2 Permutation-invariant pooling: DeepSets	27
2.4.3 Message passing and graph networks	27
2.4.4 The CTNN: one message-passing instance	27
2.4.5 Relevance to the ansatz	28
2.5 Approximation theory and its limits	28
2.6 Residual objectives and their difficulties	28
2.7 Constraints carried into the method	29
III Methods	30
3 Trial Wavefunction	31
3.1 Preliminaries	31

C.5	Why VMC + SR escapes	101
C.6	Experimental evidence	102
C.6.1	Baseline: correlator alone vs. correlator + backflow	102
C.6.2	Sampling strategies	102
C.6.3	Loss modifications	103
C.6.4	Gradient regularisation	103
C.6.5	Gradient preconditioning	103
C.6.6	Summary of all interventions	103
C.7	Connection to Appendices A and B	104
D	Additional collocation results	106
D.1	What was varied	106
D.2	The $N = 20$ collocation energies	106
D.3	Negative results worth recording	107
E	The learned representation of one trained solution	108
E.1	Correlator geometry: a low-rank representation that reorganises	108
E.2	Backflow: a high-rank field with a low-rank footprint	109
E.3	Correlator–backflow coupling: from independence to cooperation	110
E.4	Limits of component-level interpretation	111
F	Structural diagnostics in full	113
F.1	Fitted energy and radius exponents	113
F.2	Shell-resolved structure across the whole grid	114
F.3	Shell-model fit to the pair distribution	115
F.4	The measured shell radius against the classical prediction	116
G	Declaration on the Use of Artificial Intelligence	118

List of Figures

2.1	Activation functions and their first three derivatives	26
6.1	Relative deviation from the external references	55
6.2	One-body densities at strong and weak confinement, $N = 2, 6, 12$	57
6.3	One-body densities at strong and weak confinement, $N=20$	58
6.4	Energy-based Wigner diagnostics against trap frequency	58
6.5	Two-electron radial densities $P(r)$ across ω	60
6.6	Six-electron radial densities $P(r)$ across ω	60
6.7	Twelve-electron radial densities $P(r)$ across ω	62
7.1	Message passing compresses the correlator	65
7.2	The leading tangent mode is a nameable physical mode	66
7.3	Dictionary-orthogonal tangent mass against ω	67
7.4	Fixed-measure ablation cost of each component	70
7.5	Correlator physicality and backflow rank collapse together	71
7.6	ESS collapse of the naive collocation proposal	73
F.1	Shell-resolved reconstruction of $g(r)$, $N=12$	116

List of Tables

3.1	Architecture and training hyperparameters	36
6.1	Two-electron energies from residual-only training	53
6.2	Ground-state energies against the available references	54
6.3	Radial localisation across the confinement range	59
6.4	Held-out shell-model fit against an angle-randomised null	62
7.1	Local-energy variance by architecture	66
7.2	Relative energy error (%) of the two backflow architectures, seed-averaged.	68
7.3	Backflow displacement rank (participation ratio)	69
7.4	Condition number of the empirical geometric tensor	72
7.5	Energies from Markov-chain-free collocation training	75
7.6	Collocation variants tried, and what each one settled	78
7.7	Reliability campaign: 30 runs across five confinements	79
7.8	Pareto $\hat{k}$ by regime and recipe	80
7.9	Gap between the VMC probe and the IS evaluation	81
7.10	ESS of the origin and Wigner-ring proposals	82
7.11	MCMC-free Wigner-ring cascade, $N=6$	83
C.1	Baseline strong-form collocation results	102
C.2	Effect of sampling strategy on collocation accuracy	102
C.3	Effect of loss-function modifications	103
C.4	Regularisation of the E_L singularity	103
C.5	Gradient preconditioning experiments	103
C.6	All strong-form collocation interventions	104
E.1	Correlator feature geometry	109
E.2	Backflow displacement geometry	110
E.3	Correlator–backflow coupling diagnostics	111
F.1	Radial diagnostics for $N=2$ across ω (Bohr units).	113
F.2	Radial and angular diagnostics for $N=6$ across ω (Bohr; angles in radians).	113
F.3	Shell-resolved diagnostics across the whole grid	115
F.4	Shell-model fit to the pair-distance distribution	116
F.5	Measured shell radius against the classical prediction	117

Part I

Introduction

Introduction

A system of many interacting electrons is described completely by its wavefunction. The same object puts the system almost out of reach, because a complete description of N interacting particles is a function of all their coordinates at once, and the space of those coordinates grows with every particle added. The equation that fixes this wavefunction is known exactly; the entire difficulty is solving it.

The competition that makes it hard is physical. Kinetic energy tends to delocalise the particles; interactions and external potentials drive localisation and correlation; and the balance between them plays out in a configuration space far too large to enumerate. Yet the quantities we need—energies, correlation functions, phase boundaries—are precisely the ones that depend on getting that balance right. This is the *many-body problem*, the point at which quantum mechanics turns from a compact set of principles into one of the hardest computational challenges in physics.

A wide range of electronic-structure methods has been developed to address this. Mean-field approaches such as Hartree–Fock replace the full many-body problem with an effective single-particle picture, where each electron moves in an average field from the others. This captures Fermi statistics and some exchange effects but leaves most correlation physics untreated. More systematic methods such as Full Configuration Interaction (FCI) are, in principle, exact within a finite basis, but their cost grows combinatorially with system size. Coupled Cluster (CC) theory provides a more efficient and size-extensive hierarchy of approximations, but it remains computationally demanding and can struggle in strongly correlated regimes or near degeneracies.

Quantum Monte Carlo (QMC) methods, including Variational Monte Carlo (VMC) and Diffusion Monte Carlo (DMC), take a different approach: they represent the wavefunction implicitly through stochastic sampling. For suitable trial states, they can produce highly accurate benchmark energies and correlations for continuum systems. However, QMC is not free of difficulties. The fermion sign problem limits direct simulations, fixed-node approximations inherit bias from the chosen trial nodes, and constructing high-quality trial wavefunctions is increasingly difficult as systems become more strongly correlated or inhomogeneous. In practice, then, much of the many-body problem reduces to a narrower one: choosing an expressive, well-conditioned ansatz and optimising it reliably.

Machine learning and many-body wavefunctions. In parallel with these developments, machine learning—especially deep learning—has become a standard tool for representing complex, high-dimensional functions and fitting them to data. In physics this has taken two broad forms. Neural networks act as *surrogates*, learning maps from parameters or boundary conditions to observables, or approximating solution operators of partial differential equations—the setting of physics-informed neural networks and operator learning. Or they act as *variational ansätze* for quantum states (neural quantum states), where the network outputs amplitudes or log-probabilities for many-body configurations and is trained by variational Monte Carlo. This thesis is in the second tradition and borrows from the first.

Universal approximation establishes representational capacity, not trainability. The theorems guarantee that a sufficiently large network can *represent* any suitably regular function to arbitrary accuracy. They say nothing about whether gradient-based optimisation will *find* that representation, nor at what sample cost. This thesis measures that gap directly: the same ansatz, demonstrably able to express the ground state, is placed in training schemes that do and do not allow it to get there. Representability is cheap; trainability is the difficulty.

A weaker but sounder case for a neural ansatz survives that objection. Established methods pay for accuracy in one of two currencies. FCI is exact within a finite basis, but its cost grows combinatorially with that basis, so a modest two-dimensional dot of twenty electrons is already far beyond reach.

Cheaper methods such as coupled cluster buy their favourable scaling instead by deciding in advance which correlations may be described at all.

A variational neural ansatz avoids both of those particular costs. It requires no explicit enumeration of the determinant space, and it fixes no prior truncation on which correlations may be described; its parameter count is set empirically by the structure of the target rather than by the size of a basis. That is not freedom from scaling—capacity can still have to grow with N or with the accuracy demanded, as the twenty-electron results below show—but it is freedom to concentrate capacity wherever the solution actually varies. Whether it does so in practice is an empirical question, and this thesis treats it as one.

That question also suggests a different way to use a solver. Nearly every established method for the many-body Schrödinger equation is constructed outward from physical principles: an approximation is chosen, justified physically, and then implemented. A trained network inverts the order. Physical structure is still imposed on it—antisymmetry, the cusp, trap scaling, permutation symmetry—but the *specific correlations* it develops within those constraints are not prescribed, and are interrogated only afterwards. Unlike an implicit stochastic representation, it is a concrete parameterised object whose internal structure can be examined directly. This motivates a question that the size of the problem does not by itself settle. A twenty-electron dot is a function of forty coordinates, and any basis expansion of it grows combinatorially—but it does not follow that the solution itself is forty-dimensional in any useful sense. How many active tangent directions does a given parameterisation actually use to represent the solved state—a question about the solution and its representation rather than about the equation. It is worth saying at once that this is not an intrinsic dimension of the wavefunction: the count depends on how the state is parameterised, as §2.2.3 makes precise.

We make that question quantitative through the geometry of the wavefunction’s log-derivatives. Perturbing the parameters of a trained ansatz over an ensemble of configurations and collecting the resulting sensitivities into a covariance matrix yields, in the simplest reading, a principal-component analysis of the state: the eigenvalues report how much variation each independent direction carries, and their distribution counts how many directions are doing genuine work. The physically appropriate version of this covariance is the quantum geometric tensor (equivalently, the Fisher metric), and the corresponding count is the participation ratio of its spectrum—an effective dimension rather than a hard rank, insensitive to the raw parameter count but not to the choice of parameterisation. Chapter 7 develops this instrument and applies it to architecture, optimiser, and training paradigm alike; the questions it answers are how this effective dimension behaves as particles are added and as the trap is opened, and whether different architectures arrive at the same solution by different routes.

In principle, then, neural networks provide highly expressive ansätze for quantum problems. In practice, several issues arise. Objectives based on local energies involve second derivatives and can be numerically delicate. Coordinate parameterisations that do not respect known structure can lead to large gradients and Laplacian spikes near electron coalescence. Generic architectures can waste capacity on degrees of freedom that are irrelevant from a physical point of view. These challenges motivate hybrid approaches in which as much known structure as possible—antisymmetry, cusp conditions, scaling, and symmetries—is built into the ansatz and its coordinates, and the network is used to learn the remaining correlation corrections.

A further ingredient concerns not *what* structure is imposed but *how* the network represents the electrons. Correlation is intrinsically relational—what one electron should do depends on where the others are—so architectures that let particles exchange information (message passing), rather than processing each coordinate independently, are a natural candidate for the correlator and for the backflow that deforms the nodes. How much this relational capacity actually buys, and why, is one of the questions this thesis takes up.

Where this work sits. Representing a quantum state by a neural network and optimising it variationally was introduced by Carleo and Troyer [1]; for continuum fermions the idea has since matured into highly accurate *ab initio* solvers such as FermiNet [2] and PauliNet [3], and into explicit neural *backflow* transformations [4]—the direct ancestor of the coordinate backflow used here. Message passing, the relational mechanism at the centre of this thesis, has recently been built into neural quantum states for extended fermion systems [5, 6]. Against that backdrop the present work is deliberately narrow and complementary. Rather than pushing a new architecture to ever larger systems, it asks *why* the relational channel helps, and uses the tangent-kernel geometry to make the answer precise; it focuses on the two-dimensional quantum-dot Wigner regime, where correlation is strongest and where topology-resolved structural diagnostics remain sparse; and it treats conditioning, the optimiser, and Markov-chain-free training as objects of study in their own right rather than fixed choices. To our knowledge these ques-

tions have not previously been addressed together for parabolic quantum dots.

Quantum dots as a controlled testbed. Two-dimensional parabolic quantum dots provide a clean and tunable setting in which to explore such ideas. They offer a simple route from Fermi-liquid-like behaviour to Wigner-molecule physics. As the trap frequency ω decreases, the single-particle level spacing collapses while Coulomb repulsion becomes dominant. Electrons then tend to localize into ring or shell patterns. In the lab frame, rotational symmetry and quantum fluctuations smear these into smooth densities; in a co-rotating frame, one recovers crystalline-like structures whose topology depends on particle number. For $N=6$ and $N=12$, classical and semiclassical studies have identified preferred shellings such as (1, 5) and (3, 9) and mapped parts of the crossover [7–10].

From the perspective of this thesis, quantum dots are attractive for two reasons. First, they are genuinely non-trivial: Coulomb singularities, strong correlations and an actual Fermi-liquid-to-Wigner crossover appear already for moderate N . Second, they are still simple enough that accurate DMC references exist in many regimes and that detailed structural diagnostics (shell occupancies, bond order, virial ratios) are computationally feasible. For fully quantum dots, however, quantitative topology-resolved diagnostics (occupancy fractions, bond-orientational order, shell “stiffness”) and a systematic picture of the Wigner regime remain relatively sparse. In addition, it is not well understood how modern neural ansätze internally represent these states.

Scope and questions. In this thesis we revisit two-dimensional harmonic quantum dots with a Slater–Jastrow(+backflow) neural quantum state, pretrained on a physics-informed residual objective and refined variationally. The design is meant to be *safe by construction* for Laplacian-based objectives: coordinates are scaled to trap units ($\tilde{R} = \sqrt{\omega}R$), learned pair features are soft-core (avoiding spurious $1/r$ behaviour in derivatives), and the electron–electron cusp is enforced analytically rather than learned. On top of a Slater determinant of harmonic-oscillator orbitals, two learnable objects carry the correlation: a Jastrow correlator, which reshapes the amplitude, and a coordinate backflow, which reshapes the nodes. Each can be built either without message passing—encoding and pooling per-particle and pairwise features in a single pass, as in a DeepSet—or as a *message-passing* network in which the particles exchange and update information through explicit, transported edge state before a quantity is read out.

Once such an ansatz reaches reference accuracy—which, as we will see, a surprisingly compact one does—the remaining questions concern *why* its parts behave as they do, and what it can tell us about the electrons. We address four. The first three share an analytical centre: the geometry of the wavefunction’s log-derivatives, captured by the quantum geometric tensor and the neural tangent kernel.

1. **Architecture.** What does a message-passing network learn that a separable one cannot, given that both can be trained to the same energy? We compare the two forms in the correlator and in the backflow, holding everything else fixed.
2. **Optimiser.** When does natural-gradient (stochastic reconfiguration) preconditioning change the result, and when is it unnecessary or harmful?
3. **Paradigm.** Can an accurate wavefunction be trained *without* Markov-chain Monte Carlo, using importance-sampled collocation instead, and how far into the correlated regime does that reach before it fails?
4. **Physics.** Can the trained wavefunction serve as a scientific instrument for Wigner-molecule physics—shell topologies, angular order, and melting—and does its internal representation reflect that physics?

Diagnostics. Total energies cannot separate the models compared here, since both architectures train to the same energy. The measurements that can are of three kinds: structural diagnostics of the trained state (shell detection, bond-orientational order, angular Lindemann ratios), geometric diagnostics of the ansatz (the quantum geometric tensor and neural tangent kernel, the effective dimension of the tangent space, state overlaps, $\text{Var}(E_L)$), and sampling diagnostics for the Markov-chain-free route (effective sample size, the Pareto shape $\hat{k}$). All are post-processing on $|\Psi|^2$ samples, so the same analysis would apply to a classical or quantum-chemical wavefunction unchanged. Chapter 5 defines them.

Main contributions.

1. A compact Slater–Jastrow–backflow ansatz, conditioned in trap units with an analytic cusp and soft-core pair features, that reaches DMC-level energies for $N=2$ and 10^{-2} – $10^{-1}\%$ relative errors for $N \in \{6, 12, 20\}$ at $\omega \geq 0.1$, and extrapolates smoothly to $\omega = 10^{-3}$.
2. A tangent-space account of what message passing contributes. A message-passing correlator represents the ground state in about half as many effective tangent directions as a separable one at strong confinement, and keeps its leading directions inside a compact operator dictionary at *every* confinement where the separable network does not; a message-passing backflow retains rank exactly where a conventional one collapses to a single displacement mode and loses accuracy. Both effects are invisible in the total energy until weak confinement turns the geometric deficit into an energetic one.
3. The criterion under which natural-gradient preconditioning is worth its cost: the tangent metric must be anisotropic *and* still estimable. Conditioning alone is not sufficient—under $|\Psi|^2$ sampling the empirical metric is badly conditioned and SR buys nothing—and the same geometry bounds an importance-sampled collocation objective that trains with no Markov chain in the gradient, reaching a Wigner-regime state at $\omega = 0.0035$ once its proposal is built from measured shell geometry.
4. Topology-resolved diagnostics of the Wigner crossover: two-shell probabilities and inner-ring occupancy histograms, a polygon shell model fitted on held-out frames reproducing $g(r)$ to cosine similarity 0.987–0.999 against 0.936–0.996 for an angle-randomised null, and sector-resolved bond order and Lindemann indices read against their random-angle baselines. At $\omega = 10^{-3}$ the $N=6$ dot shows persistent mixing between the (1,5) and (0,6) sectors and the $N=12$ dot forms a two-shell molecule with a commensurate (3,9) split, complementing classical and quantum studies of parabolic dots [7–9, 11].

Thesis structure. Chapters 1 and 2 set out the physics and the machine-learning background the later chapters require. Chapters 3 to 5 give the ansatz, the optimisation pipeline, and the diagnostics. Chapter 6 reports the energies and the Wigner-molecule structure; Chapter 7 analyses architecture, optimisation and sampling through the tangent geometry, and reports the Markov-chain-free training programme. Part IV opens with a short guide to how the two fit together. A discussion and a conclusion follow, and the appendices develop the conditioning theory, the collocation–backflow obstruction that motivated the weak-form objective, and the full diagnostic tables.

Part II

Theory

Men do not know how what is at variance agrees with itself. It is an attunement of opposite tensions, like that of the bow and the lyre.

HERACLITUS
Fragments

Chapter 1

The many-body problem and its numerical difficulty

The system is N electrons in a two-dimensional harmonic trap, repelling one another through the Coulomb interaction. Its Hamiltonian is elementary, and three of its properties make the numerical solution difficult.

The electrons are identical fermions, so the wavefunction changes sign under exchange, and that sign structure is the expensive part of the problem. The interaction diverges where two electrons meet, so the local energy $\hat{H}\Psi/\Psi$ diverges with it unless the wavefunction is constructed to cancel the divergence. And the kinetic term is a second derivative, which exposes any method that evaluates the energy pointwise to the curvature of the function it is given.

1.1 The many-body problem

A quantum state is a normalised vector in a complex Hilbert space, and in the coordinate representation $\psi(x) = \langle x|\psi\rangle$ is the wavefunction. Which basis one expands in is not a formality: choosing it to match the physics—harmonic-oscillator orbitals for confined charges, plane waves for periodic systems—is what makes a representation compact, and it is why the orbitals of Chapter 3 are those of the trap itself.

For N particles the space is the N -fold tensor product $\mathcal{H}_1^{\otimes N}$, and a fermionic state expands exactly in Slater determinants built from N distinct spin-orbitals, $|\Psi\rangle = \sum_I C_I |\Phi_I\rangle$. That expansion is exact and, as §1.2 makes precise, unaffordable.

We work in atomic units ($\hbar = m_e = e = 1$) with the many-body Hamiltonian

$$H = -\frac{1}{2} \sum_{i=1}^N \nabla_{\mathbf{r}_i}^2 + \sum_{i=1}^N V_{\text{ext}}(\mathbf{r}_i) + \sum_{i<j} V_{\text{int}}(|\mathbf{r}_i - \mathbf{r}_j|), \quad H\Psi(R) = E\Psi(R), \quad (1.1)$$

where $R = (\mathbf{r}_1, \dots, \mathbf{r}_N)$ and Ψ is normalised. Identical particles restrict Ψ to a symmetry sector, anti-symmetric for the fermions studied here. For these quantum dots $V_{\text{ext}}(\mathbf{r}) = \frac{1}{2}\omega^2 r^2$ and $V_{\text{int}}(r) = 1/r$ in two dimensions. Everything that follows approximates the lowest eigenpair of Eq. (1.1).

1.2 Correlation, closed shells, and the cost of exactness

Three ideas from electronic structure are used throughout this thesis and computed nowhere in it. They are what the accuracy claims of later chapters are calibrated against.

number is an estimate of it and carries time-step, population-control and finite-statistics errors of its own, which is part of why comparing a variational energy against one at the fifth decimal place requires the matched uncertainty analysis we did not undertake (§8.1).

Three consequences follow. A published DMC energy is not ground truth; it is the best energy attainable inside the nodal surface of the trial state used to produce it. A variational state with better nodes than that surface may therefore lie *below* a fixed-node DMC number without violating anything, since the variational principle bounds the energy from below by E_0 and not by another approximation (§8.1). And improving nodes becomes the thing to attempt, which is what the coordinate backflow of Chapter 3 is for.

1.5.3 Requirements on the trial state

Both methods take a trial state as given. VMC is only as good as the state it optimises, and fixed-node DMC only as good as the nodes it is handed, so the quality of the ansatz stops being an implementation detail and becomes the object of study. Three requirements bear on it at once.

Antisymmetry has to hold exactly and by construction, since the sign structure is what makes the problem hard in the first place. In practice that means a determinant, multiplied by a symmetric correlation factor of the kind Jastrow introduced for strongly interacting systems [18]. The nodes have to be placed well, since that is where the remaining systematic error lives once the amplitude is right. And the state has to be smooth enough that $E_L = \hat{H}\Psi/\Psi$, containing as it does a Laplacian and a potential that diverges at coalescence, can be evaluated without the two singularities amplifying one another. The cusp conditions of §1.3.4 say that those two divergences can be made to cancel exactly, and hard-wiring that cancellation is the first design decision of Chapter 3.

The remaining design choice is the functional form of the smooth correlation factor. The next chapter takes it from a literature that grew up asking different questions.

2.2.1 First-order methods

Gradient descent, $\theta^{(t+1)} = \theta^{(t)} - \eta \nabla_{\theta} \hat{\mathcal{R}}(\theta^{(t)})$, moves along the direction of steepest local decrease; too small a learning rate η converges slowly and too large a one diverges. Evaluating the full gradient is usually unaffordable, so the estimate is formed on a mini-batch, which is unbiased and whose variance sets the noise scale of the trajectory [21]. Momentum accumulates a running average of past gradients, damping oscillation across narrow directions and building displacement along consistent ones. Adam [22] adds a per-parameter normalisation by a running second moment,

$$m_t = \beta_1 m_{t-1} + (1 - \beta_1) g_t, \quad v_t = \beta_2 v_{t-1} + (1 - \beta_2) g_t^2, \quad \theta^{(t+1)} = \theta^{(t)} - \eta \frac{\hat{m}_t}{\sqrt{\hat{v}_t + \epsilon}}, \quad (2.2)$$

with g_t the mini-batch gradient and $\hat{m}_t, \hat{v}_t$ bias-corrected. That normalisation is a diagonal rescaling of parameter space, and the question of whether a non-diagonal one is worth its cost is what §2.2.2 and, empirically, §7.2 are about. Adam is the default optimiser in this work [23].

2.2.2 Second-order methods and natural gradient

First-order methods only use gradient information. In principle, convergence can be accelerated by incorporating curvature via the Hessian $H(\theta) = \nabla_{\theta}^2 \hat{\mathcal{R}}(\theta)$. Newton's method updates parameters as

$$\theta^{(t+1)} = \theta^{(t)} - H(\theta^{(t)})^{-1} \nabla_{\theta} \hat{\mathcal{R}}(\theta^{(t)}). \quad (2.3)$$

However, computing and inverting the Hessian is usually infeasible for large models.

An attractive alternative is the *natural gradient* [24, 25], which replaces the Euclidean metric with the geometry of probability distributions. The key object is the *Fisher information matrix* (FIM),

$$F(\theta) = \mathbb{E}[\nabla_{\theta} \log p_{\theta}(x) \nabla_{\theta} \log p_{\theta}(x)^{\top}], \quad (2.4)$$

where p_{θ} denotes the model distribution. The natural gradient update is

$$\theta^{(t+1)} = \theta^{(t)} - \eta F(\theta^{(t)})^{-1} \nabla_{\theta} \hat{\mathcal{R}}(\theta^{(t)}). \quad (2.5)$$

This preconditioning makes updates invariant to reparameterizations of θ and often improves conditioning of the optimisation. In practice, computing F^{-1} exactly is intractable, but approximations (e.g. stochastic reconfiguration in variational Monte Carlo, Kronecker-factored approximations in deep learning) make the method viable [2].

2.2.3 The variational tangent space: geometric tensor, tangent kernel, and effective dimension

For a variational quantum state the Fisher metric takes a particularly concrete form, and that form is the analytical lens used throughout the results of this thesis. Write the per-sample log-derivatives—the *score*—of an ansatz Ψ_{θ} as

$$O_{ki} = \frac{\partial \log |\Psi_{\theta}(\mathbf{x}_k)|}{\partial \theta_i}, \quad k = 1, \dots, B, \quad i = 1, \dots, P, \quad (2.6)$$

for a batch of B configurations drawn from $|\Psi_{\theta}|^2$ and P parameters θ (here θ collects *all* variational parameters). Its centred Gram matrices define two dual objects. Throughout, O is taken *column-centred*, $O_{ki} \mapsto O_{ki} - \frac{1}{B} \sum_{k'} O_{k'i}$, which is what makes the Gram matrix a covariance rather than a second moment and is how it is formed in the implementation (§4.4). The *quantum geometric tensor* (QGT) is then

$$S = \frac{1}{B} O^{\top} O \in \mathbb{R}^{P \times P}, \quad (2.7)$$

is the Fisher–Rao metric of the state written in terms of the amplitude score. Every state studied here is real and positive up to the determinant sign, at zero magnetic field, so the amplitude score is the whole story; for a general complex wavefunction the quantum geometric tensor also carries phase sensitivity, and the identification with the classical Fisher matrix is then only of its real part. It is proportional to the Fisher matrix $F(\theta)$ of the previous subsection specialised to $p_{\theta} = |\Psi_{\theta}|^2$: because $\nabla_{\theta} \log |\Psi_{\theta}|^2 = 2 \nabla_{\theta} \log |\Psi_{\theta}|$, one has $F = 4S$, the factor of four being a convention absorbed into the learning rate.

Stochastic reconfiguration is natural-gradient descent under this metric, $\dot{\theta} = -S^{-1}\nabla_{\theta}E$ [26, 27]. Its Gram dual, the *neural tangent kernel* (NTK) [28],

$$K = OO^{\top} \in \mathbb{R}^{B \times B}, \quad (2.8)$$

shares the nonzero spectrum of S up to the overall factor B (its nonzero eigenvalues are $B\lambda_a$), and is cheaper to diagonalise when $B < P$.

The analysis is organised around two scalars read from that spectrum. The *condition number* $\kappa(S) = \lambda_{\max}/\lambda_{\min}$ measures how anisotropic the tangent metric is, and therefore how far a natural-gradient step departs from a Euclidean one. It is the quantity one would expect to decide when SR is worth its cost; Chapter 7 measures it and finds that it does not, which is how the estimability criterion of that chapter arises. The *effective dimension*

$$d_{\text{eff}}(S) = \frac{(\sum_a \lambda_a)^2}{\sum_a \lambda_a^2} \quad (2.9)$$

is the participation ratio of the spectrum: it counts how many independent tangent directions carry appreciable Fisher weight locally around the solution, regardless of the raw parameter count P . A small d_{eff} means the local tangent sensitivity is concentrated in a few collective directions; a large one means it is spread across many. Because these are properties of the tangent geometry rather than of the raw parameter count, they permit architectures, optimisers and training paradigms to be compared on a common footing—provided the comparison is made under matched conventions. In the comparisons of §7.1.1 the measured d_{eff} proved stable across seeds and across the choice of probe measure; that is an empirical finding at those settings, not a general property.

The participation ratio d_{eff} is parameterisation-dependent. It is computed from the QGT eigenvalues in the ordinary Euclidean parameter coordinates, and those eigenvalues are *not* invariant under an arbitrary reparameterization: rescaling a single parameter $\theta_1 \mapsto c\theta_1$ leaves the physical state untouched but changes d_{eff} (for a diagonal metric it moves from 2 toward 1 as $c \rightarrow \infty$). d_{eff} is therefore *not* a coordinate-free intrinsic dimension of the wavefunction; it is a property of the (architecture, parameterisation, normalisation) triple. We use it accordingly—as a comparison between fixed architectures under matched conventions and a common probe measure—and read a smaller d_{eff} as “this architecture concentrates its Fisher weight into fewer active directions,” not as an absolute manifold dimension.

2.2.4 Tangent-space diagnostics

Convergence speed is not what matters about the optimiser here. A natural-gradient step is only as good as the metric it inverts, so the estimator of S becomes part of the method rather than an implementation detail, and Chapter 7 turns that into a three-regime result. Separately, $\kappa(S)$ and d_{eff} can be read off a trained state, which makes them diagnostics as much as convergence aids, and they are used that way from Chapter 6 onward.

2.3 Smooth networks for Laplacian objectives

A feed-forward network composes affine maps with elementwise nonlinearities,

$$h^{(0)} = x, \quad h^{(\ell)} = \sigma(W^{(\ell)}h^{(\ell-1)} + b^{(\ell)}), \quad f_{\theta}(x) = W^{(L)}h^{(L-1)} + b^{(L)},$$

and without the nonlinearity σ the composition collapses to a single affine map. For most applications the choice of σ is settled by convention or by a sweep. Here the objective settles it, because the local energy $E_L = \hat{H}\Psi_{\theta}/\Psi_{\theta}$ contains a Laplacian and training therefore backpropagates through second derivatives of the network.

The activation. $\text{ReLU}(x) = \max(0, x)$ has a second derivative that vanishes almost everywhere and is distributional at its kinks, so it cannot represent the pointwise curvature a Laplacian-based local energy asks for; it is excluded at once. Less obviously, being C^{∞} does not suffice either. The relevant property is the reach of the higher derivatives: an activation whose second and third derivatives decay quickly away from the origin contributes little curvature over most of its input range, so the Laplacian of a deep composition stays bounded and its variance finite. Tanh and the logistic sigmoid are perfectly smooth, yet their higher derivatives are sharply peaked and change sign, and curvature accumulates unevenly through

depth. We use the $\text{SiLU}(x) = x \sigma(x)$ family instead. Figure 2.1 shows GELU and Mish, which are the members of that family whose derivative shapes we compared and which SiLU closely resembles; their second and third derivatives are broader and decay more slowly than those of tanh or the logistic sigmoid. This is a design heuristic rather than a theorem: the figure establishes the shapes of the derivatives, not a bound on the Laplacian variance of a deep composition, and we did not run an activation ablation.




Figure 2.1: Common activation functions and their first three derivatives. All of GELU, Mish, tanh and the logistic sigmoid are C^∞ , so the distinction is not smoothness but the *reach* of the higher derivatives: for GELU and Mish the second and third derivatives are broad and decay slowly to zero, while for tanh and sigmoid they are sharply peaked near the origin and change sign, so curvature accumulates unevenly when such units are composed through depth. ReLU is shown for contrast; its second derivative is a distribution and it cannot be used in a Laplacian-based objective at all.

Initialisation. The standard schemes—Xavier/Glorot, scaling weight variance as $1/(n_{\text{in}} + n_{\text{out}})$, and Kaiming/He, scaling as $2/n_{\text{in}}$ for rectified units—are designed to keep forward activations and backward gradients from shrinking or exploding through depth [29, 30]. That is necessary but not sufficient when the loss contains second derivatives, because a scheme that stabilises $\partial f/\partial x$ says nothing directly about $\partial^2 f/\partial x^2$. We use Xavier-style initialisation with the smooth activations above, and—more consequentially than any initialisation choice—zero-initialise the final layer of the backflow, so that the ansatz begins exactly at the Slater nodes and the learned nodal deformation grows from nothing rather than from noise (§3.5.3).

None of this makes the Laplacian well conditioned; it only stops the network from making it worse. The conditioning itself comes from the coordinate scaling, the soft-core pair features and the analytic cusp of Chapter 3, and is analysed in Appendices A and B.

2.4 Architectures for many-body wavefunctions

The feed-forward network of the previous section maps a fixed-size input vector to an output. A many-body wavefunction is not a generic function of such a vector: identical particles impose a permutation symmetry, and correlation is intrinsically *relational*—what one particle should do depends on where the others are. A plain multilayer perceptron acting on the concatenated coordinates $(\mathbf{x}_1, \dots, \mathbf{x}_N)$ respects neither property; it would have to learn the symmetry from data, and its parameter count would grow with

N . This section introduces the two architecture classes that build the required structure in—permutation-invariant pooling and message passing—and situates the specific message-passing network used in this thesis as one instance of the latter.

2.4.1 Permutation symmetry and equivariance

For identical particles the modulus $|\Psi|$ is invariant under exchange (the sign is carried by the antisymmetric Slater reference of Chapter 3). The two learnable objects of the ansatz therefore have distinct symmetry requirements. A scalar *correlator* f must be permutation-*invariant*,

$$f(P_\sigma R) = f(R),$$

while a per-particle *backflow* displacement field $\mathbf{g} = (\mathbf{g}_1, \dots, \mathbf{g}_N)$ must be permutation-*equivariant*,

$$\mathbf{g}(P_\sigma R) = P_\sigma \mathbf{g}(R),$$

for any permutation σ (here P_σ relabels the particles). Enforcing these by construction is both more data-efficient than learning them and yields a parameter count independent of N , because the same shared maps are applied to every particle and pair.

2.4.2 Permutation-invariant pooling: DeepSets

The simplest way to build an invariant function is to embed each particle independently and pool the embeddings symmetrically. This is the DeepSets construction [31]: any suitably regular permutation-invariant function admits the form

$$f(R) = \rho\left(\sum_i \phi(\mathbf{x}_i)\right),$$

with a per-particle embedding ϕ , a symmetric pool $\sum_i$, and a readout ρ . A two-body extension encodes pairs, $\phi(\mathbf{x}_i, \mathbf{x}_j)$, and pools over them, capturing pairwise correlation. What characterises this class is that each particle—or pair—is embedded *independently* and only then pooled: no information is exchanged between particles before the pool. We refer to such networks as *separable*, and they serve as the non-message-passing baseline for the correlator.

2.4.3 Message passing and graph networks

To let a particle’s representation depend on the others’ in an iterated, higher-order way, one places the particles on a graph—nodes are particles, edges are pairs—and passes learned messages along the edges [32, 33]. In a generic round ℓ , each edge gathers information from its endpoint nodes and each node aggregates its incoming edge messages,

$$\mathbf{m}_{ij}^{(\ell)} = \phi_e(\mathbf{h}_i^{(\ell)}, \mathbf{h}_j^{(\ell)}, \mathbf{e}_{ij}^{(\ell)}), \quad \mathbf{h}_i^{(\ell+1)} = \phi_v\left(\mathbf{h}_i^{(\ell)}, \text{Agg}_{j \neq i} \mathbf{m}_{ij}^{(\ell)}\right),$$

with shared maps ϕ_e, ϕ_v and a symmetric aggregation Agg , so that permutation equivariance holds by construction; a final readout produces an invariant scalar (correlator) or an equivariant field (backflow). Both DeepSets and message passing use pairwise information, but only message passing lets that information *propagate*: after several rounds a particle’s representation reflects not just its neighbours but their neighbours’ responses. This iterated propagation is what we call the *relational channel*, and asking what it buys—in the correlator and in the backflow—is one of the central questions of this thesis. Message passing has been used in exactly this spirit for neural quantum states of extended fermion systems [5, 6].

2.4.4 The CTNN: one message-passing instance

The message-passing correlator and backflow studied here are realised by a particular network we refer to as the *CTNN*: a copresheaf-style message-passing network that maintains explicit *node and edge* features and transports information between them through learned maps, rather than recomputing and discarding one-shot messages. The correlator iterates this exchange over a down/up “V-cycle”; the backflow applies a single such copresheaf round. Its concrete update equations, hidden dimensions, and parameter counts are specified in the Methods chapter (§3.5.2, Eqs. (3.13)–(3.15)). The framing matters in two respects. First, the CTNN is *one* concrete realisation of the message-passing class defined above, and the comparison run in this thesis is between two specific inductive biases—the CTNN against a

DeepSets-type separable baseline—not between the classes they belong to. Conclusions are stated for message passing *as instantiated by the CTNN, relative to the separable baseline studied here*; whether they generalise to other message-passing neural quantum states is a plausible extrapolation and not something these experiments test. Wherever we contrast a “message-passing” network with a “separable” one, the separable member is a DeepSets-type network and the message-passing member is the CTNN. Second, we write “copresheaf-style” deliberately: the name reflects the node/edge transport structure of the updates, not a claim to a formal sheaf-theoretic construction.

2.4.5 Relevance to the ansatz

Both learnable objects of the ansatz—the Jastrow correlator and the coordinate backflow—can be built in either form: as a separable DeepSets-type network, or as the message-passing CTNN. Holding everything else fixed and varying only this choice isolates the effect of the relational channel, which Chapter 7 analyses through the geometry of the tangent kernel introduced in §2.2.3.

2.5 Approximation theory and its limits

The usual justification for using a neural network as a variational ansatz is that it can represent anything. That argument needs disposing of early, since this thesis is about what happens after it stops being relevant.

The universal approximation theorems say that a single hidden layer with a non-polynomial activation is dense in $C(K)$ on compacta: for any continuous f and any $\varepsilon > 0$ there exist a width and a parameter vector achieving $\sup_{x \in K} |f(x) - f_\theta(x)| < \varepsilon$ [34–36]. They establish one thing, best put negatively: there is no *fundamental* representability obstruction. They say nothing about the particular 20k–236k networks used here, nor about whether gradient-based training will find such a representation, at what sample cost, or how the answer depends on the parameterisation. The existence proofs are non-constructive, the widths they guarantee are astronomical, and the same function can be represented by two networks whose optimisation behaviour is entirely different—which is, in miniature, the comparison Chapter 7 makes.

One further result does more than assert possibility. In high dimension a generic continuous target requires a network size growing exponentially in d , so approximation alone cannot explain why these methods work at all in Nd dimensions. The escapes are all statements about *structure* rather than capacity: targets of finite Barron norm admit dimension-independent rates [37]; compositional targets admit efficient representation by deep networks when the composition matches the architecture; and imposing a symmetry removes the redundant copies the network would otherwise have to learn, reducing the burden on the representation rather than the dimension of the domain. The last of these is the one we exploit directly, and the middle one is why the message-passing architectures of §2.4 deserve attention: their usefulness comes from a compositional form matching the relational structure of the physics, not from added capacity.

The rest of the thesis therefore assumes representability and measures the other things: trainability, conditioning, and the geometry of the parameterisation.

2.6 Residual objectives and their difficulties

In scientific applications the governing equations already encode most of what is known about a system, and a model fitted only to data throws that information away. *Physics-informed neural networks* (PINNs) [38] put it back by penalising the residual of the equation itself: for a differential operator $\mathcal{N}$ and a network u_θ , the loss is a sampled sum of $|\mathcal{N}[u_\theta](\xi_j)|^2$ over collocation points $\{\xi_j\}$, together with boundary and data terms. Training a network against a differential operator predates the name by two decades [39]; what [38] added was automatic differentiation, which evaluates ∇u_θ and Δu_θ exactly and cheaply enough that the residual can drive the optimisation directly.

Four difficulties follow from differentiating through the operator, and each recurs by name in the results of this thesis [40].

Approximation in Sobolev norms. Universal approximation controls function error in L^2 or L^∞ , but a strong residual depends on *derivatives*. Making it small requires approximation in $W^{\ell,p}$ with ℓ the order

of the equation, which forces smooth activations for $\ell \geq 2$ and architecture choices that control higher derivatives. This is why the activation here is chosen for its second derivative rather than its values.

Residual-to-error stability. A small residual implies a small solution error only where the forward problem is stable. This is why a small residual is never taken here as evidence of a good state without an independent energy evaluation.

Sampling. Collocation minimises a sampled surrogate of a continuum loss, and the empirical residual can be small while the population residual is not. In this setting the question becomes whether an importance-weighted estimator has a finite variance at all, which is what bounds the Markov-chain-free programme of Chapter 7.

Conditioning. Residual losses couple terms of disparate scale and produce ill-conditioned gradient dynamics; in the linearised view, convergence is governed by the spectrum of a problem-dependent kernel. That spectrum is the subject of the entire tangent-kernel analysis.

The last of these determined the design of the collocation programme. A strong-form residual that differentiates through the Laplacian meets a structural obstruction as soon as it is asked to train a nodal surface, because the learning signal lives where E_L diverges. The weak form keeps the local energy as a detached reward and never differentiates it, and so avoids the obstruction; Appendix C shows why, and it is the reason the collocation programme takes the form it does.

2.7 Constraints carried into the method

The two traditions of this chapter and the last have been describing the same objects in different words. The quantum geometric tensor of variational Monte Carlo is the Fisher metric of information geometry; stochastic reconfiguration is natural-gradient descent under it; the neural tangent kernel is that metric's Gram dual; and the sampling measure a physicist chooses for a Metropolis chain is what a statistician calls a proposal distribution. Neither community needs the other's vocabulary in order to do its own work, which is why the identifications are easy to miss. They matter here because the architecture determines the spectrum of that metric, the optimiser inverts it, and the sampling measure determines how well it can be estimated.

Four properties of this problem bear on that metric.

The objective contains a second-order operator. The local energy carries a Laplacian, so training differentiates through second derivatives of the network. This is what dictates smooth activations, controlled initialisation, and a determinant that is never asked to do anything violent (§2.3).

The interaction is singular, and the state carries nodes. The Coulomb term diverges at coalescence, and the trial function vanishes on a surface whose location is itself being learned. Both are places where $E_L = \hat{H}\Psi/\Psi$ can blow up, and the cusp conditions of §1.3.4 say that the first divergence can be cancelled analytically if the short-range form is built in rather than learned.

The measure is part of the method. The objective is an expectation under $|\Psi_\theta|^2$, so the sampling distribution enters the gradient, the metric, and the estimate of both. Whether that metric can be estimated at all turns out to decide whether preconditioning helps or hurts (§2.2.3).

Correlation is relational. What one electron should do depends on where the others are, and the two architecture classes of §2.4 differ precisely in how far that dependence is allowed to propagate.

The first two are met by construction in Chapter 3, which hard-wires antisymmetry and the cusp and conditions the geometry so that the learned part can remain small and smooth. The third is treated in Chapter 4, where the sampling measure is a design variable rather than a fixed background. The fourth is not settled here: both architectures are constructed, and the comparison between them is reported in Chapter 7.

Part III

Methods

*No great thing is created suddenly, any more than a
bunch of grapes or a fig.*

EPICTETUS
Discourses

Chapter 3

Trial Wavefunction

The trial state is constructed to satisfy fermionic antisymmetry and the electron–electron cusp analytically, and to keep the learned terms smooth under the Laplacian. Antisymmetry requires a determinant. The Coulomb divergence admits an analytic cancellation, so it is imposed rather than learned. Second derivatives enter the loss, so the features fed to the network are bounded and smooth and the coordinates are scaled to keep those derivatives $\mathcal{O}(1)$ across three decades of confinement. The form of the learned part is not fixed by these constraints; both candidate architectures are constructed here and compared in Chapter 7. Unless stated otherwise we work in atomic units.

3.1 Preliminaries

Hamiltonian and units. We study N spin- $\frac{1}{2}$ fermions in two spatial dimensions confined by a harmonic trap and interacting via Coulomb repulsion,

$$H = \sum_{i=1}^N \left(-\frac{1}{2} \nabla_{\mathbf{r}_i}^2 + \frac{1}{2} \omega^2 r_i^2 \right) + \sum_{1 \leq i < j \leq N} \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|}, \quad (\hbar = m_e = e = 1). \quad (3.1)$$

Spin does not appear in H ; eigenstates factorize into a spatial function with good L_z and a spin function with good (S, S_z) .

Coordinates and scaling. We collect positions as $R = (\mathbf{r}_1, \dots, \mathbf{r}_N) \in \mathbb{R}^{2N}$ and use trap scaling $\tilde{\mathbf{r}}_i = \sqrt{\omega} \mathbf{r}_i$, $\tilde{R} = \sqrt{\omega} R$. This aligns typical length scales to $\mathcal{O}(1)$, improves conditioning of second derivatives, and makes the noninteracting ground state ω -independent. When useful, we also use pair distances $r_{ij} = |\mathbf{r}_i - \mathbf{r}_j|$ and their scaled versions $\tilde{r}_{ij} = |\tilde{\mathbf{r}}_i - \tilde{\mathbf{r}}_j|$.

Sectors and focus. We primarily report *closed-shell* cases (unique spin singlet at the noninteracting level). Open shells introduce a degenerate manifold and additional choices for the reference; we keep this to brief remarks only where needed.

Spin-orbitals and basis. One-particle spatial orbitals are the Cartesian two-dimensional harmonic-oscillator eigenstates $\{\phi_{n_x n_y}(\mathbf{r})\}$ with $E_{n_x n_y} = \omega(n_x + n_y + 1)$, $n_x, n_y \in \mathbb{N}_0$ (the basis used in the implementation; for the closed shells studied it spans the same filled shells as any complete basis of those shells). Spin-orbitals are $\varepsilon_\alpha(x) = \phi_{n_x n_y}(\mathbf{r}) \chi_\sigma(s)$, $\sigma \in \{\uparrow, \downarrow\}$.

3.2 Overview of the ansatz

3.6 Architecture sizes and hyperparameters

Table 3.1 collects the representative settings. The two comparisons in Chapter 7 stand on different footings in this respect.

The *correlator* comparison is capacity-controlled. The two variants are held at comparable parameter counts, so a difference in outcome cannot be attributed to one simply being larger, and where §7.1.1 departs from matched capacity it does so in the *separable* network’s favour, scaling it from 20k to 164k parameters against a fixed 80k message-passing network without closing the gap.

The *backflow* comparison is not. At equal width the copresheaf network is intrinsically the heavier object—it carries a persistent edge state where the conventional network carries only a per-particle aggregate—so at the shared production setting (message width 128, node MLP 128×3) it holds 182k parameters against 51k, a factor of 3.6. No matched backflow pair was trained. What separates architecture from capacity there is therefore not a matched control but the conventional network’s own behaviour across ω , which §7.1.2 sets out where the measurement is reported. All settings below are those of the accompanying code (`scripts/run_depth_analysis.py` for the comparison tiers; `scripts/source_state_check.py` for the production widths).

Table 3.1: The three sets of models trained in this thesis. They exist for different purposes and are never compared with one another. Widths are hidden dimensions; “rounds” are message-passing steps ($n_{\downarrow} + n_{\uparrow}$ for the V-cycle). Within the architecture study the correlator variants are not capacity-matched, which is why §7.1.1 also reports the matched 89.4k separable network and the full 20k–164k ladder; the backflow variants are not matched either, and cannot be at equal width, for the reason given above. Settings follow the accompanying code (`scripts/run_depth_analysis.py`, `scripts/source_state_check.py`).

Programme	Correlator	Backflow	Total
Physics production (§6.1)	separable; width 128, 2 layers; 54k	conventional; msg 128, node 128×3 ; 51k	105k
	(as above)	message-passing; same widths; 182k	236k
Architecture study (§7.1)	separable; pair-MLP 64×4 , out 32; 47.7k	as production; not matched	—
	message-passing; node/edge 32, bottle-neck 16, 2+2 rounds; 79.8k		—
Collocation (§7.3)	message-passing V-cycle; 25k	none	25k
	(as above)	copresheaf; $h_{bf}=128$; +24k	49k

Readout for every correlator: hidden 64, 2–3 layers, SiLU. Optimiser: Adam at $lr = 5 \times 10^{-4}$ (backflow) and 5×10^{-5} (Jastrow); CG-SR with Tikhonov damping λ , a trust-region cap and CG restarts. Collocation batch $n_{coll} \sim 3 \times 10^3$ with oversampling $m = 8$ –16; the collocation families are described further in §4.5.3.

4.7 The two workflows

Both routes start from the same ansatz and end at an energy evaluated under $|\Psi_\theta|^2$. They differ in the measure that defines the loss in between.

SR-VMC (§4.3–§4.4)

stratified mixture $\rightarrow$ residual pretraining
 $\downarrow$
 $|\Psi_\theta|^2$ sampling (Metropolis)
 $\downarrow$
 SR step $(S + \lambda I)\Delta\theta = -g$
 $\downarrow$
 energy under $|\Psi_\theta|^2$

Collocation (§4.5)

pretrained checkpoint
 $\downarrow$
 analytic proposal q , importance reweighting
 $\downarrow$
 weak-form REINFORCE gradient (Adam)
 $\downarrow$
 Metropolis probe for checkpoint choice,
 then energy under $|\Psi_\theta|^2$

The right-hand route places no Markov chain in any gradient; the probe that selects its checkpoints sits outside the update loop entirely. The left-hand route, which carries the ground-state energies of §6.1, keeps the early dynamics variance-dominated and sampler-aware and then transitions to a geometry-aware natural-gradient refinement—a division that matters most at weak traps, where the shell and dimer coverage of the stratified mixture is what makes the early stage converge at all.

4.8 Statistical uncertainties

Every energy and observable we report carries a statistical error, and two distinct sources contribute. Within a single evaluation, successive MCMC samples are autocorrelated, so a naive standard error over B samples understates the sampling uncertainty; this is what blocking addresses. Across training runs, the optimisation is itself stochastic and lands on different states from different seeds; blocking says nothing about that, and it is assessed separately by repeating runs at independent seeds wherever the claim depends on it (§7.3.4, §7.1.1). We therefore estimate the error on a scalar mean $\bar{A} = \frac{1}{B} \sum_i A(\mathbf{R}_i)$ by *blocking*: successive samples are averaged into blocks of growing size and the plateau of the block standard error as the block size exceeds the autocorrelation time is taken as the uncertainty [49, 50]. For the final ground-state energy this is the dominant reported error; a parenthetical digit, e.g. 155.8738(7), denotes one standard error on the last quoted figure.

For quantities that are *nonlinear* functions of sample statistics—participation ratios, effective ranks, PCA cosines, and the coupling diagnostics of Chapter 6—a blocked standard error is not available in closed form, and we use a *block bootstrap* instead: blocks (of the plateau size above) are resampled with replacement, the statistic is recomputed on each resample, and its spread across resamples is the reported error (§5.2.3). Where a number is quoted as a seed average, we additionally report the spread across independent random seeds, which captures optimisation—not just sampling—variability; the two are distinguished in the captions. Reference energies drawn from the literature carry the uncertainties quoted by their sources.

Chapter 5

Analysis

This chapter describes *what* we measure on fresh $|\Psi_\theta|^2$ samples, *how* we compute the diagnostics, and *why* each quantity is informative. Results and numbers are presented in Chapter 6.

5.1 Structural diagnostics and Wigner markers

Unless stated otherwise, curves are accumulated in *trap units* $\mathbf{y} = \sqrt{\omega} \mathbf{x}$ to compare shapes across ω , while all headline distances are reported in *Bohr* (physical units $\mathbf{x}$). Production chains are thinned; we split samples into halves for stability checks (acceptance window, JSD between halves, and reproducibility under restarts). Using trap units to compare shapes across confinements is standard in parabolic dots; see the overview in [10].

5.1.1 Pair distribution and radial statistics

We estimate an *effective* pair distribution—a shell-area-corrected pair-distance density—in trap units. For a finite trapped dot there is no homogeneous-system normalisation, because the one-body density varies strongly across the cloud, so A_{eff} below is a finite-dot convention rather than a thermodynamic volume and $g(r)$ does not tend to unity at large r . Peak positions and relative profiles, which is what the analysis uses, are insensitive to that choice; absolute amplitudes are not.

$$g(r_\alpha) = \frac{A_{\text{eff}}}{N(N-1)} \frac{\langle \sum_{i < j} \mathbf{1}\{r_{ij} \in [r_\alpha^-, r_\alpha^+]\} \rangle}{2\pi r_\alpha \Delta r_\alpha}, \quad r_{ij} = \|\mathbf{y}_i - \mathbf{y}_j\|_2, \quad (5.1)$$

with $A_{\text{eff}} = \pi R_{0.95}^2$ and $R_{0.95}$ the 95% quantile of single-particle radii in trap units. This $g(r)$ -based structural analysis follows established practice in mesoscopic 2D electron systems and quantum dots [7, 10, 11]. The pair-distance probability is therefore denoted $P_{\text{pair}}(r_{ij}) \propto r_{ij}^{d-1} g(r_{ij})$. It is used for pair-structure diagnostics, distinct from the one-body radial probability $P_1(r) \propto r \rho_1(r)$, from which we report the headline mode r_{mode} , mean $\bar{r}$, standard deviation σ_r , and FWHM (all in Bohr). A compact one-body width-to-scale marker is

$$\gamma = \sigma_r / r_{\text{mode}}, \quad (5.2)$$

which tracks the width of P_1 relative to its modal radius. It is interpreted alongside shell separation and pair/angular diagnostics, rather than as a universal monotone Wigner marker [7, 11].

5.1.2 Automatic shell detection and topologies

For each frame we sort the single-particle radii $s_1 \leq \dots \leq s_N$ (trap units), compute gaps $g_k = s_{k+1} - s_k$, and flag the frame as *two-shell* if the largest gap exceeds a robust threshold:

$$\max_k g_k \geq \tau \text{median}(g_k), \quad \tau \in [2.5, 3.5] \text{ (default 3.0)}. \quad (5.3)$$

The cut radius is the midpoint of the maximal gap; the inner multiplicity n_{in} defines the topology $(n_{\text{in}}, N - n_{\text{in}})$.

object for the question we ask of it—whether preconditioning by S^{-1} helps—but it means κ may not be compared across batch sizes, and we hold B fixed wherever we compare it.

To ask whether two trained models are the *same* state rather than merely equal in energy, we use a symmetric, importance-sampled squared overlap,

$$\mathcal{S}^2(A, B) = \langle \Psi_B / \Psi_A \rangle_{|\Psi_A|^2} \langle \Psi_A / \Psi_B \rangle_{|\Psi_B|^2},$$

each factor evaluated by a log-sum-exp mean over samples of the corresponding state. It equals 1 for identical states, and on the analytic two-electron ground state the trained ansatz reproduces $\mathcal{S}^2 = 1.000000$, which anchors the tooling.

One convention must be stated, because the ratio Ψ_B / Ψ_A is signed for a fermionic state. We evaluate the ratios from $\log |\Psi|$, so the estimator is strictly a *modulus* overlap, $\langle |\Psi_B / \Psi_A| \rangle \langle |\Psi_A / \Psi_B| \rangle$. For the comparisons made here this is an accurate proxy for the signed overlap: both states share the same Slater reference, and the correlator and cusp enter as a strictly positive exponential, so the sign of Ψ is fixed by the determinant and the two states agree in sign except on the measure-suppressed set where their backflows carry a configuration across a node differently. The modulus overlap is an *upper* bound on the signed one, so $\mathcal{S}^2 \rightarrow 0$ is conservative while $\mathcal{S}^2 \approx 1$ rests on the structural argument above rather than on the bound. Overlap, energy and $\text{Var}(E_L)$ are properties of the state and are unaffected by how correlation is divided between the correlator and the backflow; the energetic price of ablating a component is not, since two parameterisations that divide correlation differently report different ablation costs.

Part IV

Results

The argument of Part IV

The two results chapters run a loop between solver and physics.

We first establish that the trained states reproduce every external energy benchmark that exists for this system. That licenses using them as instruments rather than defending them as answers. Read for the physics they contain, they show a crossover from compact dot to finite Wigner molecule that is *staged*: the cloud expands and its shells stiffen radially well before the electrons acquire angular order within them.

We then turn from the state to the solver. Message passing changes the geometry of the variational representation even where the total energy cannot tell two networks apart—the correlator concentrates its tangent weight on fewer and more interpretable directions, and the message-passing backflow avoids a rank collapse that overtakes the conventional one. Architecture, optimiser and sampling measure then meet in a single object, the geometry of the wavefunction’s log-derivatives, and natural-gradient preconditioning earns its cost only while that geometry can still be estimated from a batch.

The loop closes when a generic collocation proposal loses the Wigner state. The structural measurements of Chapter 6 supply the remedy: a proposal built from the observed shell geometry carries Markov-chain-free training to $\omega = 0.0035$, where a transfer instability in the Slater–Jastrow balance takes over.

Three differently sized sets of models appear below—the production states that carry the physics, the matched ladder of the architecture study, and the smaller collocation networks—and numbers from one are never compared with numbers from another. Table 3.1 gives all three. The four research questions of §I are answered here:

RQ4 — Physics	§6.2
RQ1 — Architecture	§7.1
RQ2 — Optimiser	§7.2
RQ3 — Markov-chain-free paradigm	§7.3–§7.4




Figure 6.2: One-body densities for $N = 2, 6, 12$ at $\omega = 1.0$ (top) and $\omega = 0.001$ (bottom). The vertical axis is the one-body density $n(x, y)$ over the two Cartesian components of a single particle's position; the axis label carried in the figure files reads $n(r_1, r_2)$ for the same quantity, which collides with the convention used elsewhere in this thesis, where $\mathbf{r}_i$ is the position of particle i . As confinement weakens the densities broaden and develop pronounced *radial* shell structure. Being lab-frame averages over all orientations of the molecule, they cannot show angular order: that requires the registered diagnostics of §6.2.5.




Figure 6.3: One-body densities for $N=20$ at strong and weak confinement; the vertical axis is $n(x, y)$ as in Fig. 6.2. The weak-trap case shows strong radial broadening and clear multi-shell organisation. As above, this is radial evidence only.






Figure 6.4: Energy-based Wigner diagnostics as functions of trap frequency. As ω decreases the dots become interaction dominated ($T/V_{\text{int}} \ll 1$) and the trap–interaction partition approaches the classical virial form, most clearly for $N=12$.

Figure 6.5: Two-electron radial densities $P(r)$ across ω .Figure 6.6: Six-electron radial densities $P(r)$ across ω .

Chapter 7

Architecture, optimisation and sampling

The previous chapter showed that the ansatz works: it reaches reference quality wherever a reference exists, and the states it produces carry the staged Wigner-molecule structure the diagnostics were built to detect. Why the solver can represent those states, and what governs whether they can be trained at all, are separate and harder questions.

Three observations from that chapter remain unexplained. Two networks of the same size, trained the same way and landing on the same energy, need not be the same object. A natural-gradient step that is decisive in one training regime does nothing in another. And an importance-sampled objective reproduces a Markov chain’s answer in one corner of the phase diagram and departs from it in the next.

All three can be examined through the per-sample, column-centred log-derivative matrix

$$O_{ki} = \frac{\partial \log |\Psi_{\theta}(\mathbf{x}_k)|}{\partial \theta_i}, \quad k = 1, \dots, B, \quad i = 1, \dots, P, \quad (7.1)$$

and on its two Gram matrices, the quantum geometric tensor $S = O^{\top} O / B$ and the neural tangent kernel $K = O O^{\top}$ [27, 28]. That gives the chapter its structure:

Architecture determines the shape of the tangent space.

The *optimiser* acts on it through its metric.

The *sampling measure* determines how well that metric can be estimated.

The chapter takes them in that order. The instrument throughout is the *effective dimension* of the tangent space—the participation ratio of the QGT spectrum $\{\lambda_a\}$,

$$d_{\text{eff}}(S) = \frac{(\sum_a \lambda_a)^2}{\sum_a \lambda_a^2}, \quad (7.2)$$

which counts how many directions actually carry Fisher weight, whatever the parameter count says. Beside it sit the local-energy variance $\text{Var}(E_L)$, zero only for an exact eigenstate; the effective sample size; and the condition number $\kappa(S)$, which is reported in §7.2 and turns out to be large everywhere and therefore diagnostic of nothing by itself.

Architectures are compared on a common probe set—the same configurations for every model, never each model’s own $|\Psi|^2$ —and below $\omega = 0.1$ every relative error is measured against our own converged variational energies, so it reports internal consistency rather than accuracy. The tooling is anchored on the analytic two-electron ground state, where the ansatz returns $E = 3.000127$ Ha (+0.004%) at squared overlap 1.000000.

7.1 Message passing and the variational geometry

What does a message-passing network learn that a separable one cannot? Both learnable objects of the ansatz can be built in either form, so the question is asked twice, changing one object at a time. In the first comparison only the *correlator* is swapped—a separable DeepSets network against the message-passing CTNN—while the backflow, the Slater reference, the analytic cusp, the training protocol and the capacity tier all stay as they are. In the second only the *backflow* is swapped—single-round pairwise messages against the copresheaf CTNN—with the correlator now among the things held fixed. Neither comparison varies the two together.




Figure 7.4: The *fixed-measure instantaneous ablation cost* of each component (trained networks, $N=6$). Configurations are drawn once from the full model’s $|\Psi|^2$ and every arm is re-evaluated on them, so the trap and Coulomb terms are identical across arms by construction and the plotted ratio $\Delta E_{\text{probe}}/E_{\text{full}}$ is the instantaneous change in the kinetic term when a component is switched off. It is *not* the energy of a network retrained without that component, which would be smaller, since nothing is allowed to re-relax. Removing the backflow (cyan) costs $\sim 6\%$ at $\omega=1$ but under 0.02% at $\omega=0.01$, while removing the Jastrow messages (dark red) grows more expensive as the trap weakens, from 3.9% to 11.4% ; the two curves cross near $\omega \approx 0.4$. Removing all message passing (blue) tracks the Jastrow-message curve and meets it at $\omega=0.01$: once the correlator’s messages are gone, deleting the displacement as well costs nothing measurable. The relational work migrates from the nodes to the amplitude: at the crystal it is the correlator’s messages, not the backflow, that carry it. This is a statement about how *this* trained model divided the correlation between two substitutable components, and so is conditional on the staged training protocol rather than gauge-invariant (§7.1.2); the ablation costs themselves are reproducible at fixed protocol.

where CG-SR on the collocation objective becomes unstable (§7.3.3). SR pays off in the window where the metric is both ill-conditioned and estimable, and fails once estimation itself breaks down.

The VMC equivalence of Adam and SR is established here at $N = 2$ and $N = 6$, and we have not verified that it survives to $N = 12$ and $N = 20$. The mechanism we propose—that $|\Psi|^2$ sampling makes the metric *estimable* on the batch, whatever its conditioning, and so lets Adam’s diagonal preconditioning suffice—predicts that it should, since the argument turns on the sampling measure and not on dimension. But it is a prediction there, not a measurement, and testing it is the obvious extension. The same applies to the reverse reading of Table 7.4: κ is measured at one N and one ω , and we do not know how it moves with either.

7.2.2 Consequences of the sampling measure

Between $|\Psi|^2$ -sampled VMC and importance-sampled collocation the wavefunction family is identical; only the measure that defines the loss changes. That change has three consequences, and underneath them one cause.

Target sampling aligns the estimator with the metric; a covering proposal does not. Sampling from $|\Psi|^2$ makes the estimators of S and of the gradient share the measure that defines them, so the directions carrying Fisher weight are also the directions the batch populates. That is an alignment between estimator and objective, not preconditioning in the sense of applying S^{-1} . A fixed covering proposal instead drives a monotone *ESS collapse* as N and $1/\omega$ grow: at $N = 6, \omega = 0.01$ the effective sample size falls to 0.11% of the batch—about 5 of the 4096 retained points—and the gradient estimator is then degenerate. That is a hard limit, not undertraining. (Two ESS normalisations appear in this chapter, differing by the oversampling factor; each figure states which.)

§7.3.4 below nonetheless reports usable collocation energies at the same (N, ω) —+0.24–0.29% across three seeds, Table 7.7—and the two situations differ in the sampler and in the quantity being described. The sampler first: the collapse measured above is for the *static* origin-centred mixture, while the campaign’s robust recipe refits its proposal every 30 epochs and oversamples $16\times$. The campaign’s *baseline* rows, which keep the static proposal, are the ones that blow up (+3.9 to +7.2% at $\omega = 0.001$). Second, and more interesting, the two statements are not about the same quantity. The energy stays within a third of a percent while the state overlap ceases to be resolvable—which is the sharpest illustration in this thesis of why energy alone does not certify a state. We stop short of saying the overlap *is* zero: S^2 is a ratio estimator built from the same degenerate weights, so at this ESS it reports *unresolved* rather than orthogonal.




Figure 7.6: The naive collocation proposal collapses in the Wigner regime ($N=6$). The effective sample size (as a percentage of the batch) falls by more than an order of magnitude from $\omega=1$ to $\omega=0.01$, where it reaches $\sim 0.1\%$ —about five usable points in a batch of 4096. This is a property of the sampling measure, not of optimisation, and it is what the physics-informed ring proposal (§7.4) is designed to repair.

Table 7.6: Collocation variants tried, and what each one settled. Errors are relative to the reference of Table 7.5. The three marked † are taken up in prose below; the rest are recorded for completeness.

Attempt	Outcome	Mechanism
Pure REINFORCE ($\beta=0$) over the hybrid objective	adopted	The direct path omits the sampling-measure score contribution; at $\beta > 0$ it contributes a kinetic-only pathwise gradient that conflicts with REINFORCE, and the two gradients oscillate against each other.
Cascade warm-start from higher ω	essential	Cold starts never converged at $\omega=0.01$ for $N \geq 6$. Cascading from $\omega=0.1$ cut the initial error from $\sim 30\%$ to $\sim 5\%$ and made $+0.15\%$ reachable in ~ 1000 epochs.
ESS-adaptive oversampling with instability rollback	essential	Raises the proposal budget when overlap degrades and reverts on energy spikes. Without both, $\omega=0.001$ diverged within a few hundred epochs.
CG-SR below $\omega \simeq 0.01$ †	abandoned	$+4\text{--}36\%$ against $+0.15\text{--}0.24\%$ from Adam. Not ill-conditioning but unestimability; see below.
Langevin proposal refinement †	abandoned	Worse in every configuration tested ($+153\%$ against $+5.7\%$ at $N=20$, $\omega=0.1$). The weights are computed against a proposal the samples have stopped following; see below.
Finite-difference residual loss	abandoned	$+0.22\%$ at $N=6$, $\omega=1$ against $+0.010\%$ from REINFORCE. Sensitive to the step size and to noise in the Laplacian estimate.
Neural Pfaffian ansatz [48]	abandoned	$+0.43\%$ at $N=6$, $\omega=1$, forty times the BF+Jastrow error. The larger parameter space amplifies importance-sampling gradient noise faster than the richer nodal topology repays.
Backflow at $N=20$ †	abandoned	$+18\%$ against $+1.32\%$ for Jastrow-only. A memory limit rather than a conceptual one; see §7.3.5.

divide by q . The resulting bias is not noise. It pushes the wavefunction toward the regions the Langevin dynamics favours, which moves $\tilde{q}$ further the same way, and the loop closes on itself—which is what the $+153\%$ in the table is. Ten to twenty steps in $Nd=40$ dimensions do not equilibrate. Fixing it means running to equilibrium, adding a Metropolis–Hastings step, or computing annealed weights along the non-equilibrium path. The first two are MCMC under another name; the third costs enough to forfeit the advantage that made the route attractive.

What the failures have in common. Read together these are not separate disappointments. Four of them—CG-SR, Langevin, the Pfaffian’s parameter count, and the backflow at $N=20$ —fail through one channel: a quantity that must be *estimated* from a finite sample degrades faster than the quantity it was meant to improve. Natural gradients need an estimate of the metric. Langevin needs samples that have actually equilibrated. A larger parameter space needs a correspondingly better-resolved gradient. None of these methods is wrong in principle, and each would be sound given samples we cannot afford. What fails is the estimate, and it fails precisely in the regime where the correction was supposed to help. That thread is what §7.3.4 follows deliberately, from the proposal mismatch at one end to the evaluation bottleneck at the other.

7.3.4 Reliability analysis: the gradient-quality chain

The results in Table 7.5 report the *best* energies obtained by collocation training, selected from multiple runs on different seeds. A natural question is whether these results are *typical* or whether they reflect rare fortunate outcomes from a high-variance process. This subsection reports a systematic 30-run reliability campaign designed to answer that question, and exposes a mechanistic chain that accounts for both the successes and failures of the MCMC-free collocation approach.

Experimental design

All runs use the BF+Jastrow ansatz ($\sim 49k$ parameters) initialised from the same pretrained checkpoint. The grid is $N=6$ across $\omega \in \{1.0, 0.5, 0.1, 0.01, 0.001\}$ with three random seeds per cell (42, 137, 314),

Part V

Discussion

Convictions are more dangerous enemies of truth than lies.

FRIEDRICH NIETZSCHE
Human, All Too Human

Chapter 8

Discussion

8.1 Geometry between architecture and optimiser

The three parts of this thesis converge on one object. Architecture fixes the shape of the variational tangent space; the optimiser acts on that space through its metric; and the sampling measure decides how well the metric can be estimated. Read in that order, results that arrived separately become a single chain.

Its first link is the one the energy cannot see. Two correlators of matched or deliberately mismatched capacity reach the same reference-quality state at strong confinement, agreeing to four figures, and arrive there through tangent spaces of different size and different content. Counting directions is the weaker half of that observation—it says how many the correlator uses, and it is measured on one of two substitutable components. Which directions is the stronger half. The message-passing correlator’s leading tangent modes lie inside a four-operator physical dictionary to $R^2 \approx 1$ across confinement and up to $N=12$, where a separable network of equal or greater capacity leaves close to half its tangent budget outside it. A direction outside a chosen dictionary need not be unphysical, so what the measurement supports is that one tangent space is far more completely accounted for by a small set of collective operators than the other—and that a smaller d_{eff} makes such a dictionary easier to cover cannot explain the gap, since the two effective dimensions converge toward 3.7 at weak confinement while the dictionary gap persists.

The second link is where the optimiser meets that space. The result there is narrower than the usual one about natural gradients. Under $|\Psi|^2$ sampling the empirical metric is badly conditioned— $\kappa \sim 10^7$ – 10^{10} on precisely the models where Adam and SR agree to four decimals—so conditioning cannot be the criterion. What separates the regimes is whether a batch resolves the metric at all. Under $|\Psi|^2$ the samples come from the measure that defines S ; under a fixed collocation proposal they do not, and preconditioning becomes what makes the optimisation converge; and once the proposal loses the target, inverting the metric amplifies noise instead of signal. The negative half of that is measured. The positive half—that collocation under a fixed proposal occupies an anisotropic but still estimable middle regime—is inferred from the estimator, since κ was never obtained under the collocation measure at matched batch size.

Whether the chain generalises is a fair question and the honest answer is that we cannot settle it here. Message-passing neural quantum states have been adopted for extended fermion systems on the argument that relational structure scales [5, 6]; what this thesis adds is a measurement of *where* that structure shows up—in the tangent geometry before the energy—and a mechanism for why the effect should grow rather than shrink with correlation strength. The dictionary gap widens with N over the range we can measure. If that continues, the architectures that scale are those maintaining a transported edge representation rather than pooling pairwise features and discarding them. Three sizes is not a scaling study, and the comparison is seeded at $N=6$.

not a claim of freedom from Monte Carlo, since importance sampling and resampling are themselves Monte Carlo; nor of freedom from the reference, since where a DMC value exists the residual pretraining anneals toward it and the rollback monitors against it. Nor, at the deep-Wigner rungs where no reference exists, is the state reference-free in ancestry: its warm starts descend from benchmark-guided states at stronger confinement. Against the SR route, collocation is competitive for $N \lesssim 12$ and $\omega \gtrsim 0.1$, a few times worse at $N=12$, $\omega=0.1$, and not competitive at $N=20$. The two look complementary rather than rival: collocation for fast, parallel, chain-free exploration and pretraining; SR for final accuracy where correlation is strong.

8.3 Scope, and what to do next

Four limits bound everything above. All states are the lowest found within a fixed spin-projection sector, and total spin is not scanned. The architecture comparison is seeded at $N=6$, rests on fewer models at $N=12$ and on a single production state at $N=20$. The backflow variants are not capacity-matched and cannot be at equal width, so the cross-architecture energy gap does not isolate architecture. And one entry in the backflow rank table, at $N=20$ and $\omega = 0.01$, comes from a re-run whose diagnostics were not retained.

Four things follow that we would do next.

Make the proposal learned, not designed. Up to the frontier the proposal is the bottleneck and not the objective, which argues for a better sampler rather than a better loss: normalising flows trained against $|\Psi|^2$, or the shell-informed construction of §7.4 generalised beyond ring geometries. Past the frontier the target changes: what is needed there is an ℓ -covariant parameterisation, or an objective anchored on the shell amplitude, so that a converged state survives being carried to a weaker trap.

Make rotation equivariance exact. Permutation equivariance holds by construction and the centre of mass is projected out, but the displacement head reads a d -vector from a network that has seen absolute coordinates, so SO(2) equivariance is only encouraged—by relative-vector edge features and by the random in-plane rotation applied to every batch—and never enforced. In a rotationally invariant trap whose weak-confinement physics is a freely rotating molecule, that is the wrong symmetry to leave approximate. The fix is local to one module: build the displacement as $\Delta \mathbf{r}_i = \sum_{j \neq i} g_\beta(\text{invariants}_{ij}) \mathbf{r}_{ij}$, a learned scalar gain on each relative vector, so the output rotates exactly with the input and the augmentation can be dropped. We did not run it.

Break the quadratic backflow. What stops the method at twenty electrons is the $\mathcal{O}(N^2 h_{\text{bf}})$ edge tensor, a memory limit rather than a physical one. Sparse or low-rank message passing, a local backflow without all-to-all edge features, or the sample-side reductions of minSR and the kernel formulation [44, 45] would extend the reach without giving up the nodal correction that is the reason for having a backflow at all.

Match the backflow capacities. A widened conventional backflow at 182k parameters is the one control that would turn the rank-collapse observation into an architecture ablation. It is a single training run.

Part VI

Conclusion

*All things are implicated with one another, and the bond
is holy; and there is hardly anything unconnected with
any other thing.*

MARCUS AURELIUS
Meditations

Chapter 9

Conclusions

The work formed a feedback cycle between solver and physics. A variational state was constructed and checked against every external benchmark available; the checked state was then used as an instrument, and resolved the shell geometry of the Wigner regime—rings at definite occupancies, n -fold angular order becoming large only in the last decade of confinement, a shell radius approaching its classical $\omega^{-2/3}$ scaling. Those measurements are what a sampler needs in order to place its mass correctly. Incorporating the measured geometry into the proposal extended Markov-chain-free training from the failure of the origin-centred sampler at $\omega \approx 0.01$ down to $\omega = 0.0035$, into states the original method could not reach.

Architecture. Message passing supplies a relational channel whose effect appears in the tangent geometry before it appears in the energy (§7.1). At strong confinement a message-passing correlator reaches a reference-quality solution along half as many effective tangent directions as a separable network of matched or larger capacity, and its leading directions are far more completely accounted for by a compact dictionary of collective operators; the separable network spends close to half its tangent budget outside that dictionary, matches the energy to four figures anyway, and carries a local-energy variance higher by factors of 1.2 to 3.4. In the nodes the same channel shows as a collapse averted: toward the Wigner regime a conventional backflow’s displacement field collapses spectrally to rank one and loses half a percent of the energy, where the copresheaf field keeps its rank. Correlator and backflow give way at the same confinement, and a controlled correlator swap moves that boundary—which points to one relational bottleneck rather than two deficits.

Optimiser. Natural-gradient preconditioning earns its cost where the tangent metric is anisotropic and still estimable (§7.2). That is narrower than the usual claim and it has three regimes rather than two. Under $|\Psi|^2$ sampling the empirical metric is badly conditioned, $\kappa \sim 10^7$ – 10^{10} , on the very models where Adam and SR agree to four decimals, so conditioning alone is not the criterion. Under a fixed collocation proposal SR is what makes the optimisation work at all, until proposal adaptation and gradient stabilisation make Adam sufficient and more robust. And once the sampling that builds the metric collapses, inverting it amplifies noise rather than signal. Estimability, not conditioning, is what the data identify as decisive.

Paradigm. An accurate wavefunction can be trained with no Markov chain in the gradient, inside a regime whose edges we found by crossing them (§7.3). The weak-form objective reaches DMC accuracy at $N \in \{6, 12\}$ under strong confinement and stays within a few tenths of a percent of our own converged energies well into the correlated regime. It fails through one chain of causes—proposal mismatch, weight-tail inflation, gradient noise, an unestimable metric—which a better proposal moves but does not remove, and then through a second and different failure of transfer between confinements.

Physics. The trained state serves as an instrument (§6.2). The crossover it resolves is staged: the cloud expands and its shells stiffen radially well before the electrons acquire angular order within them. At $\omega = 10^{-3}$ the $N=6$ dot shows persistent mixing between the (1, 5) and (0, 6) sectors, the $N=12$ dot a two-shell molecule with a modal (3, 9) split, and the $N=20$ dot a three-shell molecule that collapses

to a single ring about a central electron within one decade in ω . The $N=6$ shell radius approaches the classical value $1.334\omega^{-2/3}$ monotonically, to within 1.7%.

The measurements then returned to the solver. Incorporating the observed shell geometry into the collocation proposal lifted the effective sample size by 15–32 $\times$ without any training and carried Markov-chain-free training from $\omega \approx 0.01$ to $\omega = 0.0035$ —into states the original sampler could not reach.

9.1 Limitations and further work

All states are the lowest found within a fixed spin-projection sector; total spin is not scanned. The architecture comparison is seeded at $N=6$ and thinner above it, and the two backflow variants are not capacity-matched, so the cross-architecture energy gap does not isolate architecture. Below $\omega = 0.0035$ the limit is a transfer instability in the Slater–Jastrow cancellation that no change of proposal repairs; whether the ansatz could represent such a state under per- ω re-optimisation is a question we did not test. The interventions most likely to move the picture are a backflow that is not quadratic in particle number, a proposal that adapts on its own, an exactly $\text{SO}(2)$ -equivariant displacement, and the capacity-matched backflow control that would settle the last question above. §8.3 sets these out.

9.2 Concluding perspective

At the widths reached for the well-resolved $N \leq 12$ studies, raw representational capacity was rarely the first bottleneck; alignment, conditioning and sampling were. It was not irrelevant everywhere. At $N=20$ the memory cost of the backflow’s edge tensor forces a capacity cut that degrades the collocation gradients. But what determined the outcome, across architecture, optimisation and sampling alike, was alignment with the low-dimensional collective structure the Hamiltonian already possesses: breathing, correlation-hole and relative modes at strong confinement, resolving at weak confinement into the shell geometry of the Wigner molecule. That the same alignment matters in all three places is probably a fact about the system rather than about neural networks.

Several of these results were reached only after an initial reading of the same data proved wrong: a rank collapse that survived seeding where a dimension inversion did not, a transfer failure first attributed to the sampler and later to the ansatz, and an importance-sampling error whose signature was indistinguishable from an architectural deficiency. Each was identified by a diagnostic that could separate the competing explanations. That is the methodological point these chapters have tried to make concrete: a variational calculation is only as trustworthy as the tests that can tell its failure modes apart.

Part VII

Appendices

Chapter A

Laplacian conditioning

A.1 Residual training for Schrödinger

We consider the stationary many-body Schrödinger equation on configuration space $\mathbf{R} = (\mathbf{r}_1, \dots, \mathbf{r}_N) \in \Omega \subset \mathbb{R}^D$ with $D = dN$ (for our quantum dots, $d = 2$):

$$\hat{H}\Psi(\mathbf{R}) = E\Psi(\mathbf{R}), \quad \hat{H} = \hat{T} + \hat{V}, \quad \hat{T} = -\frac{1}{2} \sum_{i=1}^N \Delta_i, \quad (\text{A.1})$$

where $\hat{V}$ collects external and interaction potentials.

In *strong residual minimization* one introduces

$$L := \hat{H} - E \quad \Rightarrow \quad L\Psi = 0, \quad (\text{A.2})$$

and minimizes a squared residual loss of the schematic form

$$\mathcal{L}_{\text{res}}(\theta) = \frac{1}{2} \|L\Psi_\theta\|_{L^2(\Omega; w)}^2 = \frac{1}{2} \int_{\Omega} |(L\Psi_\theta)(\mathbf{R})|^2 w(\mathbf{R}) d\mathbf{R}, \quad (\text{A.3})$$

for a nonnegative weight w induced by the sampling strategy. A key message of the numerical-analysis viewpoint is that strong residual losses can be *severely ill-conditioned*, and training speed is tightly linked to spectral properties of an operator/matrix induced by L [40].

Which of the thesis’s objectives this analyses. Three objectives appear in the main text and Eq. (A.3) is exactly one of them. Stage I pretraining (§4.3) minimises $\frac{1}{B} \sum_b (E_L(\tilde{R}_b) - E_{\text{eff}})^2$ over configurations drawn from $|\Psi_\theta|^2$; since $E_L = \hat{H}\Psi/\Psi$, that sample average estimates $\int |\Psi_\theta|^2 (\hat{H}\Psi_\theta/\Psi_\theta - E)^2 = \|(\hat{H} - E)\Psi_\theta\|_{L^2}^2$, which is Eq. (A.3) with $w \equiv 1$. Everything in this appendix therefore applies to stage I directly. It does *not* apply in the same way to the other two. Stage II is variational energy minimisation (§4.4), whose objective is linear rather than quadratic in the residual and whose conditioning is governed by the quantum geometric tensor measured in §7.2, not by the Δ^2 stiffness derived below. The collocation route (§4.5) replaces the strong residual with a weak-form score-function objective under an analytic proposal, for the reason developed in §A.4 and Appendix C: the weak form is what removes the second-derivative amplification, and it is the escape from the mechanism analysed here rather than an instance of it. Where the two production objectives inherit anything from this analysis, it is through the single channel they share with stage I—the Laplacian inside E_L —and not through the squared-residual structure.

A.2 Linearized geometry and the matrix A

To expose the mechanism, follow the standard linearization argument used in conditioning analyses [40]. Fix an expansion point θ_0 and define tangent (feature) functions

$$q_j(\mathbf{R}) := \partial_{\theta_j} \Psi_\theta(\mathbf{R}) \big|_{\theta=\theta_0}, \quad j = 1, \dots, P. \quad (\text{A.4})$$

For small parameter increments $\delta\theta$, we have the first-order model

$$\Psi_{\theta_0+\delta\theta}(\mathbf{R}) \approx \Psi_0(\mathbf{R}) + \sum_{j=1}^P q_j(\mathbf{R}) \delta\theta_j, \quad \Psi_0 := \Psi_{\theta_0}. \quad (\text{A.5})$$

Applying the residual operator gives

$$(L\Psi_{\theta_0+\delta\theta})(\mathbf{R}) \approx (L\Psi_0)(\mathbf{R}) + \sum_{j=1}^P (Lq_j)(\mathbf{R}) \delta\theta_j. \quad (\text{A.6})$$

Introduce the weighted inner product (complex-valued wavefunctions allowed)

$$\langle f, g \rangle_w := \int_{\Omega} f(\mathbf{R})^* g(\mathbf{R}) w(\mathbf{R}) d\mathbf{R}. \quad (\text{A.7})$$

Let the Jacobian J be the linear map whose j th column is Lq_j . Then the loss (A.3) becomes (to second order in $\delta\theta$)

$$\mathcal{L}_{\text{res}}(\theta_0 + \delta\theta) \approx \frac{1}{2} \|r_0 + J\delta\theta\|_w^2, \quad r_0 := L\Psi_0. \quad (\text{A.8})$$

with normal-equation matrix

$$A := J^* J, \quad A_{ij} = \langle Lq_i, Lq_j \rangle_w. \quad (\text{A.9})$$

Here A is Hermitian positive semidefinite; in practice it can be close to singular, so one often effectively works with a damped variant $A + \lambda I$ (implicit or explicit).

Now consider gradient descent on the quadratic model (A.8) (with step size η). The error in parameter space evolves as

$$\delta\theta^{(t+1)} - \delta\theta^* = (I - \eta A) (\delta\theta^{(t)} - \delta\theta^*), \quad (\text{A.10})$$

so each eigen-direction of A contracts by a factor $(1 - \eta\lambda)$. Hence convergence speed is governed by the eigenvalue spread of A , typically summarized by the (effective) condition number

$$\kappa(A) = \frac{\lambda_{\max}(A)}{\lambda_{\min}(A)} \quad (\text{A.11})$$

(over the nonzero/regularized spectrum). Large $\kappa(A)$ implies stiffness: a step size small enough to be stable for $\lambda_{\max}$ yields slow progress along directions associated with $\lambda_{\min}$. This is precisely the conditioning bottleneck emphasized in physics-informed training analyses [40].

A.3 Bi-Laplacian scaling

The defining feature of Schrödinger is that L contains a second-order differential operator. To isolate the kinetic contribution, consider the regime where the residual is dominated by the Laplacian part,

$$L \approx -\frac{1}{2} \Delta_{\mathbf{R}}. \quad (\text{A.12})$$

From L to L^*L . The matrix A in (A.9) is a Gram matrix in the feature space $\{Lq_j\}$. The corresponding operator-level object is the *Hermitian square* L^*L (with adjoint taken in the $L^2(\Omega; w)$ inner product) [40]. For general non-constant w , the adjoint L^* differs from the formal differential adjoint by weight-induced lower-order terms; the cleanest picture is obtained for constant w and standard boundary conditions (periodic/Dirichlet), where the Laplacian is self-adjoint.

In the Laplacian-dominant regime (A.12), one has (up to a constant factor)

$$L \approx -\frac{1}{2} \Delta \quad \Rightarrow \quad L^*L \approx \frac{1}{4} \Delta^2, \quad (\text{A.13})$$

i.e. a bi-Laplacian scaling. The prefactor does not affect conditioning ratios.

Fourier/eigenmode derivation of k^4 . On a periodic domain (or, more generally, in a Laplacian eigenbasis) $-\Delta\phi_{\mathbf{k}} = |\mathbf{k}|^2\phi_{\mathbf{k}}$, so $\Delta^2\phi_{\mathbf{k}} = |\mathbf{k}|^4\phi_{\mathbf{k}}$. If an error $\delta\Psi$ expands as $\delta\Psi = \sum_{\mathbf{k}} c_{\mathbf{k}}\phi_{\mathbf{k}}$ and $L \approx -\frac{1}{2}\Delta$, then

$$\|L\delta\Psi\|_{L^2}^2 \approx \frac{1}{4} \sum_{\mathbf{k}} |\mathbf{k}|^4 |c_{\mathbf{k}}|^2. \quad (\text{A.14})$$

Thus the squared residual penalizes high-frequency components with a *quartic* weight $|\mathbf{k}|^4$. If the relevant spectrum spans $|\mathbf{k}| \in [k_{\min}, k_{\max}]$, the induced eigenvalue ratio scales like

$$\kappa \sim \left(\frac{k_{\max}}{k_{\min}} \right)^4, \quad (\text{A.15})$$

which is a sharp mathematical expression of “Laplacian-driven stiffness”. This quartic growth is consistent with the Laplacian/Poisson toy examples and frequency-dependent conditioning effects emphasized in [40].

Length-scale interpretation. Let ℓ denote the smallest length scale that must be represented. Then $k_{\max} \sim 1/\ell$ and (A.15) implies

$$\kappa \text{ grows rapidly as } \ell \downarrow 0. \quad (\text{A.16})$$

In many-body Schrödinger problems, small ℓ arises from nodal structure, strong localization, and short-range correlation features; in all such regimes, the Laplacian makes the strong residual extremely curvature-sensitive.

A.4 Barron/spectral perspective: the Laplacian

The conditioning story above can be rephrased in approximation terms. In spectral Barron-type viewpoints [37], function classes are controlled by norms that weight the Fourier transform by powers of frequency (or related spectral weights). A key implication is that applying derivatives systematically increases the importance of high-frequency tails [40]. Since Δ multiplies a Fourier mode by $|\mathbf{k}|^2$, and the squared residual effectively involves L^*L (hence Δ^2 in Laplacian-dominant regimes), the residual can require substantially more “spectral budget” than the wavefunction itself. In practical terms: even if Ψ is approximable in a given network class, *approximating its strong residual* can be much harder, because the loss implicitly demands accurate high-frequency curvature (cf. (A.14)). This is one of the motivations, in numerical analysis of PINNs, for considering weak/variational formulations that reduce derivative order when the strong form is too stiff [40].

A.5 Implications for Stochastic Reconfiguration

Stochastic reconfiguration (SR) updates parameters using a natural-gradient-like preconditioner [24, 26, 57]. Writing $O_i = \partial_{\theta_i} \log \Psi_{\theta}$, one common SR update is

$$S(\theta) \delta\theta = -g(\theta), \quad S_{ij} = \text{Cov}_{p_{\theta}}[O_i, O_j], \quad g_i = 2 \text{Cov}_{p_{\theta}}[E_L, O_i], \quad (\text{A.17})$$

where $p_{\theta} \propto |\Psi_{\theta}|^2$ and $E_L = (\hat{H}\Psi_{\theta})/\Psi_{\theta}$. SR improves conditioning in parameter space by using the (quantum) information geometry of the variational family [24, 57]. However, the Laplacian-induced stiffness described above is an *operator-level* phenomenon: strong residuals magnify curvature errors via Δ and Δ^2 (cf. (A.14)), which remains a fundamental challenge even when parameter updates are geometrically preconditioned.

Chapter B

Coulomb conditioning

B.1 Coulomb structure and singularity

For our two-dimensional electronic systems ($\mathbf{r}_i \in \mathbb{R}^2$), the interaction contains Coulomb terms of the form

$$\hat{V}_C(\mathbf{R}) = \sum_{1 \leq i < j \leq N} \frac{1}{r_{ij}}, \quad r_{ij} = |\mathbf{r}_i - \mathbf{r}_j|. \quad (\text{B.1})$$

This is a *singular multiplication operator*: it diverges at particle coalescence $r_{ij} \rightarrow 0$. The exact eigenfunction remains finite (or vanishes by antisymmetry) but develops non-smooth short-distance structure (a cusp) so that divergences in kinetic and potential contributions cancel in physically meaningful quantities such as the local energy [26].

Coulomb does not merely add “another term” to the residual; it introduces *singular short-range physics* that forces delicate cancellations with the Laplacian. Strong residual objectives square these effects and become extremely sensitive to tiny cusp errors.

B.2 2D cusp conditions: antiparallel vs parallel spins

We present the 2D cusp logic in a way that separates the two spin channels. The cleanest statement (especially for practical ansätze) is in terms of the short-distance behaviour of a *Slater–Jastrow* factorization

$$\Psi(\mathbf{R}) = D(\mathbf{R}) J(\mathbf{R}), \quad J(\mathbf{R}) = \exp\left(\sum_{i < j} u(r_{ij})\right), \quad (\text{B.2})$$

because (i) the Slater part D enforces antisymmetry and determines whether Ψ vanishes at coalescence, and (ii) the Jastrow slope $u'(0)$ is what cancels the $1/r$ Coulomb singularity in the local energy [26].

Fix a pair (i, j) and introduce relative/centre-of-mass coordinates in 2D:

$$\mathbf{R}_{ij} := \frac{\mathbf{r}_i + \mathbf{r}_j}{2}, \quad \mathbf{r} := \mathbf{r}_i - \mathbf{r}_j, \quad r := |\mathbf{r}|. \quad (\text{B.3})$$

A direct computation gives, in any dimension,

$$\Delta_i + \Delta_j = \frac{1}{2} \Delta_{\mathbf{R}_{ij}} + 2 \Delta_{\mathbf{r}}, \quad (\text{B.4})$$

hence the singular short-distance kinetic behaviour is controlled by the relative term $-\Delta_{\mathbf{r}}$.

A key 2D identity is

$$\Delta_{\mathbf{r}} r = \frac{1}{r} \quad (r > 0, \mathbf{r} \in \mathbb{R}^2). \quad (\text{B.5})$$

Antiparallel spins (distinguishable in the Slater part). If the two electrons have opposite spin, they enter different determinants in D and generically D does *not* vanish at $\mathbf{r} \rightarrow 0$. Then $\Psi(0) \neq 0$ and a direct cusp expansion in 2D is legitimate:

$$\Psi(\mathbf{r}, \dots) = \Psi(0, \dots) \left(1 + a r + \mathcal{O}(r^2) \right), \quad r \rightarrow 0. \quad (\text{B.6})$$

Using (B.5), the singular part of the relative Laplacian is

$$\Delta_{\mathbf{r}} \Psi \approx \Psi(0) a \Delta_{\mathbf{r}} r = \Psi(0) \frac{a}{r}, \quad \frac{\Delta_{\mathbf{r}} \Psi}{\Psi} \approx \frac{a}{r}. \quad (\text{B.7})$$

Since the pair kinetic contribution contains $-\Delta_{\mathbf{r}}$, the singular part of the local energy is

$$E_L(\mathbf{R}) = \frac{\hat{H}\Psi}{\Psi} \supset \left(-\frac{a}{r} + \frac{1}{r} \right) = \frac{1-a}{r}. \quad (\text{B.8})$$

Thus finiteness of E_L at coalescence requires

$$a = 1, \quad \text{i.e.} \quad \partial_r \log \Psi|_{r=0} = 1 \quad (2\text{D, antiparallel spins}). \quad (\text{B.9})$$

Equivalently, in a Slater–Jastrow form with $D(0) \neq 0$, this is the same as $\partial_r \log J|_0 = 1$ for the ij -pair.

Parallel spins (identical fermions, node at coalescence). For same-spin electrons, antisymmetry forces D (and hence Ψ) to vanish as $\mathbf{r} \rightarrow 0$. Generically one has a *linear* zero:

$$D(\mathbf{r}, \dots) = \boldsymbol{\eta}(\dots) \cdot \mathbf{r} + \mathcal{O}(r^2), \quad r \rightarrow 0, \quad (\text{B.10})$$

with some nonzero vector $\boldsymbol{\eta}$ depending on the other coordinates. In this case, the cancellation of the Coulomb singularity is controlled by the Jastrow factor through cross terms in $-\Delta_{\mathbf{r}}(DJ)/(DJ)$. A standard short-distance analysis (carried out explicitly for 2D quantum dots) shows that the divergent kinetic contribution behaves like

$$-\frac{\Delta_{\mathbf{r}}(DJ)}{DJ} \supset -\frac{(2k+d-1)}{r} \partial_r \log J \quad \text{with} \quad d=2, \quad k=1 \text{ (identical particles)}, \quad (\text{B.11})$$

while the Coulomb potential contributes $+1/r$ to E_L as before. Hence finiteness of the local energy requires

$$\partial_r \log J|_{r=0} = \frac{1}{2k+d-1} = \frac{1}{3} \quad (2\text{D, parallel spins}). \quad (\text{B.12})$$

This 1 (antiparallel) versus 1/3 (parallel) 2D cusp statement is the standard quantum-dot/Jastrow–Slater cusp condition [58].

What to remember. In 2D:

$$\text{antiparallel spins:} \quad \partial_r \log \Psi|_0 = 1 \iff \partial_r \log J|_0 = 1, \quad (\text{B.13})$$

$$\text{parallel spins:} \quad \Psi(0) = 0 \text{ (linear node)}, \quad \partial_r \log J|_0 = \frac{1}{3}. \quad (\text{B.14})$$

B.3 Why Coulomb is ill-conditioned (2D emphasis)

The cusp conditions above show *how* Coulomb creates stiffness: it forces kinetic and potential contributions to be individually large and singular, while their *sum* is finite only under a short-distance constraint. Strong residual minimization is therefore a *cancellation problem*.

Sensitivity amplification in the local energy. If the learned antiparallel-spin cusp slope is $a_\theta = 1 + \delta a$, then (B.8) implies

$$E_L(\mathbf{R}) \supset -\frac{\delta a}{r}, \quad (\text{B.15})$$

so an arbitrarily small cusp mismatch produces an arbitrarily large local-energy spike as $r \rightarrow 0$.

For parallel spins, if the learned Jastrow slope is $\partial_r \log J|_0 = \frac{1}{3} + \delta b$, then the same short-distance analysis leading to (B.12) implies

$$E_L(\mathbf{R}) \supset -\frac{3\delta b}{r}. \quad (\text{B.16})$$

Thus both channels generate $1/r$ spikes unless the cusp is satisfied.

Why squaring is especially dangerous in 2D. A $1/r$ defect in E_L (or in $(\hat{H} - E)\Psi$) is concentrated near coalescence. But in 2D the squared singularity is not integrable—it diverges logarithmically:

$$\int_{r < \varepsilon} \left(\frac{1}{r}\right)^2 d^2\mathbf{r} = \int_0^\varepsilon \frac{1}{r^2} (2\pi r dr) = 2\pi \int_0^\varepsilon \frac{dr}{r} = +\infty.$$

Therefore, without effectively resolving the cusp, *strong* squared-residual objectives can be dominated (or even made ill-defined in the continuum limit) by the coalescence region. In Monte Carlo practice, this manifests as heavy-tailed fluctuations and extreme sensitivity to rare close-pair samples.

Residual expansion and cancellation structure. Writing $L = \hat{T} + (\hat{V} - E)$ and expanding the squared residual norm:

$$\|L\Psi\|_w^2 = \|\hat{T}\Psi\|_w^2 + \|(\hat{V} - E)\Psi\|_w^2 + 2 \operatorname{Re}\langle \hat{T}\Psi, (\hat{V} - E)\Psi \rangle_w, \quad (\text{B.17})$$

we see that near coalescence both $\hat{T}\Psi$ and $\hat{V}_C\Psi$ are large. The exact solution relies on cancellation between these contributions (cf. (B.8) and (B.12)), which means (B.17) demands high *relative* accuracy in the cusp region: if either term is off by a small fraction, the net residual can be large in absolute magnitude. This is a classic recipe for ill-conditioning: the loss landscape contains directions where small parameter changes cause huge residual changes in a small region.

Operator-level picture (L^*L). From the same viewpoint as Appendix A, strong residual training induces matrices/operators tied to L^*L [40]. For $L = \hat{T} + (\hat{V} - E)$, one has (formally)

$$L^*L = \hat{T}^2 + (\hat{V} - E)^2 + \hat{T}(\hat{V} - E) + (\hat{V} - E)\hat{T}. \quad (\text{B.18})$$

Coulomb enters here in two destabilizing ways:

1. the positive term $(\hat{V} - E)^2$ involves a $1/r^2$ -type singular weight;
2. the mixed terms contain derivatives of V through product rules, e.g. $\hat{T}(V\Psi)$ generates $\nabla V \cdot \nabla\Psi$ and ΔV contributions, which are even more singular/distributional at coalescence.

This formal expansion explains why the Coulomb singularity can severely inflate the effective spectral spread of the training operator, aligning with the operator-conditioning viewpoint [40].

B.4 Barron/spectral perspective: the Coulomb singularity

Coulomb forces the *true* 2D solution to have cusp-like, non-smooth short-range structure, so approximating the *strong residual* requires representing high-frequency content that is not well captured by globally smooth/low-complexity function classes.

In spectral/Barron-style approximation theories [37], approximation quality is governed by spectral decay and norms that weight the Fourier tail [40]. A cusp (or a linear node times a cusp in J) implies slower spectral decay than a globally smooth function. Strong residual training then *further* amplifies the high-frequency tail because it applies Δ (and effectively L^*L), which multiplies Fourier modes by powers of $|\mathbf{k}|$ (Appendix A). Hence, even if Ψ can be approximated reasonably in function value, the strong residual can remain large unless the cusp structure is represented accurately. This is exactly the type of regularity mismatch that motivates weak/variational formulations in numerical analysis of PINN-type methods, since weak forms reduce the derivative order demanded of the approximation [40].

Practically, QMC wavefunctions therefore incorporate explicit short-range correlation structure (e.g. cusp-enforcing Jastrow factors), precisely to remove these singular residual pathologies and reduce the variance of local quantities [26, 58].

B.5 Implications for Stochastic Reconfiguration

SR performs a natural-gradient step using a (quantum) Fisher/metric matrix [24, 26, 57]. In one common formulation,

$$S_{ij} = \operatorname{Cov}_{p_\theta}[O_i, O_j], \quad g_i = 2 \operatorname{Cov}_{p_\theta}[E_L, O_i], \quad S \delta\theta = -g, \quad (\text{B.19})$$

with $O_i = \partial_{\theta_i} \log \Psi_\theta$.

Coulomb-induced cusp mismatch produces E_L spikes of the form (B.15)–(B.16), which has two consequences:

1. **Heavy-tailed Monte Carlo forces.** The estimator of $g_i = \text{Cov}[E_L, O_i]$ becomes noisy when E_L has rare but extreme spikes (short-range events). This increases variance and can destabilize the linear solve unless one uses damping/regularization, as standard in SR practice [26].
2. **Coupling to curvature sensitivity.** Even if SR corrects anisotropy in parameter space (natural gradient) [24, 57], it does not remove the underlying operator-driven stiffness: the spikes originate from imperfect kinetic–potential cancellation enforced by Coulomb cusp physics.

Summary. Laplacian dominance creates stiffness through k^4 -type amplification of curvature errors; Coulomb creates stiffness by enforcing cusp constraints (in 2D: 1 vs 1/3 depending on spin channel) and cancellation against the Laplacian. Both mechanisms worsen conditioning of strong residual objectives and increase statistical difficulty of SR through heavy-tailed local-energy fluctuations [26, 40, 58].

Chapter C

The collocation–backflow catch-22

This appendix formalizes a structural difficulty encountered when training backflow transformations under the *strong-form* collocation objective—the one that minimises the variance of the pointwise local energy E_L —and presents extensive numerical evidence for the $N=6$, $\omega=0.5$ quantum dot. The core observation is a *catch-22*: backflow’s learning signal is concentrated near the nodal surface, but the local energy diverges there for any imperfect wavefunction, and every sampling strategy we tried that includes those points produced gradients whose variance we could not control. The result, across every intervention below, is an accuracy floor at roughly 0.3%—set by the Jastrow alone—that is absent under VMC + SR training of the same ansatz family. We call the floor irreducible *within this pipeline*: it is an empirical bound established by exhaustion over the interventions we tested, not a proof that no strong-form scheme can do better.

The scope of this result needs two qualifications. First, it is a property of the *strong form* specifically: the weak-form (score-function / REINFORCE) collocation objective used for the results in the main text never differentiates through the Laplacian and does not inherit this pathology, which is why it trains the same ansatz to far below the 0.3% floor. The value here is therefore a diagnosis of one objective’s limits, not a measure of what MCMC-free training can achieve. Second, this analysis is what motivated the shift to the weak form, and is reported as a negative result rather than as a headline number.

C.1 Setup and notation

We consider a Slater–Jastrow–backflow ansatz of the form

$$\Psi_\theta(\mathbf{R}) = \det \tilde{S}_\uparrow(\mathbf{R}; \theta_{\text{BF}}) \det \tilde{S}_\downarrow(\mathbf{R}; \theta_{\text{BF}}) \exp(J(\mathbf{R}; \theta_J)), \quad (\text{C.1})$$

where $\tilde{S}_\sigma$ is the Slater matrix evaluated at backflow-shifted coordinates $\tilde{\mathbf{r}}_i = \mathbf{r}_i + \Delta \mathbf{x}_i(\mathbf{R}; \theta_{\text{BF}})$, J is a neural Jastrow correlator, and $\theta = (\theta_J, \theta_{\text{BF}})$ are trainable parameters. A CTNN message-passing network produces the backflow shifts $\Delta \mathbf{x}_i$.

We train by minimizing the *screened collocation* objective

$$\mathcal{L}(\theta) = \text{Var}_S[E_L(\mathbf{R}; \theta)], \quad E_L = \frac{\hat{H}\Psi_\theta}{\Psi_\theta}, \quad (\text{C.2})$$

where the collocation set S is obtained by importance-sampling from a Gaussian proposal q and retaining the top- K points ranked by $|\Psi_\theta|^2/q$ (“screened collocation”). This top- K selection naturally suppresses near-node configurations where $|\Psi|^2 \approx 0$.

C.2 The divergence of E_L near nodes

The local energy decomposes as

$$E_L = -\frac{1}{2} \frac{\nabla^2 \Psi}{\Psi} + V = -\frac{1}{2} \left[\nabla^2 J + |\nabla J|^2 + 2 \nabla J \cdot \frac{\nabla D}{D} + \frac{\nabla^2 D}{D} \right] + V, \quad (\text{C.3})$$

where $D := \det \tilde{S}_\uparrow \det \tilde{S}_\downarrow$ and the last term $\nabla^2 D/D$ contains second derivatives of the backflow shifts $\nabla^2 \Delta \mathbf{x}_i$ amplified by the inverse determinant.

For the exact wavefunction, E_L is constant. If Ψ is an eigenstate, $E_L = E$ everywhere. The cancellation is a property of the *full* local energy $\hat{H}\Psi/\Psi$ with $\Psi = D e^J$, in which the Jastrow cross term $2\nabla J \cdot \nabla D/D$ of Eq. (C.3) participates alongside $\nabla^2 D/D$; we isolate the determinant terms below because they are the ones the backflow parameters move, not because the others are absent. For an exact eigenstate, the potentially singular contributions in the full expression $\nabla^2(D e^J)/(D e^J)$ cancel at a simple node. This is not a property of $\nabla^2 D/D$ in isolation: for a generic smooth determinant with $D \sim d$ near the nodal surface $\mathcal{N}$, $\nabla^2 D$ may remain $\mathcal{O}(1)$. The exact eigenstate satisfies

$$\frac{\nabla^2(D e^J)}{D e^J} \text{ finite on } \mathcal{N}, \quad (\text{C.4})$$

through cancellation among the determinant, Jastrow-cross, and remaining Jastrow terms in Eq. (C.3).

For an approximate wavefunction, the cancellation generically fails. During training, the backflow parameters θ_{BF} are imperfect: the *learned* nodal surface $\mathcal{N}_\theta$ (where $D(\cdot; \theta) = 0$) does not coincide with the *true* nodal surface. Let d_θ denote the signed distance from the learned node and d^* that from the true node. Near a point where $d_\theta \approx 0$ but $d^* \neq 0$, the determinant $D \sim d_\theta \rightarrow 0$ while the kinetic contributions from the Jastrow and potential terms remain finite:

$$E_L(\mathbf{R}) \supset \frac{\nabla^2 D(\mathbf{R}; \theta)}{D(\mathbf{R}; \theta)} \sim \frac{c}{d_\theta} \rightarrow \pm\infty \quad \text{as } d_\theta \rightarrow 0, \ d^* \neq 0, \quad (\text{C.5})$$

for some configuration-dependent constant $c \neq 0$. Conversely, near a point on the true node ($d^* = 0$) but off the learned node ($d_\theta \neq 0$), $\Psi \sim d^* \rightarrow 0$ and the wavefunction assigns vanishing weight. The pathology (C.5) is therefore not an artifact of the nodal structure itself—it is a property of *mismatched* nodes, and is worst precisely when backflow still has something to learn.

C.3 The geometric mismatch argument

We now locate the divergence. It is *not* in the variance of the local energy, which the calculation below shows to be finite for a simple node; it is one derivative later, in the parameter gradient.

Consider the contribution to $\text{Var}[E_L]$ from a neighborhood U_ε of the learned nodal surface:

$$\text{Var}[E_L] \geq \int_{U_\varepsilon} |\Psi(\mathbf{R}; \theta)|^2 (E_L(\mathbf{R}; \theta) - \bar{E})^2 d\mathbf{R}. \quad (\text{C.6})$$

When the node is “simple” (D vanishes linearly), we have

$$|\Psi|^2 \sim d_\theta^2, \quad E_L^2 \sim d_\theta^{-2\alpha}, \quad (\text{C.7})$$

where α characterises the severity of the mismatch. For the generic case where the numerator $\nabla^2 D$ does *not* vanish on $\mathcal{N}_\theta$ (because $\mathcal{N}_\theta \neq \mathcal{N}^*$), one has $\alpha = 1$ so that $E_L^2 \sim d_\theta^{-2}$.

In D -dimensional configuration space, the volume element in a tubular neighborhood of a codimension-1 surface scales as $dV \sim d_\theta^0 d\sigma dd_\theta$. Thus the integrand in (C.6) behaves as

$$|\Psi|^2 \cdot E_L^2 \cdot dV \sim d_\theta^2 \cdot d_\theta^{-2} \cdot dd_\theta = dd_\theta, \quad (\text{C.8})$$

which integrates to $\mathcal{O}(\varepsilon)$ and is therefore *finite*: a simple node does not, by itself, make $\text{Var}[E_L]$ diverge under the $|\Psi|^2$ measure. The instability enters one derivative later, in the *parameter gradient*. But the key observation is that this is the *best case*. In practice, the mismatch between $\mathcal{N}_\theta$ and $\mathcal{N}^*$ is not uniform: there exist regions where the surfaces cross at angles, creating pockets where $|d^*|/|d_\theta|$ is large (the true wavefunction is non-negligible while the learned determinant is near-zero), and there the E_L^2 spike is not compensated by $|\Psi|^2$ suppression.

More concretely, the gradient $\partial \mathcal{L} / \partial \theta_{\text{BF}}$ involves terms of the form

$$\frac{\partial E_L}{\partial \theta_{\text{BF}}} \supset \frac{1}{D} \frac{\partial (\nabla^2 D)}{\partial \theta_{\text{BF}}} - \frac{\nabla^2 D}{D^2} \frac{\partial D}{\partial \theta_{\text{BF}}}, \quad (\text{C.9})$$

both of which diverge as $D \rightarrow 0$. Even in the marginal case where $\text{Var}[E_L]$ is finite, the *gradient variance* (which involves these higher-order poles) generically diverges, making gradient-based optimisation impossible at points near the learned node.

C.4 The catch-22

The preceding analysis reveals a three-sided dilemma:

1. **Backflow learns at the node.** The purpose of backflow is to reshape the nodal surface of the Slater determinant. The gradient signal $\partial E_L / \partial \theta_{\text{BF}}$ is largest where the backflow shifts actually change which side of the node a configuration point lies on, i.e., where $D \approx 0$.
2. **E_L diverges at the node for imperfect θ .** As shown in §C.2–C.3, the local energy and its parameter gradient contain poles of the form D^{-1} and D^{-2} whenever the learned node does not coincide with the exact node. The resulting gradient variance is unbounded.
3. **Good collocation sampling must avoid the node.** Under screened collocation, the top- K selection ranks points by $|\Psi|^2/q \propto D^2 e^{2J}/q$. Since $D^2 \rightarrow 0$ at the node, near-node points are systematically excluded from the training batch. This is *protective*—it prevents gradient explosion—but it also removes the points where backflow’s learning signal lives.

A “catch-22” emerges:

Including near-node points gives divergent, noisy gradients and backflow shrinks defensively or diverges. Excluding them gives stable training but removes backflow’s learning signal. In either regime, collocation cannot effectively train backflow beyond an accuracy floor set by the correlator quality alone.

C.5 Why VMC + SR escapes

Under VMC, the sampling distribution is $|\Psi_\theta|^2$, which assigns vanishing measure to the nodal surface. Near-node configurations are strongly *suppressed*—for a simple node the probability of lying within a distance ϵ scales as ϵ^3 —rather than strictly absent, which is enough to keep the D^{-1} singularity out of the gradient estimator in practice. However, VMC *still optimises the node position* through a subtle mechanism:

Indirect nodal information through bulk correlations. The variational energy $\langle E_L \rangle_{|\Psi|^2}$ is sensitive to nodal position because shifting the node changes the wavefunction *everywhere* (through normalisation and through the functional form of D). The SR force vector $g_i = \text{Cov}_{|\Psi|^2}[E_L, O_i]$ (where $O_i = \partial_{\theta_i} \log \Psi$) receives contributions from the *bulk*—regions where both $|\Psi|^2$ and E_L are well-behaved—that are *correlated with* nodal position but do not require evaluating E_L at the node.

SR amplifies the signal. The Fisher metric $S_{ij} = \text{Cov}[O_i, O_j]$ captures how sensitive $\log \Psi$ is to each parameter direction. By solving $(S + \epsilon I) \delta \theta = -g$, SR amplifies exactly those subtle bulk signals that encode nodal information, compensating for the fact that the node itself has zero weight under $|\Psi|^2$.

Core asymmetry. The asymmetry needs stating carefully, because a power count that is too quick proves too much. Under $|\Psi|^2$ the SR force $g_i = \text{Cov}[E_L, O_i]$ has a finite *mean*: both E_L and $O_i = \partial_{\theta_i} \log \Psi$ carry a d_θ^{-1} pole near a mismatched node, their product goes as d_θ^{-2} , and the measure supplies $|\Psi|^2 \sim d_\theta^2$, so the integral converges. Its *variance* does not, by the same counting—so VMC is not immune to near-node pathology either, and in practice this is what the damping and trust region of §4.4 are for.

The real distinction is one derivative higher. VMC and SR never differentiate E_L : the local energy enters only as a multiplicand in a covariance, so the D^{-2} pole of Eq. (C.9)—which comes from ∂_θ acting on $\nabla^2 D/D$ —never appears. Strong-form collocation differentiates E_L by construction, and inherits it. That is why the same node is survivable in one scheme and not in the other, and it is the argument

§7.3.1 makes for the weak form as well. None of the sampling, loss, or preconditioning strategies we tested (§C.6) overcame this asymmetry within the strong-form collocation framework; we present it as a structural obstruction of the tested pipeline rather than a theorem that no collocation method can ever avoid it.

C.6 Experimental evidence

We present a systematic study for $N=6$ electrons in a 2D harmonic trap with $\omega=0.5$ (DMC reference: $E_{\text{DMC}} = 11.78484(6)$ [12]). The ansatz uses a neural Jastrow correlator ($\sim 15,000$ parameters) and a CTNN backflow network ($\sim 8,600$ parameters) with `bf_scale` = 0.7. All energies are evaluated via MCMC with 15 000 samples after burn-in.

C.6.1 Baseline: correlator alone vs. correlator + backflow

Table C.1: Baseline collocation results for $N=6$, $\omega=0.5$. “Correlator only” trains without backflow; “joint” trains both networks from scratch for 300 epochs with `bf_scale` = 0.7. The backflow provides only a marginal improvement of ~ 0.1 percentage points.

Configuration	E (Ha)	σ_E	err (%)
Correlator only (300 ep)	11.834	0.003	0.42
Correlator + BF, joint (300 ep)	11.823	0.003	0.33
Correlator + BF, separate opt	11.822	0.003	0.32
DMC reference	11.78484	—	0.00

The backflow reduces the error from 0.42% to 0.32–0.33%, far short of the $<0.01\%$ reached by VMC+SR on the same ansatz *family* (Table 6.2). The two are not matched in size: the networks here total $\sim 24\text{k}$ parameters against $\sim 236\text{k}$ for the production model, so the comparison bounds what this objective achieves at this capacity rather than isolating the objective alone. The floor is nonetheless a property of the objective, because it does not move under any of the interventions below, including ones that add capacity.

C.6.2 Sampling strategies

We tested multiple strategies to bring near-node information into the collocation batch, hoping to give the backflow a stronger learning signal.

Table C.2: Effect of sampling strategy on collocation accuracy. All experiments use 50-epoch fine-tuning from the best collocation checkpoint (0.35%), except “hard-smooth (scratch)” which trains 300 epochs from random initialisation.

Method	Description	err (%)	Outcome
Top- K baseline	standard screened collocation	0.35	stable
Hard mining + Huber	top- K of residuals	0.41	neutral
Hard mining + smoothness	+ $\ \nabla^2 \Delta x\ ^2$ penalty	0.38	slight help
SIR (single proposal)	$ \Psi ^2$ -reweighted resampling	0.47	worse
SIR (mixture)	Gaussian mixture proposal	0.44	worse
Gumbel top- K [59]	stochastic top- K selection	5.00	catastrophic

Every strategy that includes more near-node points (SIR, Gumbel) produced *worse* results—consistent with the catch-22: the additional points carry divergent E_L values that corrupt the gradient. Gumbel top- K (which replaces deterministic selection with stochastic Gumbel-softmax) was catastrophic, with early stopping triggered by rapid energy divergence.

C.6.3 Loss modifications

Table C.3: Effect of loss function modifications. The hybrid loss adds a mean-energy (variational) term; the det filter discards points with small $|\det \tilde{S}|$.

Method	Description	err (%)	Outcome
Variance loss (baseline)	Huber on $E_L - \bar{E}$	0.33	best collocation
Hybrid anneal	$\beta \cdot \text{Var} + (1-\beta) \cdot \text{Mean}$, $\beta:0.7 \rightarrow 0.15$	2.59	catastrophic
Gumbel + hybrid	stochastic top- K + hybrid	5.00	catastrophic
Hybrid + det filter	$\beta=0.2$ fixed + $ \det $ filter	24.75	broken

The mean-energy gradient $\nabla_\theta \langle E_L \rangle_S$ under biased collocation sampling does *not* equal the VMC gradient—the sampling distribution is not $|\Psi|^2$ —so the variational principle does not hold, and the backflow drifts to large, unphysical shifts ($|\Delta \mathbf{x}| \approx 0.83$, versus ~ 0.23 under variance-only training). The variance loss’s implicit constraint on backflow magnitude is *protective*, not pathological.

C.6.4 Gradient regularisation

We attacked the divergence sources in E_L directly:

Table C.4: Regularisation of the E_L singularity (50-epoch fine-tune from 0.35% checkpoint). Each modification smooths or clips one of the divergent contributions to E_L .

Method	Modification	err (%)	Outcome
Soft Coulomb	$1/r \rightarrow 1/\sqrt{r^2 + \varepsilon^2}$, $\varepsilon=0.2$	1.41	much worse
Regularised det	$\log D \rightarrow \text{logaddexp}(\log D , \log \delta)$	0.49	worse
T clipping	$\max(T , T_{\max})$, $T_{\max}=30$	0.43	neutral

Soft Coulomb trains on wrong physics (V is no longer the true potential). Regularising the determinant distorts the $\log \Psi$ landscape. Kinetic energy clipping is neutral because the clipped points are exactly those that carried the (noisy) backflow signal. None of the regularisations improve upon the baseline.

C.6.5 Gradient preconditioning

Table C.5: Gradient preconditioning experiments (300-epoch scratch training unless noted). K-FAC [60] uses block-diagonal Kronecker-factored Fisher on the backflow network only.

Method	err (%)	Outcome
Vanilla Adam	0.33	baseline
K-FAC on CTNN	0.34	no improvement
K-FAC + smoothness	0.51	worse
Variance-minimising pretrain	0.54	worse
Adam fine-tune (30 ep, lr=1e-4)	0.34	no improvement
Adam burst (15 ep, lr=3e-4)	0.56	worse

K-FAC provides a block-diagonal approximation to the Fisher matrix and should, in principle, help with the ill-conditioned parameter-space geometry. In practice, it does not: the conditioning problem is *operator-level* (Appendices A–B), not merely parameter-space anisotropy, and K-FAC cannot address the fundamental divergence in E_L .

C.6.6 Summary of all interventions

The pattern is striking:

- Every intervention that *includes* more near-node points (SIR, Gumbel) *increases* the error—backflow shrinks defensively under gradient noise or diverges entirely.

Table C.6: Complete summary of all collocation-based backflow experiments for $N=6$, $\omega=0.5$. No intervention breaks the $\sim 0.32\%$ accuracy floor. The final row is the same ansatz *family* under VMC + SR at production width (Table 6.2), an order of magnitude larger in parameter count; it is included to show the size of the gap, not as a matched comparison.

Experiment	err (%)	Category
Correlator + BF, sep opt (best)	0.32	baseline
Correlator + BF, joint 300 ep	0.33	baseline
K-FAC on CTNN	0.34	preconditioner
Adam fine-tune 30 ep	0.34	optimiser
Top- K baseline (retrained)	0.35	sampling
Hard mining + smoothness	0.38	sampling
Hard mining + Huber	0.41	sampling
Correlator only	0.42	no backflow
T clipping	0.43	regularisation
SIR (mixture)	0.44	sampling
SIR (single)	0.47	sampling
Regularised det	0.49	regularisation
K-FAC + smoothness	0.51	preconditioner
Var-min pretrain	0.54	pretrain
Adam burst lr=3e-4	0.56	optimiser
Soft Coulomb	1.41	regularisation
Hybrid loss (annealed)	2.59	loss
Gumbel top- K + hybrid	5.00	sampling + loss
Hybrid + det filter	24.75	loss + filter
Same ansatz, VMC + SR	<0.01	variational

- Every intervention that *changes* the loss to tolerate near-node contributions (hybrid mean-energy, det filter) produces catastrophic results because the collocation sampling is not $|\Psi|^2$ and the variational principle does not apply.
- The smoothness penalty $\lambda\|\nabla^2\Delta\mathbf{x}\|^2$ helps modestly (from 0.59% to 0.38% without it vs. with it) by damping the $\nabla^2 D/D$ poles, but cannot eliminate the fundamental divergence.
- Ironically, the only method that trains backflow effectively is MCMC-based VMC + SR—which sidesteps the entire collocation framework.

C.7 Connection to Appendices A and B

The catch-22 is a specific manifestation of the general conditioning difficulties analysed in Appendices A–B:

Laplacian stiffness (Appendix A). The k^4 -type amplification of curvature errors (§A.3) is compounded by backflow, because $\nabla^2(\det \tilde{S})$ introduces second derivatives of $\Delta\mathbf{x}$ that are not present in a fixed-node ansatz. These additional high-frequency modes inflate the spectral spread of the training operator L^*L .

Coulomb singularity (Appendix B). The $1/r$ Coulomb potential (§B.1) adds a singular multiplication operator to $\hat{H}$. Near both the coalescence point ($r_{ij} \rightarrow 0$) and the node ($D \rightarrow 0$), E_L contains products of these two singularities, and the cusp cancellation (§B.2) must be maintained simultaneously with the nodal cancellation (C.4). In the strong residual squared $\|LE_L\|^2$, these compound singularities create an extremely stiff landscape.

The backflow-specific pathology. What distinguishes the backflow catch-22 from the general conditioning problem is the *self-referential* nature of the singularity: the backflow parameters θ_{BF} determine *where* the node is, so updating θ_{BF} moves the singularity in configuration space. Under the collocation loss, the gradient pushes θ_{BF} in a direction determined by E_L values at the current collocation points, but the resulting node shift invalidates those very points. VMC+SR avoids this because it samples from the *current* $|\Psi_\theta|^2$ at every step, so the sampling distribution co-adapts with the node position.

Chapter D

Additional collocation results

This appendix records the collocation results that Chapter 7 refers to but does not tabulate: the $N=20$ energies, and the variants that were tried and failed. Both bear on the same question—what limits the Markov-chain-free route—from the side of what does not work.

D.1 What was varied

Four axes were explored. The *sampling*: Gaussian and Gaussian-mixture proposals scaled by $1/\sqrt{\omega}$, oversample-then-resample with ESS diagnostics, ESS-adaptive oversampling, weight tempering and log-weight clipping. The *objective*: finite-difference collocation residuals, a hybrid weak form, and pure REINFORCE, which became the default, with the Laplacian kept in the reward path but never differentiated. The *preconditioner*: Adam, diagonal-Fisher natural gradient, and full SR in Woodbury and CG modes with damping anneals and trust-region caps. And the *training style*: single sweeps, wave-based cascades with checkpoint inheritance, and asynchronous campaigns relaunching from the best available checkpoint.

Two results held across the search. Optimisation quality dominated architectural novelty: moderately sized backflow+Jastrow models with stable training beat more ambitious variants with weaker optimisation geometry. And at low ω , regularising the resampling distribution (ESS control, top-mass reduction) helped where hard cusp gating did not—a controlled $2 \times 2 \times 3$ matrix over gating, resampling regularisation and three seeds showed no measurable gain from gating and a consistent gain from regularisation.

Warm-starting along the confinement cascade $\omega : 1.0 \rightarrow 0.5 \rightarrow 0.1 \rightarrow 0.01 \rightarrow 0.001$ was necessary throughout; cold starts at low ω did not converge. Representative cascade results are $N=6$ at $\omega=1.0$ ($E = 20.16167 \pm 0.00181$, +0.012%), $\omega=0.5$ ($E = 11.78466 \pm 0.00199$, -0.002%) and $\omega=0.1$ ($E = 3.56525 \pm 0.00046$, +0.32%), and $N=12$ at $\omega=1.0$ ($E = 65.70699 \pm 0.00595$, +0.010%). For $N=2$ the route covers the whole confinement range: +0.012%, +0.023%, -0.004%, +0.002% and +0.154% at $\omega = 1, 0.5, 0.1, 0.01, 0.001$, zero-variance extrapolated.

D.2 The $N = 20$ collocation energies

At $N=20$ the backflow’s edge-feature tensor, which scales as $\mathcal{O}(N^2 h_{\text{bf}})$, forces the hidden dimension down to $h_{\text{bf}}=48\text{--}64$ to fit in memory. The best backflow run at $\omega=1.0$ then reached only $\sim+18\%$ error, while a Jastrow-only ansatz—freed of that tensor, and so able to afford a larger collocation budget and higher oversampling—reached +1.32% ($E = 157.934 \pm 0.018$). The more expressive architecture is beaten by the simpler one, which is consistent with the memory-forced capacity cut degrading gradient quality more than its extra flexibility repays.

Polishing the Jastrow checkpoints at reduced learning rate improved the $\omega=1.0$ error from $\sim+2.6\%$ to +1.32% over ~ 400 epochs, and the $\omega=0.5$ error from $\sim+7.0\%$ to +2.74% ($E = 96.447 \pm 0.015$); the best $\omega=0.1$ result was +5.53% ($E = 31.637 \pm 0.008$). The gap to the sub-percent accuracy reached at $N \in \{6, 12\}$ tracks the capacity cut and the gradient noise it brings. A more memory-frugal backflow is

the direct next intervention; these experiments do not isolate the collocation objective as the source of the remaining gap.

D.3 Negative results worth recording

Scalar final energy is not sufficient for method development on this route: ESS, top-mass concentration, rollback counts and the probe-to-final gap are what distinguish a run that converged from one that stopped moving. Several intuitively attractive interventions delivered nothing in the tested range—hard cusp gating most notably—and recording that was necessary to keep the search from circling. Stricter SR trust constraints, heavier damping, ESS-adaptive sampling and rollback controls improved operational stability without closing the accuracy gap at the hardest targets. What eventually closed it at low ω and $N=6$ was not a stabilisation measure but a change of proposal (§7.4).

Chapter E

The learned representation of one trained solution

This appendix records what the internal representation of *one* high-quality trained solution looks like. It is descriptive, it is measured on a single training run, and—as §7.1.2 sets out—most of the quantities in it are component-level and therefore gauge-like: the correlator and the backflow are partial substitutes, so how the labour is divided between them records the route the training took as much as the state it reached. Nothing in the architectural argument of Chapter 7 rests on anything here. It is reported because the regularities are striking and because they are ones the network was free *not* to adopt, not because they establish a property of the architecture.

Three findings survive that qualification well enough to be worth stating. Across $N \in \{2, 6, 12, 20\}$ and $\omega \in \{1, 0.5, 0.1, 10^{-2}, 10^{-3}\}$ the correlator feature space stays on one to three effective dimensions and never expands, even where the real-space density develops elaborate shell structure; what changes across the crossover is which branch the leading axis is built from, not how many axes there are. The displacement field is the opposite kind of object, occupying nearly all the spatial directions available to it. And the coupling between the two changes character near $\omega \approx 0.1$, the same place the energy partition crosses over (§6.2.3).

Every diagnostic below is evaluated on fresh samples drawn from $|\Psi_\theta|^2$, with the estimators and stability protocol of §5.2.

E.1 Correlator geometry: a low-rank representation that reorganises

We characterise the correlator feature space $Z = \bar{\phi} \|\bar{\psi}\| \mathbf{g}$ by its entropy-based effective rank $r_{\text{eff}}(Z)$, the correlation $|\rho(\text{head}, \text{PC1})|$ between the head output and its first principal component, and the power of PC1 decomposed across architectural branches. Table E.1 reports these quantities across the grid.

The manifold is low-dimensional everywhere. The effective rank stays below 3.1 across the entire grid, and the head output is almost perfectly reconstructed from PC1 alone in all but one analysed cell ($|\rho| \geq 0.94$ except $N=20, \omega=0.01$, where $|\rho| = 0.79$). This holds even at strong correlation where the real-space density develops pronounced shell structure—the correlator does not respond by expanding its latent dimensionality. The modest peak at $N=12, \omega=0.1$ ($r_{\text{eff}} = 3.08$) coincides with the intermediate-correlation regime where neither the shell picture nor the crystalline picture provides a clean single-mode description.

A $\phi \leftrightarrow \psi$ transition in the leading axis. What varies across regimes is *which branch dominates PC1*. At moderate-to-weak correlation ($\omega \geq 0.1, N \geq 6$), PC1 is predominantly ψ -loaded: the pair branch carries 60–73% of the variance, meaning the correlator’s primary degree of freedom tracks pair correlations. At strong correlation the loading shifts toward ϕ (orbital) or $\mathbf{g}$ (auxiliary) branches for $N=6$ and $N=12$. The $N=20$ run does not follow the same branch-switching pattern: its pair branch remains dominant at both weakest confinements. Given that these are single-run, component-level diagnostics,

correlator's gradient; the change is sharp and coincides with the crossover in Γ .

These measurements describe one trained solution and do not by themselves establish which properties are architecture-specific. A different training route could divide the same labour differently and arrive at the same energy, which for two partial substitutes is what one should expect. Deciding the question requires holding everything else fixed and varying only the architecture.

F.2 Shell-resolved structure across the whole grid

Chapter 6 quotes shell occupancies, radii, bond order and Lindemann numbers for individual (N, ω) cells in running prose. Table F.3 collects them for the whole grid from a single analysis pass, so that the trends can be read off rather than assembled from the text. Shell radii are given in trap units $\tilde{r} = \sqrt{\omega} r$; multiply by $\ell = \omega^{-1/2}$ for Bohr.

Reading $|\Phi_n|$ against its null. One column of that table needs a baseline to be interpretable, and its absence would make the bond-order numbers read far more favourably than they should. For n particles placed at *independent uniform* angles on a ring, $\Phi_n = n^{-1} \sum_k e^{in\phi_k}$ is a two-dimensional random walk of n unit steps, so

$$\langle |\Phi_n| \rangle_{\text{random}} = \frac{\sqrt{\pi}}{2\sqrt{n}} \approx \frac{0.886}{\sqrt{n}}, \quad \langle |\Phi_n|^2 \rangle_{\text{random}} = \frac{1}{n}. \quad (\text{F.1})$$

A finite $|\Phi_n|$ is therefore *not* by itself evidence of angular order: a five-particle ring with no order at all returns $|\Phi_5| \approx 0.40$. The ratio $|\Phi_n| / \langle |\Phi_n| \rangle_{\text{random}}$ is the quantity that carries information, and Table F.3 reports it.

Read that way the result is sharper than the raw values suggest, and it needs stating with the right threshold. From $\omega = 1$ down to $\omega = 0.1$ the ratio is 1.00–1.02 at every particle number and on every ring. At $\omega = 0.01$ it lifts slightly, to 1.10 at $N=6$, while $N=12$ and $N=20$ remain at 1.03 and 1.01. Only at $\omega = 10^{-3}$ does it become large: 1.65 on the $N=6$ ring, 1.19 and 1.56 on the two rings of the $N=12$ (3, 9) molecule, and 1.19 and 1.47 on the outer two rings of the $N=20$ (1, 7, 12) molecule.

Whether the small excesses are *resolved* is a separate question from whether they are *large*, and with these sample sizes the two come apart. Two caveats attach to the null before it is used quantitatively. The Rayleigh form is the large- n limit of the resultant of n unit steps; for the small rings here ($n = 3$ to 19) it is an approximation, accurate to well within the effect at 10^{-3} but of the same order as the few-percent excesses at $\omega \geq 0.1$. And the frames come from a Markov chain, so a naive standard error over K raw frames understates the uncertainty, whereas the block bootstrap of §5.2.3 is used for the reported errors elsewhere. Taking those at face value, the excess at $N=6$, $\omega = 0.1$ is of order tens of block standard errors and still only 2%.

The conclusion that survives both caveats is the one that matters: statistical detectability is not what separates the regimes here, magnitude is. On that criterion the ordering is flat through $\omega = 0.1$, weakly present at $\omega = 0.01$ and only at $N=6$, and unambiguous at 10^{-3} . A cleaner version of this comparison would generate the null numerically at each actual occupancy n , through the same block-bootstrap pipeline, rather than relying on the asymptotic form; that is a small addition we did not make.

The Lindemann numbers of the same table fall monotonically throughout, so radial stiffening precedes n -fold angular ordering rather than accompanying it—the separation between radial and angular melting anticipated in §1.4. Note that the values here are for the *modal* topology over all frames; the sector-conditioned values quoted in §6.2.5 are computed on the frames assigned to a given sector and run slightly higher.

power.

Table F.5: Fitted outer-shell radius of the $N=6$ (1, 5) molecule against the classical Wigner ring $r_{\text{cl}} = 1.334\omega^{-2/3}$. Radii in Bohr.

ω	r_{shell}	r_{cl}	ratio
1.0	1.66	1.33	1.247
0.5	2.52	2.12	1.188
0.1	6.80	6.19	1.098
0.01	29.65	28.74	1.032
0.001	135.67	133.40	1.017

The ratio falls monotonically toward unity, reaching 1.017 at $\omega = 10^{-3}$: the quantum ring sits 1.7% outside the classical one at the weakest trap, and the excess grows smoothly with confinement as zero-point motion pushes the shell outward. Fitting $r_{\text{shell}} \propto \omega^\alpha$ gives $\alpha = -0.637$ over the full grid and -0.650 over the weakest three confinements, against the classical $-2/3$.

Two remarks on why this quantity, rather than the r_{mode} of Table 6.3, is the right one to compare. First, r_{mode} is the mode of the *single-particle* radial density, which for a (1, 5) topology mixes the central electron with the ring and is therefore not the ring radius; fitting it gives $\alpha \approx -0.58$ at $N=6$ and -0.54 at $N=2$ over the full grid, and approaches $-2/3$ only on the weak-trap half. Second, the shell-resolved radius is the quantity the physics-informed proposal of §7.4 actually needs, so the agreement recorded here is what licenses building a sampler out of it.

Chapter G

Declaration on the Use of Artificial Intelligence

The collocation approach in this thesis was developed over several months of failed experiments. Its central difficulty was not merely that a network did not converge, but that the parts of the Schrödinger residual that made the method physically meaningful also made its gradients unstable near the wavefunction's nodes. During that period, AI systems were used as sounding boards: to test interpretations of unexpected results, to formulate possible explanations, and to decide which experiment should come next. The exchange below is a short sample from one longer conversation at a point where the negative evidence had begun to make abandoning collocation seem plausible.

Author

I am trying to develop a PINN to find ground-state solutions of the Schrödinger equation. I have tried multiple approaches, but I am not succeeding. ... What do you think about this? What should we do from here?

AI response 1

... Anything that injects near-node points makes gradients explode or become uselessly noisy. ...

⋮

AI response 2

... This asymmetry is fundamental and cannot be resolved by sampling tricks, loss modifications, or gradient preconditioning within the collocation framework. ...

⋮

AI response 3

... Stick with SR + energy (or energy+variance) VMC. ...

Artificial-intelligence systems were therefore used for discussion, planning, and review, rather than as a source of results. They helped turn observations into hypotheses that could be checked, suggested alternative diagnostics when a run failed, and reviewed the final prose for clarity and mistakes. The code, the experiments, the scientific judgments, and the text of this thesis were written by the author. In particular, the recommendation shown above was not accepted as a methodological conclusion: it became one claim among several to investigate. The work that followed led to the collocation analysis reported in this thesis.

- [15] Nicholas Metropolis, Arianna W. Rosenbluth, Marshall N. Rosenbluth, Augusta H. Teller, and Edward Teller. Equation of state calculations by fast computing machines. *The Journal of Chemical Physics*, 21:1087, 1953.
- [16] Malvin H. Kalos, Dominique Levesque, and Loup Verlet. Helium at zero temperature with hard-sphere and other forces. *Physical Review A*, 9:2178, 1974.
- [17] Gaopei Pan and Zi Yang Meng. The sign problem in quantum Monte Carlo simulations. In *Encyclopedia of Condensed Matter Physics*, pages 879–893. Elsevier, Oxford, 2 edition, 2024. doi: 10.1016/B978-0-323-90800-9.00095-0.
- [18] Robert Jastrow. Many-body problem with strong forces. *Physical Review*, 98(5):1479–1484, 1955. doi: 10.1103/PhysRev.98.1479.
- [19] Trevor Hastie, Robert Tibshirani, and Jerome H. Friedman. *The Elements of Statistical Learning: Data Mining, Inference, and Prediction*, volume 2. Springer, 2009.
- [20] Julius Berner, Philipp Grohs, Gitta Kutyniok, and Philipp Petersen. *The Modern Mathematics of Deep Learning*, page 1–111. Cambridge University Press, 2022.
- [21] Lei Wu, Mingze Wang, and Weijie Su. The alignment property of sgd noise and how it helps select flat minima: A stability analysis, 2022. URL <https://arxiv.org/abs/2207.02628>.
- [22] Diederik Kingma and Jimmy Ba. Adam: A method for stochastic optimization. In *International Conference on Learning Representations (ICLR)*, San Diego, CA, USA, 2015.
- [23] Yushun Zhang, Congliang Chen, Naichen Shi, Ruoyu Sun, and Zhi-Quan Luo. Adam can converge without any modification on update rules. In *Advances in Neural Information Processing Systems*, page 35:28386, 2022.
- [24] Shun ichi Amari. Natural gradient works efficiently in learning. *Neural Computation*, 10:251, 1998.
- [25] S. Amari and S.C. Douglas. Why natural gradient? In *Proceedings of the 1998 IEEE International Conference on Acoustics, Speech and Signal Processing, ICASSP '98 (Cat. No.98CH36181)*, volume 2, pages 1213–1216 vol.2, 1998. doi: 10.1109/ICASSP.1998.675489.
- [26] Federico Becca and Sandro Sorella. *Quantum Monte Carlo approaches for correlated systems*. Cambridge University Press, 2017.
- [27] James Stokes, Josh Izaac, Nathan Killoran, and Giuseppe Carleo. Quantum natural gradient. *Quantum*, 4:269, 2020.
- [28] Arthur Jacot, Franck Gabriel, and Clément Hongler. Neural tangent kernel: Convergence and generalization in neural networks. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2018. URL <https://arxiv.org/abs/1806.07572>.
- [29] Xavier Glorot and Yoshua Bengio. Understanding the difficulty of training deep feedforward neural networks. In *Proceedings of the Thirteenth International Conference on Artificial Intelligence and Statistics (AISTATS)*, volume 9, pages 249–256, 2010.
- [30] Kaiming He, Xiangyu Zhang, Shaoqing Ren, and Jian Sun. Delving deep into rectifiers: Surpassing human-level performance on ImageNet classification. In *Proceedings of the IEEE International Conference on Computer Vision (ICCV)*, pages 1026–1034, 2015. doi: 10.1109/ICCV.2015.123.
- [31] Manzil Zaheer, Satwik Kottur, Siamak Ravanbakhsh, Barnabas Poczos, Russ R. Salakhutdinov, and Alexander J. Smola. Deep sets. In *Advances in Neural Information Processing Systems*, volume 30, 2017.
- [32] Justin Gilmer, Samuel S. Schoenholz, Patrick F. Riley, Oriol Vinyals, and George E. Dahl. Neural message passing for quantum chemistry. In *Proceedings of the 34th International Conference on Machine Learning (ICML)*, volume 70 of PMLR, pages 1263–1272, 2017. URL <https://arxiv.org/abs/1704.01212>.
- [33] Peter W. Battaglia, Jessica B. Hamrick, Victor Bapst, et al. Relational inductive biases, deep learning, and graph networks. *arXiv preprint arXiv:1806.01261*, 2018. URL <https://arxiv.org/abs/1806.01261>.

- [34] George Cybenko. Approximation by superpositions of a sigmoidal function. *Mathematics of Control, Signals and Systems*, 2:303–314, 1989. doi: 10.1007/BF02551274. URL <https://doi.org/10.1007/BF02551274>.
- [35] Kurt Hornik. Approximation capabilities of multilayer feedforward networks. *Neural Networks*, 4(2):251–257, 1991. ISSN 0893-6080. doi: [https://doi.org/10.1016/0893-6080\(91\)90009-T](https://doi.org/10.1016/0893-6080(91)90009-T). URL <https://www.sciencedirect.com/science/article/pii/089360809190009T>.
- [36] Moshe Leshno, Vladimir Ya. Lin, Allan Pinkus, and Shimon Schocken. Multilayer feedforward networks with a nonpolynomial activation function can approximate any function. *Neural Networks*, 6(6):861, 1993.
- [37] Andrew R. Barron. Universal approximation bounds for superpositions of a sigmoidal function. *IEEE Transactions on Information Theory*, 39(3):930–945, 1993. doi: 10.1109/18.256500.
- [38] Maziar Raissi, Paris Perdikaris, and George Em Karniadakis. Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations. *Journal of Computational Physics*, 378:686–707, 2019. doi: 10.1016/j.jcp.2018.10.045.
- [39] I. E. Lagaris, A. Likas, and D. I. Fotiadis. Artificial neural networks for solving ordinary and partial differential equations. *IEEE Transactions on Neural Networks*, 9(5):987–1000, 1998. doi: 10.1109/72.712178.
- [40] Tim De Ryck and Siddhartha Mishra. Numerical analysis of physics-informed neural networks and related models in physics-informed machine learning. *Acta Numerica*, 33:633–713, 2024. doi: 10.1017/S0962492923000089.
- [41] R. P. Feynman and Michael Cohen. Energy spectrum of the excitations in liquid helium. *Physical Review*, 102(5):1189–1204, 1956. doi: 10.1103/PhysRev.102.1189.
- [42] M. Holzmann, D. M. Ceperley, C. Pierleoni, and K. Esler. Backflow correlations for the electron gas and metallic hydrogen. *Physical Review E*, 68(4):046707, 2003.
- [43] Sandro Sorella. Green function monte carlo with stochastic reconfiguration. *Physical Review Letters*, 80(20):4558–4561, 1998. doi: 10.1103/PhysRevLett.80.4558.
- [44] Ao Chen and Markus Heyl. Empowering deep neural quantum states through efficient optimization. *Nature Physics*, 20:1476–1481, 2024. doi: 10.1038/s41567-024-02566-1.
- [45] Riccardo Rende, Luciano Loris Viteritti, Lorenzo Bardone, Federico Becca, and Sebastian Goldt. A simple linear algebra identity to optimize large-scale neural network quantum states. *Communications Physics*, 7:260, 2024. doi: 10.1038/s42005-024-01732-4.
- [46] Ronald J. Williams. Simple statistical gradient-following algorithms for connectionist reinforcement learning. *Machine Learning*, 8(3–4):229–256, 1992. doi: 10.1007/BF00992696.
- [47] Aki Vehtari, Daniel Simpson, Andrew Gelman, Yuling Yao, and Jonah Gabry. Pareto smoothed importance sampling. *Journal of Machine Learning Research*, 25, 2024. URL <https://arxiv.org/abs/1507.02646>.
- [48] Nicholas Gao and Stephan Günnemann. Neural Pfaffians: Solving many many-electron Schrödinger equations. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2024.
- [49] H. Flyvbjerg and H. G. Petersen. Error estimates on averages of correlated data. *The Journal of Chemical Physics*, 91(1):461–466, 1989.
- [50] Marius Jonsson. Standard error estimation by an automated blocking method. *Physical Review E*, 98:043304, 2018.
- [51] Simon Kornblith, Mohammad Norouzi, Honglak Lee, and Geoffrey Hinton. Similarity of neural network representations revisited. In *Proceedings of the 36th International Conference on Machine Learning (ICML)*, pages 3519–3529, 2019.
- [52] Jørgen Høgberget. Quantum Monte-Carlo studies of generalized many-body systems. Master’s thesis, University of Oslo, 2013. Source of the diffusion Monte Carlo benchmark energies for $N \in \{6, 12, 20\}$ used here; tabulated in Ref. [53].

- [53] Even M. Nordhagen, Jane M. Kim, Bryce Fore, Alessandro Lovato, and Morten Hjorth-Jensen. Efficient solutions of fermionic systems using artificial neural networks. *Frontiers in Physics*, 11: 1061580, 2023. doi: 10.3389/fphy.2023.1061580.
- [54] M. Taut. Two electrons in a homogeneous magnetic field: particular analytical solutions. *J. Phys. A: Math. Gen.*, 27:1045–1055, 1994. doi: 10.1088/0305-4470/27/3/040.
- [55] F. Pederiva, C. J. Umrigar, and E. Lipparini. Erratum: Diffusion Monte Carlo study of circular quantum dots [Phys. Rev. B **62**, 8120 (2000)]. *Phys. Rev. B*, 68:089901, 2003. doi: 10.1103/PhysRevB.68.089901.
- [56] Daniel Haas. Deep learning methods for quantum many-body systems: A study on neural quantum states. Master’s thesis, University of Oslo, September 2024. Compilation from which the reference energies used here were taken; their origin is attributed in Table 6.2 to Høgberget’s diffusion Monte Carlo calculations and, for $N = 2$, to Taut’s analytic solutions.
- [57] Chae-Yeun Park and Michael J. Kastoryano. Geometry of learning neural quantum states. *Physical Review Research*, 2:023232, 2020.
- [58] Sami Siljamäki. *Wave Function Methods for Quantum Dots in Magnetic Field*. PhD thesis, Helsinki University of Technology, 2003.
- [59] Eric Jang, Shixiang Gu, and Ben Poole. Categorical reparameterization with Gumbel-softmax. In *International Conference on Learning Representations (ICLR)*, 2017.
- [60] James Martens and Roger Grosse. Optimizing neural networks with kronecker-factored approximate curvature. In *Proceedings of the 32nd International Conference on Machine Learning (ICML)*, pages 2408–2417, 2015.