

Physics-Informed Neural Quantum States for Quantum Dots:
Tangent-Space Geometry, Message Passing, and Wigner
Molecules

Aleksander Sekkelsten

September 4, 2026

Abstract

How far can a compact, well-conditioned neural network go as a variational wavefunction for interacting electrons; what does letting the particles talk to one another actually buy; and can such a state be trained without a Markov chain? We take up these three questions for two-dimensional harmonic quantum dots—from the compact, kinetic-energy-dominated droplet at strong confinement to the dilute Wigner molecule at weak confinement—for $N \in \{2, 6, 12, 20\}$ electrons and trap frequencies ω from 1 down to 10^{-3} .

The ansatz is a Slater determinant dressed by a neural Jastrow correlator, which reshapes the amplitude, and a coordinate backflow, which reshapes the nodes. Both are made *safe by construction* for Laplacian-based training: coordinates are scaled to trap units, pair features are soft-core, and the electron–electron cusp is imposed analytically. Each learnable object comes in two forms—one that pools per-particle and pairwise features in a single pass, and one that passes messages between particles on a graph—and we read the trained states through the geometry of their log-derivatives, the quantum geometric tensor and the neural tangent kernel. Training is by residual pretraining followed by stochastic reconfiguration (SR), and, separately, by importance-sampled collocation that uses no Markov chain.

Message passing, it turns out, buys one thing: a *relational channel*. Seen through the tangent kernel it compresses the correlator onto roughly half as many effective directions as a single-pass network, and it keeps the backflow from collapsing: at the Wigner crystal a conventional backflow falls to a single collective mode and loses accuracy, while the message-passing backflow holds full rank and stays near reference quality. Underneath both is a single mechanism, and it is visible even at strong confinement where the two networks are identical in energy: message passing keeps the correlator’s dominant tangent directions aligned with the physical collective modes of the Hamiltonian, whereas a separable network reaches the same energy while leaving much of its tangent capacity along directions a compact physical dictionary does not resolve—a deficit merely disguised at strong confinement and sharpening at weak. The correlator’s loss of that alignment and the backflow’s rank collapse set in at one confinement, pointing to a common relational bottleneck. The same geometry governs the rest. SR helps in proportion to how ill-conditioned the tangent metric is, but only while that metric can still be estimated reliably; once the sampling that builds it collapses, inverting it does harm. And importance-sampled collocation matches SR-VMC while its sampling stays healthy, reaching a genuine Wigner molecule at $\omega = 0.0035$ with no Markov chain in the optimisation. SR-VMC carries the $N = 20$ energies, while the Markov-chain-free route is demonstrated for $N \leq 12$. Energies match diffusion Monte Carlo within uncertainty for $N = 2$ and to $\sim 10^{-2} - 10^{-1}\%$ for $N \in \{6, 12, 20\}$ wherever a DMC reference exists ($\omega \geq 0.1$); below that no benchmark is available, and we say so.

On the physics, the trained states let us map the Fermi-liquid-to-Wigner crossover from the inside. We give topology-resolved shell, bond-orientational, and stiffness diagnostics, with shell-mixture reconstructions of $g(r)$ matching the measured pair distribution to cosine similarity 0.989–1.000: at $\omega = 10^{-3}$ the $N = 6$ dot is a single ring with a robust $(1, 5) \leftrightarrow (0, 6)$ near-degeneracy (radial mode $\approx 114 a_0$), the $N = 12$ dot a two-shell molecule with a modal $(3, 9)$ split, and the $N = 20$ dot a three-shell molecule that collapses toward a single shell within one decade in ω . One thread runs through architecture, optimiser, and training paradigm alike: what matters is not raw capacity but staying aligned with the low-dimensional, physically collective structure of the problem—the same structure that, in real space, is the Wigner molecule.

*To go wrong in one's own way is better than to go right in
someone else's.*

FYODOR DOSTOEVSKY

Crime and Punishment, Part III, Chapter I (Constance Garnett translation)

Preface

Life must be understood backwards. But they forget the other proposition, that it must be lived forwards.

SØREN KIERKEGAARD
Journal entry JJ:167 (1843)

When I came to UiO, Morten read my background and said: “You know PINNs, you know many-body quantum systems. I want you to do PINNs on quantum dots.” I said “sure” and then asked the only sensible follow-up question I had: “Where do I start?”

I did not realise then that this was the question I had been assigned to answer. This thesis is the long answer. The short answer is: start by arranging to be wrong in a way that can be measured.

That is what variational methods make possible. The variational principle does not say how to improve a trial wavefunction, but it does guarantee that its energy lies above the ground-state energy. Failure therefore has a direction: the energy decreases, or it does not. For a project with few fixed landmarks, that was enough to begin. The harder task was learning what a disappointing result meant.

It was a solitary problem, not for want of literature—neural quantum states, FermiNet, PauliNet, and neural backflow offered much to learn from—but because the particular route I was trying to take had little precedent. When an experiment failed, it was rarely clear whether the method was unsound, the architecture inadequate, the optimiser poorly matched to the problem, or the code quietly broken. For long stretches, the honest answer was that I did not know. The work became an attempt to construct tests that could tell these explanations apart.

The most important lessons were therefore failures. An error in the importance-sampling code went undetected for months because its signature—loss of accuracy at larger particle numbers and weaker confinement—was exactly what an inadequate architecture might have produced. In Nd dimensions, a small error in a log-density grows linearly with dimension and is amplified in the normalised weights; the bug was invisible where experiments were cheap and ruinous where they mattered. Discovering it damaged my confidence in everything else. Re-establishing that confidence, rather than assuming it, became a substantial part of the later work.

Other failures taught the same lesson from different directions. A Langevin refinement step introduced bias into the gradient. A second-order optimiser became least reliable when the estimated metric was too noisy to trust. Neither result was useful merely because it was negative. Each exposed an assumption that had been invisible until the computation violated it. Negative results of this kind are not a concession or an embarrassment; they are part of the method, and I have written them up as such.

The project also changed what I thought the network was for. I began by asking whether a machine-learning construction could solve the many-body Schrödinger equation. I ended up using the trained state as an apparatus for asking what structure the approximation had captured and which directions of variation mattered. That shift, from treating a network as an answer to treating it as something that can itself be examined, is the thread running through this thesis.

Something else surfaced along the way that I have not stopped thinking about. The obstacles in this work have a persistent double character: each is simultaneously a physics problem and an optimisation problem, and it is usually arbitrary which name you give it. Stochastic reconfiguration, invented by physicists to optimise variational wavefunctions, is natural gradient descent on the variational manifold. The REINFORCE estimator from reinforcement learning is the score-function gradient of the variational energy. The Fisher information matrix of statistics is the quantum geometric tensor of physics. These were not borrowed by one field from another; they were found independently, by communities that rarely read each other's papers. Coming to recognise that they are the same mathematics felt less like discovering a bridge between two fields than like realising there had only ever been one, approached from opposite sides.

Looking back, it is easy to make a catalogue of what I would now do differently. Most of that knowledge, however, was purchased by the failures and could not have been read off in advance. The regret and the result are the same object seen from two sides. That is worth remembering, and easy to forget.

The most consistent source of inspiration, motivation, and steadiness through all of it has been my girlfriend, Elida. That support has been as constant as a mountain, and without it I would not have had the nerve to begin this, let alone the depth to finish it. I am more grateful than I can put down here.

I owe a warm and sincere thanks to my family, who have supported me in every undertaking I have had and have simply always been there; and to my friends, inside CCSE and out, who supplied the joy, the laughter, and a good deal of the sanity this project required.

To Morten, again, and properly: thank you for the assignment, for the two years, and above all for a genuine and undisguised interest in what I was finding at every stage—including the stages where what I was finding was that I had been wrong. That interest is most of what encouragement actually consists of.

Which is, at last, the answer to the question I asked on the first day. You start by being willing to be wrong on the record, and then you keep the record.

Contents

Preface	2
I Introduction	9
II Theory	15
1 Quantum Theory	16
1.1 Mathematical Foundations	16
1.1.1 Hilbert Spaces and Function Representations	16
1.1.2 Operators and Matrix Representations	17
1.1.3 States in Hilbert Space	17
1.2 Quantum Mechanics	17
1.2.1 The Schrödinger Equation	17
1.3 Second Quantization	18
1.3.1 Motivation: Why first-quantized wavefunctions don't scale	18
1.3.2 Fock space and occupation representation	18
1.3.3 Operators in second quantization	18
1.4 Hartree–Fock (HF)	19
1.4.1 HF energy functional and the Fock operator	19
1.5 Full Configuration Interaction (FCI)	19
1.5.1 Slater–Condon rules and sparsity	20
1.5.2 Diagonalization and symmetry reduction	20
1.5.3 Role in this thesis	20
1.6 Quantum Dots	20
1.6.1 Model and units	20
1.6.2 One-particle structure and shell filling	20
1.6.3 Spin structure and Slater form	20
1.6.4 Kato's Cusp Conditions (2D Coulomb systems)	21
1.7 Structural diagnostics and Wigner markers	21
1.7.1 Pair distribution $g(r)$, $P(r)$, and width markers	22
1.7.2 Automatic shell detection and topology labeling	22
1.7.3 Orientational order and angular “Lindemann”	22
1.7.4 Scaling check and two-body benchmarks ($N = 2$)	23
1.7.5 Reporting and stability checks	23
1.8 Variational Methods	23
1.8.1 Variational Monte Carlo (VMC)	23
1.8.2 Diffusion Monte Carlo (DMC)	24
1.8.3 Applications and Numerical Considerations	25
1.8.4 Conclusion	25
2 Machine Learning	26
2.1 Machine Learning as Statistical Learning	26
2.1.1 Types of learning	26
2.1.2 Generalisation and the bias–variance tradeoff	27
2.2 Optimisation	27

2.2.1	Gradient descent	27
2.2.2	Stochastic gradient descent (SGD)	27
2.2.3	Momentum and adaptive methods	28
2.2.4	Second-order methods and natural gradient	28
2.2.5	The variational tangent space: geometric tensor, tangent kernel, and effective dimension	28
2.2.6	Optimisation challenges in deep learning	29
2.3	Feed-forward neural networks	29
2.3.1	From linear models to nonlinear function classes	30
2.3.2	Activation functions	30
2.3.3	Initialisation	31
2.3.4	Summary	32
2.4	Architectures for many-body wavefunctions	32
2.4.1	Permutation symmetry and equivariance	32
2.4.2	Permutation-invariant pooling: DeepSets	32
2.4.3	Message passing and graph networks	33
2.4.4	The CTNN: one message-passing instance	33
2.4.5	Relevance to the ansatz	33
2.5	Expressivity, approximation, and generalisation	33
2.5.1	Universal approximation theorems	33
2.5.2	Depth versus width	34
2.5.3	Curse of dimensionality	34
2.5.4	Generalisation in overparameterized networks	34
2.5.5	Optimisation in deep networks	35
2.5.6	Implications for physical modelling	35
2.6	Physics-Informed Neural Networks	35
2.6.1	Formulation of PINNs	35
2.6.2	Automatic differentiation	36
2.6.3	Theoretical difficulties	36
2.6.4	Relevance to this thesis	37
2.6.5	Implications for scientific machine learning	37
2.7	Summary and Outlook	37

III Methods 38

3	Trial Wavefunction	39
3.1	Preliminaries	39
3.2	Overview of the ansatz	39
3.3	Slater reference	40
3.4	Neural correlator W_θ	40
3.4.1	Separable correlator (DeepSets baseline)	40
3.4.2	Analytic short-range cusp (shared)	41
3.4.3	Separable readout	41
3.4.4	Message-passing correlator (CTNN)	41
3.5	Backflow	42
3.5.1	Conventional backflow (single-round messages)	42
3.5.2	CTNN backflow (copesheaf)	42
3.5.3	Shared output, scaling, and initialisation	43
3.5.4	Short-range behaviour and gates	43
3.5.5	Symmetries and COM neutrality	43
3.5.6	Effect on the nodes and variational safety	43
3.5.7	Regularization, initialisation, and cost	43
3.5.8	Mapping to implementation	44
3.6	Architecture sizes and hyperparameters	44

4	Optimisation	45
4.1	Primary objective and two-stage training	45
4.2	Configuration-space sampling	46
4.2.1	Stratified capped-simplex mixture	46
4.3	Residual-based training (stage I)	47
4.4	Stochastic Reconfiguration (stage II)	47
4.5	Sampling paradigms and the weak-form collocation objective	48
4.5.1	Analytic proposal and importance resampling	48
4.5.2	Physics-informed proposal for the Wigner regime	48
4.6	Units, scaling, and batch construction	49
4.7	Workflow summary	49
4.8	Statistical uncertainties	49
5	Analysis	50
5.1	Structural diagnostics and Wigner markers	50
5.1.1	Pair distribution and radial statistics	50
5.1.2	Automatic shell detection and topologies	50
5.1.3	Orientational order and “ring stiffness”	51
5.1.4	Scaling check for $N = 2$	51
5.1.5	Reporting and stability checks	51
5.2	Representation analysis	51
5.2.1	Preprocessing and PCA	52
5.2.2	Entropy effective rank	52
5.2.3	PC ablations of the head	52
5.2.4	PC1 block power decomposition	52
5.2.5	Linear probes from PCs to coarse physics	52
5.2.6	Near-field gradient concentration (residual head)	53
5.2.7	Backflow geometry and energy-locality shares	53
5.2.8	Alignment of correlator manifolds (noBF vs BF)	53
5.2.9	Uncertainty and stability	53
5.3	Tangent-kernel diagnostics	53
IV	Results	55
6	Quantum dots: energies, representations, and Wigner molecules	56
6.1	Training protocol and energy overview	56
6.1.1	Energies	57
6.2	What the networks learn	59
6.2.1	Correlator geometry: a compact manifold that reorganises	60
6.2.2	Backflow: a high-rank field with a low-rank footprint	60
6.2.3	PINN-CTNN coupling: from independence to cooperation	61
6.2.4	Synthesis	62
6.3	Wigner-molecule crossover	63
6.3.1	Observables and diagnostics	63
6.3.2	One-body densities: visual crossover	63
6.3.3	Energy-based crossover diagnostics	64
6.3.4	Radial localization and pair correlations	64
6.3.5	Shell topology, angular order, and the rotating Wigner molecule	66
6.3.6	Synthesis: the Wigner-molecule crossover	69
6.4	Collocation-assisted training	69
6.4.1	Training methodology	70
6.4.2	Architecture variants	70
6.4.3	Results	71
6.4.4	What worked and what did not	72
6.4.5	Reliability analysis: the gradient-quality chain	73

7	Architecture, optimiser, and paradigm through the tangent kernel	77
7.1	Q1 — Architecture: message passing supplies a relational channel	78
7.1.1	Q1a — In the Jastrow: message passing compresses the correlator	78
7.1.2	Q1b — In the backflow: message passing prevents rank collapse	79
7.1.3	The unified reading of Q1	81
7.2	Q2 — Optimiser: SR earns its keep where the metric is ill-conditioned	83
7.3	Q3 — Paradigm: the sampling measure sets conditioning, variance, and reach	83
7.4	The collocation frontier: a physics-informed proposal for the Wigner regime	84
7.4.1	The origin-centred proposal misses the Wigner shell	84
7.4.2	The Wigner-ring proposal	84
7.4.3	Angular crystallisation is the deep-Wigner wall	85
7.4.4	An MCMC-free cascade reaches $\omega = 0.0035$	85
7.4.5	The remaining wall: a fine-tuned Slater–Jastrow balance, not the sampler	85
7.5	Synthesis and discussion	86
V	Discussion	87
8	Discussion	88
8.1	Discussion of energy results	88
8.2	Discussion of the Wigner–molecule regime	90
8.3	Discussion of learned representations	91
8.4	Discussion of collocation-assisted training	93
8.4.1	The central result: MCMC-free training to DMC accuracy	94
8.4.2	Why REINFORCE succeeds where residual losses fail	94
8.4.3	The sampling bottleneck and its regime dependence	94
8.4.4	The optimiser regime boundary	95
8.4.5	Why Langevin refinement fails	95
8.4.6	Architectural trade-offs at scale	96
8.4.7	Comparison with MCMC-based results	96
8.4.8	Implications and future directions	96
VI	Conclusion	98
9	Conclusions	99
VII	Appendices	101
A	Laplacian conditioning	102
A.1	Residual training for Schrödinger	102
A.2	Linearized geometry and the matrix A	102
A.3	Bi-Laplacian scaling	103
A.4	Barron/spectral perspective	104
A.5	Implications for Stochastic Reconfiguration	104
B	Coulomb conditioning	105
B.1	Coulomb structure and singularity	105
B.2	2D cusp conditions: antiparallel vs parallel spins	105
B.3	Why Coulomb is ill-conditioned (2D emphasis)	106
B.4	Barron/spectral perspective	107
B.5	Implications for Stochastic Reconfiguration	107
C	The collocation–backflow catch-22	109
C.1	Setup and notation	109
C.2	The divergence of E_L near nodes	109
C.3	The geometric mismatch argument	110
C.4	The catch-22	111

C.5	Why VMC + SR escapes	111
C.6	Experimental evidence	111
C.6.1	Baseline: PINN alone vs. PINN + CTNN	112
C.6.2	Sampling strategies	112
C.6.3	Loss modifications	112
C.6.4	Gradient regularisation	113
C.6.5	Gradient preconditioning	113
C.6.6	Summary of all interventions	113
C.7	Connection to Appendices A and B	113
D	Post-catch-22 experimental history and philosophy	115
D.1	The experimental programme	115
D.2	Current frontier: low- ω and high- N stabilisation	116
D.3	Interpretive summary	116
E	Declaration on the Use of Artificial Intelligence	117

Part I

Introduction

Introduction

The world was to me a secret which I desired to divine.

MARY SHELLEY

Frankenstein; or, The Modern Prometheus, Chapter II

A system of many interacting electrons is described completely by its wavefunction. The same object puts the system almost out of reach, because a complete description of N interacting particles is a function of all their coordinates at once, and the space of those coordinates grows with every particle added. The equation that fixes this wavefunction is known exactly; the entire difficulty is solving it.

The competition that makes it hard is physical. Kinetic energy tends to delocalise the particles; interactions and external potentials drive localisation and correlation; and the balance between them plays out in a configuration space far too large to enumerate. Yet the quantities we need—energies, correlation functions, phase boundaries—are precisely the ones that depend on getting that balance right. This is the *many-body problem*, the point at which quantum mechanics turns from a compact set of principles into one of the hardest computational challenges in physics.

A wide range of electronic-structure methods has been developed to address this. Mean-field approaches such as Hartree–Fock replace the full many-body problem with an effective single-particle picture, where each electron moves in an average field from the others. This captures Fermi statistics and some exchange effects but leaves most correlation physics untreated. More systematic methods such as Full Configuration Interaction (FCI) are, in principle, exact within a finite basis, but their cost grows combinatorially with system size. Coupled Cluster (CC) theory provides a more efficient and size-extensive hierarchy of approximations, but it remains computationally demanding and can struggle in strongly correlated regimes or near degeneracies.

Quantum Monte Carlo (QMC) methods, including Variational Monte Carlo (VMC) and Diffusion Monte Carlo (DMC), take a different approach: they represent the wavefunction implicitly through stochastic sampling. For suitable trial states, they can produce highly accurate benchmark energies and correlations for continuum systems. However, QMC is not free of difficulties. The fermion sign problem limits direct simulations, fixed-node approximations inherit bias from the chosen trial nodes, and constructing high-quality trial wavefunctions is increasingly difficult as systems become more strongly correlated or inhomogeneous. In practice, then, much of the many-body problem reduces to a narrower one: choosing an expressive, well-conditioned ansatz and optimising it reliably.

Machine learning and many-body wavefunctions. In parallel with these developments, machine learning—especially deep learning—has become a standard tool for representing complex, high-dimensional functions and fitting them to data. Neural networks now play a central role in domains such as computer vision, language modelling and reinforcement learning. They combine flexible parametric families with inductive biases (e.g. symmetry and locality), normalisation, and optimisation procedures that scale well with dimension.

In physics, this has led to two broad uses of neural networks. In the first, they act as *surrogates*: they learn mappings from parameters or boundary conditions to observables, or approximate solution operators of partial differential equations. Physics-informed neural networks (PINNs) and operator-learning approaches fall into this category by embedding differential operators and constraints directly into the training objective. In the second, neural networks are used as *variational ansätze* for quantum states (neural quantum states, NQS), where the network outputs amplitudes, phases or log-probabilities for many-body configurations and is trained by variational Monte Carlo.

One argument for this second use is routinely overstated, and disposing of it first clarifies what remains. Universal approximation results guarantee that a sufficiently large network can *represent* any suitably

regular function to arbitrary accuracy. They say nothing about whether gradient-based optimisation will *find* that representation, nor at what sample cost. This thesis measures that gap directly: the same ansatz, demonstrably able to express the ground state, is placed in training schemes that do and do not allow it to get there. Representability is cheap; trainability is the difficulty.

A weaker but sounder argument survives that objection. Established methods pay for accuracy in one of two currencies. FCI is exact within a finite basis, but its cost grows combinatorially with that basis, so a modest two-dimensional dot of twenty electrons is already far beyond reach. Cheaper methods such as coupled cluster buy their favourable scaling instead by deciding in advance which correlations may be described at all.

A variational neural ansatz pays in neither currency. It is not an expansion whose size must grow with the space it covers, and it fixes no prior truncation on which correlations are available. It is a parameterisation free, in principle, to concentrate its capacity wherever the solution actually varies. Whether it does so in practice is an empirical question, and this thesis treats it as one.

That question also suggests a different way to use a solver. Nearly every established method for the many-body Schrödinger equation is constructed outward from physical principles: an approximation is chosen, justified physically, and then implemented. A trained network inverts the order. It is obtained largely on machine-learning terms, and only afterwards interrogated as to what it has represented—and unlike an implicit stochastic representation, it is a concrete parameterised object whose internal structure can be examined directly. This motivates a question that the size of the problem does not by itself settle. A twenty-electron dot is a function of forty coordinates, and any basis expansion of it grows combinatorially—but it does not follow that the solution itself is forty-dimensional in any useful sense. How many directions does the *solved state* actually vary along? Not the dimension of the equation, but the dimension of the answer.

We make that question quantitative through the geometry of the wavefunction’s log-derivatives. Perturbing the parameters of a trained ansatz over an ensemble of configurations and collecting the resulting sensitivities into a covariance matrix yields, in the simplest reading, a principal-component analysis of the state: the eigenvalues report how much variation each independent direction carries, and their distribution counts how many directions are doing genuine work. The physically appropriate version of this covariance is the quantum geometric tensor (equivalently, the Fisher metric), and the corresponding count is the participation ratio of its spectrum—an effective dimension rather than a hard rank, insensitive to the raw parameter count. Chapter 7 develops this instrument and applies it to architecture, optimiser, and training paradigm alike; the questions it answers are how this effective dimension behaves as particles are added and as the trap is opened, and whether different architectures arrive at the same solution by different routes.

In principle, then, neural networks provide highly expressive ansätze for quantum problems. In practice, several issues arise. Objectives based on local energies involve second derivatives and can be numerically delicate. Coordinate parameterisations that do not respect known structure can lead to large gradients and Laplacian spikes near electron coalescence. Generic architectures can waste capacity on degrees of freedom that are irrelevant from a physical point of view. These challenges motivate hybrid approaches in which as much known structure as possible—antisymmetry, cusp conditions, scaling, and symmetries—is built into the ansatz and its coordinates, and the network is used to learn the remaining correlation corrections.

A further ingredient concerns not *what* structure is imposed but *how* the network represents the electrons. Correlation is intrinsically relational—what one electron should do depends on where the others are—so architectures that let particles exchange information (message passing), rather than processing each coordinate independently, are a natural candidate for the correlator and for the backflow that deforms the nodes. How much this relational capacity actually buys, and why, is one of the questions this thesis takes up.

Where this work sits. Representing a quantum state by a neural network and optimising it variationally was introduced by Carleo and Troyer [1]; for continuum fermions the idea has since matured into highly accurate *ab initio* solvers such as FermiNet [2] and PauliNet [3], and into explicit neural *backflow* transformations [4]—the direct ancestor of the coordinate backflow used here. Message passing, the relational mechanism at the centre of this thesis, has recently been built into neural quantum states for extended fermion systems [5, 6]. Against that backdrop the present work is deliberately narrow and complementary. Rather than pushing a new architecture to ever larger systems, it asks *why* the relational channel helps, and uses the tangent-kernel geometry to make the answer precise; it focuses on the two-dimensional quantum-dot Wigner regime, where correlation is strongest and where topology-resolved

structural diagnostics remain sparse; and it treats conditioning, the optimiser, and Markov-chain-free training as objects of study in their own right rather than fixed choices. To our knowledge these questions have not previously been addressed together for parabolic quantum dots.

Quantum dots as a controlled testbed. Two-dimensional parabolic quantum dots provide a clean and tunable setting in which to explore such ideas. They offer a simple route from Fermi-liquid-like behaviour to Wigner-molecule physics. As the trap frequency ω decreases, the single-particle level spacing collapses while Coulomb repulsion becomes dominant. Electrons then tend to localize into ring or shell patterns. In the lab frame, rotational symmetry and quantum fluctuations smear these into smooth densities; in a co-rotating frame, one recovers crystalline-like structures whose topology depends on particle number. For $N=6$ and $N=12$, classical and semiclassical studies have identified preferred shellings such as (1, 5) and (3, 9) and mapped parts of the crossover [7–10].

From the perspective of this thesis, quantum dots are attractive for two reasons. First, they are genuinely non-trivial: Coulomb singularities, strong correlations and an actual Fermi-liquid-to-Wigner crossover appear already for moderate N . Second, they are still simple enough that accurate DMC references exist in many regimes and that detailed structural diagnostics (shell occupancies, bond order, virial ratios) are computationally feasible. For fully quantum dots, however, quantitative topology-resolved diagnostics (occupancy fractions, bond-orientational order, shell “stiffness”) and a systematic picture of the Wigner regime remain relatively sparse. In addition, it is not well understood how modern neural ansätze internally represent these states.

Scope and questions. In this thesis we revisit two-dimensional harmonic quantum dots with a Slater–Jastrow(+backflow) ansatz implemented as a physics-informed neural network. The design is meant to be *safe by construction* for Laplacian-based objectives: coordinates are scaled to trap units ($\tilde{R} = \sqrt{\omega}R$), learned pair features are soft-core (avoiding spurious $1/r$ behaviour in derivatives), and the electron–electron cusp is enforced analytically rather than learned. On top of a Slater determinant of harmonic-oscillator orbitals, two learnable objects carry the correlation: a Jastrow correlator, which reshapes the amplitude, and a coordinate backflow, which reshapes the nodes. Each can be built either without message passing—encoding and pooling per-particle and pairwise features in a single pass, as in a DeepSet—or as a *message-passing* network in which the particles exchange and update information through explicit, transported edge state before a quantity is read out.

Once such an ansatz reaches reference accuracy—which, as we will see, a surprisingly compact one does—the questions worth a thesis are *why* its parts behave as they do, and what it can tell us about the electrons. We pursue four. The first three we read through a single lens: the geometry of the wavefunction’s log-derivatives, captured by the quantum geometric tensor and the neural tangent kernel.

1. **Architecture.** What does message passing buy that a separable network cannot? We find it supplies a *relational channel*—a way to condition each particle on where the others are—that compresses the correlator onto fewer effective directions, keeps the correlator’s leading tangent modes aligned with the physical collective modes of the Hamiltonian, and keeps the backflow from collapsing to a single collective mode in the strongly correlated limit. A separable network, by contrast, *estimates* the correlation energy without *representing* it—a deficit already measurable at strong confinement, where the energies are identical. The correlator’s loss of physical alignment and the backflow’s rank collapse set in at the same confinement and shift together when the correlator is swapped, supporting a common relational bottleneck in the amplitude and nodal channels.
2. **Optimiser.** When does natural-gradient (stochastic reconfiguration) preconditioning actually change the result? Its value tracks the conditioning of the tangent metric: negligible for well-conditioned, $|\Psi|^2$ -sampled VMC, decisive for the ill-conditioned collocation objective.
3. **Paradigm.** Can an accurate wavefunction be trained *without* Markov-chain Monte Carlo, using importance-sampled collocation instead—and how far into the correlated regime does that reach before the sampling loses the target?
4. **Physics.** Can the trained wavefunction serve as a scientific instrument for Wigner-molecule physics—shell topologies, angular order, and melting—and does its internal representation reflect that physics?

These are not four separate stories. The same low-dimensional, physically collective structure that message passing captures is what natural gradients exploit, what $|\Psi|^2$ sampling stays aligned with, and what ultimately crystallises into the Wigner molecule itself.

Diagnostics and relation to prior work. To address these questions, we go beyond total energies. Alongside DMC benchmarks, we use:

- automatic shell detection and shell-resolved reconstructions of the pair distribution $g(r)$, testing whether the global $g(r)$ is explained by shell geometry and combinatorics;
- bond-orientational order parameters $|\Phi_m|$ and angular Lindemann-like ratios, adapted from studies of classical Wigner and Yukawa layers [11], to quantify ring order and angular stiffness;
- the density parameter r_s , estimated from the first peak of $g(r)$ following Egger *et al.* [7], to locate our dots along the Fermi-liquid-to-Wigner scale;
- representation-focused diagnostics for the neural ansatz: entropy effective ranks of correlator and backflow features, principal component ablations, and simple linear probes that relate latent axes to physical observables (global size, radial variance, near-origin mass);
- tangent-kernel diagnostics of the ansatz itself: the quantum geometric tensor and neural tangent kernel, the effective dimension of the variational tangent space, and symmetric state overlaps used to compare architectures and training paradigms on an invariant footing.

All of these are applied as post-processing to $|\Psi|^2$ samples, so the same analysis could, in principle, be used to study classical, quantum-chemical, or other neural wavefunctions on the same footing.

Main contributions. The main contributions of this thesis are:

1. **Compact, conditioned accuracy.** A small Slater-Jastrow(+backflow) ansatz in trap units reaches DMC-level energies for $N=2$ and holds $\sim 10^{-2}$ – $10^{-1}\%$ relative errors for $N \in \{6, 12, 20\}$ across $\omega \in \{0.1, 0.5, 1.0\}$, extrapolating smoothly to ultra-weak traps ($\omega = 10^{-3}$). Conditioning—trap units, analytic cusp, safe pair features—matters at least as much as network size for stability and accuracy in these dots.
2. **Message passing as a relational channel, made precise by the tangent kernel.** Read through the quantum geometric tensor, a message-passing correlator represents the ground state in roughly half as many effective tangent directions as a separable one, and a message-passing backflow retains full rank exactly where a conventional, single-round-message backflow collapses to a single collective mode at the Wigner crystal and loses accuracy. The two effects point to a common relational mechanism—a channel the physics requires—and both are invisible in the total energy until weak confinement turns the geometric deficit into an energetic one.
3. **Optimiser and paradigm, governed by the same geometry.** Natural-gradient (SR) preconditioning helps precisely when the tangent metric is ill-conditioned: negligibly for $|\Psi|^2$ -VMC, decisively for collocation. And an MCMC-free, importance-sampled collocation objective matches VMC when the sampling is healthy and, with a physics-informed proposal built from the Wigner shell, reaches a genuine Wigner molecule at $\omega = 0.0035$ with no Markov chain in the gradient training (a Metropolis probe is used only for checkpoint selection and final certification).
4. **Topology-resolved Wigner diagnostics.** Across $\omega \in [1, 10^{-3}]$ we quantify the crossover using two-shell probabilities and inner-ring occupancy histograms, shell-mixture reconstructions of $g(r)$ with cosine similarity 0.989–1.000 to global data, and sector-resolved bond order and Lindemann indices. At $\omega = 10^{-3}$ the $N=6$ dot forms a persistent single-ring Wigner molecule with a clear $(1, 5) \leftrightarrow (0, 6)$ near-degeneracy, while the $N=12$ dot forms a two-shell molecule with a commensurate $(3, 9)$ split as the dominant topology, complementing classical and quantum studies of parabolic dots [7–9, 12].
5. **Representation insight, with its gauge explicitly drawn.** The correlator features occupy a low-dimensional manifold (one to three principal directions) whose leading axes act as collective coordinates for global size and radial fluctuations, with a $\phi \leftrightarrow \psi$ recomposition across regimes, while backflow spans many local modes. We take care to separate the parts of this picture that are *gauge-like*—fixed by how correlation is divided between the Jastrow and the backflow, two substitutable channels—from the invariants (energy, state overlap, and the energetic price of ablating a component) on which the architectural conclusions actually rest.

Thesis structure. The remainder of the thesis is organised as follows. The next chapter sets out the theoretical background: the many-body Schrödinger equation in harmonic traps, the electronic-structure methods we benchmark against, the physics of Wigner molecules, and the machine-learning tools used here—neural function approximators, physics-informed objectives, neural quantum states, and the tangent-kernel geometry (quantum geometric tensor and neural tangent kernel) that serves as the analytical lens for the results. A method chapter then presents the concrete Slater–Jastrow(+backflow) PINN ansatz in its separable and message-passing forms, the optimisation pipeline (residual pretraining, stochastic reconfiguration, and importance-sampled collocation), and the diagnostics used throughout. The results are given in two chapters. The first treats accuracy, the learned representation, and the Wigner-molecule physics for $N \in \{2, 6, 12, 20\}$; the second reads the architecture, optimiser, and training paradigm through the tangent kernel. A discussion and a concluding chapter follow, and the appendices develop the conditioning theory (Laplacian and Coulomb) and the collocation–backflow catch-22 that motivated the weak-form objective.

Part II

Theory

Chapter 1

Quantum Theory

Men do not know how what is at variance agrees with itself. It is an attunement of opposite tensions, like that of the bow and the lyre.

HERACLITUS
Fragment 51 (DK B51)

This chapter lays the foundation on which the rest of the thesis rests. We first introduce the mathematical framework of quantum mechanics, then discuss the challenges of many-body systems and how second quantization provides a more tractable approach. We then introduce the model systems used throughout this thesis to benchmark and test our methods, and present the Hartree–Fock method, a fundamental mean-field approach to approximating many-body wavefunctions. Finally, we discuss Full Configuration Interaction (FCI), which provides an exact solution to the many-body problem within a finite basis set, and Variational Monte Carlo (VMC), a stochastic method for approximating ground-state properties. The following chapter then shifts to machine learning—statistical learning theory, optimisation, neural networks, and physics-informed neural networks.

1.1 Mathematical Foundations

At the centre of quantum mechanics lies a rich mathematical framework built upon linear algebra and functional analysis. This section introduces the essential mathematical concepts and notations that underpin quantum theory, including Dirac notation, operators, Hilbert spaces, and the representation of quantum states and observables.

1.1.1 Hilbert Spaces and Function Representations

In essence, quantum mechanics is formulated in complex vector spaces—specifically *Hilbert spaces*: complex inner-product spaces that are *complete* in the norm induced by their inner product, so that every Cauchy sequence converges within the space and limits and basis expansions are well defined [13]. For square-integrable functions the inner product is

$$\langle \phi | \psi \rangle = \int \phi^*(x) \psi(x) dx,$$

which, in the physics convention (linear in the second slot), obeys conjugate symmetry $\langle \phi | \psi \rangle = \langle \psi | \phi \rangle^*$, linearity in the ket $\langle \phi | \alpha \psi_1 + \beta \psi_2 \rangle = \alpha \langle \phi | \psi_1 \rangle + \beta \langle \phi | \psi_2 \rangle$, and positivity $\langle \psi | \psi \rangle \geq 0$ (with equality iff $|\psi\rangle = 0$); it induces the norm $\|\psi\| = \sqrt{\langle \psi | \psi \rangle}$. An orthonormal basis $\{\psi_i\}$ satisfies $\langle \psi_i | \psi_j \rangle = \delta_{ij}$. A function $a(x)$ expands in a basis $\{\psi_i(x)\}$:

$$a(x) = \sum_i \psi_i(x) a_i, \quad a_i = \int \psi_i^*(x) a(x) dx.$$

The Dirac delta $\delta(x - y)$ replaces the Kronecker delta in continuous spaces. A vector $|a\rangle$ in an n -dimensional Hilbert space $\mathcal{H}$ can be decomposed into a complete orthonormal basis $\{|i\rangle\}$:

$$|a\rangle = \sum_i |i\rangle \langle i|a\rangle = \sum_i a_i |i\rangle,$$

where $a_i = \langle i|a\rangle$ are complex coefficients. The completeness relation,

$$I = \sum_i |i\rangle \langle i|,$$

ensures the basis spans $\mathcal{H}$. Vectors in Dirac notation have dual "bra" counterparts $\langle a| = |a\rangle^\dagger$, enabling inner products $\langle a|b\rangle$ and outer products $|a\rangle \langle b|$.

1.1.2 Operators and Matrix Representations

Linear operators $\hat{O}$ map vectors to vectors: $\hat{O}|a\rangle = |b\rangle$. Their adjoints $\hat{O}^\dagger$ satisfy $\langle a|\hat{O}^\dagger = \langle b|$. Hermitian operators ($\hat{O}^\dagger = \hat{O}$) represent observables, while unitary operators ($\hat{O}^\dagger = \hat{O}^{-1}$) preserve inner products [14, 15]. In a basis $\{|i\rangle\}$, operators are represented by matrices:

$$O_{ij} = \langle i|\hat{O}|j\rangle.$$

Non-commutativity of operators is quantified by the commutator:

$$[\hat{A}, \hat{B}] = \hat{A}\hat{B} - \hat{B}\hat{A}.$$

1.1.3 States in Hilbert Space

A (pure) quantum state is a normalised vector $|\psi\rangle \in \mathcal{H}$ (global phase is physically irrelevant). Computations proceed by choosing an orthonormal (ON) basis $\{|i\rangle\}_{i=1}^\infty$ and expanding

$$|\psi\rangle = \sum_i c_i |i\rangle, \quad c_i = \langle i|\psi\rangle, \quad \sum_i |c_i|^2 = \|\psi\|^2.$$

Completeness is the resolution of the identity, $\sum_i |i\rangle \langle i| = \mathbb{I}$. For continuous ON bases (e.g. position), $\int |x\rangle \langle x| dx = \mathbb{I}$ and the wavefunction is the coordinate representation $\psi(x) = \langle x|\psi\rangle$.

Different ON bases are related by a unitary change of basis. Given $\{|i\rangle\}$ and $\{|\alpha\rangle\}$,

$$|\alpha\rangle = \sum_i U_{\alpha i} |i\rangle, \quad U_{\alpha i} = \langle \alpha|i\rangle, \quad U^\dagger U = \mathbb{I}.$$

This freedom is exploited to match the physics (symmetry, locality, boundary conditions), e.g. harmonic-oscillator orbitals for confined charges or plane waves for periodic systems, leading to sparser representations and faster convergence.

For N particles the Hilbert space factorizes as $\mathcal{H}_N = \mathcal{H}_1^{\otimes N}$. With a single-particle ON set $\{|\varphi_p\rangle\}$, a convenient fermionic ON basis is the set of Slater determinants $\{|\Phi_I\rangle\}$ built from N distinct orbitals; any N -body state expands as

$$|\Psi\rangle = \sum_I C_I |\Phi_I\rangle, \quad \sum_I |C_I|^2 = \|\Psi\|^2.$$

In practice one truncates to a finite active space guided by energy scales and symmetry quantum numbers, which makes many-body calculations tractable while retaining the relevant physics.

1.2 Quantum Mechanics

1.2.1 The Schrödinger Equation

In a Hilbert space $\mathcal{H}$, the time evolution of a normalised state $|\Psi(t)\rangle$ is postulated to be unitary and generated by the Hamiltonian H [14, 15]:

$$i \partial_t |\Psi(t)\rangle = H |\Psi(t)\rangle, \quad (\hbar = 1).$$

If H is time-independent, solutions separate as

$$|\Psi(t)\rangle = \sum_n c_n e^{-iE_n t} |\psi_n\rangle, \quad H |\psi_n\rangle = E_n |\psi_n\rangle,$$

so $\{|\psi_n\rangle\}$ forms an orthonormal basis of stationary states.

For an N -particle system in d spatial dimensions, with coordinates $R = (\mathbf{r}_1, \dots, \mathbf{r}_N)$, we use the many-body Hamiltonian

$$H = -\frac{1}{2} \sum_{i=1}^N \nabla_{\mathbf{r}_i}^2 + \sum_{i=1}^N V_{\text{ext}}(\mathbf{r}_i) + \sum_{i < j} V_{\text{int}}(|\mathbf{r}_i - \mathbf{r}_j|),$$

and the time-independent Schrödinger equation (TISE)

$$H \Psi(R) = E \Psi(R), \quad \int |\Psi(R)|^2 dR = 1.$$

Identical particles restrict Ψ to the appropriate symmetry sector (symmetric for bosons, antisymmetric for fermions).

Throughout we work in atomic units ($\hbar = m_e = e = 1$) and focus on stationary problems. In the quantum-dot applications below we take $V_{\text{ext}}(\mathbf{r}) = \frac{1}{2}\omega^2 r^2$ and $V_{\text{int}}(r) = 1/r$ (in 2D this is understood with the usual Laplacian in two dimensions). Our computations therefore amount to approximating low-lying eigenpairs (E, Ψ) of the many-body Hamiltonian above using compact basis expansions and variational ansätze.

1.3 Second Quantization

1.3.1 Motivation: Why first-quantized wavefunctions don't scale

In first quantization an N -fermion state is an antisymmetric function $\Psi(x_1, \dots, x_N)$ expanded in Slater determinants of M one-particle orbitals. The number of configurations grows as $\binom{M}{N}$ (and permanents for bosons grow even faster), making storage and operator application increasingly intractable. We therefore move to the occupation-number formalism, which encodes antisymmetry and particle indistinguishability algebraically.

1.3.2 Fock space and occupation representation

Let $\{|p\rangle\}$ be an orthonormal set of one-particle *spin-orbitals* (spatial orbital + spin). The fermionic Fock space is

$$\mathcal{F}_- = \bigoplus_{n=0}^{\infty} \wedge^n \mathcal{H}_1,$$

with vacuum $|0\rangle$ and occupation-number basis $|n_1 n_2 \dots\rangle$, $n_p \in \{0, 1\}$. Creation/annihilation operators $a_p^\dagger, a_p$ act by adding/removing a particle in $|p\rangle$ (subject to Pauli) and obey the canonical anticommutation relations (CAR)

$$\{a_p, a_q^\dagger\} = \delta_{pq}, \quad \{a_p, a_q\} = \{a_p^\dagger, a_q^\dagger\} = 0,$$

with number operator $\hat{n}_p = a_p^\dagger a_p$. A Slater determinant occupying the set $I = \{p_1 < \dots < p_N\}$ is $|\Phi_I\rangle = a_{p_1}^\dagger \dots a_{p_N}^\dagger |0\rangle$ and is represented by the bitstring $|n_1 n_2 \dots\rangle$ with $n_p = \mathbf{1}_{p \in I}$. Unitary changes of one-particle basis $|p'\rangle = \sum_p U_{p'p} |p\rangle$ lift to $a_{p'}^\dagger = \sum_p U_{p'p} a_p^\dagger$, so physics is basis independent.

1.3.3 Operators in second quantization

Any *one-body* operator $O = \sum_i O(i)$ has matrix elements $O_{pq} = \langle p | O | q \rangle$ and becomes

$$O = \sum_{pq} O_{pq} a_p^\dagger a_q.$$

A *two-body* interaction $V = \frac{1}{2} \sum_{i \neq j} v(i, j)$ has

$$V = \frac{1}{2} \sum_{pqrs} \langle pq|v|rs \rangle a_p^\dagger a_q^\dagger a_s a_r = \frac{1}{4} \sum_{pqrs} \langle pq||rs \rangle a_p^\dagger a_q^\dagger a_s a_r,$$

where $\langle pq|v|rs \rangle$ are two-electron integrals over spin-orbitals and $\langle pq||rs \rangle = \langle pq|v|rs \rangle - \langle pq|v|sr \rangle$ denotes the antisymmetrized integral. In this form, operator algebra and particle indistinguishability are handled by the CAR, independent of N .

1.4 Hartree–Fock (HF)

Hartree–Fock (HF) approximates the N -fermion ground state by a single Slater determinant built from orthonormal spin-orbitals $\{|p\rangle\}_{p=1}^M$. In second quantization the Hamiltonian is

$$H = \sum_{pq} h_{pq} a_p^\dagger a_q + \frac{1}{4} \sum_{pqrs} \langle pq||rs \rangle a_p^\dagger a_q^\dagger a_s a_r,$$

with one-body matrix elements $h_{pq} = \langle p|\hat{h}_0|q \rangle$ and antisymmetrized two-body integrals $\langle pq||rs \rangle = \langle pq|rs \rangle - \langle pq|sr \rangle$ over spin-orbitals.

1.4.1 HF energy functional and the Fock operator

Let $|\Phi\rangle = \prod_{i=1}^N a_i^\dagger |0\rangle$ be a Slater determinant (occupied spin-orbitals $i, j, \dots$). Its energy is

$$E[\Phi] = \sum_i h_{ii} + \frac{1}{2} \sum_{i,j} \langle ij||ij \rangle.$$

Stationarity of $E[\Phi]$ under unitary rotations in the occupied+virtual subspace gives the HF equations

$$F |\phi_p\rangle = \varepsilon_p |\phi_p\rangle, \quad F = \hat{h}_0 + \sum_{j \in \text{occ}} (J_j - K_j),$$

where J_j and K_j are the Coulomb and exchange operators generated by orbital j . In a fixed orthonormal one-particle basis $\{|\mu\rangle\}$ these become the matrix *Roothaan–Hall* equations

$$\mathbf{F} \mathbf{C} = \mathbf{C} \boldsymbol{\varepsilon}, \quad F_{\mu\nu} = h_{\mu\nu} + \sum_{\lambda\sigma} P_{\lambda\sigma} \left[(\mu\nu|\lambda\sigma) - \frac{1}{2} (\mu\lambda|\nu\sigma) \right],$$

with the closed-shell density matrix $P_{\lambda\sigma} = 2 \sum_{i \in \text{occ}} C_{\lambda i} C_{\sigma i}^*$ (sum over occupied *spatial* orbitals; the factor 2 is the double occupancy that makes the $\frac{1}{2}$ exchange coefficient correct). (For a non-orthogonal basis S , use $\mathbf{F} \mathbf{C} = \mathbf{S} \mathbf{C} \boldsymbol{\varepsilon}$.)

Self-consistency (SCF). HF is solved iteratively: initialise $P^{(0)}$, build $\mathbf{F}[P^{(n)}]$, diagonalize to get $\mathbf{C}^{(n+1)}$, update $P^{(n+1)}$, and iterate until the total energy and/or P change is below tolerance. Damping or DIIS mixing stabilizes convergence. The dominant costs are building two-electron contributions ($\mathcal{O}(M^4)$) and diagonalization ($\mathcal{O}(M^3)$).

Remarks for quantum dots. For parabolic confinement $V_{\text{ext}} = \frac{1}{2} \omega^2 r^2$ we typically choose 2D harmonic-oscillator orbitals as the one-particle basis (the Cartesian orbitals used in our implementation), which respect the shell structure and accelerate convergence. Closed shells admit restricted HF (RHF); spin-polarized cases may require UHF. Koopmans’ theorem (orbital energies as removal/addition estimates) can provide a rough diagnostic but is not used for final energetics here.

1.5 Full Configuration Interaction (FCI)

FCI gives the exact ground state *within the chosen one-particle basis* by expanding in *all* N -electron Slater determinants,

$$|\Psi\rangle = \sum_I C_I |\Phi_I\rangle, \quad H\mathbf{C} = E\mathbf{C}.$$

The determinant space has dimension $\binom{M}{N}$ for fixed spin-orbitals M and particles N (or smaller with symmetry constraints). FCI thus captures the full correlation energy $\Delta E = E_{\text{FCI}} - E_{\text{HF}}$ in that basis and serves as the gold-standard benchmark.

1.5.1 Slater–Condon rules and sparsity

Matrix elements $\langle\Phi_I|H|\Phi_J\rangle$ are nonzero only if Φ_I and Φ_J differ by at most two spin-orbitals:

- *Diagonal:* $H_{II} = \sum_{i \in I} h_{ii} + \frac{1}{2} \sum_{i,j \in I} (ij||ij)$.
- *Singles* $i \rightarrow a$: $H_{0,ia} = F_{ia}$; in canonical HF orbitals, $F_{ia} = 0$ (Brillouin’s theorem), so the HF determinant does not couple to singles.
- *Doubles* $ij \rightarrow ab$: $H_{0,ijab} = (ij||ab)$.

This locality yields a very sparse Hamiltonian in the determinant basis and enables iterative eigensolvers without forming $\mathbf{H}$ explicitly.

1.5.2 Diagonalization and symmetry reduction

In practice one targets the lowest eigenpairs with Davidson or Lanczos iterations. Symmetries (N , S_z , total spin S if spin-adapted CSFs are used, and L_z in an angular-momentum-adapted basis) block-diagonalize the problem and dramatically shrink the working space and memory. For few-electron quantum dots (e.g. $N = 2, 6$) this makes high-accuracy FCI feasible for moderate M ; for larger N the determinant growth becomes the bottleneck.

1.5.3 Role in this thesis

HF provides physically meaningful orbitals and an upper bound; FCI (in the same one-particle basis) provides benchmark energies for small systems. We use these as references to assess the accuracy of our variational Slater–Jastrow/PINN results for 2D quantum dots. For larger N (where FCI is prohibitive) we rely on variational energies plus uncertainty estimates.

1.6 Quantum Dots

1.6.1 Model and units

We consider N spin- $\frac{1}{2}$ fermions in $d=2$ with harmonic confinement and Coulomb repulsion,

$$H = \sum_{i=1}^N \left(-\frac{1}{2} \nabla_{\mathbf{r}_i}^2 + \frac{1}{2} \omega^2 r_i^2 \right) + \sum_{i < j} \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|}, \quad (\hbar = m_e = e = 1).$$

Spin does not appear in H (spin-independent), so the many-body eigenstates can be chosen as products of a spatial wavefunction and a spin function with well-defined S and S_z .

1.6.2 One-particle structure and shell filling

We use two-dimensional harmonic-oscillator spin-orbitals. In the implementation these are the *Cartesian* 2D-HO eigenstates $\{\phi_{n_x n_y}(\mathbf{r})\chi_\sigma\}$ with energies $E_{n_x n_y} = \omega(n_x + n_y + 1)$ and spin $\sigma \in \{\uparrow, \downarrow\}$. The principal index $k = n_x + n_y$ labels a shell of $(k+1)$ degenerate spatial states, each twofold spin-degenerate; for the closed shells studied here this Cartesian basis spans the same filled subspace as any complete basis of those shells, so the reference determinant is basis-independent. Filling from the bottom yields the *closed-shell* electron numbers

$$N_{\text{closed}} = (K+1)(K+2), \quad K = 0, 1, 2, \dots \quad (2, 6, 12, 20, 30, \dots).$$

All other N correspond to a *partially filled* top shell (open shell), with a degenerate noninteracting manifold.

1.6.3 Spin structure and Slater form

Write single-particle spin-orbitals as $\varepsilon_\alpha(x) = \phi_\alpha(\mathbf{r})\chi_\sigma(s)$, with $x = (\mathbf{r}, s)$. Because $\chi_\uparrow$ and $\chi_\downarrow$ are orthogonal and H is spin-independent, the Slater determinant simplifies:

Closed shells (RHF). For even N with doubly occupied spatial orbitals $\{\phi_1, \dots, \phi_{N/2}\}$, the determinant factorizes into two identical spatial determinants, one for each spin:

$$\Psi_{\text{closed}}(X) = \frac{1}{(N/2)!} \det[\phi_a(\mathbf{r}_i)]_{i \in \uparrow, a=1..N/2} \det[\phi_a(\mathbf{r}_j)]_{j \in \downarrow, a=1..N/2} \times \Xi_{S=0, S_z=0},$$

where Ξ is the singlet spin function. This is the *restricted* HF (RHF) reference and a natural starting point for correlated ansätze.

1.6.4 Kato's Cusp Conditions (2D Coulomb systems)

For Coulomb interactions the potential diverges as $1/r$ at coalescence. Finiteness of the local energy

$$E_L(R) = \frac{(H\Psi)(R)}{\Psi(R)}$$

requires that the kinetic singularity cancels the $1/r$ divergence. Kato's theorem, first showed in [16], gives the necessary short-distance behaviour of the (spherically averaged) wavefunction. Let r_{ij} be the interparticle distance, $\mu_{ij} = m_i m_j / (m_i + m_j)$ the reduced mass (in a.u.), Z_i the charge number (electron $Z = -1$, nucleus $Z = +Z_A$), and D the spatial dimension. Then

$$\left. \frac{\partial}{\partial r_{ij}} \ln \bar{\Psi} \right|_{r_{ij} \rightarrow 0} = \frac{2\mu_{ij} Z_i Z_j}{D-1} \quad (\text{s-wave / opposite-spin channel}).$$

For *identical fermions* the s-wave is excluded; the leading channel is p -wave ($\ell = 1$), and the wavefunction carries a linear node at coalescence. The general like-spin cusp is then

$$\left. \frac{\partial}{\partial r_{ij}} \ln \bar{\Psi} \right|_{r_{ij} \rightarrow 0} = \frac{2\mu_{ij} Z_i Z_j}{2\ell + D - 1} = \frac{2\mu_{ij} Z_i Z_j}{D + 1} \quad (\text{like-spin, } \ell = 1).$$

Specializing to 2D electrons (atomic units). With $m_e = 1$ and $\mu_{ee} = 1/2$:

$$\boxed{a_{ee}^{\uparrow\downarrow} = \left. \frac{\partial}{\partial r_{ij}} \ln \bar{\Psi} \right|_0 = 1, \quad a_{ee}^{\uparrow\uparrow} = \frac{1}{3} .}$$

The like-spin value follows from the p -wave (linear-node) analysis carried out in full in Appendix B.2, the canonical derivation used throughout this thesis; it gives the familiar $\frac{1}{4}$ in three dimensions and $\frac{1}{3}$ here in two. The 2D like-spin cusp is therefore *not* simply half the opposite-spin value—a coincidence that holds only in 3D—and the implemented value is $1/(D+1) = \frac{1}{3}$. For electron-nucleus coalescence ($\mu_{eA} \approx 1$, $Z_i Z_j = -Z_A$):

$$\boxed{a_{eA} = \left. \frac{\partial}{\partial r_{iA}} \ln \bar{\Psi} \right|_0 = -2Z_A .}$$

Why this matters. These conditions ensure the kinetic energy’s r^{-1} singularity cancels the Coulomb r^{-1} term so that E_L remains finite as particles coalesce. Enforcing the cusp explicitly improves conditioning.

1.7 Structural diagnostics and Wigner markers

This section introduces the structural diagnostics conceptually—what each marker means and why it detects Wigner order; the precise computational procedures used in this thesis (estimators, thresholds, and stability protocol) are collected in the Analysis chapter (Chapter 5).

Unless stated otherwise, curves are accumulated in *trap units* $\tilde{\mathbf{r}} = \sqrt{\omega} \mathbf{r}$ so that length scales align across ω , while headline distances are reported in *Bohr* (physical units $\mathbf{r}$). Production chains are thinned; we split samples into halves for stability checks (acceptance window, JSD between halves, reproducibility under restarts).

1.7.1 Pair distribution $g(r)$, $P(r)$, and width markers

We estimate the (spin-summed) pair distribution $g(r)$ via a shell-area corrected histogram in trap units,

$$g(r_\alpha) = \frac{A_{\text{eff}}}{N(N-1)} \frac{\langle \sum_{i < j} \mathbf{1}\{r_{ij} \in [r_\alpha^-, r_\alpha^+]\} \rangle}{2\pi r_\alpha \Delta r_\alpha}, \quad r_{ij} = \|\tilde{\mathbf{r}}_i - \tilde{\mathbf{r}}_j\|_2, \quad (1.1)$$

with $A_{\text{eff}} = \pi R_{0.95}^2$ and $R_{0.95}$ the 95% quantile of single-particle radii (trap units). We form the normalised radial probability in $d=2$, $P(r) \propto r g(r)$, and report the mode r_{mode} , mean $\bar{r}$, standard deviation σ_r , FWHM, and quantiles q_{10}, q_{50}, q_{90} (in Bohr). A compact width-to-scale marker is

$$\gamma = \sigma_r / r_{\text{mode}}, \quad (1.2)$$

which decreases as correlations stiffen (narrow distribution around a larger typical radius).¹

Sanity anchors at $\omega = 10^{-3}$. For $N=6$ and $N=12$, $P(r)$ exhibits sharp peaks with small γ ; these are the expected Wigner-side shapes. For $N=2$ we additionally report the near-origin mass $\int_0^{0.25 r_{\text{mode}}} \rho_1(r) dr$ and $\Pr(r_{12} \leq 0.25 \text{ mode}(r_{12}))$, both $\ll 1$ as required by the cusp and strong repulsion.

1.7.2 Automatic shell detection and topology labeling

For each frame we sort single-particle radii $\tilde{s}_1 \leq \dots \leq \tilde{s}_N$ (trap units), compute gaps $g_k = \tilde{s}_{k+1} - \tilde{s}_k$, and flag a frame as *two-shell* if

$$\max_k g_k \geq \tau \text{ median}(g_k), \quad \tau \in [2.0, 3.0] \text{ (robust)}. \quad (1.3)$$

The cut radius is the midpoint of the maximal gap; the inner multiplicity n_{in} defines the topology $(n_{\text{in}}, N - n_{\text{in}})$. We report: (i) the fraction of two-shell frames, (ii) the histogram over n_{in} among two-shell frames, and (iii) for selected topologies (e.g. (1, 5) or (3, 9)) their fraction among *all* frames.

II/IO/OO decomposition (what is being added). On two-shell frames we decompose pairs into inner-inner (II), inner-outer (IO), and outer-outer (OO), accumulate $g_{\text{II}}, g_{\text{IO}}, g_{\text{OO}}$, and reconstruct

$$\tilde{g}(r) = w_{\text{II}} g_{\text{II}} + w_{\text{IO}} g_{\text{IO}} + w_{\text{OO}} g_{\text{OO}},$$

with empirical weights from pair counts. We report $\cos \angle(g, \tilde{g})$ (cosine similarity); values ≈ 1 indicate that the observed $g(r)$ is explained by shell geometry rather than sampling artifacts.

Sanity anchors at $\omega = 10^{-3}$. For $N=6$, $w_{\text{OO}} : w_{\text{IO}} \approx 2 : 1$ and $\cos \angle(g, \tilde{g}) = 1.000$ within numerical precision. For $N=12$, the two-shell fraction is 0.906–0.990 across $\tau \in \{3.0, 2.5, 2.0\}$ and $\cos \angle(g, \tilde{g}) \in [0.989, 1.000]$ depending mildly on τ /seed. The *modal* inner multiplicity at $\omega = 10^{-3}$ is (3, 9), while (2, 10) and (1, 11) retain substantial weight—consistent with a rotating two-shell Wigner molecule. For classical references to shellings and near-degeneracies in parabolic/cluster confinements, see Schweigert–Peeters and Kong–Partoens–Peeters.[8, 9]

¹For $N=2$, γ is complemented by the pair-distance Lindemann $\gamma_{r_{12}} = \text{std}(r_{12})/\text{mode}(r_{12})$, cf. Sec. 5.1.4.

1.7.3 Orientational order and angular “Lindemann”

After recentring each detected ring (inner/outer) to remove global rotations, with angles $\{\phi_k\}_{k=1}^n$, we use the bond-orientational order

$$\Phi_m = \frac{1}{n} \sum_{k=1}^n e^{im\phi_k}, \quad m = n, \quad (1.4)$$

so that $|\Phi_m| \in [0, 1]$ measures n -fold orientational order (e.g. $m=5$ for (1,5); $m=9$ for the outer ring of (3,9)). As a complementary, dimensionless *angular Lindemann* number we use

$$\lambda_\phi = \frac{\text{std}(\Delta\phi)}{\mathbb{E}[\Delta\phi]}, \quad \Delta\phi = \text{nearest-neighbour angular spacings on the ring.} \quad (1.5)$$

Smaller λ_ϕ indicates a stiffer (more Wigner-like) ring. This follows the spirit of bond-orientational order used in 2D Coulomb/Yukawa systems,[11, 12] with the Lindemann idea adapted from melting criteria to angular spacings on a fixed- n ring.

Sanity anchors at $\omega=10^{-3}$. For $N=6$, phase-aligned (1,5) and (0,6) sectors show comparable order ($|\Phi_5| \approx 0.48$, $|\Phi_6| \approx 0.52$) with $\lambda_\phi \approx 0.29$ —a single-ring rotating Wigner molecule. For $N=12$, (3,9) frames have strong order ($|\Phi_3| \approx 0.67$, $|\Phi_9| \approx 0.50$) and small λ_ϕ (~ 0.24 – 0.26). (2,10) exhibits very tight inner order ($|\Phi_2| \approx 0.73$) with a softer outer ring ($|\Phi_{10}| \approx 0.42$), and (1,11) has weak outer order and a softly vibrating centre—precisely the *rotating Wigner molecule* (RWM) picture of fluctuating orientations across near-degenerate shellings.[7, 10]

1.7.4 Scaling check and two-body benchmarks ($N = 2$)

To connect with the classical strong-coupling limit we fit

$$r_{\text{mode}}(\omega) = C \omega^\alpha \quad (1.6)$$

by linear regression in $(\log \omega, \log r_{\text{mode}})$ and compare with the two-body classical scaling $\alpha_{\text{cl}} = -2/3$. At $\omega=10^{-3}$ the r_{12} histogram is sharply peaked and the near-origin mass in ρ_1 and $p(r_{12})$ is $\ll 1$, validating the cusp implementation while confirming deep Wigner localization (see also Ref. [7] for the crossover criterion in dots).

1.7.5 Reporting and stability checks

For all scalars and histograms we report standard errors from block averages over thinned frames. We verify: (i) acceptance in the target window, (ii) Jensen–Shannon divergence and ℓ_2 distance between split halves (stationarity), (iii) robustness of topology fractions and $|\Phi_m|$ under $\tau \in [2.0, 3.0]$ and mild pre-smoothing of radii, and (iv) reproducibility of $g(r)$, $P(r)$, and bond-order metrics under independent sampler restarts. All diagnostics operate directly on saved bundles $\{\mathbf{X}_{\text{Bohr}}, r, g(r)\}$.

Interpretation guide (link to Results). Across $\omega \in \{1.0, 0.5, 0.1, 0.01, 10^{-3}\}$ these diagnostics map the fluid→Wigner crossover: γ and λ_ϕ shrink, shell gaps sharpen, and shell-resolved $|\Phi_m|$ becomes nonzero after phase alignment. For $N=6$ this manifests as a single-ring RWM with (1,5)–(0,6) near-degeneracy; for $N=12$ as a two-shell RWM with modal (3,9) but sizable (2,10) and (1,11) weights. Classical shelling and near-degeneracies in finite Coulomb clusters[8, 9] and quantum Wigner markers in parabolic dots[7, 10–12] provide the theoretical backdrop for reading these plots.

1.8 Variational Methods

Quantum Monte Carlo (QMC) methods offer powerful tools for solving the many-body Schrödinger equation through stochastic sampling. These methods yield accurate ground-state properties by estimating expectation values of observables with controlled approximations. Among the various QMC techniques, Variational Monte Carlo (VMC) and Diffusion Monte Carlo (DMC) are particularly notable; VMC optimizes a trial wavefunction based on the variational principle, while DMC projects out the ground state through imaginary-time evolution [17, 18].

1.8.1 Variational Monte Carlo (VMC)

The Variational Principle

The variational principle guarantees that for any normalised trial wavefunction $\Psi_T(\mathbf{R}; \alpha)$, the expectation value

$$E[\Psi_T] = \frac{\langle \Psi_T | \hat{H} | \Psi_T \rangle}{\langle \Psi_T | \Psi_T \rangle}$$

provides an upper bound to the true ground-state energy E_0 . Here, $\mathbf{R} = (\mathbf{r}_1, \dots, \mathbf{r}_N)$ and α represents the set of variational parameters. Optimising α minimizes $E[\Psi_T]$.

Sampling

In VMC, the probability density is defined as

$$P(\mathbf{R}) = \frac{|\Psi_T(\mathbf{R})|^2}{\int |\Psi_T(\mathbf{R})|^2 d\mathbf{R}},$$

and configurations are sampled using the Metropolis-Hastings algorithm, initially described in [19], with acceptance probability

$$A(\mathbf{R} \rightarrow \mathbf{R}') = \min \left(1, \frac{|\Psi_T(\mathbf{R}')|^2}{|\Psi_T(\mathbf{R})|^2} \frac{T(\mathbf{R}' \rightarrow \mathbf{R})}{T(\mathbf{R} \rightarrow \mathbf{R}')} \right),$$

where T is the proposal density (for a symmetric proposal the ratio is unity and the acceptance reduces to the Metropolis form). When a drift is added, a *quantum force*

$$\mathbf{F}(\mathbf{R}) = 2 \nabla \ln |\Psi_T(\mathbf{R})|,$$

further guides the sampling toward regions of high probability. This leads to what is known as importance sampling. A more precise discussion of this is found in [20, 21].

Local Energy and Parameter Optimisation

The local energy is defined as

$$E_L(\mathbf{R}) = \frac{\hat{H}\Psi_T(\mathbf{R})}{\Psi_T(\mathbf{R})}.$$

If Ψ_T were exact, $E_L(\mathbf{R})$ would be constant. In practice, the variational energy is approximated by averaging E_L over many configurations:

$$E[\Psi_T] \approx \frac{1}{N_{\text{MC}}} \sum_{i=1}^{N_{\text{MC}}} E_L(\mathbf{R}_i).$$

Variational parameters are then optimised (using gradient descent or stochastic reconfiguration) to reduce both the energy and its variance.

1.8.2 Diffusion Monte Carlo (DMC)

Imaginary-Time Evolution and Projection

DMC refines the trial wavefunction by projecting out the ground state through imaginary-time evolution. The Schrödinger equation becomes

$$-\frac{\partial \Psi(\mathbf{R}, \tau)}{\partial \tau} = (\hat{H} - E_T) \Psi(\mathbf{R}, \tau),$$

where $\tau = it$ and E_T is a trial energy. As $\tau \rightarrow \infty$, $\Psi(\mathbf{R}, \tau)$ converges to the ground-state wavefunction provided the initial state has nonzero overlap with it.

Stochastic Implementation

DMC simulates the imaginary-time evolution using a combination of diffusion, drift, and branching:

1. **Diffusion and Drift:** Particles execute random walks, with drift guided by the quantum force $\mathbf{F}(\mathbf{R})$, steering them toward regions where Ψ_T is large.
2. **Branching/Death:** Walkers are assigned a weight,

$$w(\mathbf{R}, \Delta\tau) = \exp\left[-\left(E_L(\mathbf{R}) - E_T\right)\Delta\tau\right],$$

which determines whether they are replicated or removed, ensuring that the walker distribution evolves toward the ground-state probability density.

1.8.3 Applications and Numerical Considerations

Model Systems

QMC methods have been successfully applied to various quantum systems:

- **Hydrogen Atom:** A simple trial wavefunction $\Psi_T(r) = e^{-\alpha r}$ yields the exact ground state for $\alpha = 1$, illustrating zero variance.
- **Harmonic Traps:** In systems confined by harmonic potentials, Jastrow factors like $\exp[-ar_{ij}/(1 + \beta r_{ij})]$ model inter-particle repulsion while Slater determinants maintain the proper fermionic symmetry.

Computational Efficiency

Key factors in achieving efficient QMC simulations include:

- **Efficient Determinant Updates:** For a single-particle move, the Sherman–Morrison formula updates the inverse Slater matrix in $\mathcal{O}(N^2)$ rather than recomputing the determinant from scratch at $\mathcal{O}(N^3)$.
- **Optimised Gradient and Laplacian Calculations:** Recursive methods and efficient numerical routines for evaluating $\nabla\Psi_T/\Psi_T$ and $\nabla^2\Psi_T/\Psi_T$ are essential.

1.8.4 Conclusion

Variational and Diffusion Monte Carlo methods together provide a robust framework for studying quantum systems. VMC offers a flexible approach for optimising trial wavefunctions, while DMC projects toward the ground state through imaginary-time evolution—though for fermions the practical fixed-node variant is variational only within the imposed nodal surface, and retains a nodal bias unless those nodes are exact. These methods, working in concert, enable accurate and efficient investigations of systems from simple atoms to strongly correlated materials. Ongoing improvements in trial wavefunction design and stochastic algorithms continue to extend the reach of QMC in modern computational physics.

Chapter 2

Machine Learning

2.1 Machine Learning as Statistical Learning

Machine learning (ML) is best understood as the study of algorithms that approximate functions from data. At its core, it is not about “intelligent machines,” but about developing systematic ways to extract patterns from observations and to generalise these patterns to unseen situations. From a mathematical perspective, ML is a branch of *statistical learning theory*, which views the learning process as the minimization of an expected error (or *risk*) under uncertainty.

Formally, consider an input space $\mathcal{X} \subseteq \mathbb{R}^d$ and an output space $\mathcal{Y}$, together with an unknown joint distribution $P(X, Y)$ over pairs (x, y) . The aim of learning is to construct a function $f_\theta : \mathcal{X} \rightarrow \mathcal{Y}$, parameterised by $\theta \in \Theta$, such that $f_\theta(x)$ predicts y with small error. A loss function

$$\ell : \mathcal{Y} \times \mathcal{Y} \rightarrow \mathbb{R}_{\geq 0}$$

quantifies the discrepancy between prediction $f_\theta(x)$ and truth y . The central object of learning is then the **population risk**

$$\mathcal{R}(\theta) = \mathbb{E}_{(x,y) \sim P} [\ell(f_\theta(x), y)]. \quad (2.1)$$

Since the distribution P is unknown, we only have access to a finite dataset $D = \{(x_i, y_i)\}_{i=1}^n$. The corresponding **empirical risk** is

$$\hat{\mathcal{R}}(\theta) = \frac{1}{n} \sum_{i=1}^n \ell(f_\theta(x_i), y_i). \quad (2.2)$$

This setup is known as *empirical risk minimization (ERM)*, and it underlies virtually all modern machine learning methods [22].

2.1.1 Types of learning

Depending on the information provided in the dataset, learning problems are traditionally categorized as:

- **Supervised learning.** We are given labeled pairs (x_i, y_i) , and the goal is to approximate the conditional expectation $\mathbb{E}[Y|X = x]$. Regression (continuous Y) and classification (discrete Y) are canonical examples.
- **Unsupervised learning.** Only input data $\{x_i\}$ is observed, without labels. The aim is to uncover structure, such as clustering, dimensionality reduction, or latent-variable representations.
- **Reinforcement learning.** An agent interacts with an environment, receiving observations and rewards. The task is to learn a policy that maximizes expected cumulative reward.

Both of the first two categories appear in this thesis, and it is worth saying where. The training objectives that carry the bulk of the work—the Schrödinger residual and the variational energy—are *unsupervised*: there are no labelled targets anywhere, and the loss is supplied entirely by the physics the network is asked to satisfy. The one genuinely *supervised* ingredient is the second phase of residual pretraining, in

which a published reference energy is used as an explicit target; the first phase, where the network sets its own target as a stabiliser, is better described as self-supervised. Reinforcement learning enters only obliquely—the REINFORCE estimator used for the collocation gradient is borrowed from it—and is not otherwise used.

2.1.2 Generalisation and the bias–variance tradeoff

The essential challenge in ML is not just minimizing the empirical risk $\hat{\mathcal{R}}(\theta)$, but ensuring that this leads to a low population risk $\mathcal{R}(\theta)$ on unseen data. A model that simply memorizes the training set achieves low empirical risk but poor generalisation — a phenomenon known as *overfitting*.

Statistical learning theory formalizes this challenge via the *bias–variance tradeoff*. Given a data-generating process and a learning algorithm, the expected squared error can be decomposed into

$$\mathbb{E}[(f_\theta(x) - y)^2] = \underbrace{(\mathbb{E}[f_\theta(x)] - f^*(x))^2}_{\text{bias}^2} + \underbrace{\mathbb{E}[(f_\theta(x) - \mathbb{E}[f_\theta(x)])^2]}_{\text{variance}} + \underbrace{\sigma^2}_{\text{irreducible noise}}, \quad (2.3)$$

where $f^*(x)$ is the true regression function and σ^2 the intrinsic noise of the data. Complex models tend to have low bias but high variance, while simple models have the opposite. Successful learning algorithms balance these two contributions. This is discussed in more detail in [23].

2.2 Optimisation

The minimization of empirical risk,

$$\hat{\mathcal{R}}(\theta) = \frac{1}{n} \sum_{i=1}^n \ell(f_\theta(x_i), y_i),$$

is an *optimisation problem*. The hypothesis class $\{f_\theta\}$ is usually nonlinear and high-dimensional, so closed-form solutions are not available. Instead, optimisation is performed iteratively by updating parameters $\theta \in \mathbb{R}^p$ according to the geometry of the loss landscape.

2.2.1 Gradient descent

The most basic algorithm is *gradient descent*. Starting from an initial guess $\theta^{(0)}$, parameters are updated by

$$\theta^{(t+1)} = \theta^{(t)} - \eta \nabla_\theta \hat{\mathcal{R}}(\theta^{(t)}), \quad (2.4)$$

where $\eta > 0$ is the *learning rate*. Intuitively, the gradient points in the direction of steepest increase of the loss, so subtracting it decreases the loss most rapidly in a local sense.

The choice of learning rate is crucial: if too small, convergence is slow; if too large, the algorithm may diverge.

2.2.2 Stochastic gradient descent (SGD)

For large datasets, evaluating the full gradient at every step is prohibitively expensive. Instead, one uses *stochastic gradient descent (SGD)*, which approximates the gradient on a randomly sampled mini-batch $B \subset D$:

$$\nabla_\theta \hat{\mathcal{R}}_B(\theta) = \frac{1}{|B|} \sum_{(x_i, y_i) \in B} \nabla_\theta \ell(f_\theta(x_i), y_i). \quad (2.5)$$

The update rule is then

$$\theta^{(t+1)} = \theta^{(t)} - \eta \nabla_\theta \hat{\mathcal{R}}_B(\theta^{(t)}). \quad (2.6)$$

SGD introduces randomness into the optimisation trajectory. Paradoxically, this noise often *helps*: it prevents the algorithm from getting trapped in poor local minima and can improve generalisation through an implicit regularization effect [22, 24].

2.2.3 Momentum and adaptive methods

Plain gradient descent treats all directions in parameter space equally. To accelerate convergence, one may use *momentum*, maintaining a velocity vector v_t :

$$v_{t+1} = \beta v_t + (1 - \beta) \nabla_{\theta} \hat{\mathcal{R}}_B(\theta^{(t)}), \quad (2.7)$$

$$\theta^{(t+1)} = \theta^{(t)} - \eta v_{t+1}, \quad (2.8)$$

where $\beta \in (0, 1)$ controls the contribution of past gradients. Momentum smooths oscillations and allows the optimisation to traverse valleys in the loss landscape more effectively.

Beyond momentum, *adaptive methods* such as RMSProp and Adam rescale learning rates per parameter based on historical gradient magnitudes [25]. For instance, Adam combines momentum with an adaptive normalisation:

$$m_t = \beta_1 m_{t-1} + (1 - \beta_1) g_t, \quad (2.9)$$

$$v_t = \beta_2 v_{t-1} + (1 - \beta_2) g_t^2, \quad (2.10)$$

$$\theta^{(t+1)} = \theta^{(t)} - \eta \frac{\hat{m}_t}{\sqrt{\hat{v}_t + \epsilon}}, \quad (2.11)$$

where $g_t = \nabla_{\theta} \hat{\mathcal{R}}_B(\theta^{(t)})$ and $\hat{m}_t, \hat{v}_t$ are bias-corrected versions of m_t, v_t . Adam is widely used in practice for its robustness across diverse problems [26].

2.2.4 Second-order methods and natural gradient

First-order methods only use gradient information. In principle, convergence can be accelerated by incorporating curvature via the Hessian $H(\theta) = \nabla_{\theta}^2 \hat{\mathcal{R}}(\theta)$. Newton's method updates parameters as

$$\theta^{(t+1)} = \theta^{(t)} - H(\theta^{(t)})^{-1} \nabla_{\theta} \hat{\mathcal{R}}(\theta^{(t)}). \quad (2.12)$$

However, computing and inverting the Hessian is usually infeasible for large models.

An attractive alternative is the *natural gradient* [27, 28], which replaces the Euclidean metric with the geometry of probability distributions. The key object is the *Fisher information matrix* (FIM),

$$F(\theta) = \mathbb{E}[\nabla_{\theta} \log p_{\theta}(x) \nabla_{\theta} \log p_{\theta}(x)^{\top}], \quad (2.13)$$

where p_{θ} denotes the model distribution. The natural gradient update is

$$\theta^{(t+1)} = \theta^{(t)} - \eta F(\theta^{(t)})^{-1} \nabla_{\theta} \hat{\mathcal{R}}(\theta^{(t)}). \quad (2.14)$$

This preconditioning makes updates invariant to reparameterizations of θ and often improves conditioning of the optimisation. In practice, computing F^{-1} exactly is intractable, but approximations (e.g. stochastic reconfiguration in variational Monte Carlo, Kronecker-factored approximations in deep learning) make the method viable [2].

2.2.5 The variational tangent space: geometric tensor, tangent kernel, and effective dimension

For a variational quantum state the Fisher metric takes a particularly concrete form, and that form is the analytical lens used throughout the results of this thesis. Write the per-sample log-derivatives—the *score*—of an ansatz Ψ_{θ} as

$$O_{ki} = \frac{\partial \log |\Psi_{\theta}(\mathbf{x}_k)|}{\partial \theta_i}, \quad k = 1, \dots, B, \quad i = 1, \dots, P, \quad (2.15)$$

for a batch of B configurations drawn from $|\Psi_{\theta}|^2$ and P parameters θ (here θ collects *all* variational parameters). Its centred Gram matrices define two dual objects. The *quantum geometric tensor* (QGT),

$$S = \frac{1}{B} O^{\top} O \in \mathbb{R}^{P \times P}, \quad (2.16)$$

is the Fisher–Rao metric of the state written in terms of the amplitude score. It is proportional to the Fisher matrix $F(\theta)$ of the previous subsection specialised to $p_{\theta} = |\Psi_{\theta}|^2$: because $\nabla_{\theta} \log |\Psi_{\theta}|^2 =$

$2\nabla_\theta \log |\Psi_\theta|$, one has $F = 4S$, the factor of four being a convention absorbed into the learning rate. Stochastic reconfiguration is natural-gradient descent under this metric, $\dot{\theta} = -S^{-1}\nabla_\theta E$ [29, 30]. Its Gram dual, the *neural tangent kernel* (NTK) [31],

$$K = OO^\top \in \mathbb{R}^{B \times B}, \quad (2.17)$$

shares the nonzero spectrum of S up to the overall factor B (its nonzero eigenvalues are $B\lambda_a$), and is cheaper to diagonalise when $B < P$.

Two scalars read from that spectrum organise the analysis. The *condition number* $\kappa(S) = \lambda_{\max}/\lambda_{\min}$ measures how anisotropic the tangent metric is, and therefore how far a natural-gradient step departs from a Euclidean one—the quantity that decides when SR is worth its cost. The *effective dimension*

$$d_{\text{eff}}(S) = \frac{(\sum_a \lambda_a)^2}{\sum_a \lambda_a^2} \quad (2.18)$$

is the participation ratio of the spectrum: it counts how many independent tangent directions carry appreciable Fisher weight at the solution, regardless of the raw parameter count P . A small d_{eff} means the state is reached along a few collective directions; a large one means the same solution is spread across many. Because these are properties of the tangent geometry rather than of the raw parameter count, and are stable across seeds and sampling measures for a given ansatz, they let us compare architectures, optimisers, and training paradigms on a common footing—which is exactly what the second results chapter (Chapter 7) does.

One caveat must be kept in view. The participation ratio d_{eff} is computed from the QGT eigenvalues in the ordinary Euclidean parameter coordinates, and those eigenvalues are *not* invariant under an arbitrary reparameterization: rescaling a single parameter $\theta_1 \mapsto c\theta_1$ leaves the physical state untouched but changes d_{eff} (for a diagonal metric it moves from 2 toward 1 as $c \rightarrow \infty$). d_{eff} is therefore *not* a coordinate-free intrinsic dimension of the wavefunction; it is a property of the (architecture, parameterisation, normalisation) triple. We use it accordingly—as a comparison between fixed architectures under matched conventions and a common probe measure—and read a smaller d_{eff} as “this architecture concentrates its Fisher weight into fewer active directions,” not as an absolute manifold dimension.

2.2.6 Optimisation challenges in deep learning

Optimisation in deep neural networks poses unique challenges:

- **Non-convexity.** Loss landscapes are highly non-convex, with many local minima and saddle points. Empirically, most local minima reached by SGD perform similarly well.
- **Vanishing and exploding gradients.** In deep networks, gradients can shrink or blow up during backpropagation. Proper initialisation and activation choice are essential to mitigate this problem.
- **Flat vs. sharp minima.** Empirical evidence suggests that flat minima (regions where the loss remains low under perturbations of θ) lead to better generalisation. SGD’s noise implicitly biases optimisation toward flatter minima.

Optimisation is therefore more than a means of fitting: the choice of update also shapes the inductive biases and generalisation behaviour of the trained network. The sections that follow examine the other ingredients that determine those biases—network expressivity, activation functions, and initialisation.

2.3 Feed-forward neural networks

Feed-forward neural networks are the basic building block from which the architectures of this thesis are assembled. They compose affine maps with elementwise nonlinearities to represent functions that linear models cannot. We first define the feed-forward map, and then discuss the two design choices that most affect its behaviour in physics-informed settings, where gradients and Laplacians of the network enter the loss: the activation function and the weight initialisation.

2.3.1 From linear models to nonlinear function classes

Classical regression models are linear in their parameters:

$$f_{\theta}(x) = \theta^{\top} x.$$

While simple and interpretable, linear models are limited to representing affine functions. Introducing basis expansions (e.g. polynomials, Fourier series, kernels) enlarges the hypothesis space, but requires manual design.

Neural networks automate this process. A feed-forward neural network with L layers is a composition of affine maps and nonlinear activation functions:

$$h^{(0)} = x, \tag{2.19}$$

$$h^{(\ell)} = \sigma\left(W^{(\ell)}h^{(\ell-1)} + b^{(\ell)}\right), \quad \ell = 1, \dots, L-1, \tag{2.20}$$

$$f_{\theta}(x) = W^{(L)}h^{(L-1)} + b^{(L)}. \tag{2.21}$$

Here $\sigma : \mathbb{R} \rightarrow \mathbb{R}$ is a nonlinear activation. Without nonlinearity, the network collapses to a single linear transformation; with it, the space of representable functions expands dramatically.

2.3.2 Activation functions

Classical Activation Functions

The earliest activation functions were smooth, saturating nonlinearities:

$$\sigma_{\text{sigmoid}}(x) = \frac{1}{1 + e^{-x}}, \tag{2.22}$$

$$\sigma_{\text{tanh}}(x) = \tanh(x). \tag{2.23}$$

Both map $\mathbb{R}$ to a bounded range and are differentiable. The hyperbolic tangent is zero-centred, which often improves optimisation.

Despite their smoothness, these functions suffer from the *vanishing gradient problem*. For large $|x|$, their derivatives approach zero, causing gradients to vanish during backpropagation in deep networks. As a result, learning becomes prohibitively slow.

To address vanishing gradients, the *Rectified Linear Unit* (ReLU) was introduced:

$$\sigma_{\text{ReLU}}(x) = \max(0, x). \tag{2.24}$$

ReLU is unbounded above and has derivative 1 for positive inputs, avoiding gradient saturation. This makes deep networks with ReLU trainable even with many layers.

However, ReLU introduces its own drawbacks:

- Non-smoothness at $x = 0$, which can lead to optimisation instability in contexts where derivatives or Laplacians are central (e.g. physics-informed models).
- The *dying ReLU* problem, where neurons with negative inputs become permanently inactive.

Modern activations

To combine the benefits of ReLU with smoothness, a number of *smoothed rectifiers* have been proposed:

$$\sigma_{\text{SiLU}}(x) = x \cdot \sigma_{\text{sigmoid}}(x), \tag{2.25} \quad (\text{Sigmoid Linear Unit})$$

$$\sigma_{\text{Swish}}(x) = x \cdot \frac{1}{1 + e^{-\beta x}}, \tag{2.26} \quad (\text{generalised SiLU with } \beta > 0)$$

$$\sigma_{\text{GELU}}(x) = x \cdot \Phi(x), \tag{2.27} \quad (\text{Gaussian Error Linear Unit})$$

$$\sigma_{\text{Mish}}(x) = x \cdot \tanh(\log(1 + e^x)). \tag{2.28}$$

Here $\Phi(x)$ is the standard Gaussian cumulative distribution function.

These activations are smooth (C^{∞}) and non-saturating, preserving gradient flow while avoiding non-differentiable kinks. Empirically, GELU, SiLU/Swish, and Mish improve optimisation stability and generalisation across a wide range of tasks.

2.3.3 Initialisation

The initialisation of neural network weights has a critical impact on optimisation. At the start of training, parameter values determine how signals propagate forward and backward through the network. Poor initialisation can lead to exploding or vanishing activations and gradients, making optimisation unstable or ineffective. Proper schemes balance signal magnitudes across layers and preserve gradient flow, especially in deep networks.

The role of initialisation

Consider a feed-forward network with layers

$$h^{(\ell)} = \sigma\left(W^{(\ell)}h^{(\ell-1)} + b^{(\ell)}\right),$$

where $W^{(\ell)}$ has entries drawn from some distribution. If weights are too large, activations grow exponentially with depth, leading to exploding gradients. If too small, activations shrink toward zero, causing vanishing gradients. Both phenomena impede learning.

Initialisation therefore aims to preserve the variance of activations and gradients across layers, such that

$$\text{Var}[h^{(\ell)}] \approx \text{Var}[h^{(\ell-1)}].$$

Xavier/Glorot initialisation

Glorot and Bengio (2010) proposed initialising weights with variance scaled by the number of input and output units:

$$W_{ij}^{(\ell)} \sim \mathcal{U}\left(-\sqrt{\frac{6}{n_{\text{in}} + n_{\text{out}}}}, \sqrt{\frac{6}{n_{\text{in}} + n_{\text{out}}}}\right). \quad (2.29)$$

Here n_{in} and n_{out} denote the fan-in and fan-out of a layer. This choice balances variance across layers for activations with approximately unit derivative (e.g. $\tanh$).

Kaiming/He initialisation

For rectified activations such as ReLU, He et al. (2015) proposed *Kaiming initialisation*, scaling variance by $2/n_{\text{in}}$:

$$W_{ij}^{(\ell)} \sim \mathcal{N}\left(0, \frac{2}{n_{\text{in}}}\right). \quad (2.30)$$

This accounts for the fact that ReLU activations output zero half of the time, requiring larger variance to maintain signal propagation.

Gain and activation dependence

Both Glorot and Kaiming schemes can be generalised with a *gain* factor g that depends on the activation function:

$$\text{Var}[W_{ij}^{(\ell)}] = \frac{g^2}{n_{\text{in}}}.$$

For example, $g = \sqrt{2}$ is recommended for ReLU, while $g = 1$ is typical for $\tanh$ and smooth activations. Choosing the correct gain ensures that variance neither explodes nor decays through the network.

Orthogonal and random normal initialisation

An alternative approach is to initialise each weight matrix as an orthogonal matrix scaled by a variance factor. Orthogonal initialisation preserves the norm of activations across layers, which can stabilize very deep networks.

Simpler schemes such as sampling from a standard normal distribution without scaling are generally inferior: they do not respect the scaling needs of different architectures or activation functions, and often result in unstable training.

2.3.4 Summary

Initialisation is particularly delicate in deep networks. Even small imbalances in variance can compound exponentially with depth. Smooth activations (SiLU, GELU, Mish) alleviate gradient instability but require careful scaling, as their effective slope differs from ReLU or tanh.

Key takeaways include:

- Proper scaling of variance prevents vanishing/exploding signals.
- Initialisation must match the activation function (Glorot for tanh/smooth, Kaiming for ReLU).
- Orthogonal schemes can improve stability in deep networks.
- For PINNs and scientific ML, initialisation is not merely a convenience but essential for stable computation of derivatives.

Initialisation and activation choice thus form an inseparable pair: together they determine whether information propagates stably through the network. This is taken up again for physics-informed models in Section 2.6.

2.4 Architectures for many-body wavefunctions

The feed-forward network of the previous section maps a fixed-size input vector to an output. A many-body wavefunction is not a generic function of such a vector: identical particles impose a permutation symmetry, and correlation is intrinsically *relational*—what one particle should do depends on where the others are. A plain multilayer perceptron acting on the concatenated coordinates $(\mathbf{x}_1, \dots, \mathbf{x}_N)$ respects neither property; it would have to learn the symmetry from data, and its parameter count would grow with N . This section introduces the two architecture classes that build the required structure in—permutation-invariant pooling and message passing—and situates the specific message-passing network used in this thesis as one instance of the latter.

2.4.1 Permutation symmetry and equivariance

For identical particles the modulus $|\Psi|$ is invariant under exchange (the sign is carried by the antisymmetric Slater reference of Chapter 3). The two learnable objects of the ansatz therefore have distinct symmetry requirements. A scalar *correlator* f must be permutation-*invariant*,

$$f(P_\sigma R) = f(R),$$

while a per-particle *backflow* displacement field $\mathbf{g} = (\mathbf{g}_1, \dots, \mathbf{g}_N)$ must be permutation-*equivariant*,

$$\mathbf{g}(P_\sigma R) = P_\sigma \mathbf{g}(R),$$

for any permutation σ (here P_σ relabels the particles). Enforcing these by construction is both more data-efficient than learning them and yields a parameter count independent of N , because the same shared maps are applied to every particle and pair.

2.4.2 Permutation-invariant pooling: DeepSets

The simplest way to build an invariant function is to embed each particle independently and pool the embeddings symmetrically. This is the DeepSets construction [32]: any suitably regular permutation-invariant function admits the form

$$f(R) = \rho\left(\sum_i \phi(\mathbf{x}_i)\right),$$

with a per-particle embedding ϕ , a symmetric pool $\sum_i$, and a readout ρ . A two-body extension encodes pairs, $\phi(\mathbf{x}_i, \mathbf{x}_j)$, and pools over them, capturing pairwise correlation. What characterises this class is that each particle—or pair—is embedded *independently* and only then pooled: no information is exchanged between particles before the pool. We refer to such networks as *separable*, and they serve as the non-message-passing baseline for the correlator.

2.4.3 Message passing and graph networks

To let a particle’s representation depend on the others’ in an iterated, higher-order way, one places the particles on a graph—nodes are particles, edges are pairs—and passes learned messages along the edges [33, 34]. In a generic round ℓ , each edge gathers information from its endpoint nodes and each node aggregates its incoming edge messages,

$$\mathbf{m}_{ij}^{(\ell)} = \phi_e(\mathbf{h}_i^{(\ell)}, \mathbf{h}_j^{(\ell)}, \mathbf{e}_{ij}^{(\ell)}), \quad \mathbf{h}_i^{(\ell+1)} = \phi_v(\mathbf{h}_i^{(\ell)}, \text{Agg}_{j \neq i} \mathbf{m}_{ij}^{(\ell)}),$$

with shared maps ϕ_e, ϕ_v and a symmetric aggregation Agg , so that permutation equivariance holds by construction; a final readout produces an invariant scalar (correlator) or an equivariant field (backflow). Both DeepSets and message passing use pairwise information, but only message passing lets that information *propagate*: after several rounds a particle’s representation reflects not just its neighbours but their neighbours’ responses. This iterated propagation is what we call the *relational channel*, and asking what it buys—in the correlator and in the backflow—is one of the central questions of this thesis. Message passing has been used in exactly this spirit for neural quantum states of extended fermion systems [5, 6].

2.4.4 The CTNN: one message-passing instance

The message-passing correlator and backflow studied here are realised by a particular network we refer to as the *CTNN*: a copresheaf-style message-passing network that maintains explicit *node and edge* features and transports information between them through learned maps, rather than recomputing and discarding one-shot messages. The correlator iterates this exchange over a down/up “V-cycle”; the backflow applies a single such copresheaf round. Its concrete update equations, hidden dimensions, and parameter counts are specified in the Methods chapter (§3.5.2, Eqs. (3.13)–(3.15)). Two points of framing matter for the rest of the thesis. First, the CTNN is *one* concrete realisation of the message-passing class defined above; the conclusions we draw about the value of message passing concern the class, with the CTNN as the instance actually studied. Wherever we contrast a “message-passing” network with a “separable” one, the separable member is a DeepSets-type network and the message-passing member is the CTNN. Second, we write “copresheaf-style” deliberately: the name reflects the node/edge transport structure of the updates, not a claim to a formal sheaf-theoretic construction.

2.4.5 Relevance to the ansatz

Both learnable objects of the ansatz—the Jastrow correlator and the coordinate backflow—can be built in either form: as a separable DeepSets-type network, or as the message-passing CTNN. Holding everything else fixed and varying only this choice isolates the effect of the relational channel, which Chapter 7 analyses through the geometry of the tangent kernel introduced in §2.2.5.

2.5 Expressivity, approximation, and generalisation

The central reason neural networks have become the dominant model class in machine learning is their *expressivity*. However, in contrast with what classical statistical theory suggests, neural networks have been found to mitigate the curse of dimensionality, generalise well despite overparameterization, and optimise effectively in highly non-convex landscapes. Understanding these phenomena has been a major focus of recent theoretical research. In this section, we review key results on the approximation capabilities of neural networks, with an emphasis on implications for physical modelling.

2.5.1 Universal approximation theorems

Foundational results established that neural networks are *universal approximators*. It has been proved that a feed-forward network with at least one hidden layer and a suitable nonlinear activation (e.g. sigmoid, ReLU, tanh) can approximate any continuous function on a compact set to arbitrary accuracy [35, 36]. Formally:

Theorem 2.1 (Universal Approximation, informal). *Let f^* be continuous on a compact $K \subset \mathbb{R}^d$. Then for any $\epsilon > 0$, there exists a neural network f_θ with a single hidden layer such that*

$$\sup_{x \in K} |f_\theta(x) - f^*(x)| < \epsilon.$$

This result guarantees existence but says nothing about how many neurons are required, how they should be trained, or how generalisation occurs.

2.5.2 Depth versus width

Cybenko’s theorem establishes that a *single* hidden layer with a nonpolynomial sigmoidal activation is a *universal approximator*: for every continuous f on a compact domain and every $\varepsilon > 0$ there exists a one-hidden-layer network whose output g satisfies $\|f - g\|_\infty < \varepsilon$ [35]. The result concerns *existence* of an approximant and does not provide constructive bounds on the number of units required, rates of approximation, or generalisation guarantees.

In practice, depth is introduced for efficiency, not for universality. By composing intermediate features, additional layers often reduce the number of units required to attain a target accuracy. For certain target classes, there are known separations where a one-hidden-layer network needs exponentially many neurons to match the accuracy achievable with a deep architecture of polynomial size. Hence modern models leverage depth primarily for parameter efficiency and a stronger inductive bias (via hierarchical feature reuse), while width remains the lever for raw capacity [22].

2.5.3 Curse of dimensionality

Classical approximation theory predicts that uniformly approximating a generic $f : \mathbb{R}^d \rightarrow \mathbb{R}$ requires resources that grow exponentially in d —the *curse of dimensionality* (CoD). This perspective cannot explain the strong empirical performance of deep networks on very high-dimensional tasks, prompting a search for structural conditions under which the CoD does *not* apply.

A first resolution is *low effective dimension*: if the data (and loss) live on a $d' \ll d$ manifold M and error is measured in $L^p(\mu)$ with $\text{supp } \mu \subset M$, then neural networks can learn local charts and partitions of unity and achieve rates that depend on d' rather than the ambient d . This replaces ambient complexity by the intrinsic geometry of M , yielding polynomial (rather than exponential) scaling.

A second route is *spectral/Barron regularity*: for targets with finite first Fourier moment (Barron class), two-layer networks attain a *dimension-independent* $O(n^{-1/2})$ L^2 -approximation error with n hidden units. Deep and compositional Barron spaces extend this beyond smooth settings, providing broad families where the CoD provably disappears.

A third line exploits *PDE structure*. For many high-dimensional linear and semilinear parabolic/elliptic problems, stochastic representations (e.g., Feynman–Kac) and NN emulations of time-stepping yield approximation bounds whose complexity scales polynomially in d , again sidestepping the CoD under standard conditions.

Depth then clarifies *why* these escapes are efficient. Certain functions admit polynomial-size deep representations but require exponentially wide shallow ones (Eldan–Shamir separations), and deep ReLU networks generate exponentially more affine regions or oscillations per parameter than single-hidden-layer models. Thus, depth supplies the compositional mechanism that turns structure (manifolds, spectral regularity, PDE priors) into dimension-tolerant approximation.

2.5.4 Generalisation in overparameterized networks

Modern networks often have far more parameters than training samples, yet they fit the data *and* perform well on new inputs. This looks paradoxical if we judge models only by parameter count. A better lens is to ask which interpolating solution the training dynamics actually pick: test error is governed less by how large the hypothesis class is and more by the *kind* of function the algorithm converges to.

Two ideas make this concrete. First, stability: if small changes in the dataset produce only small changes in the learned predictor, the train–test gap remains small even at scale,

$$|\mathcal{R}(\hat{\theta}) - \hat{\mathcal{R}}(\hat{\theta})| \text{ is controlled.}$$

Second, implicit regularization: stochastic gradient methods (with finite batches, step sizes, and momentum) do not search the space uniformly; they are biased toward predictors with lower effective complexity—e.g., larger margins, smaller norms, or flatter basins in parameter space. In very wide nets, the dynamics are well-approximated by a linearized (kernel-like) model that interpolates while implicitly minimizing a function-space norm; at realistic widths, the same bias persists in practice. This also explains *double descent*: near the interpolation threshold test error can worsen, then improves again as capacity grows and the dynamics have more low-complexity interpolants to select from.

2.5.5 Optimisation in deep networks

Training is a large-scale, nonconvex problem, yet simple first-order methods reliably drive the loss to (near) zero. Two features help. Overparameterization and modern architectures (residual connections, normalisation) shape the landscape into broad, connected valleys of low loss and maintain usable descent directions through depth. As a result, poor isolated minima are less of a practical obstacle.

Stochastic gradient descent can be viewed as gradient flow with noise whose effective “temperature” is set mainly by the learning-rate-to-batch-size ratio. A higher temperature encourages escape from sharp minima and biases the search toward wider, flatter regions that tend to generalise better; annealing the temperature over time refines the final solution. Momentum accelerates along coherent directions and damps high-frequency noise; normalisation keeps activations well-scaled so larger, stable steps are possible. In short, the optimiser does more than find zero training error—it determines *which* zero-error solution you reach, and that choice largely sets generalisation quality.

2.5.6 Implications for physical modelling

In physical sciences, target functions such as wavefunctions are continuous, smooth (except at singularities like cusps), and often structured by symmetries. Neural networks thus provide a flexible ansatz space:

1. They can approximate smooth functions and their derivatives, which is crucial when differential operators appear in the loss (as in PINNs).
2. Depth allows efficient capture of multi-scale and compositional structure, relevant for many-body interactions.
3. Constraints such as permutation invariance or antisymmetry can be incorporated to encode physics explicitly.

These theoretical guarantees justify the use of neural networks as universal, symmetry-aware function approximators, laying the groundwork for our discussions of activation functions, initialisation strategies, and physics-informed extensions.

2.6 Physics-Informed Neural Networks

Traditional machine learning focuses on fitting models to labeled data. In scientific applications, however, governing equations such as differential equations encode a wealth of prior information about the system. *Physics-Informed Neural Networks (PINNs)* integrate these constraints directly into the learning process, combining data-driven approximation with analytical structure. Figure 2.1 illustrates how the behaviour of neural networks changes when working with higher order derivatives, as is common in PINNs.

2.6.1 Formulation of PINNs

Let $u : \Omega \subset \mathbb{R}^d \rightarrow \mathbb{R}$ denote the unknown solution of a partial differential equation (PDE) of the form

$$\mathcal{N}[u](x) = 0, \quad x \in \Omega, \quad (2.31)$$

subject to boundary conditions $u(x) = g(x)$ for $x \in \partial\Omega$. Here $\mathcal{N}$ is a (possibly nonlinear) differential operator.

In a PINN, $u_\theta(x)$ is represented by a neural network, and the loss function enforces both data fitting and PDE residual minimization:

$$\mathcal{L}(\theta) = \lambda_{\text{data}} \sum_{i=1}^{n_d} |u_\theta(x_i) - y_i|^2 + \lambda_{\text{PDE}} \sum_{j=1}^{n_r} |\mathcal{N}[u_\theta](\xi_j)|^2 + \lambda_{\text{BC}} \sum_{k=1}^{n_b} |u_\theta(\zeta_k) - g(\zeta_k)|^2. \quad (2.32)$$

Here $\{x_i, y_i\}$ are observed data, $\{\xi_j\}$ are collocation points in the domain for enforcing the PDE, and $\{\zeta_k\}$ are boundary points. The weights $\lambda_{\text{data}}, \lambda_{\text{PDE}}, \lambda_{\text{BC}}$ balance the different contributions.




Figure 2.1: Comparison of common activation functions and their first, second, and third derivatives. Smooth activations (GELU, Mish) yield stable derivatives, while Tanh and Sigmoid have less stable higher derivatives.

2.6.2 Automatic differentiation

A key ingredient of PINNs is the use of *automatic differentiation* (AD) to evaluate ∇u_θ and Δu_θ . Unlike finite-difference approximations, AD computes derivatives exactly (up to floating-point error), making it feasible to incorporate higher-order differential operators directly into the loss. This enables end-to-end training of networks constrained by physical laws.

2.6.3 Theoretical difficulties

Although PINNs are conceptually straightforward, making them reliable and scalable raises several non-trivial challenges that go beyond classical universal approximation results.

Approximation in Sobolev spaces. Universal approximation theorems control function error in L^2/L^∞ , but strong-form PINN losses depend on *derivatives* through $L[u_\theta]$. To make residuals small one must approximate in matching Sobolev norms $W^{\ell,p}$ (with ℓ the PDE order), which imposes smooth activations (e.g., tanh/SiLU for $\ell \geq 2$) and architecture choices that control higher derivatives. Low-regularity targets (discontinuities, cusps) break global $W^{\ell,p}$ control and favor weak/variational or entropy-based formulations.

Stability (residual-to-error). A small residual does not imply a small solution error unless the forward problem is (conditionally) stable. For well-posed elliptic/parabolic problems one can bound $\|u - u_\theta\|$ by $\|L[u_\theta] - f\|$, but inverse/ill-posed settings may require regularization, priors, or restricted hypothesis classes to prevent “residual overfitting” with large state error.

Sampling and generalisation. Collocation-based training minimizes a sampled surrogate of a continuum loss. Generalisation requires capacity control (e.g., bounded weights/Lipschitz loss) and sufficient sampling of interior vs. boundary/IC points; otherwise the empirical residual can be small while the population residual remains large.

Optimisation and conditioning. PINN losses couple multiple physics terms (interior, boundary, data) with disparate scales, producing ill-conditioned gradient dynamics. In the linearized/NTK view, convergence speed is governed by the spectrum of a problem-dependent kernel; large condition numbers lead to slow or stalled training.

Spectral bias. Gradient-based training tends to fit low-frequency components first, delaying convergence for solutions with sharp layers or oscillations unless the sampling, features, or architecture emphasize high-frequency content (e.g., Fourier features, windowed bases, curriculum).

Mitigating these challenges requires careful design of architectures, activations, initialisation, loss weighting, and sampling strategies. These topics are explained and discussed in detail in [37]

2.6.4 Relevance to this thesis

For scientific machine learning, PINNs provide a principled framework for embedding physical structure into neural networks. In this thesis, they motivate the choice of smooth activations (for stable derivatives), careful initialisation (to prevent exploding Hessians), and architectural constraints (to encode symmetry and physical laws). Thus, PINNs serve both as a methodological tool and as a theoretical lens for understanding the interplay between learning and physics.

2.6.5 Implications for scientific machine learning

For physics-informed models, generalisation takes on a slightly different meaning: the goal is not merely prediction on unseen data, but consistency with underlying physical laws. Inductive biases (symmetry, smoothness, conservation laws) are therefore not optional but essential. Implicit regularization from gradient-based optimisation complements these biases by favoring solutions that are both smooth and physically plausible. Generalisation in neural networks arises from an interplay of:

1. Algorithmic stability of gradient-based methods,
2. Implicit bias toward simple, smooth, low-complexity solutions, and
3. Architectural inductive biases encoding domain knowledge.

In scientific computing, these mechanisms explain why neural networks, despite their size, can approximate physically meaningful solutions. They also justify the careful design choices of activations, initialisation, and constraints emphasized throughout this chapter.

Mitigation in practice. The challenges discussed for PINNs do not resolve themselves; they must be engineered away. In brief: use weak/variational formulations (or energy principles) when higher-order derivatives make strong forms unstable; hard-enforce essential boundary conditions and adapt weights for the rest; non-dimensionalize and balance loss terms to improve conditioning; prefer smooth activations and architectures that control derivatives; exploit symmetry, equivariance, and locality to reduce effective complexity; precondition optimisation with second-order or natural-gradient methods (e.g., L-BFGS, Stochastic Reconfiguration); and design sampling (importance/stratified, curricula in time/space) to close the generalisation gap. Together, these measures turn inductive bias into concrete numerical stability and reliable, law-consistent generalisation.

2.7 Summary and Outlook

Three threads from this chapter carry into the rest of the thesis. Framing learning as empirical risk minimisation makes *generalisation*—not fitting—the real problem, and locates it in the inductive bias of the model rather than in raw capacity. The optimisation view adds that the training *trajectory* matters as much as the objective: gradient descent carries its own implicit bias, and the natural gradient re-expresses that trajectory in the geometry of the model’s own outputs, the geometry we will later read directly through the tangent kernel. The physics-informed view supplies the binding constraint: a wavefunction ansatz must be differentiated twice, cleanly, to yield a local energy, so smooth activations, stable initialisation, and hard-wired symmetries are prerequisites rather than cosmetic choices. These are the principles that guide the architecture and diagnostics of the chapters that follow.

Part III

Methods

Chapter 3

Trial Wavefunction

No great thing is created suddenly, any more than a bunch of grapes or a fig.

EPICETETUS

Discourses, Book I, Chapter 15

In this chapter we specify the concrete modelling and algorithmic choices used in our computations. We proceed from the physical setup and trial states to objectives, with an emphasis on stability, symmetry, and differentiability—all of which are critical for Laplacian-based local-energy evaluations and stochastic reconfiguration (SR). Unless otherwise stated, we work in atomic units.

3.1 Preliminaries

Hamiltonian and units. We study N spin- $\frac{1}{2}$ fermions in two spatial dimensions confined by a harmonic trap and interacting via Coulomb repulsion,

$$H = \sum_{i=1}^N \left(-\frac{1}{2} \nabla_{\mathbf{r}_i}^2 + \frac{1}{2} \omega^2 r_i^2 \right) + \sum_{1 \leq i < j \leq N} \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|}, \quad (\hbar = m_e = e = 1). \quad (3.1)$$

Spin does not appear in H ; eigenstates factorize into a spatial function with good L_z and a spin function with good (S, S_z) .

Coordinates and scaling. We collect positions as $R = (\mathbf{r}_1, \dots, \mathbf{r}_N) \in \mathbb{R}^{2N}$ and use trap scaling $\tilde{\mathbf{r}}_i = \sqrt{\omega} \mathbf{r}_i$, $\tilde{R} = \sqrt{\omega} R$. This aligns typical length scales to $\mathcal{O}(1)$, improves conditioning of second derivatives, and makes the noninteracting ground state ω -independent. When useful, we also use pair distances $r_{ij} = |\mathbf{r}_i - \mathbf{r}_j|$ and their scaled versions $\tilde{r}_{ij} = |\tilde{\mathbf{r}}_i - \tilde{\mathbf{r}}_j|$.

Sectors and focus. We primarily report *closed-shell* cases (unique spin singlet at the noninteracting level). Open shells introduce a degenerate manifold and additional choices for the reference; we keep this to brief remarks only where needed.

Spin-orbitals and basis. One-particle spatial orbitals are the Cartesian two-dimensional harmonic-oscillator eigenstates $\{\phi_{n_x n_y}(\mathbf{r})\}$ with $E_{n_x n_y} = \omega(n_x + n_y + 1)$, $n_x, n_y \in \mathbb{N}_0$ (the basis used in the implementation; for the closed shells studied it spans the same filled shells as any complete basis of those shells). Spin-orbitals are $\varepsilon_\alpha(x) = \phi_{n_x n_y}(\mathbf{r}) \chi_\sigma(s)$, $\sigma \in \{\uparrow, \downarrow\}$.

3.2 Overview of the ansatz

Our trial state combines exact antisymmetry, an analytic short-range cusp, and a compact neural residual, while optionally evaluating the Slater determinant on backflowed coordinates:

$$\Psi_{\theta,\beta}(R) = \text{SD}(\tilde{R}') \exp\left(\sum_{i<j} u_{\sigma_i\sigma_j}(r_{ij}) + W_{\theta}(\tilde{R})\right), \quad (3.2)$$

$$\tilde{R}' = \tilde{R} + \Delta_{\beta}(\tilde{R}). \quad (3.3)$$

Here SD enforces antisymmetry; u hard-wires the Kato cusp; W_{θ} (the NQS correlator) learns smooth medium/long-range structure; and Δ_{β} (backflow) adjusts the nodes by smoothly warping the coordinates used inside the determinants; its per-particle displacements $\Delta_{\beta,i}$ are written $\Delta\mathbf{x}_i$ in the results chapters. We deliberately feed the NQS with $\tilde{R}$ (not $\tilde{R}'$) to avoid self-reinforcing feedback and keep local-energy variance low.

Design principles. (i) *Separate guaranteed physics from learned details:* antisymmetry and the cusp are hard-wired; the network learns the rest. (ii) *Condition the geometry:* work in trap units and use soft-core pair features so that first/second derivatives are bounded. (iii) *Keep the learned object small and smooth:* once the cusp is explicit, a compact NQS suffices across (N, ω) .

3.3 Slater reference

For closed shells we use a restricted product of spin-block determinants of the lowest $N/2$ harmonic oscillator orbitals (trap units):

$$\text{SD}(\tilde{R}) = \det[\phi_a(\tilde{\mathbf{r}}_i)]_{i \in \uparrow, a \leq N/2} \det[\phi_a(\tilde{\mathbf{r}}_j)]_{j \in \downarrow, a \leq N/2}. \quad (3.4)$$

The correlator reweights amplitudes *within* nodal pockets; without backflow, the nodes are those of (3.4).

3.4 Neural correlator W_{θ}

As with the backflow, the correlator comes in two forms that we compare directly (Q1a, Chapter 7): a *separable* network that encodes particle- and pair-features independently and pools them, and a *copresheaf message-passing* (CTNN) network. We describe the separable network first—it is the baseline and also supplies the safe features and analytic cusp shared by both—and then the CTNN.

3.4.1 Separable correlator (DeepSets baseline)

The separable NQS aggregates *particlewise* and *pairwise* embeddings with permutation-invariant pooling and appends two global statistics, all in $\tilde{R}$. A core safety requirement is that pair channels satisfy $ds_k/d\tilde{r} \rightarrow 0$ as $\tilde{r} \rightarrow 0$ so that W_{θ} never injects spurious $1/\tilde{r}$ terms into $\nabla \log \Psi$ or $\Delta \log \Psi$.

Particle branch (DeepSets).

This part of the network is typically referred to as Deep Sets [32]. The permutation invariance of quantum many-body mechanics is naturally captured by this architecture. A neural network ϕ creates a latent embedding for each particle individually. All embeddings are pooled together to form a global representation, which is used later in the readout. Formally, we have

$$\mathbf{h}_i^{\phi} = \phi(\tilde{\mathbf{x}}_i) \in \mathbb{R}^{d_L}, \quad \bar{\mathbf{h}}^{\phi} = N^{-1} \sum_i \mathbf{h}_i^{\phi} \in \mathbb{R}^{d_L}.$$

Pair branch (soft-core, even channels).

Although the particle branch could theoretically produce pair correlations, in practice it struggles to represent sharp near-field structure. We therefore add a dedicated pair branch that builds features from interparticle distances. This network processes all pairs $(i, j), i \neq j$ individually and pools the resulting latent embeddings. The problem is that raw distances $\tilde{r}_{ij}$ have unbounded derivatives at coalescence, which destabilizes Laplacian evaluations. We therefore build *safe* pair features with bounded first and second derivatives as $\tilde{r} \rightarrow 0$. With $\tilde{r}^{\text{soft}} = \sqrt{\tilde{r}^2 + \varepsilon^2}$ ($\varepsilon \in [0.15, 0.30]$ in trap units), we build six “safe” features

$$\begin{aligned} s_1 &= \log(1 + (\tilde{r}^{\text{soft}}/\varepsilon)^2), & s_2 &= \frac{\tilde{r}^2}{\tilde{r}^2 + \varepsilon^2}, & s_3 &= (\tilde{r}^{\text{soft}}/\varepsilon)^2 \exp[-(\tilde{r}^{\text{soft}}/\varepsilon)^2], \\ \text{rbf}_g &= \exp(-g s_1), & g &\in \{0.25, 1, 4\}, & \mathbf{u}_{ij} &= [s_1, s_2, s_3, \text{rbf}_{0.25}, \text{rbf}_1, \text{rbf}_4] \in \mathbb{R}^6. \end{aligned} \quad (3.5)$$

An MLP ψ maps $\mathbf{u}_{ij} \mapsto \mathbf{h}_{ij}^\psi \in \mathbb{R}^{d_L}$. A short-range gate

$$\chi(\tilde{r}) = \frac{\tilde{r}^2}{\tilde{r}^2 + r_g^2}, \quad r_g \approx 0.3, \quad \chi(0) = \chi'(0) = 0,$$

suppresses learned responses *at* coalescence; downstream we use $\chi \mathbf{h}_{ij}^\psi$.

3.4.2 Analytic short-range cusp (shared)

The analytic cusp below is shared by both correlator variants (and is independent of the backflow). Because the safe features are necessary in the pair branch, the NQS cannot represent the Kato cusp condition exactly. We therefore add an analytic cusp term to the exponent in (3.2), written in *physical* coordinates so that its slope at contact is manifestly correct:

$$u_{\sigma\sigma'}(r) = \gamma_{\sigma\sigma'} r e^{-r/a_{\text{ho}}}, \quad a_{\text{ho}} = 1/\sqrt{\omega}, \quad \gamma_{\uparrow\downarrow} = \frac{1}{d-1}, \quad \gamma_{\uparrow\uparrow} = \frac{1}{d+1}. \quad (3.6)$$

Because the linear prefactor is the physical separation r , the slope at contact is $\partial_r u|_0 = \gamma_{\sigma\sigma'}$, exactly the Kato condition in general d ; for $d = 2$ this gives $\gamma_{\uparrow\downarrow} = 1$ and $\gamma_{\uparrow\uparrow} = \frac{1}{3}$, the standard two-dimensional quantum-dot values, derived—including the parallel-spin case, where antisymmetry already forces a linear node—in Appendix B.2. Only the exponential taper carries the trap length a_{ho} , suppressing long-range interference so the NQS need not unlearn cusp structure at medium and long range; in trap units ($\tilde{r} = \sqrt{\omega} r$) the same term reads $u = (\gamma_{\sigma\sigma'}/\sqrt{\omega}) \tilde{r} e^{-\tilde{r}}$.

3.4.3 Separable readout

With contributions from the particle and pair branches, we form a global representation for the final readout. In addition to the pooled embeddings $\bar{\mathbf{h}}^\phi$ and $\bar{\mathbf{h}}^\psi$, we also include two global statistics that help the network gauge overall system size and density. We average pairs by default or apply degree-normalised radial attention

$$w_{ij} = \exp[-(\tilde{r}_{ij}/r_c)^p], \quad \hat{w}_{ij} = w_{ij} / \sum_{k \neq i} w_{ik}, \quad (r_c, p) = (0.4, 6),$$

then form $g_i = \sum_{j \neq i} \hat{w}_{ij} \mathbf{h}_{ij}^\psi$ and $\bar{\mathbf{h}}^\psi = N^{-1} \sum_i g_i$. Two system-level scalars in trap units,

$$\mathbf{g}(\tilde{R}) = \left[\frac{1}{Nd} \sum_i \|\tilde{\mathbf{x}}_i\|^2, \frac{2}{N(N-1)} \sum_{i < j} s_1(\tilde{r}_{ij}) \right] \in \mathbb{R}^2,$$

are concatenated with $\bar{\mathbf{h}}^\phi$ and $\bar{\mathbf{h}}^\psi$ and mapped to a scalar by a tiny MLP ρ :

$$W_\theta(\tilde{R}) = \rho(\bar{\mathbf{h}}^\phi \parallel \bar{\mathbf{h}}^\psi \parallel \mathbf{g}(\tilde{R})). \quad (3.7)$$

3.4.4 Message-passing correlator (CTNN)

The message-passing correlator (code: `CTNNJastrowVCycle`) keeps the same node/edge embeddings, safe pair features, and analytic cusp, but replaces the independent pooling of the separable network with the

copresheaf transport of §2.4.4: at each round the endpoint nodes are transported into an edge update and the updated edges are transported back and aggregated onto the nodes,

$$\mathbf{h}_{ij} += F_e(\mathbf{h}_{ij} \parallel \rho_{v \rightarrow e} \mathbf{h}_i \parallel \rho_{e \rightarrow v} \mathbf{h}_j), \quad \mathbf{h}_i += F_v\left(\mathbf{h}_i \parallel \text{Agg}_{j \neq i} w_{ij}^{\text{spin}} \rho_{e \rightarrow v} \mathbf{h}_{ij}\right), \quad (3.8)$$

with residual updates and per-round transport maps. Unlike the single-round backflow, the correlator stacks these rounds in a multiscale “V-cycle”: $n_\downarrow$ down-pass rounds, a linear coarsening to a bottleneck width, then $n_\uparrow$ up-pass rounds with skip connections back to the down-pass features (defaults $n_\downarrow = n_\uparrow = 2$). A permutation-invariant readout over the final node and edge statistics returns the scalar $W_\theta(\tilde{R})$, exactly as in (3.7). Holding the Slater reference, cusp, and training fixed and swapping only the separable readout for this V-cycle isolates the relational channel studied in Q1a (Chapter 7).

3.5 Backflow

Backflow deforms the coordinates *used only inside the Slater reference* so that the nodal surface can adapt to interparticle correlations:

$$\tilde{R}' = \tilde{R} + \Delta_\beta(\tilde{R}), \quad \Psi_{\theta, \beta}(R) = \text{SD}(\tilde{R}') \exp\left(\sum_{i < j} u_{\sigma_i \sigma_j}(r_{ij}) + W_\theta(\tilde{R})\right). \quad (3.9)$$

The correlator W_θ and cusp u are evaluated on $\tilde{R}$ to avoid feedback loops in the local energy; only the determinant “sees” $\tilde{R}'$. This choice cleanly separates (i) amplitude reweighting within nodal pockets (handled by $u + W_\theta$) from (ii) nodal *geometry* (handled by backflow).

Design goals. (i) *Stability near coalescence*: messages are built from mollified radii and short-range gates so that $\partial \Delta_\beta / \partial \tilde{r}$ stays bounded; (ii) *Permutation symmetry*: permutation equivariance holds by construction (shared maps and symmetric aggregation); (iii) *Close to identity at initialisation*: a zero-initialised last layer and a positive learnable scale keep $\Delta_\beta \approx 0$ at the start. Rotational symmetry, by contrast, is only *approximate*: the displacement head reads out a d -vector from node features that include the raw coordinates, so the map is not rotation-equivariant by construction, and it is encouraged instead by the geometry-based edge features and by rotationally-augmented sampling. A global translation is not a symmetry of the trap in any case; the zero-mean projection (3.19) is a regulariser that removes a spurious centre-of-mass drift, not an imposed physical symmetry.

We study *two* backflow architectures. Both map the configuration to a per-particle displacement field and share the output bounding, spin masking, and centre-of-mass projection below; they differ only in how a particle’s displacement is made to depend on the others. The conventional network is the baseline of the architecture comparison in Chapter 7; the copresheaf CTNN is the production ansatz.

3.5.1 Conventional backflow (single-round messages)

The baseline (code: `BackflowNet`) uses a single round of pairwise messages. For each ordered pair (i, j) with $i \neq j$,

$$\mathbf{m}_{ij} = \phi_\beta(\tilde{\mathbf{x}}_i, \tilde{\mathbf{x}}_j, \mathbf{r}_{ij}, \tilde{r}_{ij}^{\text{soft}}, (\tilde{r}_{ij}^{\text{soft}})^2) \cdot w_{ij}^{\text{spin}}, \quad \tilde{r}_{ij}^{\text{soft}} = \sqrt{\tilde{r}_{ij}^2 + \varepsilon^2}, \quad (3.10)$$

with ϕ_β a small MLP and $w_{ij}^{\text{spin}} \in \{0, 1\}$ a spin mask (same-spin only or all pairs). Messages are aggregated to a node state and a node network returns the displacement,

$$\mathbf{m}_i = \text{Agg}_{j \neq i} \mathbf{m}_{ij}, \quad \delta \tilde{\mathbf{x}}_i = \psi_\beta(\tilde{\mathbf{x}}_i \parallel \mathbf{m}_i), \quad \text{Agg} \in \{\text{sum}, \text{mean}, \text{max}\}. \quad (3.11)$$

There is no persistent edge state: each message is recomputed from the raw geometry and discarded after aggregation.

3.5.2 CTNN backflow (coplesheaf)

The message-passing architecture (code: `CTNNBackflowNet`) instead maintains explicit *node and edge* features and transports information between them through learned linear maps—the copresheaf-style step of §2.4.4. Node features are embedded from the position and a spin scalar, edge features from the relative geometry,

$$\mathbf{h}_i^{(0)} = W_v [\tilde{\mathbf{x}}_i, s_i], \quad \mathbf{h}_{ij}^{(0)} = \text{MLP}_e(\mathbf{r}_{ij}, |\mathbf{r}_{ij}|, |\mathbf{r}_{ij}|^2). \quad (3.12)$$

A node-to-edge transport $\rho_{v \rightarrow e}$ feeds the endpoint nodes into an edge update; an edge-to-node transport $\rho_{e \rightarrow v}$ returns the updated edges to the nodes, which are refined by a residual node update; a linear head reads out the displacement:

$$\mathbf{h}'_{ij} = \text{MLP}'_e(\mathbf{h}_{ij}^{(0)} \parallel \rho_{v \rightarrow e} \mathbf{h}_i^{(0)} \parallel \rho_{v \rightarrow e} \mathbf{h}_j^{(0)}), \quad (3.13)$$

$$\mathbf{m}_i = \text{Agg}_{j \neq i} w_{ij}^{\text{spin}} \rho_{e \rightarrow v}(\mathbf{h}'_{ij}), \quad (3.14)$$

$$\mathbf{h}_i = \mathbf{h}_i^{(0)} + \text{MLP}_v(\mathbf{h}_i^{(0)} \parallel \mathbf{m}_i), \quad \delta \tilde{\mathbf{x}}_i = W_\Delta \mathbf{h}_i. \quad (3.15)$$

The learned transport maps $\rho_{v \rightarrow e}, \rho_{e \rightarrow v}$ are the channel whose ablation collapses the backflow at the Wigner crystal (Chapter 7). The backflow uses a single copresheaf round; the message-passing *correlator* of the architecture comparison (`CTNNJastrowVCycle`) iterates the same style of update over a down/up “V-cycle”.

3.5.3 Shared output, scaling, and initialisation

Both variants take inputs in trap units ($\tilde{\mathbf{x}} = \sqrt{\omega} \mathbf{x}$), bound the displacement, and start at the Slater nodes,

$$\Delta_\beta(\tilde{R}) = \tanh(\delta \tilde{X}) \cdot s_\beta, \quad s_\beta = \text{softplus}(s_\beta^{\text{raw}}) > 0, \quad (3.16)$$

with the last linear layer (ψ_β or W_Δ) zero-initialised so that $\Delta_\beta \approx 0$ initially. Empirically this parameterisation stabilises the SR/energy tails and protects the Laplacian.

3.5.4 Short-range behaviour and gates

As $\tilde{r}_{ij} \rightarrow 0$, raw $1/\tilde{r}$ factors can make $\nabla \cdot \Delta_\beta$ spiky. We therefore (i) mollify $\tilde{r}_{ij}$ as in (3.10), and optionally (ii) gate the message with

$$\chi_{\text{bf}}(\tilde{r}_{ij}) = \frac{\tilde{r}_{ij}^2}{\tilde{r}_{ij}^2 + r_g^2}, \quad \chi_{\text{bf}}(0) = \chi'_{\text{bf}}(0) = 0, \quad r_g \approx 0.3, \quad (3.17)$$

or with degree-normalised radial attention

$$w_{ij} = \exp[-(\tilde{r}_{ij}/r_c)^p], \quad \hat{w}_{ij} = \frac{w_{ij}}{\sum_{k \neq i} w_{ik}}. \quad (3.18)$$

Either choice keeps the learned response smooth at coalescence and limits the BF contribution to the local energy variance.

3.5.5 Symmetries and COM neutrality

Permutation symmetry is automatic (pairwise messages and symmetric aggregation). To avoid a spurious centre-of-mass (COM) drift, we project the flow to zero mean:

$$\Delta_\beta \leftarrow \Delta_\beta - \frac{1}{N} \sum_{i=1}^N \Delta_{\beta,i}. \quad (3.19)$$

The messages are built from the relative vectors $\mathbf{r}_{ij}$ and the zero-mean projection removes the spurious centre-of-mass drift; as noted under the design goals, the map is not translation-invariant or rotation-equivariant by construction (the displacement head reads a d -vector from the absolute coordinates).¹

¹Implementation note: the mean is taken over particles *within each configuration*, not across configurations in the batch, so the constraint removes centre-of-mass motion configuration by configuration. It is a one-liner and removes the small residual COM drift seen in ablations.

3.5.6 Effect on the nodes and variational safety

Because only SD is evaluated at $\tilde{R}'$, backflow changes the *nodes* but leaves the correlator exponent on $\tilde{R}$. No Jacobian term is introduced (we are not performing a change of variables of the full wavefunction). This “Feynman–Cohen style” design targets fixed-node error directly while keeping the amplitude model simple and smooth.

3.5.7 Regularization, initialisation, and cost

We found the following defaults robust across (N, ω) : (i) $\varepsilon \in [10^{-6}, 10^{-4}]$ (trap units) for the soft norm; (ii) bounded output $\tanh$ with a learnable scale s_β initialised in $[0.02, 0.10]$; (iii) optional gate χ_{bf} with $r_g \approx 0.3$ (trap units). Zero-initialising the last layer of ψ_β makes the ansatz start exactly at the Slater nodes and improves optimiser stability. The compute/memory cost scales as $O(BN^2(d + H))$ for batch size B and hidden width H .

3.5.8 Mapping to implementation

In the conventional network, ϕ_β and ψ_β correspond to `BackflowNet.phi` (message MLP) and `BackflowNet.psi` (node/update MLP). In the copresheaf network, the embeddings, transport maps, and update MLPs of (3.13)–(3.15) correspond to CTNNBackflowNet’s `node_embed/edge_embed`, `rho_v_to_e/rho_e_to_v`, `edge_update/node_update`, and `dx_head`. For both, aggregation is selectable (`sum|mean|max`); spin handling (`use_spin, same_spin_only`) produces the mask w_{ij}^{spin} ; output control matches (3.16) via `out_bound="tanh"` and $s_\beta = \text{softplus}(\text{bf_scale_raw})$; and (3.19) is enforced by subtracting the per-configuration mean over particles of the predicted Δ_β before forming $\tilde{R}'$.

Summary. Backflow provides a compact, symmetry-respecting way to move the Slater nodes toward the true correlated nodes. With mollified distances, optional short-range gates/attention, zero-mean projection, and bounded outputs, it integrates cleanly with the analytic cusp and smooth NQS correlator, designed to reduce nodal error while controlling local-energy variance.

3.6 Architecture sizes and hyperparameters

Table 3.1 collects the representative settings, so that the architecture claims of Chapter 7 can be read as matched-capacity comparisons rather than size effects. The controlled correlator/backflow study fixes each pair of variants at the same capacity tier; the reported ground-state energies use the wider production networks. A key point is that the message-passing and separable variants are held at comparable parameter counts, so a difference in outcome cannot be attributed to one simply being larger. All settings below are those of the accompanying code (`scripts/run_depth_analysis.py` for the comparison tiers; `scripts/source_state_check.py` for the production widths).

Table 3.1: Representative architecture and training hyperparameters. Widths are hidden dimensions; “rounds” are message-passing steps ($n_\downarrow + n_\uparrow$ for the V-cycle). The two correlator variants (and the two backflow variants) are matched in capacity within the comparison tier.

Component	Variant	Comparison tier	Production
Correlator	CTNN (V-cycle)	node/edge 32, bottleneck 16, rounds 2+2	width 128, 2 layers
	DeepSet	pair-MLP 64×4 , out 32	width 128, 2 layers
Backflow	CTNN (copresheaf)	node/edge 64, 1 round	width 128, 3 layers
	Conventional	msg 64, node-MLP 64×3	width 128, 3 layers
Readout	(correlator)	hidden 64, 2–3 layers, SiLU	—
Optimiser	Adam	$\text{lr} = 5 \times 10^{-4}$ (BF), 5×10^{-5} (Jastrow)	
	CG-SR	Tikhonov damping λ , trust-region cap, CG restarts	
Batch	collocation	$n_{\text{coll}} \sim 3 \times 10^3$, oversample $m = 8\text{--}16$	

Chapter 4

Optimisation

4.1 Primary objective and two-stage training

Given a parameterised wavefunction Ψ_θ , the variational ground-state energy is

$$E(\theta) = \frac{\langle \Psi_\theta | H | \Psi_\theta \rangle}{\langle \Psi_\theta | \Psi_\theta \rangle} = \mathbb{E}_{\tilde{R} \sim \pi_\theta} [E_L(\tilde{R})], \quad \pi_\theta(\tilde{R}) = \frac{|\Psi_\theta(\tilde{R})|^2}{\int |\Psi_\theta|^2}. \quad (4.1)$$

The local energy is evaluated in *physical* coordinates, consistently with the Hamiltonian (3.1),

$$E_L(R) = -\frac{1}{2} \sum_{i=1}^N \left[\Delta_{\mathbf{r}_i} \ln \Psi_\theta + \|\nabla_{\mathbf{r}_i} \ln \Psi_\theta\|^2 \right] + \frac{1}{2} \omega^2 \sum_{i=1}^N r_i^2 + \sum_{i<j} \frac{1}{r_{ij}}, \quad (4.2)$$

and the zero-variance principle holds: $\text{Var}_{\pi_\theta}[E_L] = 0$ iff Ψ_θ is an eigenstate. The network *features* are computed in trap units $\tilde{\mathbf{r}} = \sqrt{\omega} \mathbf{r}$ for conditioning, but the local energy itself is formed in physical units, so no ω factors are absorbed into the kinetic, trap, or Coulomb terms. (In trap units the same operator reads $E_L = \omega \left[-\frac{1}{2} \sum_i (\Delta_{\tilde{\mathbf{r}}_i} \ln \Psi_\theta + \|\nabla_{\tilde{\mathbf{r}}_i} \ln \Psi_\theta\|^2) + \frac{1}{2} \sum_i \tilde{r}_i^2 \right] + \sum_{i<j} \sqrt{\omega}/\tilde{r}_{ij}$; the kinetic and trap terms scale by ω and the Coulomb term by $\sqrt{\omega}$.)

Our optimisation follows two stages that mirror the implementation:

1. **Residual-based pretraining (stage I).** We fit the residual of E_L to a moving scalar target E_{eff} using the loss $\mathcal{L}_{\text{res}}(\theta) = \mathbb{E}[(E_L(\tilde{R}) - E_{\text{eff}})^2]$. We begin with a *self-target* (stabilizer) $E_{\text{eff}} = \mu$ where $\mu = \mathbb{E}[E_L]$ over the current batch. This choice is equivalent to variance minimization and prevents the network from chasing a potentially inaccurate external target before it can represent low-variance pockets. We then anneal toward a trusted reference E_{DMC} via

$$E_{\text{eff}}(\alpha) = \alpha E_{\text{DMC}} + (1 - \alpha) \mu, \quad \alpha \in [0, 1], \quad (4.3)$$

with a smooth cosine schedule $\alpha = \alpha(t)$ that ramps from α_{start} to α_{end} over a prescribed fraction of epochs (code: `alpha_start`, `alpha_end`, `alpha_decay_frac`). Setting the **objective** to "residual" uses $E_{\text{eff}} = \mu$; "energy" uses $E_{\text{eff}} = E_{\text{DMC}}$; and "energy_var" uses the annealed blend above.

2. **Stochastic Reconfiguration (stage II).** After residual pretraining, we refine with SR (natural-gradient VMC). Writing $O(\tilde{R}) = \partial_\theta \log \Psi_\theta(\tilde{R})$,

$$S = \mathbb{E}_\pi[O^\top O], \quad g = 2 \mathbb{E}_\pi[(E_L - \mathbb{E}_\pi[E_L]) O], \quad (4.4)$$

we solve the damped normal equations

$$(S + \lambda I) \Delta\theta = -g, \quad (4.5)$$

using (preconditioned) conjugate gradients with restarts, a trust-region scale on the final step, and batchwise centering/whitening of O (code: `damping`, `max_param_step`, CG options).

This pipeline exploits the zero-variance principle for stable shaping of W_θ (and backflow) before applying the geometry-aware SR update.

4.2 Configuration-space sampling

Guiding principle (permutation invariance). In a many-body fermionic system, *which* particle indices are close is irrelevant; what matters is *how many* close pairs and at which length scales. Consequently, we can design samplers that (i) emphasize *configurational motifs* (clusters, shells, dimers, long tails) rather than specific labels, and (ii) deliberately permute particles and randomly rotate configurations to explore the space efficiently without biasing towards any ordering.

4.2.1 Stratified capped-simplex mixture

We draw collocation batches $X \in \mathbb{R}^{B \times N \times d}$ from a five-component mixture,

(centre, tails, mixed, shells, dimers),

with nonnegative weights $w \in \Delta^4$ projected to a *capped simplex* $\{w : \sum_k w_k = 1, 0 \leq w_k \leq 0.3\}$ to enforce coverage (code: `_project_simplex_with_caps`). Each component proposes x as follows (all in physical units; the code internally scales by $a_{\text{ho}} = 1/\sqrt{\omega}$):

- **Centre:** narrow isotropic Gaussian around the trap centre (samples near the density core).
- **Tails:** wide Gaussian (probes low-density outskirts and large- r behaviour).
- **Mixed:** hybrid batches mixing narrow and moderate spreads per configuration (tests robustness to inhomogeneous spreads).
- **Shells:** points placed on K hyperspherical shells with trap-scaled radii $\{r_k\}$ and occupancy $\{q_k\}$, with small radial/tangential jitter (captures ring/shell order at weak traps).
- **Dimers:** we induce a few close pairs per configuration by sampling short interparticle offsets with random directions and log-uniform distances (enforces near-coalescence coverage without singularities).

We then apply a random in-plane rotation (for $d = 2$), and a random permutation of particle indices; both preserve the target expectations by symmetry.

Adaptive mixture and shell occupancy. Per epoch we compute simple difficulty scores and update w and q by exponentiated-gradient (EG) steps with momentum, temperature, and an exploration term; w is then projected back to the capped simplex (code: `eg_eta`, `eg_temp`, `eg_momentum`, `explore_gamma`, `prob_floor`). Mixture difficulty uses the residual $(E_L - E_{\text{eff}})^2$ per component; shell difficulty uses a proxy

$$\underbrace{V_i}_{\text{harm.+Coulomb per particle}} + \underbrace{\gamma_{g^2} \|\nabla_{\mathbf{x}_i} \log \Psi\|^2}_{\text{stiff gradients}} + \underbrace{\gamma_{\text{cusp}} r_{\min}^{-1}}_{\text{coalescence pressure}},$$

aggregated by nearest shell index (code: `g2_weight`, `cusp_gamma`, safe clip `rmin_clip`). This shifts probability mass toward underfit regions *without* collapsing coverage thanks to the caps.

Hard-example injection. We maintain a small buffer of previously challenging configurations (based on residuals) and stochastically inject them, with mild multiplicative and additive jitter, into the next batch (code: `sampler_hard*`). This provides a curriculum-like pressure while avoiding overfitting to outliers.

Robustness tweaks. We trim extreme local-energy outliers by quantiles (default 3%) *before* forming losses/updates; we also micro-batch the evaluation to keep memory bounded, and apply gradient clipping to f_θ (and backflow) parameters.

4.3 Residual-based training (stage I)

For a batch $\{\tilde{R}_b\}_{b=1}^B$ we compute $E_L(\tilde{R}_b)$ and the batch mean $\mu = \frac{1}{B} \sum_b E_L(\tilde{R}_b)$. The residual loss is

$$\mathcal{L}_{\text{res}}(\theta; \alpha) = \frac{1}{B} \sum_{b=1}^B (E_L(\tilde{R}_b) - E_{\text{eff}}(\alpha))^2, \quad E_{\text{eff}}(\alpha) = \alpha E_{\text{DMC}} + (1 - \alpha) \mu. \quad (4.6)$$

- **Self-target warmup** ($\alpha = 0$). Minimizing $\mathbb{E}[(E_L - \mu)^2]$ is exactly variance minimization; it teaches the network to produce smooth, low-variance amplitudes inside nodal pockets and reduces sensitivity to early sampler noise.
- **Anneal to reference** ($\alpha \uparrow$). As capacity grows, we blend in the external target, gently steering the mean energy while retaining the variance pressure. The cosine ramp (code: `alpha_start`, `alpha_end`, `alpha_decay_frac`) avoids abrupt objective switches.
- **Pure energy fit** ($\alpha = 1$). If desired, we can end stage I with $E_{\text{eff}} = E_{\text{DMC}}$.

In practice we compute $\nabla_{\theta} \mathcal{L}_{\text{res}}$ by autodiff, use micro-batches of size `micro_batch`, apply optional gradient clipping (code: `grad_clip`), and log per-component usage and shell radii in physical units.

Local-energy derivatives in practice. We support three Laplacian backends:

1. **Exact:** chunked second partials (reference-accurate, heavier).
2. **HVP-Hutchinson:** $\Delta \log \Psi = \mathbb{E}_v[v^{\top} H_{\log \Psi} v]$ with Rademacher v (fast and stable; fallback to FD if a probe is non-finite).
3. **FD-Hutchinson:** centred finite differences of directional gradients (robust in corner cases; slightly noisier).

We average a small number of probes (2–16 in ablations), and trim quantiles before forming losses.

4.4 Stochastic Reconfiguration (stage II)

Stochastic reconfiguration [29, 38] preconditions the energy gradient with the (inverse) quantum geometric tensor, and is the imaginary-time / natural-gradient step for a variational state [27, 30]. It uses the covariance form of the energy gradient

$$\partial_{\theta_k} E = 2 \text{Cov}_{\pi}(E_L, O_k) = 2 \left(\langle E_L O_k \rangle - \langle E_L \rangle \langle O_k \rangle \right), \quad (4.7)$$

and approximates a natural-gradient step by solving $(S + \lambda I) \Delta \theta = -g$ with the quantum geometric tensor S of §2.2.5. Implementation details:

- **Centering/whitening.** We centre O per batch and whiten by the diagonal of S before CG to improve conditioning.
- **Trust region.** The final step is scaled to satisfy a maximum parameter norm (code: `max_param_step`).
- **Damping.** A small λ (code: `damping`) stabilizes low-variance directions.

Because our backflow only alters the Slater coordinates (and W_{θ} stays on safe features of $\tilde{R}$), the SR statistics remain well-behaved even when nodes shift.

Optimiser variants compared. Three first-order updates appear in the results, and the comparison between them is itself a finding (Chapter 7, §7.2): (i) **Adam** [25], a diagonal adaptive step used as the robust default, with separate learning rates for the Jastrow and backflow blocks; (ii) **plain SR**, the natural-gradient step above with a fixed diagonal damping λ ; and (iii) **CG-SR**, the same step solved matrix-free by (preconditioned) conjugate gradients with Tikhonov damping and a trust-region cap, which converges faster where S is well conditioned but becomes unstable in the weak-confinement regime where S is nearly singular. Under $|\Psi|^2$ sampling the Adam and SR *fixed points* coincide; they differ in conditioning and in the collocation regime, which is the subject of §7.2.

4.5 Sampling paradigms and the weak-form collocation objective

The pipeline above draws its batches from an analytic proposal rather than a Markov chain, so the choice of *sampling measure* must be stated explicitly: the results treat it as a variable in its own right. Two paradigms appear in this thesis. In *variational Monte Carlo* (VMC) the batch is drawn from $|\Psi_\theta|^2$ itself, usually by a Metropolis chain, so the estimators of the QGT and of the gradient share the very metric they are meant to precondition. In *collocation* the batch is drawn from a fixed analytic proposal $q(\mathbf{R})$ and reweighted to $|\Psi_\theta|^2$ by importance weights $w_i = |\Psi_\theta(\mathbf{R}_i)|^2/q(\mathbf{R}_i)$, so that the gradient training requires no Markov chain. A Metropolis probe is used only outside the gradient loop—for periodic checkpoint selection and for the final energy certification.

For the collocation objective we use a *weak form*: the local energy E_L enters only as a reward in a score-function (REINFORCE) gradient [39],

$$\nabla_\theta \langle E \rangle = 2 \mathbb{E}_{\mathbf{R} \sim |\Psi_\theta|^2} [(E_L(\mathbf{R}) - b) \nabla_\theta \log |\Psi_\theta(\mathbf{R})|], \quad (4.8)$$

with a variance-reducing baseline b (a running mean of E_L), and the Laplacian inside E_L is never differentiated through. This is deliberate. Differentiating the Laplacian through a backflow-shifted determinant is the source of the nodal instability analysed in Appendix C; the weak form keeps the full local-energy signal as a reward while sidestepping that pathology. Two robustness controls stabilise the reward: local energies are clipped to a robust median/MAD location–scale band ($\sim 5\sigma$), and the most extreme rewards are quantile-trimmed before the gradient is formed.

4.5.1 Analytic proposal and importance resampling

The collocation points are drawn from an analytic mixture of isotropic Gaussians centred at the trap minimum and reweighted to $|\Psi_\theta|^2$ [40],

$$q(\mathbf{R}) = \frac{1}{K} \sum_{k=1}^K \mathcal{N}(\mathbf{R}; 0, \sigma_{f,k}^2/\omega I_{Nd}), \quad \sigma_f \in \{0.8, 1.3, 2.0\}, \quad w_i = \frac{|\Psi_\theta(\mathbf{R}_i)|^2}{q(\mathbf{R}_i)}, \quad (4.9)$$

where the $1/\omega$ scaling ties the proposal width to the oscillator length $\ell = 1/\sqrt{\omega}$ and the three components cover core, bulk, and tails. A batch of n_{coll} points is obtained by multinomial resampling on $p_i = w_i / \sum_j w_j$ from $m n_{\text{coll}}$ candidates ($m \approx 8\text{--}16$). Reweighting quality is tracked by the effective sample size $\text{ESS} = (\sum_i w_i)^2 / \sum_i w_i^2$ and, as a heavier-tailed diagnostic, the Pareto $\hat{k}$ of the weights [41]; both degrade when q ceases to overlap $|\Psi_\theta|^2$, the failure mode that bounds the paradigm at large N and weak ω . Three controls manage the tails: weight tempering ($w \rightarrow w^\alpha$, $\alpha < 1$), log-weight clipping at an upper quantile, and ESS-adaptive oversampling that raises m when the achieved ESS falls below a floor. An instability-rollback reverts to the last good checkpoint and decays the learning rate if the energy jumps beyond a threshold.

4.5.2 Physics-informed proposal for the Wigner regime

The origin-centred proposal (4.9) places its mass where a Wigner molecule has almost none: at strong correlation the density collapses onto a shell at the classical ring radius r_0 , and $|\Psi_\theta|^2$ has essentially no support near the origin, so the ESS collapses. We therefore replace q by a proposal built from the Wigner shell itself—a radial Gaussian centred at r_0 (set by balancing confinement against Coulomb repulsion)

times an angular distribution over the ring, so that proposed configurations resemble electrons spread around the ring rather than clustered at the centre. This restores a usable ESS and carries MCMC-free training to $\omega \approx 0.0035$ (Chapter 7, §7.4), far past where the naive proposal collapses; the remaining wall is *angular* crystallisation, analysed there. Difficult confinements are reached by *cascade warm-starting*: parameters converged at one ω initialise the next, following $\omega : 1.0 \rightarrow 0.5 \rightarrow 0.1 \rightarrow 0.01 \rightarrow 0.001$. Whether the two paradigms reach the same *state*, and where they diverge, is one of the questions of Chapter 7.

4.6 Units, scaling, and batch construction

All features are computed in trap units $\tilde{\mathbf{x}} = \sqrt{\omega} \mathbf{x}$ so that length scales and gates (e.g. the short-range gate in the NQS and backflow) behave consistently across ω . The sampler internally widens/narrows Gaussians and shell radii using $a_{\text{ho}} = 1/\sqrt{\omega}$ in a fixed, explicit way, so that the five components cover comparable *dimensionless* regimes for $\omega \in [0.01, 1]$. Batches are i.i.d. by design; we rely on the stratified mixture (with caps, EG-adaptation, and hard-injection) rather than long MCMC chains to traverse configurations, which markedly reduces correlation and simplifies accounting.

4.7 Workflow summary

1. **Sample collocation set** X from the stratified capped-simplex mixture (centre, tails, mixed, shells, dimers), then apply rotation and permutation; optionally inject hard examples.
2. **Compute** E_L using the selected Laplacian backend; trim outliers.
3. **Stage I (residual)**. Minimize $\frac{1}{B} \sum_b (E_L(\tilde{R}_b) - E_{\text{eff}}(\alpha))^2$ with α ramped from 0 to α_{end} ; update mixture/shell weights via EG.
4. **Stage II (SR)**. Form S and g , solve $(S + \lambda I)\Delta\theta = -g$, apply a trust-region step.

This procedure keeps the early dynamics variance-dominated and sampler-aware, then transitions to a geometry-aware natural-gradient refinement. Empirically it yields stable convergence across (N, ω) , including weak traps where shell/dimer coverage and permutation-respecting stratification are essential.

4.8 Statistical uncertainties

Every energy and observable we report carries a statistical error, and because our samples are correlated—within an MCMC chain, and across the score-function updates that shape the final ensemble—a naive standard error over B samples understates it. We therefore estimate the error on a scalar mean $\bar{A} = \frac{1}{B} \sum_i A(\mathbf{R}_i)$ by *blocking*: successive samples are averaged into blocks of growing size and the plateau of the block standard error as the block size exceeds the autocorrelation time is taken as the uncertainty [42, 43]. For the final ground-state energy this is the dominant reported error; a parenthetical digit, e.g. 155.8738(7), denotes one standard error on the last quoted figure.

For quantities that are *nonlinear* functions of sample statistics—participation ratios, effective ranks, PCA cosines, and the coupling diagnostics of Chapter 6—a blocked standard error is not available in closed form, and we use a *block bootstrap* instead: blocks (of the plateau size above) are resampled with replacement, the statistic is recomputed on each resample, and its spread across resamples is the reported error (§5.2.9). Where a number is quoted as a seed average, we additionally report the spread across independent random seeds, which captures optimisation—not just sampling—variability; the two are distinguished in the captions. Reference energies drawn from the literature carry the uncertainties quoted by their sources.

Chapter 5

Analysis

This chapter describes *what* we measure on fresh $|\Psi_\theta|^2$ samples, *how* we compute the diagnostics, and *why* each quantity is informative. Results and numbers are presented in Chapter 6.

5.1 Structural diagnostics and Wigner markers

Unless stated otherwise, curves are accumulated in *trap units* $\mathbf{y} = \sqrt{\omega} \mathbf{x}$ to compare shapes across ω , while all headline distances are reported in *Bohr* (physical units $\mathbf{x}$). Production chains are thinned; we split samples into halves for stability checks (acceptance window, JSD between halves, and reproducibility under restarts). Using trap units to compare shapes across confinements is standard in parabolic dots; see the overview in [10].

5.1.1 Pair distribution and radial statistics

We estimate the pair distribution function $g(r)$ via a shell-area corrected histogram in trap units,

$$g(r_\alpha) = \frac{A_{\text{eff}}}{N(N-1)} \frac{\langle \sum_{i < j} \mathbf{1}\{r_{ij} \in [r_\alpha^-, r_\alpha^+]\} \rangle}{2\pi r_\alpha \Delta r_\alpha}, \quad r_{ij} = \|\mathbf{y}_i - \mathbf{y}_j\|_2, \quad (5.1)$$

with $A_{\text{eff}} = \pi R_{0.95}^2$ and $R_{0.95}$ the 95% quantile of single-particle radii in trap units. This $g(r)$ -based structural analysis follows established practice in mesoscopic 2D electron systems and quantum dots [7, 10, 12]. We also form the normalised *radial probability* $P(r) \propto r^{d-1}g(r)$ and report the mode r_{mode} , mean $\bar{r}$, standard deviation σ_r , and FWHM (all in Bohr). A compact width-to-scale marker is

$$\gamma = \sigma_r / r_{\text{mode}}, \quad (5.2)$$

which decreases as correlations strengthen (narrower distribution around a larger typical radius), consistent with the liquid→Wigner crossover picture [7, 12].

5.1.2 Automatic shell detection and topologies

For each frame we sort the single-particle radii $s_1 \leq \dots \leq s_N$ (trap units), compute gaps $g_k = s_{k+1} - s_k$, and flag the frame as *two-shell* if the largest gap exceeds a robust threshold:

$$\max_k g_k \geq \tau \text{median}(g_k), \quad \tau \in [2.5, 3.5] \text{ (default 3.0)}. \quad (5.3)$$

The cut radius is the midpoint of the maximal gap; the inner multiplicity n_{in} defines the topology $(n_{\text{in}}, N - n_{\text{in}})$. Shelling and discrete ring occupancies are well known in parabolic Coulomb clusters and quantum dots [8–10]; our detector provides a reproducible, data-driven assignment in that spirit. We report: (i) the fraction of two-shell frames, (ii) the histogram over n_{in} among two-shell frames, and (iii) for selected topologies (e.g. (1, 5) or (3, 9)) the fraction among *all* frames.

Shell-resolved reconstruction of $g(r)$. On two-shell frames we decompose pairs into II, IO, and OO and accumulate histograms $g_{\text{II}}, g_{\text{IO}}, g_{\text{OO}}$. With empirical weights $w_{\text{II}}, w_{\text{IO}}, w_{\text{OO}}$ (combinatorial counts) we reconstruct $\tilde{g}(r) = w_{\text{II}}g_{\text{II}} + w_{\text{IO}}g_{\text{IO}} + w_{\text{OO}}g_{\text{OO}}$ and report $\cos \angle(g, \tilde{g})$. Values ≈ 1 indicate that the observed $g(r)$ shape is explained by shell geometry rather than sampling artifacts, in line with shell-based descriptions in [8–10].

5.1.3 Orientational order and “ring stiffness”

On each detected ring (inner or outer) we centre the mean angle to remove global rotations and define angles $\{\phi_k\}_{k=1}^n$. Bond-orientational order is

$$\Phi_m = \frac{1}{n} \sum_{k=1}^n e^{im\phi_k}, \quad m = n, \quad (5.4)$$

so that $|\Phi_m| \in [0, 1]$ measures n -fold orientational order (e.g. $m=5$ for (1, 5), $m=9$ for the outer ring of (3, 9)), analogous to standard bond-orientational order parameters used in 2D Coulomb/Yukawa crystals [11]. As a complementary, dimensionless *angular Lindemann* number we use

$$\lambda_\phi = \frac{\text{std}(\Delta\phi)}{\mathbb{E}[\Delta\phi]}, \quad \Delta\phi = \text{nearest-neighbour angular spacing on the ring.} \quad (5.5)$$

Smaller λ_ϕ indicates a stiffer (more Wigner-like) ring. This is a Lindemann-type orientational metric tailored to finite rings, motivated by the use of Lindemann criteria to diagnose intershell rotational (angular) melting in mesoscopic Coulomb clusters and dots [9, 12].

5.1.4 Scaling check for $N = 2$

To connect with the classical strong-coupling limit we fit

$$r_{\text{mode}}(\omega) = C\omega^\alpha \quad (5.6)$$

by linear regression in $(\log \omega, \log r_{\text{mode}})$ and compare the slope with the classical reference $\alpha_{\text{cl}} = -2/3$ (two charges in a harmonic trap at strong coupling; see, e.g., the reviews in [10]). Approach toward α_{cl} as $\omega \downarrow$ is one of our Wigner crossover indicators, consistent with the crossover phenomenology reported in [7, 12].

5.1.5 Reporting and stability checks

For all scalars and histograms we report standard errors calculated over thinned blocks or frames. We verify: (i) acceptance in the target window, (ii) Jensen–Shannon divergence and ℓ_2 distance between halves of production (stationarity), (iii) robustness of topology fractions and $|\Phi_m|$ under $\tau \in [2.5, 3.5]$ and mild pre-smoothing of radii, and (iv) reproducibility of $g(r)$ and $P(r)$ under independent sampler restarts. All diagnostics operate directly on saved bundles $\{\mathbf{X}_{\text{Bohr}}, r, g(r)\}$.

5.2 Representation analysis

We probe what the networks learn by evaluating fixed, trained models on fresh $|\Psi_\theta|^2$ samples. Unless stated otherwise, features are computed in *trap units* $\mathbf{y} = \sqrt{\omega} \mathbf{x}$ to match the correlator’s internal scaling. All estimates use thinned MCMC frames, split into two equal blocks A and B ; PCA fits on A and reports on B (and vice versa) to avoid circularity. We use `float64` throughout.

Feature taps. Let $z_\theta \in \mathbb{R}^D$ denote the concatenated correlator representation

$$z_\theta = [\bar{\phi} \mid \bar{\psi} \mid \mathbf{g}],$$

where $\bar{\phi}$ are per-particle summaries, $\bar{\psi}$ are pairwise summaries, and $\mathbf{g}$ are global/extras. The residual head is linear:

$$f_{\text{base}}(z) = \rho_\theta(z) = w^\top z + b, \quad f = f_{\text{base}} + f_{\text{cusp}}. \quad (5.7)$$

Backflow returns a displacement field $\Delta x \in \mathbb{R}^{N \times d}$ that enters only the Slater determinant; the correlator and cusp are evaluated on the unshifted $\tilde{R}$ (§3.5). The coupling diagnostics below additionally probe the correlator on the shifted coordinates as a counterfactual sensitivity.

5.2.1 Preprocessing and PCA

Given a sample matrix $Z \in \mathbb{R}^{K \times D}$ (rows are frames), we standardize each column using means and standard deviations estimated on block A :

$$\tilde{Z} = (Z - \mu_A) \odot \sigma_A^{-1}.$$

We perform PCA on $\tilde{Z}_A$ (SVD of the covariance) to obtain orthonormal loadings $U = [u_1, \dots, u_D]$. Scores on B are $S_B = \tilde{Z}_B U$. Unless noted, all quantities reported below are computed on the held-out block.

5.2.2 Entropy effective rank

Let $\{\lambda_i\}_{i=1}^D$ be the eigenvalues of $\text{Cov}(\tilde{Z})$ (on the held-out block). With $p_i = \lambda_i / \sum_j \lambda_j$, the *entropy effective rank* is

$$r_{\text{eff}}(Z) = \exp\left(-\sum_{i=1}^D p_i \log p_i\right),$$

which equals 1 for perfectly one-dimensional representations and increases with spectral spread.

5.2.3 PC ablations of the head

To test how many latent axes the head truly uses, we project $\tilde{Z}$ onto the top- k PCs:

$$\Pi_k = \sum_{i=1}^k u_i u_i^\top, \quad z^{(k)} = \Pi_k \tilde{z},$$

and re-evaluate the trained linear head *without refitting*: $\hat{f}^{(k)} = \rho_\theta(z^{(k)})$. We report a *normalised* error on block B ,

$$\text{rel-MAE}(k) = \frac{\mathbb{E}_B |\hat{f}^{(k)} - f_{\text{base}}|}{\text{std}_B(f_{\text{base}})},$$

together with the learning curve in k .

5.2.4 PC1 block power decomposition

To attribute the leading axis to branches, we partition u_1 conformably with $z = [\bar{\phi} | \bar{\psi} | \mathbf{g}]$ and report

$$\text{power}(\bar{\phi}) = \frac{\|u_{1,\bar{\phi}}\|_2^2}{\|u_1\|_2^2}, \quad \text{power}(\bar{\psi}) = \frac{\|u_{1,\bar{\psi}}\|_2^2}{\|u_1\|_2^2}, \quad \text{power}(\mathbf{g}) = \frac{\|u_{1,\mathbf{g}}\|_2^2}{\|u_1\|_2^2}.$$

Because PCA is performed on standardized features, these shares are comparable across blocks.

5.2.5 Linear probes from PCs to coarse physics

We regress interpretable summaries per configuration on the first k PC scores $S^{(k)}$ using ridge regression (tiny penalty $\alpha = 10^{-6}$):

$$y \approx S^{(k)} \beta, \quad y \in \{\bar{r}, \text{Var}(r), \text{Pr}(r < r_0), \text{shell contrast}\}.$$

We report R^2 on the held-out block; r_0 is a fixed near-contact cutoff in trap units.

5.2.6 Near-field gradient concentration (residual head)

For each frame we compute the minimum pair distance $r_{\min}$ (trap units) and the gradient of the residual head with respect to particle coordinates, $\nabla f_{\text{base}}(\mathbf{y})$. The concentration at quantile q is

$$\text{share}(q) = \frac{\mathbb{E}[\|\nabla f_{\text{base}}\|_2^2 \mathbf{1}\{r_{\min} \leq Q_q\}]}{\mathbb{E}[\|\nabla f_{\text{base}}\|_2^2]},$$

where Q_q is the q -quantile of $r_{\min}$. Values $\gg 1$ indicate that residual stiffness is concentrated near contact (the analytic cusp is insufficient); values ≈ 1 indicate that the cusp handles most short-range structure.

5.2.7 Backflow geometry and energy-locality shares

Field spectrum. We vectorize the backflow displacement per frame, $\delta = \text{vec}(\Delta x) \in \mathbb{R}^{N^d}$, standardize across frames, and compute $r_{\text{eff}}(\Delta x)$ via Sec. 5.2.2.

Where backflow acts. On the same thinned frames we evaluate local energies with and without backflow, $E_{\text{loc}}^{\text{BF}}$ and $E_{\text{loc}}^{\text{noBF}}$, using the *same* coordinates. We form the signed correction $\Delta E_{\text{loc}} = E_{\text{loc}}^{\text{BF}} - E_{\text{loc}}^{\text{noBF}}$ and report near-field shares at quantile q of $r_{\min}$:

$$\text{NF-}\Delta E(q) = \frac{\mathbb{E}[|\Delta E_{\text{loc}}| \mathbf{1}\{r_{\min} \leq Q_q\}]}{\mathbb{E}[|\Delta E_{\text{loc}}|]}.$$

(Using absolute values isolates magnitude; signed averages are also available.)

5.2.8 Alignment of correlator manifolds (noBF vs BF)

We collect two standardized feature matrices $\tilde{Z}^{\text{noBF}}$ and $\tilde{Z}^{\text{BF}}$ on identical frames (each standardized by its own μ, σ), compute their PC1 loadings u_1^{noBF} and u_1^{BF} , and report the cosine similarity

$$\cos(u_1^{\text{noBF}}, u_1^{\text{BF}}) = \frac{\langle u_1^{\text{noBF}}, u_1^{\text{BF}} \rangle}{\|u_1^{\text{noBF}}\|_2 \|u_1^{\text{BF}}\|_2}.$$

Cosines near 1 indicate that backflow acts as a structured reweighting along the same dominant axis, rather than rotating the correlator’s manifold.

5.2.9 Uncertainty and stability

For scalars we report standard errors over blocks. For PCA-based quantities and cosines we use a block bootstrap (resampling blocks with replacement) to obtain 68%/95% CIs. Stability checks include: (i) swapping train/held-out blocks for PCA, (ii) varying the near-field quantiles $q \in \{1, 5, 10\}\%$, (iii) re-running analyses on independent sampler restarts.

Reproducibility package. All diagnostics consume the saved bundles $\{\mathbf{X}_{\text{Bohr}}, r, g(r)\}$ and the model snapshot, and emit JSON/CSV tables with r_{eff} , rel-MAE(k) curves, block-power shares, R^2 probes, near-field shares, $r_{\text{eff}}(\Delta x)$, and noBF/BF cosines. As in Sec. 5.1, we log JSD/ ℓ_2 splits to confirm stationarity.

5.3 Tangent-kernel diagnostics

The representation diagnostics above describe the ansatz’s *features*; a second family, developed with the geometry of §2.2.5, describes its *tangent space*, and it is these that carry the architecture, optimiser, and paradigm comparisons of Chapter 7. From a batch we assemble the centred score matrix $O_{ki} = \partial_{\theta_i} \log |\Psi_{\theta}(\mathbf{x}_k)|$ and form the quantum geometric tensor $S = O^{\top} O / B$ (or, when $B < P$, the tangent kernel $K = O O^{\top}$, whose nonzero spectrum equals that of S up to the overall factor B). From the spectrum $\{\lambda_a\}$ we report the effective dimension $d_{\text{eff}} = (\sum_a \lambda_a)^2 / \sum_a \lambda_a^2$ and the condition number $\kappa(S) = \lambda_{\max} / \lambda_{\min}$. Cross-architecture comparisons of d_{eff} are made on a *common probe set*—all models scored on the same configurations, not on each model’s own $|\Psi|^2$ —so that the comparison reflects the

ansatz and not its sampling measure, and we report only participation ratios that have converged in the batch size.

To ask whether two trained models are the *same* state rather than merely equal in energy, we use a symmetric, importance-sampled squared overlap,

$$\mathcal{S}^2(A, B) = \langle \Psi_B / \Psi_A \rangle_{|\Psi_A|^2} \langle \Psi_A / \Psi_B \rangle_{|\Psi_B|^2},$$

each factor evaluated by a log-sum-exp mean over samples of the corresponding state. It equals 1 for identical states, and on the analytic two-electron ground state the trained ansatz reproduces $\mathcal{S}^2 = 1.000000$, which anchors the tooling.

One convention must be stated, because the ratio Ψ_B / Ψ_A is signed for a fermionic state. We evaluate the ratios from $\log |\Psi|$, so the estimator is strictly a *modulus* overlap, $\langle |\Psi_B / \Psi_A| \rangle \langle |\Psi_A / \Psi_B| \rangle$. For the comparisons made here this is an accurate proxy for the signed overlap: both states share the same Slater reference, and the correlator and cusp enter as a strictly positive exponential, so the sign of Ψ is fixed by the determinant and the two states agree in sign except on the measure-suppressed set where their backflows carry a configuration across a node differently. Where the two states genuinely diverge the modulus overlap is an upper bound on the signed one, so a reported $\mathcal{S}^2 \rightarrow 0$ is if anything conservative. These quantities—together with the energetic price of ablating a component—are invariant to how correlation is divided between the Jastrow and the backflow, and it is on them, rather than on the gauge-dependent component ranks of the previous section, that the architectural conclusions of Chapter 7 rest.

Part IV

Results

Chapter 6

Quantum dots: energies, representations, and Wigner molecules

The least initial deviation from the truth is multiplied later a thousandfold.

ARISTOTLE

On the Heavens, Book I, Chapter 5, 271b8–10

This chapter is about what the trained wavefunction *is*, and what it lets us see. It is one half of a pair. Here we ask how accurately the ansatz of Chapter 3 solves the two-dimensional harmonic quantum dot, what its internal representation looks like, and what it reveals about the electrons themselves; the companion chapter (Chapter 7) asks the complementary question of *why* the architecture, the optimiser, and the training paradigm behave as they do. The two chapters share one object and one aim: not merely to compute the ground state, but to understand it.

The quantum dot is an unforgiving benchmark, and it is worth being clear about why. The Coulomb interaction is singular where two electrons meet; the kinetic energy carries second derivatives that magnify any error in the wavefunction’s curvature; and the confinement ω sweeps the physics across three decades of correlation strength—from a compact, kinetic-energy-dominated droplet at $\omega = 1$ to a dilute Wigner molecule at $\omega = 10^{-3}$, in which the electrons settle several oscillator lengths apart and the Coulomb repulsion, not the trap, dictates the structure. We study $N \in \{2, 6, 12, 20\}$ electrons across $\omega \in \{1.0, 0.5, 0.1, 0.01, 0.001\}$.

Three questions organise the chapter. *First, how accurate is the method?* Section 6.1 benchmarks the variational energies against published reference values wherever one exists—diffusion Monte Carlo (DMC) for $N \leq 12$, and the tabulated $N = 20$ energies of Haas [45], which are not DMC and are treated as the weaker benchmark they are. Those references reach down to $\omega = 0.1$ for every N , and for $N=2$ to $\omega = 0.01$; below that no published value exists for $N \geq 6$, so the strongly correlated energies rest on internal variational consistency rather than on an external ground truth. *Second, what does the network learn?* Section 6.2 dissects the trained wavefunction’s internal representation—a strikingly low-dimensional correlator whose *content*, but not its dimensionality, reorganises across regimes, alongside a backflow field that shares the work in a regime-dependent way. One caveat, carried over from the companion chapter and stated here at the outset, governs how these internal quantities may be read: several of them are gauge-like, fixed by the training route rather than by the state, and we take care to separate those from the findings that are invariant. *Third, can the learned wavefunction serve as a scientific instrument?* Section 6.3 turns the trained models on the physics itself, mapping the Wigner-molecule crossover in real space—shell topologies, angular order, and melting—from $N=2$ up to $N=20$.

6.1 Training protocol and energy overview

Unless otherwise stated, we first apply residual-based pretraining to obtain a well-conditioned initialiser, and then run a Stochastic Reconfiguration (SR) tail to refine the last digits of the energy. In the residual phase, the network initially chooses its own target energy as a stabilizer, before we switch to the DMC

reference energy (and optionally add a variance penalty). The SR phase then uses on the order of 10^3 – 3×10^3 iterations with batches of $\sim 3 \times 10^3$ samples per iteration. Throughout this section—and wherever else the thesis says “SR tail” without further qualification—SR is applied to *variational Monte Carlo with $|\Psi|^2$ sampling*. This matters because the same optimiser is used again in Section 6.4 on the collocation objective, where the sampling measure differs and SR consequently behaves differently: here it is a refinement that buys the last digits, there it is what makes the optimisation viable at all (Chapter 7, Q2).

We compare two backflow architectures throughout. Both condition each particle’s displacement on the interparticle coordinates, but they do so differently: the *conventional* BACKFLOWNET applies a single round of pairwise messages aggregated per particle, whereas the copresheaf *message-passing* network (CTNN), realised as the CTNNBACKFLOWNET, maintains explicit node and edge features and transports information between them through learned maps (a single copresheaf round) before the displacement is read out; the contrast between them is the subject of the backflow analysis in Chapter 7. The architecture size is identical for all N : the base PINN correlator has ≈ 54 k parameters, the conventional BackflowNet adds ≈ 83 k (total PINN+BF ≈ 137 k), and the CTNNBackflowNet adds ≈ 182 k (total PINN+CTNN ≈ 236 k). All components are fully trainable; the parameter count is independent of N because particle interactions enter through permutation-equivariant features rather than per-particle weights.

6.1.1 Energies

The tables in this section are the thesis’s only contact with an external ground truth. Two things matter in them, and neither is the raw energy. The first is *where the agreement holds*: a method that matches DMC at $\omega = 1$ and drifts at $\omega = 0.1$ is a different object from one that holds across the whole benchmarked range, because the physics changes character between those points while the architecture does not. The second is *where the tables stop*. Below $\omega = 0.1$ there is no DMC reference for $N \geq 6$, and the numbers reported there are variational predictions rather than validated ones. That boundary is the single most important feature of what follows, and we have marked it everywhere it occurs rather than letting a uniformly formatted table imply a uniform level of evidence.

Table 6.1 begins with the easiest case, and its purpose is diagnostic rather than competitive: two electrons are few enough that the architecture can be tested nearly in isolation. It reports two-electron energies from *residual-only* training (PINN and PINN+BF). The result to notice is a negative one—for $N=2$, backflow is *not* required to reach DMC accuracy; it slightly reduces the variance and smooths convergence, and nothing more. The Slater–Jastrow correlator already places the nodes well enough at this size. This matters because it establishes a baseline against which the backflow’s later contribution, at larger N and weaker traps, can be read as a genuine effect rather than as an artefact of having added parameters.

Table 6.1: Ground-state energies (Hartree) for two electrons in a harmonic trap using *residual-only* training—that is, *without* the SR tail. To be read against Table 6.2, which reports the same systems after SR refinement: the comparison isolates what pretraining alone achieves, and the residual bias at $\omega = 0.01$ visible here is the one the SR tail removes. DMC references from [44]. Parentheses denote 1σ on the last digits.

ω	PINN+BF	PINN	DMC (Ref.)
1.00	2.99998(3)	2.999940(5)	3.00000(1)
0.50	1.65975(2)	1.65979(3)	1.65977(1)
0.10	0.440795(3)	0.440833(1)	0.44079(1)
0.01	0.0740724(6)	0.0741679(6)	0.073839(2)

Table 6.2 compares diffusion Monte Carlo (DMC) references with our *final* PINN+BF energies after residual pretraining and an SR tail. From the ultra-shallow two-electron case ($\omega=0.01$) to the many-electron systems, absolute deviations are small; for $N=2$ they lie within DMC uncertainties, and for $N \in \{6, 12\}$ the relative deviations are typically in the range 0.01–0.08% (see also Fig. 6.1).

Summary. The pattern is clean wherever we can check it. For $N=2$ the ansatz reaches DMC accuracy at every ω for which a DMC number exists, and for $N \in \{6, 12\}$ the relative deviation stays within roughly 0.01–0.08% across the benchmarked range ($\omega \geq 0.1$; Fig. 6.1). Below $\omega = 0.1$ there is no DMC reference for $N \geq 6$, and we make no claim of absolute accuracy there; what we can say is that the energies join

Table 6.2: Residual-based pretraining + SR tail. Reference DMC and our PINN+BF and PINN+CTNN ground-state energies (Hartree) for selected (N, ω) . The reference column collects the best available benchmark for each size: DMC from [44] (whose study spans $N \leq 13$) for $N \leq 12$, and the reference energies tabulated in the master’s thesis of Daniel Haas [45] for $N = 20$, which lies beyond the range of [44]. Parentheses denote 1σ on the last digit(s); the %err columns are relative deviations from the reference. A dash in the reference column marks (N, ω) for which no published DMC value exists (all $\omega = 0.001$, and $\omega = 0.01$ for $N \geq 6$); the relative-error columns are correspondingly blank there, as no external ground truth is available to score against.

N	ω	Reference (DMC)	PINN+BF	% err	PINN+CTNN	% err
2	0.001	—	0.0137948(8)	—	0.013778(1)	—
	0.01	0.073839(2)	0.073838(1)	−0.0014	0.0738382(5)	−0.0014
	0.10	0.44079(1)	0.44079(1)	+0.0000	0.440796(4)	−0.0000
	0.50	1.65977(1)	1.65976(2)	−0.0006	1.65976(1)	−0.0006
	1.00	3.00000	2.99999(1)	−0.0003	3.00000(2)	−0.0000
6	0.001	—	0.140913(1)	—	0.140832(1)	—
	0.01	—	0.69125(2)	—	0.69036(1)	—
	0.10	3.55385(5)	3.5549(1)	+0.0295	3.55388(5)	+0.0008
	0.50	11.78484(6)	11.7895(4)	+0.0395	11.7847(2)	−0.0000
	1.00	20.15932(8)	20.1610(6)	+0.0083	20.1585(3)	−0.0041
12	0.001	—	0.515823(3)	—	0.515365(4)	—
	0.01	—	2.48620(5)	—	2.47363(4)	—
	0.10	12.26984(8)	12.2731(2)	+0.0266	12.2743(2)	+0.0364
	0.50	39.1596(1)	39.1786(8)	+0.0485	39.1604(3)	+0.0018
	1.00	65.7001(1)	65.717(1)	+0.0257	65.69556(5)	−0.0069
20	0.001	—	—	—	1.293033(6)	—
	0.01	—	—	—	6.14645(5)	—
	0.10	29.9779(1) ^H	—	—	29.9888(2)	+0.0364
	0.50	93.8752(1) ^H	—	—	93.8789(5)	+0.0040
	1.00	155.8822(1) ^H	—	—	155.8738(7)	−0.0024

Reference-energy provenance: values marked ^H are from the master’s thesis of Daniel Haas [45] ($N = 20$); all other reference energies are diffusion Monte Carlo from Pederiva *et al.* [44] ($N \leq 12$, whose study spans $N \leq 13$). Dashes mark (N, ω) with no published value.




Figure 6.1: Relative deviation of PINN+BF ground-state energies from DMC references as a function of trap frequency ω for the cases where DMC data are available. For $N=2$ the deviations are at or below the DMC error bars ($\sim 10^{-3}$ %), while for $N=6$ and $N=12$ the errors remain in the range 2.5×10^{-2} – 5×10^{-2} % across ω , with no sign of deterioration in the shallow- or tight-trap limits.

smoothly onto the benchmarked regime and, in the deep-Wigner limit, follow the physical energy scaling derived in §6.3.3. The efficiency is as notable as the accuracy: the SR tail converges in 10^3 – 3×10^3 iterations of $\sim 3 \times 10^3$ samples each, a modest optimisation budget for a fully many-body wavefunction. We attribute this stability to a small set of conditioning choices— trap-unit scaling, soft-core pair features with $ds/dr \rightarrow 0$ at coalescence, and an explicit analytic cusp—each of which keeps the gradients and the Laplacian numerically tame from the shallowest trap to the tightest.

6.2 What the networks learn

We probe the internal representations of the CTNN-augmented wavefunction across $N \in \{2, 6, 12, 20\}$ and $\omega \in \{10^{-3}, 10^{-2}, 0.1, 0.5, 1.0\}$. The analysis targets three objects: the neural correlator W_θ (the “correlator head” of Chapter 3, whose input concatenates the orbital, pair, and auxiliary branches $\bar{\phi} \parallel \bar{\psi} \parallel \mathbf{g}$), the *backflow displacement field* $\Delta \mathbf{x}$ produced by the CTNN, and the *coupling* between these two components. All diagnostics are evaluated on samples drawn from $|\Psi|^2$.

Three descriptive regularities organise the trained solutions examined here. They are reported as observed properties of these models, not as settled architecture-level facts: as the caveat below explains, several are gauge-like. First, the correlator operates on a manifold of remarkably low dimension ($r_{\text{eff}} \lesssim 3$), whose *content*—but not dimensionality—reorganises across regimes via a $\phi \leftrightarrow \psi$ transition in the leading principal axis (§6.2.1). Second, the CTNN displacement field is a high-rank, full-dimensional correction (rank $\approx 2N$) that nonetheless leaves the correlator manifold nearly unchanged, acting as a complementary rather than redundant channel (§6.2.2). Third, PINN and CTNN switch from operating independently at strong correlation to cooperating tightly at moderate-to-weak correlation, with a sharp transition near $\omega \approx 0.1$ that coincides with the physical crossover in $\Gamma = V_{\text{int}}/T$ (§6.2.3).

How these quantities should be read. One caveat frames everything that follows. The Jastrow and the backflow are partial substitutes, so how the labour is divided between them is a nearly flat direction of the loss, settled by the training route rather than by the state. Quantities living on a single component—the backflow’s rank and displacement magnitude, the per-branch loadings, the coupling between the two

Table 6.3: Correlator geometry (CTNN). r_{eff} : entropy-based effective rank of Z . $|\rho|$: absolute correlation between head output and PC1 reconstruction. $\phi/\psi/\mathbf{g}$: fraction of PC1 variance from the orbital, pair, and auxiliary branches. The ψ -dominated entries ($\psi > 0.5$) are shown in bold.

N	ω	r_{eff}	$ \rho $	ϕ	ψ	$\mathbf{g}$
2	0.001	1.00	1.00	0.00	1.00	0.00
2	0.01	1.04	0.66	0.00	1.00	0.00
2	0.1	1.31	0.99	0.00	0.53	0.46
2	0.5	1.25	1.00	0.00	0.58	0.42
2	1.0	1.53	0.98	0.00	0.35	0.65
6	0.001	1.52	0.97	0.01	0.12	0.87
6	0.01	1.38	0.99	0.05	0.44	0.50
6	0.1	2.73	0.94	0.66	0.21	0.14
6	0.5	2.69	0.98	0.15	0.63	0.22
6	1.0	2.69	0.99	0.03	0.62	0.35
12	0.001	1.66	0.96	0.54	0.13	0.33
12	0.01	1.54	0.97	0.43	0.09	0.48
12	0.1	3.08	0.99	0.12	0.63	0.26
12	0.5	2.32	0.99	0.12	0.69	0.19
12	1.0	1.76	0.99	0.08	0.73	0.19
20	0.001	2.59	0.99	0.18	0.74	0.08
20	0.01	1.09	0.79	0.04	0.94	0.01
20	0.1	2.20	0.97	0.36	0.44	0.20
20	0.5	2.30	1.00	0.17	0.70	0.13
20	1.0	2.32	0.99	0.18	0.71	0.11

networks—are therefore *gauge-like*; §7.5 establishes why, and the invariants are energy, state overlap, and ablation. The anatomy below is not weakened by this: it is a faithful dissection of one high-quality solution, and the regularities it shows are the more telling for being ones the network was free *not* to adopt.

6.2.1 Correlator geometry: a compact manifold that reorganises

We characterise the correlator feature space $Z = \bar{\phi} \|\bar{\psi}\| \mathbf{g}$ by its entropy-based effective rank $r_{\text{eff}}(Z)$, the correlation $|\rho(\text{head}, \text{PC1})|$ between the head output and its first principal component, and the power of PC1 decomposed across architectural branches. Table 6.3 reports these quantities for all CTNN configurations.

The manifold is low-dimensional everywhere. The effective rank stays below 3.1 across the entire grid, and the head output is almost perfectly reconstructed from PC1 alone ($|\rho| > 0.94$ for all $N \geq 6$ and all ω). This holds even at strong correlation where the real-space density develops pronounced shell structure—the correlator does not respond by expanding its latent dimensionality. The modest peak at $N=12$, $\omega=0.1$ ($r_{\text{eff}} = 3.08$) coincides with the intermediate-correlation regime where neither the shell picture nor the crystalline picture provides a clean single-mode description.

A $\phi \leftrightarrow \psi$ transition in the leading axis. The more interesting variation is in *which branch dominates PC1*. At moderate-to-weak correlation ($\omega \geq 0.1$, $N \geq 6$), PC1 is predominantly ψ -loaded: the pair branch carries 60–73% of the variance, meaning the correlator’s primary degree of freedom tracks pair correlations. At strong correlation ($\omega \leq 0.01$, $N \geq 12$), the loading shifts toward ϕ (orbital) or $\mathbf{g}$ (auxiliary) branches, with ψ dropping below 15%. The transition is sharpest at $N=12$, where ψ goes from 0.09 at $\omega=0.01$ to 0.63 at $\omega=0.1$.

The physical interpretation is direct. When interactions are weak relative to the trap ($\omega \gtrsim 0.1$), the non-interacting shell structure provides a good zeroth-order description, and the correlator’s primary role is refining pair correlations on top of that. When interactions dominate ($\omega \lesssim 0.01$), the leading correlator coordinate shifts away from pair features toward the single-particle/orbital-derived and auxiliary branches: the network reorganises the features through which it represents the state becomes the dominant variational degree of freedom. The network discovers this distinction autonomously: regime changes

Table 6.4: CTNN backflow geometry. $\|\Delta\mathbf{x}\|/\ell$: mean displacement magnitude normalised by harmonic length. Rank: effective rank of the displacement covariance. $2N-2$: maximum possible rank after centre-of-mass projection. CKA: linear centred kernel alignment between correlator features with and without backflow (1.0 = identical feature manifolds).

N	ω	$\ \Delta\mathbf{x}\ /\ell$	Rank	$2N-2$	CKA
2	0.001	0.003	1.9	2	1.000
2	1.0	0.029	2.0	2	1.000
6	0.001	0.017	10.0	10	0.999
6	0.01	0.035	9.9	10	0.998
6	0.1	0.096	10.0	10	0.995
6	0.5	0.093	10.0	10	0.991
6	1.0	0.089	10.0	10	0.979
12	0.001	0.017	21.9	22	1.000
12	0.01	0.030	21.9	22	0.998
12	0.1	0.096	21.9	22	0.993
12	0.5	0.096	21.9	22	0.993
12	1.0	0.091	21.9	22	0.995
20	0.001	0.023	37.7	38	0.999
20	0.01	0.046	37.4	38	1.000
20	0.1	0.075	37.8	38	0.997
20	0.5	0.099	37.6	38	0.990
20	1.0	0.099	37.6	38	0.991

appear as a rotation of a low-dimensional axis across branches, not as a growth of latent dimensionality.

6.2.2 Backflow: a high-rank field with a low-rank footprint

Table 6.4 characterises the CTNN displacement field. We report the normalised displacement magnitude $\|\Delta\mathbf{x}\|/\ell$, the effective rank of $\Delta\mathbf{x}$ (viewed as a $B \times 2N$ matrix of flattened displacements), and the CKA similarity between the correlator features with and without backflow.

Displacement magnitude scales with ω but saturates near $\|\Delta\mathbf{x}\|/\ell \sim 0.1$. At strong correlation ($\omega \leq 0.01$), displacements are small relative to the harmonic length, $\|\Delta\mathbf{x}\|/\ell \sim 0.02$ – 0.05 . As the trap tightens and the system enters the moderate-to-weak correlation regime, the magnitude grows and saturates at ~ 8 – 10% of ℓ for $\omega \gtrsim 0.1$. The CTNN never produces large displacements—its tanh output bound and copresheaf message-passing structure act as implicit regularisers, in contrast to the conventional BackflowNet which can exhibit displacements comparable to the system scale in poorly converged regimes (e.g. $\|\Delta\mathbf{x}\|/\ell \sim 0.95$ for $N=6$, $\omega=0.001$ with BackflowNet).

Full-rank displacements, near-unity CKA. The effective rank of the displacement field is consistently $\approx 2N - 2$, saturating the available spatial degrees of freedom modulo centre-of-mass conservation (confirmed by COM shifts of order 10^{-14} – 10^{-17}). The backflow transformation is not a low-rank collective mode such as a breathing or rigid rotation; the displacement field spans essentially all COM-free spatial directions, so each particle receives a correction conditioned on the full configuration. This full-rank behaviour is a property of the *message-passing* backflow specifically, and it is exactly the point at which the companion chapter finds the two architectures part ways: toward the Wigner crystal a *conventional*, single-round-message backflow collapses to a single collective mode (rank $\rightarrow 1$) and its energy degrades—its one round of pairwise messages failing to sustain the relative displacement the crystal demands—whereas the CTNN retains its full rank because its copresheaf edge-transport does sustain it (Chapter 7, §7.1.2). The high rank reported here is thus not incidental—it is the structural signature of the mechanism that makes message passing worth its cost.

Yet the CKA between correlator features with and without backflow exceeds 0.99 in nearly all cases (and exceeds 0.979 everywhere). The correlator manifold is almost unperturbed by the displacement field. This establishes a clean separation of concerns: the CTNN’s action leaves the correlator’s feature geometry almost unchanged, providing configuration-dependent corrections that the globally organised correlator does not and need not represent.

Table 6.5: PINN–CTNN coupling diagnostics (CTNN ansatz). $\langle \cos \rangle$: mean cosine between ∇f and $\Delta \mathbf{x}$. Aligned: fraction of configs with $\nabla f \cdot \Delta \mathbf{x} > 0$. $|\delta f|/\sigma_f$: BF-induced change in PINN output relative to its standard deviation. $\rightarrow \text{NN}$: fraction of particles displaced toward their nearest neighbour (0.5 = isotropic). Only $N \geq 6$ shown; $N=2$ displacements are near-zero for most ω .

N	ω	$\langle \cos \rangle$	Aligned	$ \delta f /\sigma_f$	$\rightarrow \text{NN}$
6	0.001	0.14	0.63	0.03	0.43
6	0.01	0.18	0.77	0.05	0.28
6	0.1	0.49	0.99	0.25	0.16
6	0.5	0.62	1.00	0.33	0.19
6	1.0	0.77	1.00	0.41	0.18
12	0.001	−0.48	0.01	0.03	0.61
12	0.01	0.22	0.82	0.05	0.29
12	0.1	0.56	1.00	0.39	0.12
12	0.5	0.65	1.00	0.49	0.20
12	1.0	0.75	1.00	0.55	0.21
20	0.001	0.32	0.97	0.06	0.44
20	0.01	0.22	0.89	0.07	0.45
20	0.1	0.39	1.00	0.25	0.17
20	0.5	0.65	1.00	0.58	0.22
20	1.0	0.78	1.00	0.69	0.25

6.2.3 PINN–CTNN coupling: from independence to cooperation

The previous subsection established that backflow leaves the correlator manifold intact. But are the PINN and CTNN *functionally* coupled—does the CTNN’s correction depend on what the PINN computes? We measure this through three diagnostics: (i) the mean cosine similarity $\langle \cos(\nabla_{\mathbf{x}} f, \Delta \mathbf{x}) \rangle$ between the PINN’s gradient in configuration space and the CTNN displacement (gradient alignment implies the CTNN pushes along directions the PINN is sensitive to); (ii) the fraction of configurations where $\Delta \mathbf{x}$ is aligned with ∇f ; and (iii) $|\delta f|/\sigma_f$, where $\delta f = f(\tilde{R} + \Delta \mathbf{x}) - f(\tilde{R})$ is the change in the correlator output when it is evaluated on the backflow-shifted coordinates, in units of the correlator’s own standard deviation σ_f . Two cautions attach to this last quantity. It is a *counterfactual sensitivity*: in the production ansatz the correlator sees the unshifted $\tilde{R}$ (only the determinant sees $\tilde{R} + \Delta \mathbf{x}$; §3.5), so δf measures how much the correlator *would* respond to the displacement, not an actual term in the wavefunction. And it is a typical relative change, not a variance decomposition: $|\delta f|/\sigma_f = 0.7$ means the counterfactual change is 0.7 standard deviations, not that 70% of the variance is attributable to backflow. Table 6.5 reports these alongside the fraction of particles pushed toward their nearest neighbour.

A transition at $\omega \approx 0.1$. The data reveal a sharp regime change. At $\omega \leq 0.01$, the PINN and CTNN are weakly coupled: $\langle \cos \rangle \sim 0.1$ –0.2, the alignment fraction is well below unity, and backflow barely affects the PINN output ($|\delta f|/\sigma_f < 0.07$). In this regime the CTNN operates in a subspace largely orthogonal to the PINN’s gradient—it modifies the wavefunction in directions that W_θ is insensitive to, consistent with its role being to adjust the Slater determinant’s orbital positions rather than the correlator.

At $\omega \geq 0.1$, the picture changes qualitatively. The gradient cosine jumps to 0.5–0.8, the alignment fraction reaches 1.0 for $N \geq 6$, and the counterfactual output change reaches $|\delta f|/\sigma_f = 0.25$ –0.69 (i.e. up to ~ 0.7 standard deviations of the correlator). Here the CTNN acts as a cooperative refinement: it pushes configurations along directions the PINN is maximally sensitive to, systematically correcting the correlator’s residual errors. The transition is sharpest between $\omega = 0.01$ and $\omega = 0.1$, coinciding with the crossover in $\Gamma = V_{\text{int}}/T$ from $\Gamma > 10$ (interaction-dominated) to $\Gamma \sim 3$ –6 (moderate correlation).

Directional structure: correlation-hole deepening. The nearest-neighbour direction fraction provides a complementary signature. At $\omega \geq 0.1$ for $N \geq 6$, only 12–25% of particles are displaced toward their nearest neighbour. The CTNN preferentially pushes particles *apart*, actively deepening the correlation hole. This directional bias is strongest at $N=12$, $\omega=0.1$ ($\rightarrow \text{NN} = 0.12$), exactly where the correlator effective rank peaks and the $\phi \leftrightarrow \psi$ transition occurs. At strong correlation ($\omega \leq 0.01$, large N), the fraction rises toward 0.5 (isotropic), because the large equilibrium inter-particle separations leave little room for further correlation-hole deepening.

6.2.4 Synthesis

The representation analysis supports a coherent picture of how the CTNN-augmented architecture partitions the variational problem across its components.

The **correlator head** W_θ forms a $\sim 1\text{--}3$ dimensional manifold whose leading axis tracks the dominant global coordinate of the electron cloud. The identity of that coordinate shifts across regimes: pair correlations (ψ branch) dominate at moderate-to-weak coupling ($\omega \gtrsim 0.1$), while orbital structure (ϕ) or auxiliary features ($\mathbf{g}$) take over at strong coupling ($\omega \lesssim 0.01$). This $\phi \leftrightarrow \psi$ recomposition is the primary way the architecture adapts to changing physics; dimensionality expansion plays a negligible role.

The **CTNN backflow** provides a full-rank ($\sim 2N$), configuration-dependent displacement field that is implicitly regularised to $\sim 10\%$ of the harmonic length. Despite its geometric complexity, it leaves the correlator manifold almost invariant ($\text{CKA} > 0.98$), confirming that it operates in a complementary subspace. At strong correlation, the CTNN and PINN function independently: backflow adjusts orbital positions via the Slater determinant while the correlator tracks global structure. At moderate-to-weak correlation, the two components become tightly coupled: the CTNN’s displacements align with the PINN’s gradient ($\langle \cos \rangle \sim 0.5\text{--}0.8$), and the counterfactual output change reaches $|\delta f|/\sigma_f$ up to 0.70 (about 0.7 standard deviations of the correlator). The transition between these regimes is sharp and coincides with the physical crossover at $\omega \approx 0.1$.

This division of labour is architecturally desirable. It means the base correlator can robustly learn compact collective coordinates across regimes, while backflow provides an optional, fine-grained correction channel whose character—*independent adjustment versus cooperative refinement*—adapts automatically to the strength of electron–electron correlation. These regularities are, once again, features of one well-trained solution rather than theorems about every possible one. What the companion chapter supplies is their invariant counterpart: the same division of labour priced in energy, where deleting the backflow’s inter-particle messages costs as much as deleting the displacement field entirely (Chapter 7, §7.1.2).

6.3 Wigner–molecule crossover

Up to this point the trained wavefunction has been the thing under examination. In this section it becomes the instrument. Having established that the ansatz is accurate wherever accuracy can be checked, we now turn it on the physics and ask what the electrons are actually doing—and the answer is only reachable this way, because the structure we are looking for is invisible in the observable that would most naturally be plotted. That is the first thing to understand before any of the diagnostics below make sense.

In a finite, rotationally symmetric harmonic trap the appropriate picture for the strongly correlated limit is a *Wigner molecule*: electrons localize in *relative* coordinates—forming concentric rings with polygon-like angular correlations—while the entire pattern may rotate freely. Consequently, lab-frame densities can appear nearly uniform even when strong internal order is present [7, 10]. The following analysis combines one-body density visualization, label-invariant shell topology, rotation-registered angular order, shell radii and separation gaps, pair-statistics validation, and Lindemann-type disorder to establish a coherent, quantitative picture of the Wigner-to-liquid crossover across $N = 2, 6, 12, 20$ and $\omega \in \{1, 0.5, 0.1, 0.01, 0.001\}$.

6.3.1 Observables and diagnostics

For each (N, ω) we draw $\mathbf{X} \sim |\Psi|^2$ and report (i) the pair distribution $g(r)$, (ii) the single-particle radial density $P(r) \propto r \rho(r)$, and summary statistics: r_{mode} , $\langle r \rangle$, σ_r , FWHM, quantiles q_{10}, q_{50}, q_{90} , and a localization ratio $\gamma = \sigma_r/r_{\text{mode}}$ (smaller $\gamma \Rightarrow$ stiffer localization). For multi-electron systems we additionally monitor shell-resolved bond-orientational order $|\Phi_m|$ and a dimensionless Lindemann-type ratio for positional fluctuations on each ring [11].

Density parameter r_s . Following Egger *et al.* [7], we estimate the Wigner–Seitz parameter from the first maximum r^* of the spin-summed pair correlation, $r_s \equiv r^*/a_B^*$. The Fermi–liquid $\rightarrow$ Wigner–molecule crossover occurs near $r_s \simeq 4$ in parabolic dots; our weak-trap cases lie well beyond this threshold [7, 12].

6.3.2 One-body densities: visual crossover

Figures 6.2 and 6.3 compare one-body densities at strong confinement ($\omega = 1$) and weak confinement ($\omega = 10^{-3}$) for $N = 2, 6, 12, 20$. At $\omega = 1$ the density remains compact and the dominant change with N is progressive shell filling under the harmonic trap. At $\omega = 10^{-3}$ the density broadens dramatically and develops pronounced radial structure for all N , consistent with Coulomb-dominated localization and the onset of Wigner-molecule ordering.




Figure 6.2: One-body densities for $N = 2, 6, 12$ at $\omega = 1.0$ (top) and $\omega = 0.001$ (bottom). As confinement weakens, the densities broaden and develop pronounced shell structures characteristic of Wigner molecules.

6.3.3 Energy-based crossover diagnostics

Log-log fits $E(\omega) \propto \omega^\alpha$ on the grid $\omega \in \{10^{-3}, 10^{-2}, 10^{-1}, 0.5, 1\}$ give $\alpha_T \simeq 1.00$, $\alpha_{V_{\text{int}}} \simeq 0.66$, and $\alpha_{V_{\text{trap}}} \simeq 0.81, 0.73, 0.72$ for $N = 2, 6, 12$ respectively. The kinetic energy thus scales essentially linearly with ω , while both potential terms grow sublinearly with nearly N -independent exponents. Across $\omega \in [1, 10^{-3}]$ the interaction-to-kinetic ratio $\Gamma = \langle V_{\text{int}} \rangle / \langle T \rangle$ increases from $\Gamma \simeq 0.9 \rightarrow 8.7$ ($N=2$), $2.4 \rightarrow 25.4$ ($N=6$), and $3.6 \rightarrow 39.9$ ($N=12$), while T/V_{int} falls correspondingly (Fig. 6.4a).

The partitioning between trap and interaction energies becomes quasi-classical at weak confinement. At $\omega = 10^{-3}$ we find $2\langle V_{\text{trap}} \rangle / \langle V_{\text{int}} \rangle \simeq 1.25, 1.12, 1.02$ for $N = 2, 6, 12$, so that $N=12$ already satisfies the classical virial condition $V_{\text{int}} \approx 2V_{\text{trap}}$ to within 2% (Fig. 6.4b). Using $V_{\text{trap}} = \frac{1}{2}\omega^2 \sum_i \langle r_i^2 \rangle$ we extract an rms radius $r_{\text{rms}} = \sqrt{2V_{\text{trap}}/(N\omega^2)}$, which expands from $\approx 1.1\text{--}2.0 a_B^*$ at $\omega = 1$ to $\approx 70\text{--}169 a_B^*$ at $\omega = 10^{-3}$. Fitting $r_{\text{rms}} \propto \omega^{\alpha_r}$ yields $\alpha_r \simeq -0.60$ to -0.64 , i.e. $\langle r^2 \rangle \propto \omega^{-\beta}$ with $\beta \simeq 1.19\text{--}1.28$: the cloud expands *faster* than the non-interacting ω^{-1} scaling, consistent with interaction-driven swelling and shell formation.




Figure 6.3: One-body densities for $N=20$ at strong and weak confinement. The weak-trap case shows strong radial broadening and clear multi-shell organization, consistent with deep Wigner-molecule ordering.




Figure 6.4: Energy-based Wigner diagnostics as functions of trap frequency. As ω decreases the dots become interaction dominated ($T/V_{\text{int}} \ll 1$) and the trap–interaction partition approaches the classical virial form, most clearly for $N=12$.

6.3.4 Radial localization and pair correlations

Table 6.6 summarizes the single-particle radial law for $N=2$ and Table 6.7 for $N=6$ across all five trap strengths. In both cases r_{mode} grows as a power law ($r_{\text{mode}} \sim \omega^\alpha$ with $\alpha \simeq -0.62$ for $N=2$, close to the classical two-body scaling $-2/3$) while the localization ratio γ shrinks, signaling stiffening of the single-particle shell.

Table 6.6: Radial diagnostics for $N=2$ across ω (Bohr units).

ω	r_{mode}	$\langle r \rangle$	σ_r	FWHM	q_{10}	q_{50}	q_{90}	γ
1.00	1.435	1.632	0.705	1.743	0.735	1.581	2.581	0.491
0.50	2.303	2.475	1.007	2.466	1.209	2.409	3.829	0.437
0.10	3.601	3.830	1.728	4.449	1.640	3.714	6.127	0.480
0.01	14.191	14.593	5.615	13.602	7.390	14.322	22.038	0.396
0.001	64.292	63.953	18.479	43.977	40.392	63.814	87.714	0.287

Table 6.7: Radial and angular diagnostics for $N=6$ across ω (Bohr; angles in radians).

ω	r_{mode}	σ_r	γ	$\sigma(\text{ring angles})$	$\gamma_{r_{ij}}$
1.00	2.0907	0.9249	0.4424	0.7046	0.3538
0.50	3.0702	1.3648	0.4445	0.6943	0.3397
0.10	8.1686	3.5025	0.4288	0.6842	0.3149
0.01	33.1538	13.9094	0.4195	0.6497	0.2889
0.001	113.5542	47.8076	0.4210	0.6368	0.2664

At $\omega = 10^{-3}$, the $N=2$ pair has mode separation $r_{12} \approx 122$, Lindemann ratio $\gamma_{r_{12}} \approx 0.20$, and a relative-angle distribution sharply peaked near π ($\sigma(\pi - \Delta\phi) \approx 0.35$), establishing a dilute, strongly anti-aligned Wigner dimer. For $N=6$ the ring angular fluctuations and pair Lindemann both decrease monotonically, but γ plateaus near 0.42: the single shell stiffens internally even though the radial profile broadens due to the growing equilibrium radius.




Figure 6.5: Two-electron radial densities $P(r)$ across ω .




Figure 6.6: Six-electron radial densities $P(r)$ across ω .

6.3.5 Shell topology, angular order, and the rotating Wigner molecule

This subsection resolves an apparent contradiction: the densities plotted above look like smooth rings, yet we are about to claim the electrons form a crystal. Both are true, and the reason is symmetry. Because the Hamiltonian is rotationally invariant, so is its ground state, and any lab-frame average necessarily smears an angularly ordered configuration into a featureless ring. The order is real; it simply lives in relative coordinates, and looking for it in the lab frame is looking in the one place it is guaranteed not to be. Everything below follows from taking that seriously.

Concretely, a Wigner molecule’s internal angular order can only be observed after registering frames to a co-rotating reference. For a shell s with n_s particles we compute a rotation-registered bond-orientational parameter $|\Phi_{n_s}|$ and a corresponding Lindemann-type ratio quantifying angular stiffness. Lab-frame densities remain ring-like (the “rotating Wigner molecule” scenario), while the co-rotating frame reveals crystal-like polygon order [7, 10].

$N=6$: single ring with sector mixing. At $\omega = 0.01$ the system fluctuates between a (1,5) sector (fraction ≈ 0.69) and a (0,6) single-ring sector (fraction ≈ 0.31), with bond order $|\Phi_5| = 0.453 \pm 0.214$ and $|\Phi_6| = 0.466 \pm 0.206$ respectively. At $\omega = 10^{-3}$ the ring stiffens further: (1,5) fraction ≈ 0.80 , (0,6) ≈ 0.20 , with $|\Phi_5| = 0.597 \pm 0.224$ (Lind ≈ 0.225) and $|\Phi_6| = 0.663 \pm 0.183$ (Lind ≈ 0.205). Classically (1,5) is the ground-state topology and (0,6) a proximate competitor; the observed quantum mixture mirrors this hierarchy [8, 9]. In both weak-trap cases the global $g(r)$ is reproduced to numerical precision by the appropriate inner–inner / inner–outer / outer–outer (II/IO/OO) mixtures.

$N=12$: two shells and commensurability. At $\omega = 0.01$ two-shell formation is already frequent (fraction 0.761 at radial-gap threshold $\tau=3.0$), distributed across sectors (1,11) = 32.6%, (2,10) = 25.7%, (3,9) = 13.7%. At $\omega = 10^{-3}$ shelling becomes ubiquitous ($f_{2\text{shell}} \approx 0.99$ –0.92 for $\tau = 2.0$ –3.0) and the commensurate (3,9) split emerges as modal (43.3% of quality-cut frames), with strong bond order on both rings: $|\Phi_3| = 0.687 \pm 0.229$ (Lind_{in} = 0.223), $|\Phi_9| = 0.526 \pm 0.194$ (Lind_{out} = 0.224). A shell-resolved $g(r)$ reconstruction (II/IO/OO mixture with the combinatorial weights 0.045:0.409:0.545 of the ideal (3,9) topology) reproduces the global $g(r)$ with cosine similarity 0.9982 (Fig. 6.7). Since the sector histograms and the total are built from the same samples, this is a test of *decomposability* rather than of sampling correctness: it shows the pair structure is cleanly explained by the identified shell sectors and, because the weights are the idealised (3,9) combinatorics and not empirical sector fractions, that this topology accounts for it. Sampling correctness itself rests on the independent restarts, threshold robustness, and

chain-stationarity diagnostics reported above.




Figure 6.7: Shell-resolved reconstruction of $g(r)$ at $\omega = 10^{-3}$ for $N=12$. The global pair distribution is reproduced by a convex mixture of inner–inner (II), inner–outer (IO) and outer–outer (OO) histograms with weights set by the observed ($n_{\text{in}}, n_{\text{out}}$) statistics.

$N=20$: three-shell molecule and rapid topology collapse. At $\omega = 10^{-3}$ the dominant shell topology is (1, 7, 12)—a clear three-shell Wigner molecule with widely separated rings at radii $r_0 \approx 38$, $r_1 \approx 132$, $r_2 \approx 247$ (gaps ≈ 94 and ≈ 115). Rotation-registered angular order is strong ($\tilde{C}_{\text{global}} \approx 0.79$), while the lab-frame pinning metric remains small ($C_{\text{global}} \approx 0.06$), as expected for a freely rotating molecule. Lindemann disorder on the two occupied outer shells is low: $L_{7,\text{reg}} \approx 0.35$ and $L_{12,\text{reg}} \approx 0.27$.

Already at $\omega = 10^{-2}$, the three-shell pattern collapses to (1, 19): the shell radii contract by an order of magnitude ($r_0 \approx 10$, $r_1 \approx 45$), the registered angular order drops to $\tilde{C}_{\text{global}} \approx 0.20$, and the outer-shell Lindemann rises to ≈ 0.57 . At $\omega = 1$ the same (1, 19) topology persists with compact radii ($r_0 \approx 0.55$, $r_1 \approx 2.3$), negligible internal order ($\tilde{C}_{\text{global}} \approx 0.20$), and high disorder ($L_{\text{out}} \approx 0.69$). The sharpness of this transition—a single decade in ω separates a stiff three-shell crystal from a disordered single-shell state—is the most dramatic structural collapse observed in our system-size series.

6.3.6 Synthesis: the Wigner–molecule crossover

Across $N = 2, 6, 12, 20$ we observe a consistent, internally validated structural evolution as ω decreases from 1 to 10^{-3} :

1. **One-body densities** show dramatic spatial expansion and ring formation at $\omega = 10^{-3}$ (Figs. 6.2, 6.3).
2. **Shell topology** reveals discrete finite- N Wigner patterns at $\omega = 10^{-3}$ —(1, 5) for $N=6$, (3, 9) for $N=12$, (1, 7, 12) for $N=20$ —consistent with classical ground-state analyses [8, 9]. Sharp topology collapses occur between 10^{-3} and 10^{-2} for the larger systems.
3. **Shell radii and separation gaps** provide a quantitative real-space counterpart: shells are widely separated and stable in the Wigner regime, while both radii and gaps shrink rapidly with increasing ω .




Figure 6.8: Twelve-electron radial densities $P(r)$ across ω .

4. **Rotation-registered angular order** demonstrates strong polygon-like internal localization (including multi-shell ordering for $N=12$ and $N=20$). Lab-frame measures remain comparatively small, consistent with free global rotation of the molecule.
5. **Lindemann disorder** rises strongly with ω —independently confirming progressive melting—while the Wigner point supports simultaneously stiff shells (e.g. $L_{\text{out}} \approx 0.22\text{--}0.27$ for $N=12, 20$).
6. **Energy ratios** show the interaction-to-kinetic ratio Γ growing to ~ 40 for $N=12$ at $\omega = 10^{-3}$, and the virial condition $V_{\text{int}} \approx 2V_{\text{trap}}$ approaching its classical value.

The commensurate (3, 9) sector for $N=12$ aligns with known classical ground states, but with finite weight in competing (1, 11) and (2, 10) sectors reflecting quantum rotational mixing [10, 12]. Our sector-resolved ring-order parameters $|\Phi_m|$ are extracted *within* a fully variational quantum ansatz and measured conditionally on shell occupancy, which—to our knowledge—has not been reported before for $N \geq 12$ parabolic dots. The $N=20$ results extend this picture to three concentric shells and demonstrate that topology collapse, angular-order loss, and Lindemann melting all coincide within the same narrow ω window, providing a sharp quantitative trajectory from localized Wigner molecule to the denser, more weakly ordered liquid state [7–9, 12].

From physics back to method. These structural facts are also what make part of the method possible. The shell radius that grows as $\omega^{-2/3}$ and the angular order that sharpens as the trap weakens are exactly the quantities a sampler must respect to reach the Wigner regime without MCMC. Chapter 7 uses them directly: an origin-centred proposal aims where the wavefunction no longer has any mass, whereas a proposal built from the measured shell radius and angular order tracks it, and carries MCMC-free training far into the correlated regime (§7.4). What the trained model reveals about the physics is, in turn, what lets us train it more cheaply—the instrument and the method inform one another.

6.4 Collocation-assisted training

The results in Sections 6.1–6.3 rely on a two-phase protocol in which a residual-loss pretraining stage conditions the ansatz, followed by a stochastic-reconfiguration (SR) refinement phase that uses Markov-chain Monte Carlo (MCMC) sampling from $|\Psi|^2$ to optimise the variational energy. This section reports

a complementary investigation: can an accurate many-body wavefunction be trained *without any MCMC sampling during training*, using instead a purely importance-sampled collocation approach?

The motivation is threefold. First, MCMC introduces autocorrelation, burn-in overhead, and mixing timescales that grow with particle number and correlation strength—particularly problematic at low ω where the wavefunction support is diffuse. Second, a collocation-only pipeline is embarrassingly parallel: each batch is drawn independently from a fixed analytic proposal, eliminating the sequential dependence inherent in MCMC chains. Third, collocation exposes the optimisation geometry more directly, making it a useful laboratory for understanding which architectural and sampling choices are load-bearing.

Two ways of posing the collocation problem must be separated at the outset, because they behave very differently. The *strong form* asks the wavefunction to satisfy the local Schrödinger equation pointwise: it minimises the variance of the local energy E_L , whose evaluation requires the Laplacian. The *weak form* instead uses a score-function (REINFORCE) gradient in which the Laplacian enters only as a reward and is never differentiated through. The distinction is not cosmetic. Appendix C shows that the strong form meets a structural obstruction as soon as it is asked to train backflow: the learning signal lives at the nodal surface, but E_L diverges there for any imperfect node, so no combination of sampling, loss, or preconditioning breaks an accuracy floor of $\sim 0.3\%$ —and that floor is set by the Jastrow alone, with the backflow contributing almost nothing. This is a property of the strong-form objective, not of the method as a whole; it is precisely why the experiments in this section use the weak form throughout, and why the $\sim 0.3\%$ number should not be read as the accuracy of collocation training.

Read in that light, the finding is clean. The *weak-form* objective, on the same Slater–Jastrow–backflow ansatz, recovers the variational energy to within DMC uncertainty at strong confinement and to a few tenths of a percent of our variational gold standard well into the correlated regime, for $N \in \{6, 12\}$; $N=20$ remains harder. And in the deep-Wigner limit a physics-informed proposal carries the same MCMC-free objective all the way to $\omega = 0.0035$ (Chapter 7, §7.4)—far past the point where a naive proposal collapses. What follows sets out what makes the weak form work, and where it, too, eventually breaks.

6.4.1 Training methodology

The collocation training used here is exactly the MCMC-free pipeline specified in the Methods: a weak-form (REINFORCE) objective in which the local energy enters as a detached reward (§4.5, Eq. (4.8)), an analytic Gaussian-mixture proposal reweighted to $|\Psi_\theta|^2$ with ESS / Pareto- $\hat{k}$ monitoring and tail controls (§4.5.1), and—in the Wigner regime—the physics-informed ring proposal with cascade warm-starting (§4.5.2). We do not repeat those definitions; this section reports what the pipeline achieves and where it fails. Two methodological findings shape the results below. First, among the optimisers (§4.4), **Adam** was the robust default across all regimes, whereas **CG-SR** converged faster at $\omega \geq 0.1$ but diverged for $\omega \leq 0.01$, where the geometric tensor becomes ill-conditioned. Second, **cascade warm-starting**—initialising each confinement from the converged checkpoint of a stronger one—was decisive for reaching the deep-Wigner regime, exploiting the transfer of correlation structure across ω .

6.4.2 Architecture variants

Three ansatz families were explored within the collocation framework, all built on the same Slater–Jastrow core: $\Psi(\mathbf{R}) = \det[\phi_i(\mathbf{r}_j + \Delta\mathbf{r}_j)] \cdot e^{J(\mathbf{R})}$.

Jastrow-only (no backflow). The Jastrow factor is parameterised by a graph neural network (CTNNJastrowVCycle) with $\sim 25\text{k}$ parameters. It operates on node features (single-particle orbital values) and edge features (pairwise distances with soft-core regularisation). The V-cycle variant uses a coarse-to-fine message-passing scheme. This family was the fastest to train and the most robust, serving both as a standalone ansatz and as an initialiser for backflow runs.

Coordinate backflow + Jastrow (BF). The CTNNBackflowNet adds a copresheaf-style message-passing network that predicts electron displacement vectors $\Delta\mathbf{r}_j$ fed into the Slater determinant. Edge features in the backflow network scale as $\mathcal{O}(N^2 \times h_{\text{bf}})$ where h_{bf} is the hidden dimension, creating a memory bottleneck at large N . For $N=6$ and $N=12$, the default $h_{\text{bf}}=128$ was used (total $\sim 49\text{k}$ parameters); for $N=20$, reduced variants with $h_{\text{bf}} \in \{48, 64\}$ were necessary. The BF+Jastrow family produced the best results at $N=6$ and $N=12$.

Neural Pfaffian. The Pfaffian ansatz replaces the Slater determinant with a more general antisymmetric form $\text{Pf}[A_{ij}]$, which in principle offers richer nodal topology. Representative results: $E=20.261 \pm 0.004$ and $E=20.245 \pm 0.005$ at $N=6$, $\omega=1.0$ (both $\sim 0.4\%$ above DMC), compared to $+0.009\%$ for BF+Jastrow. The Pfaffian’s greater expressivity was not exploitable within the current training pipeline; the simpler BF+Jastrow family consistently outperformed it.

6.4.3 Results

Table 6.8 presents the best energies achieved by collocation-assisted training across all (N, ω) combinations studied. All energies are evaluated by independent VMC with $\sim 30\text{k}$ MCMC samples after training (a “heavy evaluation”), ensuring that the reported numbers are not biased by the training-time importance sampling.

Table 6.8: Best ground-state energies from collocation-assisted training. “Multi-stage” denotes the best result from earlier campaigns using hand-tuned continuation chains; “Campaign (best)” denotes the best single result from the 30-run reliability campaign for $N=6$ (Table 6.9), and the best result across all independent training runs for $N=12$ (no dedicated reliability campaign was conducted for $N=12$). The $N=2$ rows report a single collocation run each (the zero-variance-extrapolated energy under Multi-stage; the campaign column does not apply), reaching DMC accuracy for $\omega \geq 0.01$ and degrading only in the deep-Wigner limit ($+0.15\%$ at $\omega = 0.001$). All values are VMC-evaluated ($\geq 20\text{k}$ samples) at the best checkpoint. The reference column is the DMC energy where one exists and, for (N, ω) with no published DMC value ($N=6$ at $\omega \leq 0.01$; see Table 6.2), our own converged variational (PINN+CTNN) energy; the corresponding %err is then a measure of internal consistency, not of absolute accuracy. The tabulated $N=6$ low- ω errors are the *reliability-campaign* bests and are deliberately conservative: the best single runs we observed reached $+0.06$ – 0.13% at $\omega \leq 0.1$ (§6.4.5). All tabulated rows use the backflow+Jastrow (BF) architecture; the memory-limited $N=20$ case is discussed separately in Appendix D.2. Parentheses denote 1σ statistical uncertainty on the last digit(s).

N	ω	Reference	Multi-stage	Campaign (best)	%err	Arch
2	1.0	3.00000	3.00035	—	+0.012	BF
	0.5	1.65977(1)	1.66014	—	+0.023	BF
	0.1	0.44079(1)	0.44077	—	−0.004	BF
	0.01	0.073839(2)	0.073840	—	+0.002	BF
	0.001	0.013778	0.013799	—	+0.154	BF
6	1.0	20.15932(8)	20.1613(23)	20.1620(56)	+0.013	BF
	0.5	11.78484(6)	11.7847(20)	11.7850(48)	+0.002	BF
	0.1	3.55385(5)	3.5571(4)	3.5632(7)	+0.263	BF
	0.01	0.69036	0.69141(8)	0.69200(1)	+0.237	BF
	0.001	0.140832	0.14118(1)	0.14118(1)	+0.250	BF
12	1.0	65.7001(1)	65.712(4)	65.706(4)	+0.009	BF
	0.5	39.1596(1)	39.171(4)	39.169(4)	+0.024	BF
	0.1	12.2698(1)	12.285(1)	12.2824(6)	+0.102	BF

$N=6$: near-reference accuracy across all ω . For six electrons, collocation training reaches DMC-level accuracy where DMC exists—at $\omega \geq 0.5$ the errors are $\lesssim 0.01\%$, inside the DMC statistical uncertainty. At the most extreme confinements, $\omega \in \{0.01, 0.001\}$, where the electrons are separated by ~ 30 – $120 a_0^*$ (a few oscillator lengths) and no DMC reference is available, the energies stay within 0.1–0.25% of our converged variational gold standard; this is agreement between two independent training paradigms on the same ansatz, not a certificate of absolute accuracy. The best result at $\omega=1.0$ is $E=20.1613 \pm 0.0023$, an error of $+0.009\%$, indistinguishable from DMC.

These results are achieved without any Markov chain in the training gradients. MCMC enters only outside the gradient loop: a small Metropolis probe periodically selects the best checkpoint, and the final heavy evaluation certifies the energy. The training itself uses 4096 importance-resampled collocation points per epoch with an oversampling factor of 8, requiring $\sim 32\text{k}$ proposal evaluations per gradient step—modest compared to the $\sim 3 \times 10^3$ MCMC samples per step in the SR-based approach.

The best $N=6$ result at $\omega=1.0$ was obtained through a three-stage continuation chain (see §4.5.2): an initial BF+Jastrow joint REINFORCE run (900 epochs, $\text{lr}=5\times 10^{-4}$), followed by a low-LR continuation (750 epochs, $\text{lr}=3\times 10^{-4}$), and a final hard-focus stage (500 epochs, $\text{lr}=2\times 10^{-4}$). Each stage inherited the full checkpoint from the previous one. The monotonic improvement across stages—+0.08%, +0.05%, +0.009%—demonstrates that patient multi-stage polishing is effective within the collocation framework.

$N=12$: sub-0.03% at strong confinement. For twelve electrons, the BF+Jastrow collocation results track DMC closely at $\omega \geq 0.5$, with errors of +0.018% and +0.028% at $\omega=1.0$ and 0.5 respectively. At $\omega=0.1$ the error increases to +0.122%, reflecting the growing difficulty of importance sampling in 24 dimensions at weak confinement. The ESS at $\omega=0.1$ is typically 5–15 per epoch (out of 4096 kept points), indicating that the Gaussian mixture proposal still maintains usable overlap with $|\Psi|^2$.

Attempts to extend to $\omega \leq 0.01$ at $N=12$ were unsuccessful: both direct starts and cascade warm-starts from $\omega=0.1$ failed to converge, with the ESS collapsing to 1 and the rollback mechanism rejecting all gradient updates. This marks the boundary of the current collocation approach for $N=12$.

$N=20$ exposes the memory–sampling trade-off. Twenty electrons in two dimensions push the collocation pipeline against a hardware wall rather than a conceptual one. The backflow’s edge-feature tensor scales as $\mathcal{O}(N^2 \times h_{\text{bf}})$, which at $N=20$ forces the hidden dimension, the collocation budget, or the oversampling factor down far enough that the gradient quality suffers—so much so that a Jastrow-only ansatz, freed of that cost, trains better than the backflow ansatz it should in principle beat. Because the limitation is one of memory and gradient noise rather than of the method, we report the $N=20$ collocation results, and the reversal that exposes the bottleneck, in Appendix D.2 rather than here.

6.4.4 What worked and what did not

The collocation programme explored a large space of methods, and most of that space did not work. We report the successful and the failed variants together, because the failures identify which assumptions are load-bearing. Each failure below was, at the time, an intuitively reasonable idea; each turned out to violate a specific assumption that the method had never been forced to state. A working configuration tells you that some set of choices is sufficient. A failure, if you chase it far enough to name the mechanism, tells you which choice was load-bearing and why—and that is the kind of statement that transfers to a system other than this one. We therefore report the mechanism in each case, not just the fact that the attempt was abandoned.

Positive: pure REINFORCE over mixed objectives. Setting $\beta=0$ in the hybrid loss (pure REINFORCE, no direct weak-form term) was consistently more stable and produced better final energies than any nonzero β . The direct term $\tilde{E}=\frac{1}{2}\|\nabla \log \Psi\|^2+V$ provided a cheaper but noisier gradient signal; at $\beta>0$ it interfered with the REINFORCE gradient by pulling toward a local-kinetic minimum that did not correspond to the variational minimum.

Positive: cascade warm-start for low ω . Cold-start training at $\omega=0.01$ never converged for $N\geq 6$. Cascade initialisation from $\omega=0.1$ made the $\omega=0.01$ regime accessible, reducing the initial error from $\sim 30\%$ (random init) to $\sim 5\%$ (warm-started) and enabling subsequent optimisation to reach +0.15% within ~ 1000 epochs.

Positive: ESS-adaptive oversampling + rollback. The combination of ESS-adaptive oversampling (automatically increasing the proposal budget when overlap is poor) and instability rollback (reverting to a good checkpoint when the energy spikes) was essential for the ultra-low- ω results. Without these mechanisms, training at $\omega=0.001$ diverged within a few hundred epochs.

Negative: CG-SR at ultra-low ω . Natural-gradient preconditioning via CG-SR produced excellent results at $\omega \geq 0.1$ (fastest convergence, best final energies for $N=6, 12$). However, at $\omega \leq 0.01$ the Fisher information matrix becomes extremely ill-conditioned: the eigenvalues span many orders of magnitude because the wavefunction’s sensitivity to parameters varies enormously across the diffuse, weakly confined state. CG-SR updates in this regime produced errors of +4–36% for $N=6$ at $\omega=0.01$ and $\omega=0.001$, compared to +0.15% and +0.24% from Adam with per-parameter adaptive learning rates. Ill-conditioning alone does not decide whether SR helps; what matters is whether the metric is *also* statistically resolvable. This gives a three-regime picture (developed in Chapter 7, Q2): where the metric is well-conditioned,

Adam already suffices and SR buys little; where it is anisotropic but still estimable, SR is most valuable; and where importance-sampling collapse makes the empirical metric itself unreliable, inverting it amplifies sampling noise and does harm. The failures above are the third regime, not the second.

Negative: Langevin proposal refinement. To address the ESS catastrophe at high N , we implemented overdamped Langevin dynamics on the proposal samples: $\mathbf{R}' = \mathbf{R} + \varepsilon \nabla_{\mathbf{R}} \log |\Psi(\mathbf{R})|^2 + \sqrt{2\varepsilon} \boldsymbol{\eta}$, applied for $K=10$ – 20 steps before importance resampling. The intent was to push proposal samples toward high- $|\Psi|^2$ regions without running a full MCMC chain.

This approach was systematically worse than standard importance resampling in every configuration tested:

- $N=20$, $\omega=0.1$: Langevin VMC error +153% vs. standard +5.7% (catastrophic divergence).
- $N=20$, $\omega=1.0$: Langevin +5.2% vs. standard +1.5%.
- $N=6$, $\omega=0.001$: Langevin +0.30% vs. Adam +0.24%.

The failure mechanism is that $K=10$ – 20 Langevin steps from a Gaussian starting point in $Nd=40$ dimensions do not equilibrate to $|\Psi|^2$. The resulting non-equilibrium sample distribution biases the REINFORCE gradient, which in turn pushes the wavefunction to concentrate further around the biased distribution—a positive-feedback loop that rapidly destroys the ansatz. Langevin refinement would require either much longer chains ($K \gtrsim 100$, negating the computational advantage over MCMC) or a Metropolis–Hastings acceptance step (reintroducing MCMC).

Negative: finite-difference residual loss. The FD collocation loss, which computes E_L via graph-safe finite differences and penalises its spread, was tested extensively. Representative BF results at $N=6$, $\omega=1.0$: $E=20.204 \pm 0.002$ (+0.22%), compared to +0.009% from REINFORCE. FD training was more sensitive to step-size h and noise in the Laplacian estimate, and never matched the REINFORCE objective for final accuracy.

Negative: Neural Pfaffian. Despite its richer antisymmetric structure, the Pfaffian ansatz underperformed the simpler BF+Jastrow family in all collocation experiments. The best Pfaffian result at $N=6$, $\omega=1.0$ was $E=20.245 \pm 0.005$ (+0.43%), more than $40\times$ worse than BF+Jastrow. The Pfaffian’s larger parameter space amplified the gradient noise from importance sampling, making it harder to optimise reliably.

Negative: backflow at $N=20$. Backflow training at $N=20$ was severely limited by memory. The edge-feature tensor scales as $\mathcal{O}(N^2 \times h_{\text{bf}})$, requiring h_{bf} to be reduced from 128 to 48–64 for $N=20$. This reduced-capacity backflow achieved only +18% error at $\omega=1.0$, compared to +1.45% for the Jastrow-only ansatz. The gradient signal at $N=20$ is too noisy for the additional backflow parameters to be trained effectively; the simpler Jastrow network with its smaller parameter count permits higher oversampling and better gradient quality.

What the failures have in common. Read together, these five negatives are not five unrelated disappointments. Four of them—CG-SR, Langevin refinement, the Pfaffian’s parameter count, and the backflow at $N=20$ —fail through the same channel: a quantity that must be *estimated* from a finite sample degrades faster than the quantity it is meant to improve. Natural gradients need an estimate of the metric; Langevin needs samples that have actually equilibrated; a larger parameter space needs a correspondingly better-resolved gradient. In each case the method is not wrong in principle, and would be perfectly sound given samples we cannot afford. What fails is the estimate, and it fails precisely in the regime where the correction was supposed to help. This is the thread that §6.4.5 follows deliberately, from the proposal mismatch at one end to the evaluation bottleneck at the other. Stated here, it can be carried into that analysis rather than rediscovered at its conclusion.

6.4.5 Reliability analysis: the gradient-quality chain

The results in Table 6.8 report the *best* energies obtained by collocation training, selected from multiple runs on different seeds. A natural question is whether these results are *typical* or whether they reflect rare fortunate outcomes from a high-variance process. This subsection reports a systematic 30-run reliability

campaign designed to answer that question, and reveals a simple causal chain that explains both the successes and failures of the MCMC-free collocation approach.

Experimental design

All runs use the BF+Jastrow ansatz (~ 49 k parameters) initialised from the same pretrained checkpoint. The grid is $N=6$ across $\omega \in \{1.0, 0.5, 0.1, 0.01, 0.001\}$ with three random seeds per cell (42, 137, 314), giving $5 \times 2 \times 3 = 30$ runs. Two training recipes are compared head-to-head:

- **Config A (baseline):** Adam optimiser with $\text{lr}=5 \times 10^{-4}$, $\text{lr}_{\text{jas}}=5 \times 10^{-5}$, 4096 collocation points, $8\times$ oversampling, local-energy clipping at 5σ , no reward normalisation, no rollback, fixed Gaussian mixture proposal (Eq. (4.9)).
- **Config B (robust):** Identical to Config A plus four stabilisation mechanisms: $16\times$ oversampling (doubling the proposal budget), reward normalisation (standardising the REINFORCE reward to zero mean and unit variance before the gradient step), instability rollback (reverting to the previous checkpoint when the energy jumps exceed 3σ , with learning-rate decay of 0.95), and an adaptive Gaussian mixture proposal (8 GMM components refitted every 30 epochs to track the evolving $|\Psi|^2$).

All runs train for 1200 epochs. Evaluation uses a 20 000-sample VMC probe every 50 epochs, with the best-VMC checkpoint restored for a final 20 000-sample IS evaluation.

Main results

Table 6.9 presents the complete results of the 30-run campaign. For each (ω, recipe) combination, three seeds provide a mean, standard deviation, and worst-case error.

Table 6.9: Reliability campaign: relative energy errors (% above DMC) for $N=6$ across five trap frequencies. Each cell shows the final IS-evaluated error for one $(\omega, \text{recipe}, \text{seed})$ combination. Bold entries mark the best result at each ω . “Mean” and “CV” are the mean error and coefficient of variation across three seeds.

ω	Recipe	Seed 42	Seed 137	Seed 314	Mean	CV
1.0	Baseline	+0.073	+0.045	+0.060	0.059	0.24
	Robust	+0.032	+0.013	+0.105	0.050	0.97
0.5	Baseline	+0.106	+0.060	+0.108	0.091	0.30
	Robust	+0.002	+0.112	+0.047	0.054	1.04
0.1	Baseline	+0.419	+0.547	+0.427	0.464	0.15
	Robust	+0.325	+0.263	+0.270	0.286	0.12
0.01	Baseline	+0.500	+0.379	+0.746	0.542	0.34
	Robust	+0.289	+0.282	+0.237	0.269	0.10
0.001	Baseline	+3.876	+7.187	+4.733	5.265	0.33
	Robust	+0.250	+0.331	+0.263	0.281	0.15

Several patterns are immediately apparent. First, the robust recipe dramatically improves the low- ω regimes: at $\omega=0.001$ the mean error drops from +5.3% (baseline) to +0.28% (robust), a factor of 19 improvement. Second, the best single result at every ω is a robust run, achieving +0.013% at $\omega=1.0$, +0.002% at $\omega=0.5$, +0.263% at $\omega=0.1$, +0.237% at $\omega=0.01$, and +0.250% at $\omega=0.001$. Third, the robust recipe exhibits higher seed-to-seed variance ($\text{CV} \sim 1.0$) at high ω but dramatically lower variance at low ω , suggesting that its stabilisation mechanisms have the most impact precisely where sampling is hardest.

The gradient-quality chain

The reliability campaign exposes a simple causal chain that governs the quality of collocation-trained wavefunctions:

proposal mismatch $\xrightarrow{\text{causes}}$ $\hat{k}$ inflation $\xrightarrow{\text{causes}}$ gradient noise

$\xrightarrow{\text{causes}}$ optimisation stochasticity $\xrightarrow{\text{causes}}$ run-to-run variance

(6.1)

Each link in this chain can be verified quantitatively from the training logs. We now explain each component and the evidence supporting each causal link.

Step 1: proposal–target mismatch. As described in §4.5.1, the REINFORCE estimator in Eq. (4.8) requires samples from $|\Psi_\theta|^2$, but collocation draws from the analytic Gaussian mixture proposal $q(\mathbf{R})$ in Eq. (4.9) and reweights via importance sampling. The quality of this reweighting depends on how well q overlaps with $|\Psi_\theta|^2$.

For the harmonic quantum dot, $|\Psi|^2$ concentrates on a manifold whose geometry depends on ω : at $\omega=1.0$, electrons occupy a compact cloud of radius of order the oscillator length $\ell = 1/\sqrt{\omega}$, and the three-component Gaussian mixture in Eq. (4.9) provides adequate coverage. At $\omega=0.001$, the electron cloud extends to $\sim 100 a_0^*$ (several oscillator lengths, $\ell \approx 31.6 a_0^*$), the pair correlation function develops sharp shell structure (Section 6.3), and $|\Psi|^2$ concentrates on a thin correlated submanifold that the isotropic Gaussian proposal covers inefficiently.

The mismatch manifests directly in the importance weights. At $\omega=1.0$, weights are relatively uniform: the effective sample size is typically $\text{ESS} \sim 5000\text{--}6000$ out of $\sim 65\text{k}$ proposal samples ($16\times$ oversampling of 4096 points), and the Pareto shape parameter $\hat{k}$ of the weight tail is 0.4–0.5, indicating a well-behaved thin-tailed distribution. At $\omega=0.001$, the ESS collapses to 1–6 (out of the same 65k proposal), and $\hat{k}$ rises above 1.0, entering the regime of infinite-variance importance weights.

Step 2: $\hat{k}$ inflation and gradient quality. The Pareto shape parameter $\hat{k}$ of the importance weight distribution, estimated via the PSIS diagnostic of Vehtari et al. (2024), provides a precise measure of gradient reliability. When $\hat{k} < 0.7$, the importance-weighted gradient estimator has finite variance and the central limit theorem applies normally. When $\hat{k} > 1.0$, the variance of the estimator is infinite and the gradient signal is dominated by a few extreme samples.

Table 6.10 summarises the average $\hat{k}$ across the campaign. The correspondence between $\hat{k}$ and final accuracy is striking.

Table 6.10: Average Pareto $\hat{k}$ of the importance weight distribution during training, by regime and recipe. Lower is better; $\hat{k} < 0.7$ is considered reliable, $\hat{k} > 1.0$ indicates infinite-variance weights.

ω	$\hat{k}$ (avg)		Mean error (%)	
	Baseline	Robust	Baseline	Robust
1.0	0.75	0.50	0.059	0.050
0.5	0.83	0.54	0.091	0.054
0.1	1.48	0.70	0.464	0.286
0.01	2.63	1.39	0.542	0.269
0.001	3.13	1.86	5.265	0.281

The robust recipe reduces $\hat{k}$ by approximately a factor of 2 at every ω . At $\omega=1.0$ this takes $\hat{k}$ from 0.75 (borderline) to 0.50 (reliable), resulting in a modest improvement. At $\omega=0.1$ it takes $\hat{k}$ from 1.48 (broken) to 0.70 (borderline), corresponding to the transition from mediocre (+0.46%) to marginal (+0.29%) accuracy. At $\omega=0.001$ even the robust recipe has $\hat{k}=1.86$, but the adaptive proposal and rollback mechanisms prevent the catastrophic divergence seen in the baseline (+5.3%).

Step 3: from gradient noise to optimisation stochasticity. When $\hat{k}$ is large, most of the gradient signal comes from a few samples with extreme importance weights. Which samples are extreme depends on the random proposal draw at each epoch—a function of the random seed. Two seeds that happen to draw different extreme samples will follow different optimisation trajectories, even though they share the same architecture and hyperparameters.

This is not the usual seed dependence of stochastic gradient descent, where the law of large numbers ensures that gradient estimates are unbiased and the trajectory variance decreases with batch size. When $\hat{k} > 1$, the variance of the gradient estimator is *infinite*: increasing the batch size reduces the variance sublinearly (or not at all), and extreme samples can dominate the update indefinitely. The result is optimisation trajectories that diverge qualitatively across seeds, explaining the large run-to-run variance observed at low ω .

Step 4: the evaluation bottleneck. An additional subtlety is that the final energy evaluation itself uses importance sampling from the same proposal distribution. Even if the model has found an excellent wavefunction, a noisy evaluation can report a misleading energy.

The data expose this directly. Table 6.11 compares the VMC-best energy seen during training (a 20 000-sample Metropolis probe from the model’s own $|\Psi|^2$, used for checkpoint selection) with the final IS evaluation that uses the Gaussian proposal. At high ω where IS is reliable, the two estimates agree closely. At low ω , the gap between them grows, and its sign is unpredictable—the IS evaluation can overestimate or underestimate the true energy.

Table 6.11: Gap between VMC-best checkpoint energy (Metropolis sampling from $|\Psi|^2$) and final IS evaluation, for selected runs. Positive gap means IS reports higher (worse) energy than the VMC probe saw.

ω	Recipe	Seed	VMC-best (%)	Final IS (%)	Gap (%)
1.0	Baseline	314	+0.001	+0.060	+0.059
1.0	Robust	137	+0.002	+0.013	+0.011
0.5	Robust	42	+0.037	+0.002	−0.035
0.1	Robust	137	+0.136	+0.263	+0.127
0.01	Baseline	42	+0.320	+0.500	+0.180
0.001	Baseline	314	+4.114	+4.733	+0.619

The first row is particularly instructive: this baseline run with seed 314 achieved a VMC-best energy of +0.001%—essentially exact—during training, but the IS evaluation reports +0.060%. The wavefunction is excellent; the evaluation is noisy. This demonstrates that some nominally “bad” runs may contain excellent wavefunctions measured poorly, and some nominally “good” results may benefit from evaluation noise in the favorable direction.

Summary of the chain. The causal chain in Eq. (6.1) provides a unified explanation for the full landscape of collocation results. At high ω , the proposal matches the target well ($\hat{k} \sim 0.5$), gradients are reliable, optimisation is stable, and all seeds converge to near-DMC accuracy. At low ω , the proposal-to-target mismatch grows, $\hat{k}$ inflates, gradients become noisy, optimisation becomes seed-dependent, and the evaluation itself becomes unreliable. The robust recipe acts on multiple links of this chain simultaneously: higher oversampling improves the raw ESS, reward normalisation stabilises the gradient scale, rollback prevents catastrophic divergence, and the adaptive proposal reduces the fundamental mismatch. No single mechanism is sufficient; each addresses a different link in the chain.

Chapter 7

Architecture, optimiser, and paradigm through the tangent kernel

Our ansatz is a closed-shell Slater determinant dressed by two learnable objects: a *Jastrow*-type neural correlator W_θ , which reshapes the amplitude, and a coordinate *backflow* $\Delta\mathbf{x}$, which reshapes the nodes. Each can be built in one of two ways—without message passing, encoding and pooling per-particle and pairwise features in a single feed-forward pass (as in a DeepSet), or as a copresheaf *message-passing* (CTNN) network that maintains explicit node and edge features and transports information between them through learned maps before a quantity is read out (the correlator iterates this exchange over a V-cycle; the backflow applies a single round). The previous chapter showed that the message-passing ansatz reaches diffusion Monte Carlo (DMC) quality wherever a DMC reference exists ($\omega \geq 0.1$), and matches our own variational gold standard in the strongly correlated regime below it, where no DMC benchmark is available. That settles *whether* the ansatz works and leaves the question worth a chapter: *why* message passing helps at all. We answer it for both learnable objects through a single lens, and then turn the same lens on the optimiser and on the training paradigm.

One idea, one lens. The thesis of this chapter is that message passing buys one thing—a **relational channel**, a way to condition each particle’s contribution on where the others are—and that this single mechanism explains three superficially separate design choices. We read all three through the geometry of the per-sample log-derivative (score) matrix

$$O_{ki} = \frac{\partial \log |\Psi_\theta(\mathbf{x}_k)|}{\partial \theta_i}, \quad k = 1, \dots, B, \quad i = 1, \dots, P, \quad (7.1)$$

whose Gram matrices are the quantum geometric tensor (QGT / Fisher metric) $S = O^\top O / B$ and the neural tangent kernel $K = OO^\top$. The central diagnostic is the *effective dimension* of the variational tangent space, the participation ratio of the QGT spectrum $\{\lambda_a\}$,

$$d_{\text{eff}}(S) = \frac{(\sum_a \lambda_a)^2}{\sum_a \lambda_a^2}, \quad (7.2)$$

which counts how many independent tangent directions carry Fisher weight at the solution, regardless of the parameter count P . We read it alongside the local-energy variance $\text{Var}(E_L)$ (zero only for an exact eigenstate, hence a measure of wavefunction quality), the sampling efficiency (effective sample size ESS), and the condition number $\kappa(S)$.

The three questions. **Q1 (architecture).** What does a message-passing correlator/backflow learn that a separable one cannot? **Q2 (optimiser).** What does stochastic reconfiguration (SR, natural gradient) do to the update, and where does it beat Adam? **Q3 (paradigm).** What changes between $|\Psi|^2$ -sampled VMC and importance-sampled collocation?

Discipline. Cross-architecture d_{eff} comparisons use a *common probe set* (all models scored on the same configurations, not on each model’s own $|\Psi|^2$); we report only participation ratios converged in the sample count B ; and we anchor the tooling on the analytic two-electron ground state, where the ansatz reproduces $E = 3.000127$ Ha (+0.004%) with squared overlap 1.000000. Backflow comparisons

are seeded (≥ 2 seeds) and held at reference quality. Every relative error in this chapter depends on that reference, and an external DMC benchmark exists only for $\omega \geq 0.1$. In the strongly correlated regime below, we quote errors against our own converged variational energies (the $|\Psi|^2$ -sampled SR result), and, in the deep-Wigner cascade of §7.4, cross-check against the physical energy-scaling law rather than any ground truth. Relative errors below $\omega = 0.1$ are therefore measures of internal consistency, not of absolute accuracy, and we read them as such.

Headline findings. **(Q1a, Jastrow)** A message-passing correlator compresses the ground state into $\sim 2\times$ fewer effective tangent directions than a separable DeepSet at weak coupling (1.40 ± 0.17 vs 3.25 ± 0.03 at $\omega = 1$, $\sim 11\sigma$, seeded), and those directions are the physical collective modes of H (breathing, correlation-hole, relative excitation)—to $R^2 = 1.00$ at the exact $N = 2$ anchor, and $R^2 \gtrsim 0.93$ at $N = 6$ rising to ≈ 1.00 at $N = 12$ across the confinement range. A separable network of matched or larger capacity instead leaves nearly half its tangent budget along directions a compact physical dictionary does not resolve—already measurable at strong confinement, where the two networks are identical in energy. **(Q1b, backflow)** At the Wigner crystal the *conventional* backflow collapses to a single collective displacement mode (rank $10 \rightarrow 1$ at $N = 6$, $22 \rightarrow 1$ at $N = 12$) and its energy degrades to $+0.5\%$, while the message-passing backflow keeps full rank $\sim N$ and stays at $\sim 0.02\%$; the gain there is Coulomb (not kinetic) correlation, and the inter-particle messages account for the entire backflow contribution. Q1a and Q1b are not independent: holding the backflow fixed and varying only the correlator, its loss of physical alignment and the backflow’s rank collapse set in at the *same* confinement, pointing to a common relational bottleneck seen in the amplitude and in the nodes. **(Q2)** SR whitens the tangent kernel; its value tracks $\kappa(S)$ —negligible for the well-conditioned VMC metric, decisive for the ill-conditioned collocation objective. **(Q3)** VMC and collocation reach the *same state* when the importance sampling is healthy ($N = 2$: overlap > 0.997) and *diverge* when it is not ($N = 6$ Wigner: ESS $\rightarrow 0.11\%$ of the batch, overlap $\rightarrow 0$ under Adam). A cross-cutting caveat, itself a result: component-level quantities (rank, $|\Delta\mathbf{x}|$) are gauge-like; energy, overlap, and ablation are the invariants.

7.1 Q1 — Architecture: message passing supplies a relational channel

Message passing helps in both learnable objects, in two complementary ways that are the same statement about the tangent kernel: in the Jastrow it *compresses* the correlation into fewer effective directions; in the backflow it keeps the displacement from *collapsing* to a single direction. We take them in turn.

7.1.1 Q1a — In the Jastrow: message passing compresses the correlator

At $N = 6$, $\omega = 1$ the two Jastrow families are *energetically indistinguishable*: trained to DMC quality across a 20k–164k parameter ladder, every model lands within $\pm 0.1\%$ of the reference, and the best raw energy belongs to a 164k DeepSet—twice the CTNN parameter count. Total energy is a blunt discriminator; both have saturated the variational floor.

The difference is geometric. On a common probe set the CTNN reaches the ground state in $d_{\text{eff}} \approx 1.2$ – 1.7 (three seeds), while every DeepSet needs $d_{\text{eff}} \approx 2.9$ – 3.6 —a factor of ~ 2 . The gap is robust to the probe measure ($\lesssim 0.2$ whether scored on CTNN, DeepSet, or pooled points) and does not shrink as the DeepSet is given more capacity (the 164k DeepSet sits at 2.9, no lower than the 20k one). The compression is architectural, not a matter of size or sampling mass. It is also a well-posed comparison despite d_{eff} not being a reparameterization-invariant absolute dimension (§2.2.5): both families are compared under matched normalisation conventions and a common probe measure, so the factor-of-two gap reflects how each architecture spends its tangent directions, not a change of coordinates.

One subtlety should be dispatched here, because the synthesis (§7.5) will caution that quantities split between the Jastrow and the backflow are gauge-like. That caution does not reach this comparison: it is between Jastrow families with *no backflow present*, so there is no correlation to divide between two substitutable channels. What remains is a comparison of how two architectures spend their tangent directions on the same ground state, and it is stable across seeds and across the probe measure—a property of the correlator, not an accounting artefact.

Seeded, at weak coupling the separation is $d_{\text{eff}} = 1.40 \pm 0.17$ (CTNN) vs 3.25 ± 0.03 (DeepSet), $\sim 11\sigma$. The gap is a *weak-coupling* effect: it closes toward the Wigner crystal, where both families reach ~ 3.7




Figure 7.1: Message passing compresses the correlator (Q1a, $N=6$, $\omega=1$). *Left*: the effective rank of the quantum geometric tensor on a common pooled probe, converged in the number of probe samples. *Right*: the same pooled measure per model. The CTNN (blue) reaches the ground state in $d_{\text{eff}} \approx 1.2$ – 1.7 , while every DeepSet (red) needs ≈ 2.9 – 3.6 —and the gap does not close as the DeepSet is scaled from 20k to 164k parameters. The compression is architectural, not a matter of capacity.

but occupy *different* tangent subspaces, with the CTNN’s carrying the lower $\text{Var}(E_L)$. A single-seed dimension “inversion” at the crystal, reported in an earlier draft, did *not* survive seeding and is retracted here as a worked cautionary result (see §7.5).

Naming the manifold, at every size. The leading tangent directions are not abstract. At the exact $N = 2$ anchor an operator decomposition identifies them, to $R^2 = 1.00$, as the physical collective modes of H —a breathing mode (∂_ω), a correlation-hole mode, and the first relative excitation. This is not special to two bodies. Projecting the leading tangent mode of trained correlators onto a dictionary of physical operators (breathing/monopole, quadrupole, correlation-hole, pair-linear), seed-averaged over well-trained checkpoints (energy within 1.5% of reference), the CTNN reaches $R^2 \gtrsim 0.93$ at $N = 6$ and $R^2 \approx 1.00$ at $N = 12$ across the whole confinement range $\omega \in [0.01, 1]$, and both architectures reproduce the $N = 2$ naming at strong confinement (Fig. 7.2). The message-passing correlator finds the ground state by moving along the handful of coordinates the Hamiltonian actually couples.

The missing correlation channel is visible at *high* ω too. The leading-mode score is generous to the DeepSet: at $\omega = 1$ it is nominally the more physical of the two ($R^2 \approx 0.996$ vs. 0.991), with energies identical to four figures. Yet a correlation-specific probe exposes a deficit the leading mode hides. Write the eigenvalue-weighted fraction of the top tangent modes lying *outside* the physical-operator span as the *dictionary-orthogonal tangent mass*. The DeepSet carries a large excess at *every* confinement—0.44 against the CTNN’s 0.06 at $\omega = 1$, a gap of ~ 0.3 – 0.5 that persists from the strongly-confined limit down to the crystal (Fig. 7.3). The reading is that a separable network reaches the same energy—correlation being a small fraction of the total at strong confinement, so the energy looks converged—while leaving nearly half its tangent budget along directions a compact physical dictionary does not resolve. A direction outside the dictionary need not be unphysical; the robust statement is that the CTNN’s tangent space is far more completely accounted for by a small set of physical operators than the DeepSet’s. The deficiency is intrinsic and present throughout; weak confinement merely removes the disguise, because there correlation *is* the energy. At $N = 12$ the contrast is starker still: the CTNN’s dictionary-orthogonal mass stays $\lesssim 0.02$ across ω , while a matched DeepSet (trained to +0.2%) sits at ≈ 0.77 at $\omega = 1$ —the gap *widens* with system size rather than closing.

7.1.2 Q1b — In the backflow: message passing prevents rank collapse

The same relational channel is decisive, in a sharper form, in the backflow. Here we fix everything but the backflow architecture (conventional single-round messages vs. the copresheaf CTNN) and train both to a Woodbury-SR endpoint. At $\omega = 1$ both reach reference quality, but they diverge as the trap weakens (Table 7.1): the CTNN backflow holds $\sim 0.02\%$ error across all N, ω , while the conventional backflow




Figure 7.2: The leading tangent mode is a nameable physical mode. R^2 of the correlator’s leading tangent-kernel eigenmode onto the physical-operator dictionary, seed-averaged over well-trained checkpoints. The CTNN stays physical at every ω and both sizes; the DeepSet matches it down to $\omega \approx 0.1$ and then loses the leading mode entirely in the deep-Wigner regime. Three seeds at $\omega = 0.01$.




Figure 7.3: A limitation visible where the energy cannot see it. Dictionary-orthogonal tangent mass—the eigenvalue-weighted tangent variance outside the span of a compact physical-operator dictionary—versus ω . At $\omega = 1$ the two correlators are indistinguishable in energy (-0.001%) and in leading-mode physicality, yet the DeepSet already spends 44% of its tangent variance on directions not resolved by the chosen physical-operator dictionary, against the CTNN’s 6%. The excess is roughly constant in ω : the correlation deficit is intrinsic, disguised at high ω only because correlation is a small slice of the energy there.

degrades to +0.5% at $N = 6, \omega = 0.01$ and $\sim 6\times$ the CTNN error at $N = 20, \omega = 0.01$ (+0.69% vs. +0.12%).

Table 7.1: Q1b. Relative energy error (%), seed-averaged.

ω	$N = 6$		$N = 12$		$N = 20$	
	CTNN	conv	CTNN	conv	CTNN	conv
1.0	-0.03	+0.06	-0.01	+0.04	+0.06	+0.07
0.1	-0.05	+0.16	+0.01	+0.16	+0.02	+0.18
0.01	-0.04	+0.58	-0.01	+0.54	+0.12	+0.69

The energy gap has a sharp structural signature. Let the *backflow rank* be the participation ratio of the displacement field $\Delta\mathbf{x}$ (how many independent directions the backflow moves electrons along). At large ω both architectures use $\sim N$ modes. Toward Wigner the *conventional* backflow collapses to a single collective mode (rank $\rightarrow 1$), while the CTNN backflow retains full rank $\sim N$ (Table 7.2); the collapse sharpens with size ($10 \rightarrow 1$ at $N = 6$, $22 \rightarrow 1$ at $N = 12$, $35 \rightarrow 1$ at $N = 20$). The tangent dimension tells the same story from the other side: toward Wigner d_{eff} grows for CTNN ($\sim 4 \rightarrow 10$) and collapses for the conventional backflow ($\sim 6 \rightarrow 1$). The collapse to *exactly* rank 1 is seed-robust at every size, and the CTNN backflow holds full rank $\sim N$ throughout ($\sim 10/21/36$ at $N = 6/12/20$).

Table 7.2: Q1b. Backflow displacement rank (participation ratio), seed-averaged.

ω	$N = 6$		$N = 12$		$N = 20$	
	CTNN	conv	CTNN	conv	CTNN	conv
1.0	8.8	10.4	20.8	22.2	36.9	35.4
0.1	7.7	10.4	18.2	21.7	33.8	37.6
0.01	9.8	1.0	20.8	1.0	35.2	1.0

Why, mechanistically. Ablating the backflow ($\Delta\mathbf{x} = 0$, sampler re-optimised) and splitting the recovered energy into kinetic and Coulomb parts gives two seed-stable facts. (i) A **kinetic $\rightarrow$ Coulomb crossover**: at $\omega = 1$ the backflow buys kinetic energy; at $\omega = 0.01$ the gain is entirely Coulomb ($\Delta V_{\text{Coul}} \approx +0.017$, $\Delta T \approx 0$). “Message passing buys kinetic energy” is a high- ω statement only. (ii) **At Wigner the transport maps are the backflow**: within the CTNN, zeroing the inter-particle transport maps ($\rho_{v \rightarrow e}, \rho_{e \rightarrow v}$) costs as much as deleting the whole displacement. Both architectures condition each particle on the relative coordinates $\mathbf{r}_{ij}$, so neither is blind to inter-particle structure; the difference is that the conventional backflow uses a single round of pairwise messages aggregated per particle, whereas the CTNN maintains and transports explicit edge state across the graph. Empirically the single-round backflow does not *sustain* a full-rank relative displacement at the crystal—it collapses to a single collective displacement mode (rank 1)—while the CTNN’s copresheaf transport holds the full-rank response, and the ablation shows that this transport carries essentially the entire effect.

7.1.3 The unified reading of Q1

Q1a and Q1b are one statement about how relational information is *routed*. The baselines are not blind to inter-particle structure—the DeepSet pools independently-encoded pairs, and the conventional backflow aggregates a single round of pairwise messages—but neither maintains a persistent, transported edge representation: each pairwise term is computed and discarded. The copresheaf CTNN does maintain one, and moves it back and forth between nodes and edges through learned transport maps ($\rho_{v \rightarrow e}, \rho_{e \rightarrow v}$); the message-passing correlator iterates this exchange over a V-cycle, while the backflow applies a single such copresheaf round. Where a compact representation exists this *compresses* the correlation onto fewer tangent directions (Jastrow: smaller d_{eff}); where the physics demands a genuinely collective, full-rank response the single-round baseline instead collapses to a single collective displacement mode (backflow: rank 1), which the copresheaf transport avoids. Both effects are visible only in the tangent-kernel geometry, not in the total energy, until weak confinement makes the geometric deficit an energetic one.

The two channels fail together. That last sentence is not a metaphor: the correlator deficit and the backflow deficit collapse at the *same* confinement. Holding the backflow architecture fixed (the conventional network, in both cases) and varying only the correlator, we track the correlator’s correlation-hole




Figure 7.4: What message passing buys, by ablation (trained networks, $N=6$; energy relative to the full model). Removing the backflow (orange) costs almost nothing deep in the Wigner regime ($\omega=0.01$) but $\sim 6\%$ at $\omega=1$, while removing the Jastrow messages (red) or all message passing (blue) is expensive throughout. The gain from message passing is a high- ω , kinetic-energy statement; at the crystal the relational work has moved into the backflow’s transport maps.

capture and the backflow’s displacement rank down the ω -sweep (Fig. 7.5). With a DeepSet correlator both give way together at $\omega \approx 0.05$ —the hole capture falls from 0.68 to 0.07 as the backflow rank drops from ~ 10 to 1. With a CTNN correlator both hold until $\omega \approx 0.01$. The correlator’s grip on the physical correlation modes and the backflow’s ability to sustain a full-rank displacement are the same capability seen from two sides: when the amplitude model can no longer resolve the correlation hole, the nodal deformation can no longer stay high-rank, and a better correlator postpones *both*. The evidence—coincident transitions that shift together under a correlator swap—supports a common relational bottleneck underlying Q1a and Q1b rather than proving them literally identical.

or physicality and backflow rank collapse together




Figure 7.5: Correlator physicality and backflow rank collapse together ($N = 6$; the backflow architecture is held fixed, only the correlator changes). For each correlator, the correlation-hole capture (left axis) and the backflow displacement rank (right axis) give way at the same ω : $\omega \approx 0.05$ for the DeepSet, $\omega \approx 0.01$ for the CTNN. A correlator that keeps its physical structure lets the backflow keep its rank.

7.2 Q2 — Optimiser: SR earns its keep where the metric is

ill-conditioned

Stochastic reconfiguration preconditions the gradient by the inverse QGT, $\theta \leftarrow \theta - \eta S^{-1} \nabla_{\theta} E$; geometrically it *whitens* the tangent kernel. At $N = 2$ this is exact: the SR step aligns with the imaginary-time direction to numerical precision. The question is where whitening changes the outcome.

Fixing the ansatz and varying only the optimiser from a common stage-3 base: **for $|\Psi|^2$ -VMC, SR and Adam are indistinguishable**—exact at $N = 2$ ($\text{var} \sim 10^{-9}$), and at $N = 6$ the energy gap $E_{\text{Adam}} - E_{\text{SR}}$ is $\lesssim 0.002$ across ω and both seeds. The VMC metric is well enough conditioned ($\kappa(S)$ moderate, and $|\Psi|^2$ sampling is self-preconditioning) that Adam already moves along the right directions. **For weak-form collocation, SR is essential**: there $E_{\text{Adam}} - E_{\text{SR}}$ is large and positive (e.g. +2.9 at $N = 6, \omega = 1$), because the importance weights make the objective badly conditioned and plain Adam fails to converge it.

The verdict is therefore not “SR wins” or “SR ties” but **SR matters when the tangent metric is ill-conditioned and still estimable**. Its value is a function of $\kappa(S)$, set by the ansatz and the sampling measure, not an intrinsic property of the optimiser—which is exactly why an earlier low- ω VMC sweep saw a modest SR advantage that never grew into a decisive win.

The claim needs one sharpening, because “more ill-conditioned $\Rightarrow$ more useful SR” cannot hold without bound: inverting S requires S itself to be *reliably estimated*. SR helps when the tangent metric is anisotropic but its empirical estimate is trustworthy; once the sampling that builds S collapses—as the importance weights do deep in the Wigner regime—the empirical S becomes noisy and rank-deficient, and preconditioning by its inverse amplifies that noise rather than the signal. This is exactly the behaviour seen at ultra-weak confinement, where CG-SR on the collocation objective becomes unstable (Chapter 6, §6.4.4): not a counterexample to the conditioning story but its completion—SR pays off in the window where the metric is both ill-conditioned and estimable, and fails once estimation itself breaks down.

One further boundary deserves to be drawn explicitly. The VMC equivalence of Adam and SR is established here at $N = 2$ and $N = 6$; we have not verified that it survives to $N = 12$ and $N = 20$, where the conditioning of the VMC metric could in principle worsen with dimension. The mechanism we propose—that $|\Psi|^2$ sampling keeps $\kappa(S)$ moderate and so lets Adam’s diagonal preconditioning suffice—is a prediction there, not yet a measurement, and testing it is the obvious extension.

7.3 Q3 — Paradigm: the sampling measure sets conditioning, variance, and reach

Between $|\Psi|^2$ -VMC and importance-sampled collocation the wavefunction family is identical; only the measure that defines the loss changes. Three consequences, one statement.

$|\Psi|^2$ is self-preconditioning; the covering proposal is not. Sampling from $|\Psi|^2$ makes the estimator of S and of the gradient share the same measure, which conditions the update. A fixed covering proposal instead drives a monotone *ESS collapse* as N and $1/\omega$ grow: at $N = 6, \omega = 0.01$ the effective sample size falls to 0.11% of the batch (about 5 of 4096 points). When ESS collapses the gradient estimator is degenerate—a hard limit, not undertraining.

Same state, until the sampling loses the target. We test whether the two paradigms reach the same *state* (not merely the same energy) via the symmetric overlap $\mathcal{S}^2 = \langle \Psi_B / \Psi_A \rangle_{|\Psi_A|^2} \langle \Psi_A / \Psi_B \rangle_{|\Psi_B|^2}$. At $N = 2$ they are identical ($\mathcal{S}^2 > 0.997$ for every ω , both optimisers, both seeds): the paradigm is efficiency-only, collocation paying $\sim 5\times$ the variance for the same wavefunction. At $N = 6$ they *diverge*: under Adam $\mathcal{S}^2$ falls $0.85 \rightarrow 0.77 \rightarrow \approx 0$ from $\omega = 1$ to 0.01 (both seeds), tracking the ESS collapse; SR partially rescues it ($\mathcal{S}^2 \approx 0.95$) but does not close the gap. Weak-form collocation thus has a **domain of validity**—equivalent to VMC when the importance sampling is healthy, degrading as correlation strengthens. This couples Q3 to Q2: the paradigm that needs SR is the one that breaks down at scale. An adaptive proposal that tracks $|\Psi|^2$ is the natural remedy, pursued in §7.4.




Figure 7.6: The naive collocation proposal collapses in the Wigner regime ($N=6$). The effective sample size (as a percentage of the batch) falls by more than an order of magnitude from $\omega=1$ to $\omega=0.01$, where it reaches $\sim 0.1\%$ —about five usable points in a batch of 4096. This is a property of the sampling measure, not of optimisation, and it is what the physics-informed ring proposal (§7.4) is designed to repair.

Why the collocation energy readout is biased. For an exact eigenstate the local energy $E_L = \hat{H}\Psi/\Psi$ is constant over configuration space, so $\text{Var}(E_L) \rightarrow 0$ as the ansatz approaches the ground state—the zero-variance principle, and the reason $\text{Var}(E_L)$ is a sharp quality measure. That cancellation is a property of the *full* local energy, in which the kinetic (Laplacian) and potential terms combine to a constant; it is not a property of the Laplacian term alone. The collocation objective does *not* discard the Laplacian: it is a score-function (REINFORCE) estimator that *forms* the full E_L as a detached reward and differentiates only $\log|\Psi|$ (§6.4.1). What it gives up is the *measure*: the reward is averaged over the analytic proposal with importance weights (and, in the hardest regimes, tempered weights), so the importance-sampled energy *readout* is a biased estimate of $\langle E_L \rangle_{|\Psi|^2}$ and must not be used as the convergence criterion. The reliable criteria are $\text{Var}(E_L)$ and the state overlap, both evaluated on $|\Psi|^2$ samples.

7.4 The collocation frontier: a physics-informed proposal for the Wigner regime

The ESS collapse of §7.3 is a limitation of the *proposal*, not of collocation itself, and it can be substantially pushed back. This section reports a physics-informed proposal that extends MCMC-free collocation deep into the Wigner regime, the mechanism that limits it, and the wall that remains.

7.4.1 The origin-centred proposal misses the Wigner shell

Every proposal in the standard pipeline is a mixture of *origin-centred* Gaussians $q(\mathbf{R}) = \frac{1}{K} \sum_k \mathcal{N}(\mathbf{R}; 0, \sigma_{f,k}^2/\omega)$. But as $\omega \rightarrow 0$ the electrons localise into a Wigner molecule on a *shell* far from the origin: the classical $N = 6$ minimum is the (1,5) configuration—one central electron, five on a ring at radius $1.334\omega^{-2/3}$ [8, 9]—which in oscillator units sits at $1.96/2.87/4.22\ell$ for $\omega = 0.1/0.01/0.001$, moving outward as the trap weakens, past the reach of a width- 2ℓ origin Gaussian. The proposal aims where $|\Psi|^2$ has no mass; hence the collapse.

7.4.2 The Wigner-ring proposal

We replace the origin mixture with a *Wigner-ring* proposal: a radial Gaussian at the shell radius times an angular distribution for the ring electrons, plus a central Gaussian. On trusted states this lifts the effective sample size by 15–47 $\times$ over the origin mixture (Table 7.3), turning a collapsed sampler into a

usable one—without any training, a cheap decisive test. Fitting the ring to $|\Psi|^2$ samples (the simplest adaptive form) doubles the ESS again; a $(1, 5) + (0, 6)$ mixture *hurts* at $N = 6$, since the state is $(1, 5)$ and the extra channel dilutes the proposal mass.

Table 7.3: ESS (of 4096) against a trusted $N = 6, \omega = 0.01$ state. The origin mixture is already collapsed; the ring proposal covers the shell.

proposal	ESS
origin-Gaussian mixture (standard)	1.5
Wigner ring $(1, 5)$, hand-tuned	22.5
Wigner ring $(1, 5)$, fitted to $ \Psi ^2$	47.5

7.4.3 Angular crystallisation is the deep-Wigner wall

Why does the ring work at $\omega = 0.01$ but fail at $\omega = 0.001$? The molecule *crystallises angularly* as the trap weakens. The nearest-neighbour angular-gap standard deviation of the five ring electrons is $40.8^\circ/33.3^\circ/25.5^\circ/6.7^\circ$ at $\omega = 1/0.1/0.01/0.001$: a “ring liquid” at moderate coupling becomes an angular crystal (electrons pinned near 72° apart) at $\omega = 0.001$. A proposal with independent, uniform angles almost never lands on the crystalline configuration—so the angular distribution must crystallise too. This is captured MCMC-free by modelling the five gaps (which sum to 2π) as Dirichlet(α): an exact tractable density whose single concentration α interpolates liquid (α small) to crystal (α large), fitted to the measured gap variance. This diagnosis *explains* the collocation wall that is otherwise only observed.

7.4.4 An MCMC-free cascade reaches $\omega = 0.0035$

Combining the ring proposal (radial Gaussian $\times$ Dirichlet gaps) with importance oversampling, a *continuous* adaptive re-fit of the proposal to the current importance-weighted batch (every few steps, no MCMC), and a gentle ω -cascade seeded from a converged $\omega = 0.01$ state, MCMC-free collocation produces states that lie on the Wigner energy-scaling curve down to $\omega = 0.0035$ —four cascade rungs, with heavy-VMC energies within 1–2% of the scaling estimate (Table 7.4), well past where the origin proposal collapses (ESS ~ 1 even at $\omega = 0.01$). The essential ingredient is the continuous re-fit: it keeps ESS healthy ($\sim 10^2$) *while* the energy descends, where a static or slowly-updated proposal collapses to ESS ~ 1 as the wavefunction moves away from it.

Table 7.4: MCMC-free Wigner-ring cascade, $N = 6$: heavy-VMC energy vs. the Wigner-scaling estimate $E(\omega) \approx 0.690 (\omega/0.01)^{0.689}$. Good to $\omega = 0.0035$.

ω	E	scaling est.	ratio
0.0077	0.577	0.577	1.00
0.0059	0.481	0.480	1.00
0.0045	0.400	0.398	1.01
0.0035	0.340	0.335	1.02

7.4.5 The remaining wall: a fine-tuned Slater–Jastrow balance, not the sampler

Below $\omega = 0.0035$ the cascade becomes unstable, and—informatively—the instability is *non-monotonic* across proposal variants: wide-exploratory, tempered-tracking, and outward-anchored proposals each converge some rungs and diverge others. That pattern locates the obstacle not in the proposal but in the warm-start, and four transfer diagnostics (taking the converged $\omega = 0.0035$ rung down to $\omega = 0.0027$, a 23% step) pin down its mechanism precisely.

First, the frontier state is genuine: at its own $\omega = 0.0035$ the rung is a bona-fide Wigner molecule, with radial density peaked at 3.34ℓ (classical ring 3.42ℓ) and a *single* basin—initialising the sampler on the shell converges to the same 3.34ℓ . Transferred to $\omega = 0.0027$, however, the wavefunction collapses to $\sim 1\ell$, and two tests show *where* the collapse lives. It is not the sampler: initialising the Metropolis walkers directly on the classical shell (3.57ℓ) still relaxes inward to 1.1ℓ , so the collapsed density is a property of the ansatz, not of sampler initialisation. And it is not the backflow: its displacement is small

($|\Delta \mathbf{x}| \approx 0.13\ell$, radially inward) and essentially *preserved* across the transfer ($0.15\ell \rightarrow 0.13\ell$)—it scales correctly with ℓ but is far too small to bridge the 2.4ℓ gap between the Slater core ($\sim 1\ell$) and the Wigner shell.

The mechanism is a delicate cancellation. The bare harmonic-oscillator determinant suppresses the shell by $\exp(-\omega r^2/2) \sim \exp(-5.6)$ in amplitude ($\sim 10^{-5}$ in density); the Wigner peak exists only because the Jastrow amplifies it back by a comparable factor, with the backflow contributing a small inward nudge. A 23% change in ω perturbs both the Slater confinement and the Jastrow correlation length, the fine-tuned balance tips, and the density falls back to the non-interacting shell—from which the collocation re-fit cannot reliably climb out (hence the non-monotonic proposal behaviour). This rules out the natural fix of an ω -robust warm-start by coordinate rescaling: because the ansatz relaxes inward even when handed shell configurations, no coordinate transform can preserve the state. Reaching $\omega < 0.0035$ would require per- ω re-optimisation against a shell-*amplitude*-anchored objective, or a genuinely ℓ -covariant parametrisation—a distinct problem left to future work. The contribution here is concrete: a physics-informed proposal that carries MCMC-free collocation from the origin-proposal wall down to a verified Wigner molecule at $\omega = 0.0035$, and a precise account of the angular-crystallisation and Slater–Jastrow-balance effects that bound it.

7.5 Synthesis and discussion

The three axes are one geometry. Architecture sets the shape of the tangent space (message passing compresses it in the Jastrow, and keeps it from collapsing in the backflow); the optimiser inverts the metric on that space (SR pays off only when it is ill-conditioned); and the paradigm chooses which measure defines the metric ($|\Psi|^2$ self-preconditions and stays on target, while the covering proposal loses conditioning and eventually the state). The single cross-cutting discriminator, robust where dimension and rank are not, is $\text{Var}(E_L)$, lower for the message-passing ansatz at every ω . What this geometry means for the physics, and how the threads combine, is taken up in the Discussion (Chapter 8) and Conclusion.

A gauge caveat, itself a result. Component-level quantities—backflow rank, displacement magnitude $|\Delta \mathbf{x}|$, per-component d_{eff} —are **gauge-like**. The Jastrow and backflow are substitutes, so how correlation is split between them is a nearly-flat, path-dependent direction of the loss; those quantities depend on the training route, not on the state. Every claim that reversed during this project was of that kind (the retracted d_{eff} inversion at the crystal; a mislabelled backflow; an artefactual rank collapse from joint-from-scratch training). The reliable measurements—energy, state overlap, and ablation (delete a component, price it in energy)—are invariant to the gauge, and the conclusions above rest only on those.

Part V

Discussion

Chapter 8

Discussion

Convictions are more dangerous enemies of truth than lies.

FRIEDRICH NIETZSCHE

Human, All Too Human, Vol. I, aphorism 483

This chapter asks what the preceding results mean, which is a different question from what they are. Four threads run through it, and they are ordered by how much interpretation they require: the energies, where an external benchmark does part of the arguing for us; the Wigner regime, where the trained state is used as an instrument and the burden of proof shifts onto internal consistency; the learned representations, where the quantities are informative but partly gauge-dependent and must be read with that in mind; and the collocation programme, where the central claim—that a Markov chain is not necessary—needs its boundaries drawn as carefully as its successes.

8.1 Discussion of energy results

The energies are the least interesting result in this thesis and the most necessary one. They establish that the ansatz works; everything downstream—every claim about what the network learned, every structural diagnostic read off the trained state—is worthless if it does not. So the question to press here is not whether the numbers are good, but whether they are good *for the right reason*, and whether the regimes where no benchmark exists inherit any of that credibility.

The benchmarks in Tables 6.1 and 6.2 show that the Slater–Jastrow PINN, with and without backflow, achieves DMC-level accuracy for two-electron dots and maintains sub-0.1% errors for $N \in \{6, 12\}$ across a wide range of trap frequencies. At the same time, it extrapolates in a controlled way into ultra-weak traps ($\omega = 10^{-3}$) where no DMC data are available. That last clause carries the weight, and we return to what it can and cannot support below.

Two electrons: architecture test and SR refinement. For $N=2$, residual-only training provides a clean test of the bare architecture. Across $\omega \in \{1.0, 0.5, 0.1, 0.01\}$, both the plain PINN and PINN+BF track DMC references at the $\sim 10^{-4}$ Ha level or better (Table 6.1), with deviations fully compatible with quoted uncertainties except at the shallowest pretraining point $\omega = 0.01$. There, residual-only PINN+BF remains variational but sits $\sim 3 \times 10^{-4}$ Ha above DMC, indicating a small residual bias from the PDE-based training alone.

The SR tail in Table 6.2 removes this bias. For $N=2$ and all $\omega \geq 0.01$, the SR-refined PINN+BF energies are statistically indistinguishable from DMC, with relative deviations $\lesssim 10^{-3}\%$. Thus, once the ansatz is well conditioned, a short SR phase suffices to close the last few 10^{-4} Ha of variational gap without changing the qualitative structure learned during residual training.

The comparison between PINN and PINN+BF in Table 6.1 also confirms that backflow is *not necessary* to reach DMC accuracy for two electrons. Energies with and without backflow agree within error bars at all traps, and the only visible effect of backflow in this case is a mild variance reduction and smoother optimisation in shallow traps. For $N=2$ the nodal structure is adequately captured by the Slater–Jastrow correlator alone.

Scaling to $N=6$ and $N=12$. The more demanding tests are the many-electron dots in Table 6.2. For $N \in \{6, 12\}$ and the traps with DMC references ($\omega \in \{0.1, 0.5, 1.0\}$), the final PINN+BF energies are consistently close to DMC and always slightly higher, as required by variationality. Relative deviations fall in the band 0.026–0.048%, corresponding to absolute errors of order 10^{-3} – 10^{-2} Ha for the interacting cases, with no sign of catastrophic growth in error with either particle number or trap strength. The worst relative deviations occur at intermediate ω and remain safely below the percent level.

One subtlety matters here, because the message-passing (CTNN) ansatz occasionally sits *below* the quoted DMC reference (for example $N=12$, $\omega=1$: 65.6956 vs. 65.7001 Ha). This does not violate the variational principle, which bounds the energy only from below by the *exact* ground state, $E_{\text{VMC}} \geq E_0$. The DMC benchmarks are *fixed-node* calculations and therefore carry their own nodal (systematic) bias; a neural ansatz with a better nodal surface can legitimately lie below a fixed-node DMC value obtained with poorer nodes. We therefore do not treat sitting above the published DMC number as a requirement, and where the CTNN falls below it we read this as a plausibly improved nodal structure rather than an error — though establishing that rigorously would require the DMC trial nodes and a matched uncertainty analysis, which we have not undertaken.

This is non-trivial for two reasons. First, the grid spans qualitatively different physical regimes: compact, kinetic-dominated states at $\omega = 1.0$ and swollen, interaction-dominated states at $\omega = 0.1$. Second, the same architecture and training protocol are used across all (N, ω) , aside from trivial scaling in trap units. The fact that a single Slater–Jastrow+backflow ansatz maintains sub-0.1% accuracy across this grid suggests that the chosen combination of (i) trap-unit normalisation, (ii) soft-core pair features, and (iii) an explicit analytic cusp is robust with respect to both particle number and confinement.

Ultra-weak traps and extrapolation to $\omega = 10^{-3}$. At $\omega = 10^{-3}$ there are no DMC references in Table 6.2, so the PINN+BF values ($E \approx 0.0138$ Ha for $N=2$, 0.1409 Ha for $N=6$, 0.5158 Ha for $N=12$) should be viewed as variational predictions rather than benchmarked numbers. Several internal checks, however, support their reliability.

First, the energies connect smoothly to the DMC-validated points at $\omega = 0.01$ under the log–log scaling analysis of T , V_{int} and V_{trap} in Sec. 6.3, with exponents and virial ratios consistent with classical expectations in the Wigner regime. Second, for $N=2$ at $\omega = 10^{-3}$, independent MCMC diagnostics (stable r_{12} and $P(r)$ histograms, smooth local-energy distributions, consistent block estimates) indicate that sampling noise is well below the quoted uncertainties in Table 6.2. Third, for $N=6$ and $N=12$, the structural and energy-based diagnostics in Sec. 6.3 confirm that the states lie on the Wigner side of the crossover (correct shell topologies, near-classical virial ratios), with no indication of a mis-set energy scale.

Within these caveats, the $\omega = 10^{-3}$ energies can be regarded as working variational anchors, supported by the internal checks above, for the Wigner-molecule analysis in Sec. 6.3 and for the representation analysis in Sec. 6.2.

Efficiency and conditioning. From an algorithmic point of view, the benchmarks highlight the importance of conditioning. The SR tail uses only $\mathcal{O}(10^3)$ iterations with $\sim 3 \times 10^3$ samples each, i.e. a few million local-energy evaluations per (N, ω) point. This is modest compared to some earlier neural approaches to quantum dots that require orders of magnitude more samples to reach similar accuracy.

The key design choices are: (i) working in trap units, which keeps typical coordinates $\mathcal{O}(1)$ from tight to shallow traps; (ii) soft-core pair features with $ds/dr \rightarrow 0$ at coalescence, which control gradients and Laplacian spikes near $r_{ij} = 0$; and (iii) an explicit analytic cusp that imposes the correct short-range behaviour analytically rather than learning it numerically. Together, these keep local energies numerically tame even in ultra-weak traps where electrons are far apart and the wavefunction varies on long length scales.

Energetic role of backflow. Finally, the tables clarify the energetic role of backflow across regimes. For $N=2$, backflow is not needed to attain DMC-level accuracy. For $N \in \{6, 12\}$ at intermediate traps ($\omega \in \{0.1, 0.5\}$), backflow provides a modest but systematic improvement over residual-only baselines (not shown in Table 6.2), bringing final errors into the $\mathcal{O}(10^{-2}$ – $10^{-1})\%$ band. In the deepest Wigner cases ($\omega = 10^{-3}$), however, the backflow correction becomes energetically tiny (Sec. 6.2): the Slater–Jastrow correlator already captures the dominant structure and the optimiser effectively *turns off* backflow. This regime dependence is mirrored in the representation analysis, where backflow appears as a high-dimensional, predominantly local correction channel that is only variationally active away from the ultra-classical limit.

8.2 Discussion of the Wigner–molecule regime

Having established the wavefunction’s accuracy wherever a benchmark exists, we now use it to probe Wigner-molecule structure. Everything up to this point asked whether the trained state is accurate; here we take that as given and ask what it can be used to see. That shift raises the evidential stakes considerably, because it is exactly the regime in which no DMC reference exists—so any structural claim has to be carried by internal consistency, by agreement between independent diagnostics that could easily have disagreed, and by the fact that the answers reproduce known classical topologies nobody trained the network to reproduce. The phrase “strongly correlated” is a label; what follows is an attempt to say something falsifiable instead.

The structural diagnostics in Sec. 6.3 show that the weakest traps considered place the dots firmly on the Wigner side of the crossover, in a sense that goes beyond that loose label. Three strands of evidence support this: (i) localization and energy partitioning, (ii) shell geometry and bond order, and (iii) topology statistics and reconstruction of $g(r)$.

A caveat on total spin. Our reference states are the closed-shell singlets of the non-interacting problem, with $N_\uparrow = N_\downarrow$ held fixed throughout. This is safe at weak coupling, but in the deep-Wigner regime the ground-state *spin* can change: for parabolic dots the interaction can drive spin transitions as it strengthens (for instance an $S = 0 \rightarrow S = 1$ change for $N = 6$ at strong coupling) [7]. Fixing $N_\uparrow = N_\downarrow$ does not by itself pin the total spin, since an $S = 1$ state also has an $S_z = 0$ component. We do not scan total spin here, so the weak-trap energies and structures below should be read as the lowest states we find *within the fixed spin-projection sector*, not as proven global-spin ground states. Determining the true spin ordering across the crossover, by comparing sectors of different S , is left to future work.

Localization and energy partitioning. On the one-body level, the weak-trap radial laws $P(r)$ are sharply peaked at large radii with relatively small fluctuations. For the two-electron dot at $\omega = 10^{-3}$ we find $r_{\text{mode}} \approx 64.3$, $\langle r \rangle \approx 64.0$ and $\gamma = \sigma_r / r_{\text{mode}} \approx 0.29$, with only $\sim 4.6 \times 10^{-3}$ of the single-particle mass inside $r \leq 0.25 r_{\text{mode}}$ (Table 6.6). The pair distribution $p(r_{12})$ is peaked at $r_{12}^{\text{mode}} \approx 122$ with Lindemann ratio $\gamma_{r_{12}} \approx 0.20$, and the relative angle is sharply anti-aligned, $\sigma(\pi - \Delta\phi) \approx 0.35$, with $p(r_{12} \leq 0.25 r_{12}^{\text{mode}}) \approx 2.6 \times 10^{-3}$. These numbers are characteristic of a dilute, essentially two-site Wigner dimer rather than a delocalized Fermi liquid.

For $N \in \{6, 12\}$ the same pattern appears in the aggregate. Both r_{mode} and r_{rms} grow roughly as $\omega^{-0.6}$, the localization ratio γ decreases only mildly with N and ω , and the extracted $\langle r^2 \rangle \propto \omega^{-\beta}$ exponents $\beta \simeq 1.2\text{--}1.3$ exceed the non-interacting value $\beta = 1$. At the same time, the energy partitioning approaches the classical virial limit: at $\omega = 10^{-3}$,

$$\frac{2\langle V_{\text{trap}} \rangle}{\langle V_{\text{int}} \rangle} \simeq 1.25, 1.12, 1.02 \quad \text{for } N = 2, 6, 12,$$

with the $N=12$ dot satisfying $V_{\text{int}} \approx 2V_{\text{trap}}$ to within 2% (Fig. 6.4). Combined with the log–log scaling exponents $\alpha_T \approx 1$ and $\alpha_{V_{\text{int}}} \approx 0.66$, this indicates that the kinetic energy has become a small correction on top of a nearly classical potential-energy landscape, as expected in the Wigner-molecule limit.

Shell geometry and bond order. Beyond global localization, the weak-trap states exhibit robust shell structure with non-trivial bond order. For $N=6$ the dot retains a single pronounced shell across all $\omega \in [1, 10^{-3}]$; what evolves is the stiffness of angular correlations. The ring Lindemann index decreases from ~ 0.70 at $\omega = 1$ to ~ 0.64 at $\omega = 10^{-3}$, while the pair-distance Lindemann ratio drops from ~ 0.35 to ~ 0.27 (Table 6.7). After angular registration, the electrons form a rotating Wigner ring with clear sixfold order.

At $\omega = 10^{-3}$, the $N=6$ ensemble is dominated by two symmetry classes, (1, 5) and (0, 6), with fractions ≈ 0.801 and ≈ 0.199 . Both show strong ring order, with $|\Phi_5| \approx 0.60$, $|\Phi_6| \approx 0.66$ and ring Lindemann indices ≈ 0.21 . The shell itself never breaks; quantum fluctuations manifest primarily as rotations and occasional transitions between (1, 5) and (0, 6). This matches classical analyses where (1, 5) is the ground-state geometry and (0, 6) a proximate competitor.

For $N=12$ the structure is richer but equally ordered. At $\omega = 10^{-3}$ the two-shell fraction is very high for a broad range of radial-gap thresholds, confirming that shelling is robust rather than a threshold artifact. Within the two-shell subset the inner-ring occupancy histogram is sharply peaked at (3, 9),

with (2, 10) and (1, 11) as the leading subdominant topologies. On (3, 9) frames we find $|\Phi_3| \approx 0.69$, $|\Phi_9| \approx 0.53$ and Lindemann indices ≈ 0.22 on both rings, i.e. clear threefold and ninefold bond order with small angular fluctuations. This is precisely the commensurate, multi-shell crystalline order expected for a twelve-electron Wigner molecule.

Topology statistics and reconstruction A more stringent test of the Wigner picture comes from shell-resolved reconstruction of the pair distribution. For $N=12$ at $\omega = 10^{-3}$ we partition two-shell configurations into II/IO/OO pairs, form their individual histograms and weight them by the observed topology mixture (0.045:0.409:0.545). The resulting synthetic $g(r)$ has cosine similarity 0.9982 to the global $g(r)$ (Fig. 6.7), i.e. is indistinguishable at our statistical resolution. The same holds for $N=6$, where the global $g(r)$ is reproduced essentially exactly by an OO/IO mixture of (1, 5) and (0, 6) sectors.

Two implications follow, and the first needs its scope stated precisely. Because the sector histograms and the global $g(r)$ are built from the *same* samples, this is a test of *decomposability*, not of sampling correctness: it shows that once the shell occupancies and topology statistics are fixed, the global $g(r)$ is determined by shell geometry and combinatorics alone, leaving no residual structure unaccounted for. It cannot, and does not, rule out a sampling artefact common to both—that case rests independently on the restart, threshold and chain diagnostics. Second, subject to that scope, the ansatz is not only reproducing the energy but also the detailed two-body correlation structure expected from classical and quantum studies of parabolic Coulomb clusters.

From dimer to multi-shell Wigner molecules. Viewed across $N = 2, 6, 12$ and decreasing ω , a coherent Wigner-molecule picture emerges. The two-electron dot evolves into a dilute, strongly anti-aligned dimer with negligible overlap near the origin. The six-electron dot forms a single, stiff Wigner ring whose lab-frame symmetry is restored by rotation but whose co-rotating frame exhibits clear sixfold order and a quantum mixture of classically known (1, 5) and (0, 6) topologies. The twelve-electron dot, finally, develops a robust two-shell structure with a commensurate (3, 9) split as the modal configuration and topology fractions that track classical ground-state expectations.

All of this structure is obtained from samples drawn from the same variational neural ansatz whose energies match or closely track DMC benchmarks. This gives strong evidence that the states analysed in Sec. 6.2 are bona fide Wigner molecules, and not artefacts of an over-flexible wavefunction. The next section discusses how this rich real-space structure is compressed into a remarkably low-dimensional latent geometry by the learned correlator, and how backflow behaves across the same regimes.

8.3 Discussion of learned representations

There is no guarantee that a trained network organises itself in a way anyone can interpret. High-dimensional function approximators are entitled to be incomprehensible, and frequently are. So the finding that structures this section is a surprising one: the internal organisation is not merely interpretable but *simple*, and simple in a way that matches the physics rather than the parameterisation. The qualification, which follows immediately and matters, is that not every quantity we can measure on this state is a property of the state.

The representation analysis in Sec. 6.2 reveals a surprisingly simple internal organization of the ansatz: the correlator W_θ lives on a low-dimensional manifold whose leading axes track global size and fluctuations of the electron cloud, while the BackflowNet implements a high-dimensional but predominantly local correction whose net energy becomes small in the deepest Wigner regimes. This mirrors the physical picture developed in Sec. 6.3. As established in §7.5, component-level quantities are gauge-like; the conclusions drawn below therefore rest on the invariants—energy, overlap, and ablation.

Low-dimensional correlator manifold. Across all (N, ω) , the entropy effective rank $r_{\text{eff}}(Z)$ of the correlator features stays low across all (N, ω) , spanning roughly 1 to 3 (Table 6.3). In the most extreme case, $N=2$ at $\omega = 10^{-3}$, we find $r_{\text{eff}}(Z) \approx 1.00$ with head-PC1 correlation ≈ 0.998 and PC1 block power almost entirely in the pair branch ψ . PC ablations show that projecting onto the top- k PCs rapidly recovers the head, while random k -dimensional subspaces leave most of the error intact: for $(N, \omega) = (2, 10^{-3})$ the relative MAE drops from $\sim 4.5 \times 10^{-2}$ with $k=1$ to $\sim 2.0 \times 10^{-5}$ with $k=12$, whereas random projections of the same dimension do not achieve comparable reductions.

For $N \in \{6, 12\}$ and for both quantum and Wigner regimes, the picture is similar. Effective ranks cluster between ~ 1.3 and ~ 2.5 , head-PC1 correlations remain $\gtrsim 0.95$, and a small number of principal directions captures the bulk of sensitivity. From a representation-learning perspective, the correlator behaves more like a low-dimensional order-parameter model than like a generic high-dimensional feature extractor.

What the leading axes encode. Linear probes from the correlator PCs to coarse physical summaries clarify what these axes represent. For $N=2$ (both at $\omega = 0.01$ and $\omega = 10^{-3}$), the first few PCs predict the mean radius and its variance essentially perfectly ($R^2(r_{\text{mean}}) \approx 1$, $R^2(r_{\text{var}}) \approx 1$), and even near-origin masses such as $\text{Pr}(r \leq 0.25 r_{\text{mode}})$ are accurately captured. For $N \in \{6, 12\}$, the leading PCs still track global size with $R^2(r_{\text{mean}})$ close to unity and radial fluctuations with $R^2(r_{\text{var}})$ in the range 0.4–0.85 across regimes, while shell contrast and angular order remain only weakly linearly encoded. This is consistent with the structural analysis: the correlator primarily encodes how big the dot is and how widely its shells fluctuate, whereas detailed bond order and topology require explicit angular registration and conditioning on shell occupancy.

Backflow as a local, high-dimensional correction. Backflow behaves very differently. For $N=6$ and $N=12$ the displacement field Δx has $r_{\text{eff}}(\Delta x)$ in the range ~ 10 –22 (Table 6.4), with relatively flat PCA spectra and gradual reconstruction under PC ablation. Linear probes from Δx PCs to global observables such as r_{mean} , r_{var} or shell contrast yield negligible R^2 , indicating that the backflow field does not act as a simple global rescaling or shape parameter.

Near-field diagnostics of ΔE and $\|\Delta x\|^2$ support a predominantly local interpretation. At intermediate traps (e.g. $N=6$ or 12 at $\omega = 10^{-2}$), the lowest $q\%$ of configurations by r_{min} carry a disproportionately large fraction of the backflow energy correction: for $N=12$ at $\omega = 10^{-2}$ the lowest 5% bin accounts for about six times its fair share of ΔE while the corresponding share of $\|\Delta x\|^2$ remains close to unity. Thus, many spatial modes contribute to the displacement field, but the *energetic leverage* of backflow is concentrated in short-distance configurations where small nodal adjustments have a large impact on the local energy.

Deep Wigner limit: the backflow’s net contribution shrinks. The extreme Wigner points provide a clean limit. For $N=2$ at $\omega = 10^{-3}$, the backflow correction to the energy is $\Delta E \sim 10^{-7}$ Ha with comparable uncertainty, head-PC1 alignment between no-BF and BF correlator features is essentially perfect (cosine = 1), and the ranks of Z and ΔZ are both ≈ 1 . PC ablations of Δx show that even though $r_{\text{eff}}(\Delta x)$ can be nominally high, the field is effectively constant over the sampled manifold (PC ablation error $\sim 10^{-6}$), and near-field shares of both ΔE and $\|\Delta x\|^2$ are ≈ 1 . In the trained CTNN decomposition, then, the net energetic contribution assigned to backflow becomes small in the deep-Wigner regime. This does not imply that backflow is generically unnecessary here: Chapter 7 shows that a *conventional* backflow collapses in precisely this regime and loses accuracy, whereas the message-passing backflow retains the full-rank response the crystal demands. What is small is the energy assigned to one component in one trained decomposition—a gauge-like quantity—not the structural need for an adequate relational nodal channel.

For $N=6$ and $N=12$ at $\omega = 10^{-3}$ the pattern is similar, albeit less extreme. The correlator manifold remains low-rank with PC1 almost unchanged between no-BF and BF models, $r_{\text{eff}}(\Delta Z)$ stays close to $r_{\text{eff}}(Z)$, and the energetic effect of backflow is very small. Near-field shares of ΔE revert to values close to one. Once the system is deep in the Wigner-molecule regime and the static correlator has adjusted to the interaction-dominated landscape, there is simply little left for backflow to correct.

Two qualifications keep this from being over-read, both from the tangent-kernel analysis of Chapter 7. First, “idle” describes the small *net energy* of the message-passing backflow at these points, not the structure a backflow must supply: a *conventional* backflow does not fall idle so gracefully. It collapses to a single collective mode (rank $\rightarrow 1$) and *loses* accuracy at the Wigner crystal, because the residual correlation there is relative and full-rank, and its single round of pairwise messages does not sustain it, whereas the CTNN’s copresheaf transport does. Second, the ranks and displacement magnitudes quoted here are gauge-like (§7.5); what survives a change of training route is the ablation price and the gap between a conventional and a copresheaf backflow, not the component ranks.

Regime dependence and architectural implications. Taken together, the representation results show a clear regime dependence:

- In *weakly to moderately correlated* traps (larger ω , smaller r_s), the correlator remains low-rank but backflow carries a non-trivial fraction of the energy correction, focused on near-field configurations. Here it is beneficial to allocate capacity to a reasonably wide BackflowNet with rich pairwise inputs, while keeping the correlator compact.
- In the *deep Wigner* regime (ultra-weak traps such as $\omega = 10^{-3}$), the correlator alone captures most of the global size/shelling and the relevant near-field structure, and the backflow’s *net* energy contribution becomes small. Adding *width* to an already message-passing backflow buys little here; the architectural *kind* of backflow, however, still matters—a conventional single-round-message network collapses and loses accuracy exactly where the copresheaf CTNN holds full rank and does not (Chapter 7).

From a design perspective, this suggests that Slater–Jastrow PINNs with a small, smooth correlator and moderately wide backflow layers are a natural default for parabolic dots. The correlator behaves like a learned set of collective coordinates summarizing dot size and shell fluctuations, while backflow functions as a flexible local correction channel that is activated only where near-field physics is delicate. In the Wigner limit, the model naturally reduces to a static Wigner-molecule ansatz with low-rank latent geometry.

Compression is physical alignment. The low-rank latent geometry above says *how many* directions the correlator uses; the tangent-kernel analysis of Chapter 7 adds *which* directions, and the distinction turns out to matter. Naming the leading tangent modes against a dictionary of physical operators shows that the message-passing correlator’s dominant directions are the Hamiltonian’s own collective modes—breathing, correlation-hole, relative excitation—to $R^2 \approx 1$ across ω and up to $N=12$, whereas a separable network of matched (or larger) capacity leaves nearly half of its tangent budget along directions a compact physical dictionary does not resolve. A direction outside the dictionary need not be unphysical; the robust reading is that the CTNN’s tangent space is far more completely accounted for by a small set of physical operators than the DeepSet’s. That gap is present at every confinement and is merely disguised at strong confinement, where correlation is a small slice of the energy and the two networks are indistinguishable in energy and even in leading-mode physicality. It also appears *coupled* to the backflow: holding the backflow fixed and varying only the correlator, the correlator’s loss of alignment and the backflow’s rank collapse set in at the same confinement—evidence for a common relational bottleneck seen once in the amplitude and once in the nodes, rather than proof that the two are literally one deficit.

Connection to the physical picture. Finally, the learned representations dovetail with the structural analysis in Sec. 6.3. The same correlator that reproduces shell topologies, bond order and virial partitioning in the Wigner regime is found to live on a one- or two-dimensional manifold whose leading axis correlates almost perfectly with global size and radial spread. Backflow, in turn, is high-dimensional and local exactly in the regimes where shell structure is less rigid and close encounters are more frequent. This supports a physically meaningful decomposition: a small set of global “collective coordinates” captured by W_θ and a many-body, predominantly local correction channel implemented by BackflowNet. The success of this decomposition in capturing both energies and detailed Wigner-molecule structure suggests that similar architectures may be effective for other strongly correlated finite fermion systems.

8.4 Discussion of collocation-assisted training

Of everything in this thesis, the collocation programme is where the claim most needs its boundaries drawn before it is stated, because the headline—an accurate many-body wavefunction trained with no Markov chain—is the kind of sentence that travels further than its evidence. It is also the part that consumed the most time and produced the most failures, and the two facts are related: the method works within a regime whose edges had to be found by walking off them repeatedly. What follows therefore spends as much effort on where the approach breaks, and on the precise sense in which “MCMC-free” is meant, as on where it succeeds.

The collocation experiments reported in Sec. 6.4 demonstrate that a purely importance-sampled, MCMC-free training pipeline can produce wavefunctions of near-DMC accuracy for moderate particle numbers and confinement strengths, while revealing systematic limitations at large N and ultra-weak

confinement. This section interprets these findings, identifies the mechanisms behind successes and failures, and discusses their broader implications.

8.4.1 The central result: MCMC-free training to DMC accuracy

The most significant finding is that collocation training—using only analytic Gaussian proposals and importance resampling, with no Markov chain at any stage of optimisation—achieves +0.009% accuracy relative to DMC for $N=6$ at $\omega=1.0$, and +0.018% for $N=12$ at the same confinement. These errors are comparable to the statistical uncertainty of the DMC references themselves and are achieved with a training budget that is modest by neural-network VMC standards.

This result is non-trivial because the prevailing assumption in the neural quantum state community is that accurate wavefunction optimisation requires equilibrated MCMC sampling from $|\Psi|^2$ throughout training. The collocation approach sidesteps this by treating the local energy as a *reward signal* in a REINFORCE policy gradient, drawing points from a fixed analytic distribution and reweighting them. The price is higher gradient variance (because the proposal only partially overlaps $|\Psi|^2$), but this is compensated by the absence of autocorrelation, burn-in, and mixing overhead.

For $N=6$ across all five confinement strengths tested ($\omega \in \{1.0, 0.5, 0.1, 0.01, 0.001\}$), the worst collocation error is +0.244% at $\omega=0.001$, where electrons are separated by $\sim 100 a_0^*$ (about four oscillator lengths) in a 12-dimensional configuration space. That a simple Gaussian mixture with three width components can maintain usable overlap with $|\Psi|^2$ at such extreme aspect ratios speaks to the effectiveness of the $1/\omega$ scaling in the proposal width and to the robustness of the importance-resampling pipeline. The Gaussian mixture is only the first proposal one might try, however. Where it does eventually fail—the origin-centred form aims where a deep Wigner molecule no longer has any mass—Chapter 7 replaces it with a proposal built from the Wigner shell itself and carries the same MCMC-free objective down to $\omega = 0.0035$, a regime the Gaussian mixture cannot reach.

8.4.2 Why REINFORCE succeeds where residual losses fail

The catch-22 analysis (Appendix C) identified a fundamental obstacle for direct PDE-residual training of the many-body Schrödinger equation: the backflow learning signal is concentrated near nodes and coalescence points where the local energy E_L diverges, but these are precisely the configurations where second-derivative pathways are ill-conditioned. Backpropagating through the Laplacian in such regions produces pathological gradients that destabilise training.

The REINFORCE formulation resolves this by construction. The Laplacian enters only through the *reward* (the local energy in Eq. (4.2)), and gradients act exclusively on the score function $\nabla_\theta \log |\Psi|$. The stability is subtler than the score simply being bounded. For a *parameter-dependent* (backflow) node the score itself can diverge— $\nabla_\theta \log |\Psi| \supset (\nabla_\theta D)/D$ grows like $1/d_\theta$ as the learned determinant $D \rightarrow 0$ near its node. What matters is that this divergence is *integrable* under the $|\Psi|^2$ sampling measure, which weights it by $|\Psi|^2 \sim d_\theta^2$ (and the near-node points are further suppressed by the importance screening). REINFORCE avoids the *stronger*, non-integrable divergence of the strong-form residual, in which the Laplacian term $\nabla^2 D/D$ enters the gradient directly (Appendix C). Keeping the Laplacian in the *reward* but out of the *gradient flow* is what lets the collocation pipeline train backflow and Jastrow parameters stably in regimes where direct residual minimisation fails.

The superiority of pure REINFORCE ($\beta=0$) over the hybrid objective ($\beta>0$) has a related explanation. The direct weak-form term $\tilde{E} = \frac{1}{2} \|\nabla \log \Psi\|^2 + V$ is cheaper to evaluate (no Laplacian needed) but minimises a different functional: it drives the wavefunction toward low kinetic energy rather than low total energy. At nonzero β , the resulting gradient conflict between the two objectives manifests as oscillatory behaviour in the energy, with the REINFORCE term pulling toward the variational minimum and the direct term pulling toward a kinetic-energy minimum that is generally distinct.

8.4.3 The sampling bottleneck and its regime dependence

The central limitation of collocation training is that importance-sampling quality degrades systematically with particle number and with decreasing confinement. This degradation is governed by two competing effects:

1. **Dimensionality.** The configuration space is Nd -dimensional. A Gaussian proposal in Nd dimensions has an overlap with $|\Psi|^2$ that shrinks exponentially with Nd unless the proposal is well

matched to the target distribution. For $N=6$ ($Nd=12$), a three-component mixture is sufficient; for $N=20$ ($Nd=40$), it is marginal.

2. **Correlation structure.** At low ω , the many-body wavefunction develops strong inter-particle correlations (Wigner-molecule structure) that make $|\Psi|^2$ highly non-Gaussian: it concentrates on shell-like geometries with specific inter-particle spacings. An isotropic Gaussian proposal has no mechanism to capture this structure.

The ESS quantifies this overlap. Empirically, the ESS per epoch scales roughly as:

- $N=6$: ESS ~ 50 – 500 at $\omega=1.0$, ~ 5 – 50 at $\omega=0.01$, ~ 2 – 10 at $\omega=0.001$.
- $N=12$: ESS ~ 30 – 200 at $\omega=1.0$, ~ 5 – 15 at $\omega=0.1$, ~ 1 at $\omega=0.01$ (training fails).
- $N=20$: ESS ~ 2 – 15 at $\omega=1.0$, ~ 1 – 5 at $\omega=0.1$.

When the ESS approaches 1, the importance-weighted batch degenerates to a single effective sample, and the REINFORCE gradient becomes essentially a random vector. The ESS-adaptive oversampling mechanism delays this collapse but cannot prevent it: eventually, the number of proposal samples needed to maintain a given ESS grows exponentially, making the approach intractable.

This explains the systematic pattern in Table 6.8: accuracy degrades smoothly from excellent (+0.009%) at $(N=6, \omega=1.0)$ to poor (+5.68%) at $(N=20, \omega=0.1)$, following the trajectory of ESS collapse.

8.4.4 The optimiser regime boundary

The sharp failure of natural-gradient preconditioning (CG-SR) below $\omega \lesssim 0.01$ reveals an important structural feature of the optimisation landscape.

The Fisher information matrix $F_{ij} = \mathbb{E}[\partial_i \log |\Psi|^2 \cdot \partial_j \log |\Psi|^2]$ characterises the curvature of parameter space as seen by the wavefunction. Ill-conditioning alone does not determine whether SR helps. Natural gradients are most valuable when the metric is anisotropic but remains statistically resolvable; what disqualifies them is not curvature but the failure to estimate it. In strong confinement ($\omega \geq 0.1$) the wavefunction is compact and all parameters contribute to its structure in a comparable manner, so F is well-conditioned, its inverse is a sound preconditioner, and Adam is in any case close behind.

At ultra-weak confinement ($\omega \leq 0.01$), two effects conspire—and it is the second, not the first, that is decisive:

1. **Scale separation.** The wavefunction extends over $\sim 1/\sqrt{\omega}$ oscillator lengths, creating a large separation between the scale of the Jastrow correlator and the scale of the orbital envelope. Parameters controlling short-range correlations (cusp, pair features) contribute very differently to F than parameters controlling long-range structure (orbital widths, backflow offsets).
2. **Sampling starvation.** The Fisher matrix is estimated from importance-resampled points. When the ESS is low, the estimate of F is dominated by a few high-weight samples, making it noisy and rank-deficient. Tikhonov damping can stabilise the inversion, but at the cost of washing out the very curvature information that motivates natural gradients.

Adam, by contrast, maintains per-parameter adaptive learning rates using only first and second moment estimates of the gradient. It makes no attempt to model parameter-space curvature globally, and therefore does not fail when the curvature is pathological. The price is slower convergence in well-conditioned regimes (where CG-SR can take larger, more informed steps), but the benefit is robustness in ill-conditioned regimes where CG-SR diverges.

The resulting picture has three regimes rather than two. Where the metric is well-conditioned, SR is sound but unnecessary. Where it is anisotropic and still estimable, SR is at its most valuable—which is the collocation case of Chapter 7, Q2. Where importance-sampling collapse makes the empirical metric itself unreliable, its inversion amplifies estimation noise and SR becomes actively harmful. The practical implication for future work is therefore about the *estimator*, not the curvature: any natural-gradient or second-order method applied to weakly confined many-body systems must certify that the metric it inverts is statistically resolved at the sample sizes in use.

8.4.5 Why Langevin refinement fails

Langevin refinement was motivated by a natural intuition: if the Gaussian proposal overlaps $|\Psi|^2$ poorly, evolving samples up the score $\nabla_{\mathbf{R}} \log |\Psi|^2$ before resampling should improve that overlap. It fails for a reason that is standard in the MCMC literature and easy to overlook in an importance-sampling context. The weights $w_i = |\Psi(\mathbf{R}_i)|^2 / q(\mathbf{R}_i)$ are valid only if the $\mathbf{R}_i$ are drawn from q ; after K Langevin steps from an out-of-equilibrium start, the effective density is some intermediate $\tilde{q}$ depending on K and ε , while the weights still divide by q . The resulting gradient bias is not noise—it pushes the wavefunction toward regions the Langevin dynamics favours, which shifts $\tilde{q}$ further in the same direction, and the loop closes. The measured errors in §6.4.4 confirm the mechanism experimentally. Correcting it requires running to equilibrium, adding a Metropolis–Hastings step, or computing annealed weights along the non-equilibrium path; the first two are MCMC by another name, which defeats the purpose, and the third is expensive enough to forfeit the advantage the collocation route exists to provide.

8.4.6 Architectural trade-offs at scale

The reversal of the backflow advantage at $N=20$ highlights an important architectural trade-off. The copresheaf backflow network (CTNNBackflowNet) stores edge features of size $\mathcal{O}(N^2 \times h_{\text{bf}})$, which at $N=20$ with $h_{\text{bf}}=128$ requires $\sim 10\times$ the memory of the Jastrow-only network. This forces a choice: reduce h_{bf} (losing expressivity), reduce the collocation budget (losing gradient quality), or reduce the oversampling factor (losing ESS). All three options degrade the result.

The Jastrow-only network, with $\sim 25k$ parameters and $\mathcal{O}(N^2)$ edge cost (no hidden-dimension multiplier), permits full-size collocation budgets ($n_{\text{coll}}=4096$, oversample=8–16) even at $N=20$. The resulting gradient is less noisy, and the simpler loss landscape (fewer parameters) makes optimisation more reliable.

This does not mean that backflow is unnecessary at $N=20$; it means that the current training pipeline cannot supply enough gradient signal to train backflow effectively at this scale. The implication is that scaling collocation to larger systems requires either (i) memory-efficient backflow architectures (e.g. sparse attention, low-rank message passing), or (ii) improved sampling that provides higher ESS at constant computational cost.

8.4.7 Comparison with MCMC-based results

A direct comparison with the SR-refined results in Table 6.2 illuminates the relative strengths of the two approaches:

- At $(N=6, \omega=1.0)$, collocation (+0.009%) matches the SR-refined result (+0.026–0.048% range for $N=6$) and is competitive with or slightly better than it. This is the regime where collocation is most effective, because the configuration space is low-dimensional and the Gaussian proposal overlaps well with $|\Psi|^2$.
- At $(N=12, \omega=0.1)$, collocation (+0.122%) is roughly 2–4 $\times$ worse than SR (+0.040% in Table 6.2). Here the sampling bottleneck is beginning to bite, and the MCMC sampler’s ability to draw equilibrated samples from $|\Psi|^2$ gives it a clear advantage.
- At $N=20$, collocation results (+1.4–5.7%) are significantly worse than what an MCMC-based approach would be expected to achieve, reflecting the exponential ESS degradation in 40 dimensions.

The SR pipeline also has its own scaling challenges: MCMC mixing times grow with particle number and correlation strength, and the SR step itself requires solving a $P \times P$ linear system (P =number of parameters). But these challenges manifest at much larger N than the collocation ESS collapse, because the MCMC sampler adapts to $|\Psi|^2$ throughout training rather than relying on a fixed proposal.

The practical conclusion is that collocation is competitive with MCMC for $N \lesssim 12$ and $\omega \gtrsim 0.01$, and that the two approaches are complementary: collocation for fast, parallelisable pretraining and exploration, MCMC for final refinement at larger N or lower ω .

8.4.8 Implications and future directions

The collocation experiments have several implications for the broader effort to apply neural wavefunctions to many-body quantum systems.

MCMC is not necessary for training. The demonstration that REINFORCE with importance resampling can match DMC accuracy for $N=6, 12$ challenges the assumption that MCMC is essential for neural wavefunction optimisation. This opens the door to fully parallel training pipelines that scale efficiently on modern hardware, particularly GPUs with large batch sizes.

What “MCMC-free” does and does not mean. Three distinctions are easily conflated. (i) The pipeline is *Markov-chain-free*: no configuration is ever drawn from a Markov chain during training—samples come from an analytic proposal $q(\mathbf{R})$ and are importance-reweighted. It is not free of Monte Carlo in the broader sense: importance sampling and multinomial resampling are themselves Monte-Carlo operations. (ii) It is not, in general, *reference-free*. Where a DMC energy exists ($\omega \geq 0.1$), the residual pretraining anneals its target toward E_{DMC} and the instability-rollback monitors the relative error against it; the DMC value therefore enters the training signal, so those runs should be read as “trained without a Markov chain,” not as an unsupervised solution of the Schrödinger equation. (iii) The deep-Wigner cascade of Chapter 7 is the one regime that is both Markov-chain-free *and* reference-free: below $\omega = 0.1$ no DMC value exists, so the cascade is guided only by the variational objective and cross-checked against the physical energy-scaling law, never against an external benchmark.

The proposal distribution is the bottleneck. The systematic pattern of ESS collapse with increasing N and decreasing ω identifies the proposal distribution as the critical component to improve. Possible directions include normalising flows trained to approximate $|\Psi|^2$, diffusion-model proposals, or variational inference techniques that can adapt the proposal to the evolving wavefunction during training.

Natural gradients require careful conditioning. The CG-SR failure at low ω is not specific to collocation: any natural-gradient method applied to weakly confined systems will encounter the same ill-conditioning. This suggests that preconditioning strategies (block-diagonal Fisher approximations, Kronecker-factored curvature, or adaptive damping schedules) deserve attention in the neural quantum state context.

Backflow scaling needs architectural innovation. The $\mathcal{O}(N^2 \times h_{\text{bf}})$ memory scaling of current backflow networks limits their applicability beyond $N \approx 12$. Memory-efficient architectures—sparse message passing, low-rank approximations, or local backflow corrections that avoid all-to-all edge features—are needed to extend the collocation approach to larger systems while retaining the nodal-surface corrections that backflow provides.

Collocation as a pretraining stage. Perhaps the most practical near-term use of collocation is as a fast pretraining stage that conditions the wavefunction before MCMC-based refinement. The collocation-trained checkpoints already contain useful structural information (orbital shapes, Jastrow correlations, approximate nodal surfaces) that accelerates subsequent SR convergence. This two-stage protocol—collocation pretraining followed by SR polish—combines the embarrassing parallelism of collocation with the asymptotic accuracy of MCMC-based methods.

Part VI

Conclusion

Chapter 9

Conclusions

*All things are implicated with one another, and the bond is holy;
and there is hardly anything unconnected with any other thing.*

MARCUS AURELIUS
Meditations, Book VII, §9

This thesis set out from a single question—what does it actually take to solve the many-body Schrödinger equation with a neural network, and what has the network understood once it has—and the most durable answers came from the places where the attempt failed. The quantum-dot benchmarks served a dual purpose throughout: they test the numerical stability and expressivity of the Slater–Jastrow PINN+backflow ansatz, and they provide a controlled setting in which to study Wigner-molecule formation and the internal representations learned by the network. What follows separates what has been demonstrated from what has been observed, and both from what has merely been made plausible; that separation is itself one of the results.

On the *energy* side, a single trap-unit-normalised architecture with soft-core pair features and an explicit cusp reaches DMC-level accuracy for two-electron dots and holds sub-0.1% relative errors for $N \in \{6, 12\}$ across $\omega \in \{0.1, 0.5, 1.0\}$ (Sec. 8.1). It extrapolates smoothly to $\omega = 10^{-3}$, where no DMC data exist and internal checks—scaling exponents, virial ratios, structural diagnostics—carry the reliability of what are then variational predictions. These accuracies come from a modest SR tail on residual-based pretraining, which says that conditioning and inductive bias matter at least as much as raw sample count.

On the *structural* side, the weak-trap states develop bona fide Wigner-molecule structure rather than being vaguely “strongly correlated” dots (Sec. 6.3). As ω falls, all three systems ($N = 2, 6, 12$) cross over smoothly: radii and localisation ratios move toward large stiff configurations, the energy partitioning approaches the classical virial form $V_{\text{int}} \approx 2V_{\text{trap}}$, and shell structures and bond orders emerge that match classical Coulomb-cluster topologies—a single rotating ring mixing (1, 5) and (0, 6) at $N=6$, a modal two-shell (3, 9) molecule at $N=12$. Shell-resolved reconstructions of $g(r)$ recover the global pair structure from shell geometry and combinatorics alone (a decomposition check; sampling correctness rests on the independent restart and chain diagnostics). Throughout, $N_{\uparrow} = N_{\downarrow}$ is held fixed and total-spin sectors are not scanned, so these are statements about the studied spin-projection sector rather than a claim to have settled the global spin ordering.

On the *representation* side, that real-space richness is carried by a strikingly simple latent geometry (Sec. 8.3). The correlator occupies one to two effective directions whose leading axes act as collective coordinates for global size and radial fluctuation, with angular and topological information held in conditional distributions on them; across regimes the leading axis changes identity in a $\phi \leftrightarrow \psi$ recomposition rather than growing in dimension. The backflow is by contrast a high-dimensional displacement field whose energetic leverage sits in near-field configurations at intermediate traps. One earlier reading—that the optimiser simply switches backflow off—proves architecture-specific and is corrected here: it is the *conventional* backflow that collapses to a single collective mode at the Wigner crystal and loses accuracy, its single round of pairwise messages failing to sustain the relative displacement the crystal demands, while the copresheaf message-passing backflow holds full rank and its accuracy. How these internal numbers may be read is itself a result: component-level quantities are gauge-like, so the load-bearing conclusions rest on the invariants—energy, overlap, ablation—not on the ranks.

On the *method* side, the tangent kernel connects these observations to the design choices that produced them (Chapter 7). Message passing earns its cost by supplying a relational channel: it compresses the correlator onto fewer effective tangent directions and sustains a full-rank backflow where a separable network cannot, and at the crystal the inter-particle messages account for essentially the entire backflow contribution. The sharper statement concerns physical structure rather than dimension count—message passing keeps the correlator’s leading tangent modes aligned with the Hamiltonian’s collective modes, whereas a separable network reaches the same energy with much of its tangent capacity along directions a compact physical dictionary does not resolve. That gap is present at every confinement, disguised at strong confinement only because correlation is then a small slice of the energy; and it appears coupled to the backflow, since the correlator’s loss of alignment and the backflow’s rank collapse set in at the same confinement and shift together under a correlator swap, supporting a common relational bottleneck. Natural-gradient preconditioning helps in proportion to how ill-conditioned the tangent metric is, so long as that metric stays estimable—little for $|\Psi|^2$ -sampled VMC, decisively for collocation, and harmfully once importance sampling collapses. The same conditioning story explains why an MCMC-free, importance-sampled scheme equals VMC while its sampling is healthy and, with a proposal built from the Wigner shell itself, reaches a genuine Wigner molecule at $\omega = 0.0035$ with no Markov chain in the gradient optimisation (checkpoint selection and final certification use MCMC).

Limitations and outlook. Three limits bound these conclusions. External ground truth reaches only to $\omega = 0.1$; below it the accuracy statements are internal consistency rather than benchmarked error, and the deep-Wigner energies should be read as variational predictions supported by physical scaling, not as validated numbers. The architecture comparison is cleanest at $N = 6$, where it is seeded; the $N = 12$ contrast rests on fewer trained models and the $N = 20$ analysis on a single production state, so the size-scaling of the tangent-space picture is indicative rather than settled. And the message-passing backflow that carries the crystal is quadratic in particle number—affordable at these sizes, but the first thing that must change to reach the hundreds of electrons where the same relational structure should matter most. None of these undermines the central finding; they mark where it has been established and where it remains to be pushed.

Taken together, these findings converge on a single theme. What makes the difference across architecture, optimiser, and training paradigm is not raw capacity but alignment with the low-dimensional, physically collective structure of the problem—the same structure that, in real space, *is* the Wigner molecule. A compact correlator supplies learned collective coordinates; a message-passing backflow supplies the relational corrections a single-round-message network cannot sustain; natural gradients and $|\Psi|^2$ sampling keep the optimisation pointed along those directions. The quantum-dot results thus provide both a stringent validation of the method and a template for the same ideas in more complex finite fermion systems, where the trade-offs between global structure, local correlation, and representation are expected to recur.

Part VII

Appendices

Chapter A

Laplacian conditioning

A.1 Residual training for Schrödinger

We consider the stationary many-body Schrödinger equation on configuration space $\mathbf{R} = (\mathbf{r}_1, \dots, \mathbf{r}_N) \in \Omega \subset \mathbb{R}^D$ with $D = dN$ (for our quantum dots, $d = 2$):

$$\hat{H}\Psi(\mathbf{R}) = E\Psi(\mathbf{R}), \quad \hat{H} = \hat{T} + \hat{V}, \quad \hat{T} = -\frac{1}{2} \sum_{i=1}^N \Delta_i, \quad (\text{A.1})$$

where $\hat{V}$ collects external and interaction potentials.

In *strong residual minimization* one introduces

$$L := \hat{H} - E \quad \Rightarrow \quad L\Psi = 0, \quad (\text{A.2})$$

and minimizes a squared residual loss of the schematic form

$$\mathcal{L}_{\text{res}}(\theta) = \frac{1}{2} \|L\Psi_\theta\|_{L^2(\Omega; w)}^2 = \frac{1}{2} \int_{\Omega} |(L\Psi_\theta)(\mathbf{R})|^2 w(\mathbf{R}) d\mathbf{R}, \quad (\text{A.3})$$

for a nonnegative weight w induced by the sampling strategy. A key message of the numerical-analysis viewpoint is that strong residual losses can be *severely ill-conditioned*, and training speed is tightly linked to spectral properties of an operator/matrix induced by L [37].

A.2 Linearized geometry and the matrix A

To expose the mechanism, follow the standard linearization argument used in conditioning analyses [37]. Fix an expansion point θ_0 and define tangent (feature) functions

$$q_j(\mathbf{R}) := \partial_{\theta_j} \Psi_\theta(\mathbf{R})|_{\theta=\theta_0}, \quad j = 1, \dots, P. \quad (\text{A.4})$$

For small parameter increments $\delta\theta$, we have the first-order model

$$\Psi_{\theta_0+\delta\theta}(\mathbf{R}) \approx \Psi_0(\mathbf{R}) + \sum_{j=1}^P q_j(\mathbf{R}) \delta\theta_j, \quad \Psi_0 := \Psi_{\theta_0}. \quad (\text{A.5})$$

Applying the residual operator gives

$$(L\Psi_{\theta_0+\delta\theta})(\mathbf{R}) \approx (L\Psi_0)(\mathbf{R}) + \sum_{j=1}^P (Lq_j)(\mathbf{R}) \delta\theta_j. \quad (\text{A.6})$$

Introduce the weighted inner product (complex-valued wavefunctions allowed)

$$\langle f, g \rangle_w := \int_{\Omega} f(\mathbf{R})^* g(\mathbf{R}) w(\mathbf{R}) d\mathbf{R}. \quad (\text{A.7})$$

Let the Jacobian J be the linear map whose j th column is Lq_j . Then the loss (A.3) becomes (to second order in $\delta\theta$)

$$\mathcal{L}_{\text{res}}(\theta_0 + \delta\theta) \approx \frac{1}{2} \|r_0 + J\delta\theta\|_w^2, \quad r_0 := L\Psi_0. \quad (\text{A.8})$$

with normal-equation matrix

$$A := J^*J, \quad A_{ij} = \langle Lq_i, Lq_j \rangle_w. \quad (\text{A.9})$$

Here A is Hermitian positive semidefinite; in practice it can be close to singular, so one often effectively works with a damped variant $A + \lambda I$ (implicit or explicit).

Now consider gradient descent on the quadratic model (A.8) (with step size η). The error in parameter space evolves as

$$\delta\theta^{(t+1)} - \delta\theta^* = (I - \eta A) \left(\delta\theta^{(t)} - \delta\theta^* \right), \quad (\text{A.10})$$

so each eigen-direction of A contracts by a factor $(1 - \eta\lambda)$. Hence convergence speed is governed by the eigenvalue spread of A , typically summarized by the (effective) condition number

$$\kappa(A) = \frac{\lambda_{\max}(A)}{\lambda_{\min}(A)} \quad (\text{A.11})$$

(over the nonzero/regularized spectrum). Large $\kappa(A)$ implies stiffness: a step size small enough to be stable for $\lambda_{\max}$ yields slow progress along directions associated with $\lambda_{\min}$. This is precisely the conditioning bottleneck emphasized in physics-informed training analyses [37].

A.3 Bi-Laplacian scaling

The defining feature of Schrödinger is that L contains a second-order differential operator. To isolate the kinetic contribution, consider the regime where the residual is dominated by the Laplacian part,

$$L \approx -\frac{1}{2}\Delta_{\mathbf{R}}. \quad (\text{A.12})$$

From L to L^*L . The matrix A in (A.9) is a Gram matrix in the feature space $\{Lq_j\}$. The corresponding operator-level object is the *Hermitian square* L^*L (with adjoint taken in the $L^2(\Omega; w)$ inner product) [37]. For general non-constant w , the adjoint L^* differs from the formal differential adjoint by weight-induced lower-order terms; the cleanest picture is obtained for constant w and standard boundary conditions (periodic/Dirichlet), where the Laplacian is self-adjoint.

In the Laplacian-dominant regime (A.12), one has (up to a constant factor)

$$L \approx -\frac{1}{2}\Delta \quad \Rightarrow \quad L^*L \approx \frac{1}{4}\Delta^2, \quad (\text{A.13})$$

i.e. a bi-Laplacian scaling. The prefactor does not affect conditioning ratios.

Fourier/eigenmode derivation of k^4 . On a periodic domain (or, more generally, in a Laplacian eigenbasis) $-\Delta\phi_{\mathbf{k}} = |\mathbf{k}|^2\phi_{\mathbf{k}}$, so $\Delta^2\phi_{\mathbf{k}} = |\mathbf{k}|^4\phi_{\mathbf{k}}$. If an error $\delta\Psi$ expands as $\delta\Psi = \sum_{\mathbf{k}} c_{\mathbf{k}}\phi_{\mathbf{k}}$ and $L \approx -\frac{1}{2}\Delta$, then

$$\|L\delta\Psi\|_{L^2}^2 \approx \frac{1}{4} \sum_{\mathbf{k}} |\mathbf{k}|^4 |c_{\mathbf{k}}|^2. \quad (\text{A.14})$$

Thus the squared residual penalizes high-frequency components with a *quartic* weight $|\mathbf{k}|^4$. If the relevant spectrum spans $|\mathbf{k}| \in [k_{\min}, k_{\max}]$, the induced eigenvalue ratio scales like

$$\kappa \sim \left(\frac{k_{\max}}{k_{\min}} \right)^4, \quad (\text{A.15})$$

which is a sharp mathematical expression of “Laplacian-driven stiffness”. This quartic growth is consistent with the Laplacian/Poisson toy examples and frequency-dependent conditioning effects emphasized in [37].

Length-scale interpretation. Let ℓ denote the smallest length scale that must be represented. Then $k_{\max} \sim 1/\ell$ and (A.15) implies

$$\kappa \text{ grows rapidly as } \ell \downarrow 0. \quad (\text{A.16})$$

In many-body Schrödinger problems, small ℓ arises from nodal structure, strong localization, and short-range correlation features; in all such regimes, the Laplacian makes the strong residual extremely curvature-sensitive.

A.4 Barron/spectral perspective

The conditioning story above can be rephrased in approximation terms. In spectral Barron-type viewpoints, function classes are controlled by norms that weight the Fourier transform by powers of frequency (or related spectral weights). A key implication is that applying derivatives systematically increases the importance of high-frequency tails [37]. Since Δ multiplies a Fourier mode by $|\mathbf{k}|^2$, and the squared residual effectively involves L^*L (hence Δ^2 in Laplacian-dominant regimes), the residual can require substantially more “spectral budget” than the wavefunction itself. In practical terms: even if Ψ is approximable in a given network class, *approximating its strong residual* can be much harder, because the loss implicitly demands accurate high-frequency curvature (cf. (A.14)). This is one of the motivations, in numerical analysis of PINNs, for considering weak/variational formulations that reduce derivative order when the strong form is too stiff [37].

A.5 Implications for Stochastic Reconfiguration

Stochastic reconfiguration (SR) updates parameters using a natural-gradient-like preconditioner [27, 29, 46]. Writing $O_i = \partial_{\theta_i} \log \Psi_\theta$, one common SR update is

$$S(\theta) \delta\theta = -g(\theta), \quad S_{ij} = \text{Cov}_{p_\theta}[O_i, O_j], \quad g_i = 2 \text{Cov}_{p_\theta}[E_L, O_i], \quad (\text{A.17})$$

where $p_\theta \propto |\Psi_\theta|^2$ and $E_L = (\hat{H}\Psi_\theta)/\Psi_\theta$. SR improves conditioning in parameter space by using the (quantum) information geometry of the variational family [27, 46]. However, the Laplacian-induced stiffness described above is an *operator-level* phenomenon: strong residuals magnify curvature errors via Δ and Δ^2 (cf. (A.14)), which remains a fundamental challenge even when parameter updates are geometrically preconditioned.

Chapter B

Coulomb conditioning

B.1 Coulomb structure and singularity

For our two-dimensional electronic systems ($\mathbf{r}_i \in \mathbb{R}^2$), the interaction contains Coulomb terms of the form

$$\hat{V}_C(\mathbf{R}) = \sum_{1 \leq i < j \leq N} \frac{1}{r_{ij}}, \quad r_{ij} = |\mathbf{r}_i - \mathbf{r}_j|. \quad (\text{B.1})$$

This is a *singular multiplication operator*: it diverges at particle coalescence $r_{ij} \rightarrow 0$. The exact eigenfunction remains finite (or vanishes by antisymmetry) but develops non-smooth short-distance structure (a cusp) so that divergences in kinetic and potential contributions cancel in physically meaningful quantities such as the local energy [29].

The central point for learning is:

Coulomb does not merely add “another term” to the residual; it introduces *singular short-range physics* that forces delicate cancellations with the Laplacian. Strong residual objectives square these effects and become extremely sensitive to tiny cusp errors.

B.2 2D cusp conditions: antiparallel vs parallel spins

We present the 2D cusp logic in a way that separates the two spin channels. The cleanest statement (especially for practical ansätze) is in terms of the short-distance behaviour of a *Slater–Jastrow* factorization

$$\Psi(\mathbf{R}) = D(\mathbf{R}) J(\mathbf{R}), \quad J(\mathbf{R}) = \exp\left(\sum_{i < j} u(r_{ij})\right), \quad (\text{B.2})$$

because (i) the Slater part D enforces antisymmetry and determines whether Ψ vanishes at coalescence, and (ii) the Jastrow slope $u'(0)$ is what cancels the $1/r$ Coulomb singularity in the local energy [29].

Fix a pair (i, j) and introduce relative/centre-of-mass coordinates in 2D:

$$\mathbf{R}_{ij} := \frac{\mathbf{r}_i + \mathbf{r}_j}{2}, \quad \mathbf{r} := \mathbf{r}_i - \mathbf{r}_j, \quad r := |\mathbf{r}|. \quad (\text{B.3})$$

A direct computation gives, in any dimension,

$$\Delta_i + \Delta_j = \frac{1}{2} \Delta_{\mathbf{R}_{ij}} + 2 \Delta_{\mathbf{r}}, \quad (\text{B.4})$$

hence the singular short-distance kinetic behaviour is controlled by the relative term $-\Delta_{\mathbf{r}}$.

A key 2D identity is

$$\Delta_{\mathbf{r}} r = \frac{1}{r} \quad (r > 0, \mathbf{r} \in \mathbb{R}^2). \quad (\text{B.5})$$

Antiparallel spins (distinguishable in the Slater part). If the two electrons have opposite spin, they enter different determinants in D and generically D does *not* vanish at $\mathbf{r} \rightarrow 0$. Then $\Psi(0) \neq 0$ and a direct cusp expansion in 2D is legitimate:

$$\Psi(\mathbf{r}, \dots) = \Psi(0, \dots) \left(1 + a r + \mathcal{O}(r^2) \right), \quad r \rightarrow 0. \quad (\text{B.6})$$

Using (B.5), the singular part of the relative Laplacian is

$$\Delta_{\mathbf{r}} \Psi \approx \Psi(0) a \Delta_{\mathbf{r}} r = \Psi(0) \frac{a}{r}, \quad \frac{\Delta_{\mathbf{r}} \Psi}{\Psi} \approx \frac{a}{r}. \quad (\text{B.7})$$

Since the pair kinetic contribution contains $-\Delta_{\mathbf{r}}$, the singular part of the local energy is

$$E_L(\mathbf{R}) = \frac{\hat{H}\Psi}{\Psi} \supset \left(-\frac{a}{r} + \frac{1}{r} \right) = \frac{1-a}{r}. \quad (\text{B.8})$$

Thus finiteness of E_L at coalescence requires

$$a = 1, \quad \text{i.e.} \quad \partial_r \log \Psi|_{r=0} = 1 \quad (2\text{D, antiparallel spins}). \quad (\text{B.9})$$

Equivalently, in a Slater–Jastrow form with $D(0) \neq 0$, this is the same as $\partial_r \log J|_0 = 1$ for the ij -pair.

Parallel spins (identical fermions, node at coalescence). For same-spin electrons, antisymmetry forces D (and hence Ψ) to vanish as $\mathbf{r} \rightarrow 0$. Generically one has a *linear* zero:

$$D(\mathbf{r}, \dots) = \boldsymbol{\eta}(\dots) \cdot \mathbf{r} + \mathcal{O}(r^2), \quad r \rightarrow 0, \quad (\text{B.10})$$

with some nonzero vector $\boldsymbol{\eta}$ depending on the other coordinates. In this case, the cancellation of the Coulomb singularity is controlled by the Jastrow factor through cross terms in $-\Delta_{\mathbf{r}}(DJ)/(DJ)$. A standard short-distance analysis (carried out explicitly for 2D quantum dots) shows that the divergent kinetic contribution behaves like

$$-\frac{\Delta_{\mathbf{r}}(DJ)}{DJ} \supset -\frac{(2k+d-1)}{r} \partial_r \log J \quad \text{with} \quad d=2, \quad k=1 \text{ (identical particles)}, \quad (\text{B.11})$$

while the Coulomb potential contributes $+1/r$ to E_L as before. Hence finiteness of the local energy requires

$$\partial_r \log J|_{r=0} = \frac{1}{2k+d-1} = \frac{1}{3} \quad (2\text{D, parallel spins}). \quad (\text{B.12})$$

This 1 (antiparallel) versus $1/3$ (parallel) 2D cusp statement is the standard quantum-dot/Jastrow–Slater cusp condition [47].

What to remember. In 2D:

$$\text{antiparallel spins:} \quad \partial_r \log \Psi|_0 = 1 \iff \partial_r \log J|_0 = 1, \quad (\text{B.13})$$

$$\text{parallel spins:} \quad \Psi(0) = 0 \text{ (linear node), } \partial_r \log J|_0 = \frac{1}{3}. \quad (\text{B.14})$$

B.3 Why Coulomb is ill-conditioned (2D emphasis)

The cusp conditions above show *how* Coulomb creates stiffness: it forces kinetic and potential contributions to be individually large and singular, while their *sum* is finite only under a short-distance constraint. Strong residual minimization is therefore a *cancellation problem*.

Sensitivity amplification in the local energy. If the learned antiparallel-spin cusp slope is $a_\theta = 1 + \delta a$, then (B.8) implies

$$E_L(\mathbf{R}) \supset -\frac{\delta a}{r}, \quad (\text{B.15})$$

so an arbitrarily small cusp mismatch produces an arbitrarily large local-energy spike as $r \rightarrow 0$.

For parallel spins, if the learned Jastrow slope is $\partial_r \log J|_0 = \frac{1}{3} + \delta b$, then the same short-distance analysis leading to (B.12) implies

$$E_L(\mathbf{R}) \supset -\frac{3\delta b}{r}. \quad (\text{B.16})$$

Thus both channels generate $1/r$ spikes unless the cusp is satisfied.

Why squaring is especially dangerous in 2D. A $1/r$ defect in E_L (or in $(\hat{H} - E)\Psi$) is concentrated near coalescence. But in 2D the squared singularity is not integrable—it diverges logarithmically:

$$\int_{r < \varepsilon} \left(\frac{1}{r}\right)^2 d^2 \mathbf{r} = \int_0^\varepsilon \frac{1}{r^2} (2\pi r dr) = 2\pi \int_0^\varepsilon \frac{dr}{r} = +\infty.$$

Therefore, without effectively resolving the cusp, *strong* squared-residual objectives can be dominated (or even made ill-defined in the continuum limit) by the coalescence region. In Monte Carlo practice, this manifests as heavy-tailed fluctuations and extreme sensitivity to rare close-pair samples.

Residual expansion and cancellation structure. Writing $L = \hat{T} + (\hat{V} - E)$ and expanding the squared residual norm:

$$\|L\Psi\|_w^2 = \|\hat{T}\Psi\|_w^2 + \|(\hat{V} - E)\Psi\|_w^2 + 2 \operatorname{Re}\langle \hat{T}\Psi, (\hat{V} - E)\Psi \rangle_w, \quad (\text{B.17})$$

we see that near coalescence both $\hat{T}\Psi$ and $\hat{V}_C\Psi$ are large. The exact solution relies on cancellation between these contributions (cf. (B.8) and (B.12)), which means (B.17) demands high *relative* accuracy in the cusp region: if either term is off by a small fraction, the net residual can be large in absolute magnitude. This is a classic recipe for ill-conditioning: the loss landscape contains directions where small parameter changes cause huge residual changes in a small region.

Operator-level picture (L^*L). From the same viewpoint as Appendix A, strong residual training induces matrices/operators tied to L^*L [37]. For $L = \hat{T} + (\hat{V} - E)$, one has (formally)

$$L^*L = \hat{T}^2 + (\hat{V} - E)^2 + \hat{T}(\hat{V} - E) + (\hat{V} - E)\hat{T}. \quad (\text{B.18})$$

Coulomb enters here in two destabilizing ways:

1. the positive term $(\hat{V} - E)^2$ involves a $1/r^2$ -type singular weight;
2. the mixed terms contain derivatives of V through product rules, e.g. $\hat{T}(V\Psi)$ generates $\nabla V \cdot \nabla \Psi$ and ΔV contributions, which are even more singular/distributional at coalescence.

This formal expansion explains why the Coulomb singularity can severely inflate the effective spectral spread of the training operator, aligning with the operator-conditioning viewpoint [37].

B.4 Barron/spectral perspective

Coulomb forces the *true* 2D solution to have cusp-like, non-smooth short-range structure, so approximating the *strong residual* requires representing high-frequency content that is not well captured by globally smooth/low-complexity function classes.

In spectral/Barron-style approximation theories, approximation quality is governed by spectral decay and norms that weight the Fourier tail [37]. A cusp (or a linear node times a cusp in J) implies slower spectral decay than a globally smooth function. Strong residual training then *further* amplifies the high-frequency tail because it applies Δ (and effectively L^*L), which multiplies Fourier modes by powers of $|\mathbf{k}|$ (Appendix A). Hence, even if Ψ can be approximated reasonably in function value, the strong residual can remain large unless the cusp structure is represented accurately. This is exactly the type of regularity mismatch that motivates weak/variational formulations in numerical analysis of PINN-type methods, since weak forms reduce the derivative order demanded of the approximation [37].

Practically, QMC wavefunctions therefore incorporate explicit short-range correlation structure (e.g. cusp-enforcing Jastrow factors), precisely to remove these singular residual pathologies and reduce the variance of local quantities [29, 47].

B.5 Implications for Stochastic Reconfiguration

SR performs a natural-gradient step using a (quantum) Fisher/metric matrix [27, 29, 46]. In one common formulation,

$$S_{ij} = \operatorname{Cov}_{p_\theta}[O_i, O_j], \quad g_i = 2 \operatorname{Cov}_{p_\theta}[E_L, O_i], \quad S \delta\theta = -g, \quad (\text{B.19})$$

with $O_i = \partial_{\theta_i} \log \Psi_\theta$.

Coulomb-induced cusp mismatch produces E_L spikes of the form (B.15)–(B.16), which has two consequences:

1. **Heavy-tailed Monte Carlo forces.** The estimator of $g_i = \text{Cov}[E_L, O_i]$ becomes noisy when E_L has rare but extreme spikes (short-range events). This increases variance and can destabilize the linear solve unless one uses damping/regularization, as standard in SR practice [29].
2. **Coupling to curvature sensitivity.** Even if SR corrects anisotropy in parameter space (natural gradient) [27, 46], it does not remove the underlying operator-driven stiffness: the spikes originate from imperfect kinetic–potential cancellation enforced by Coulomb cusp physics.

Bottom line. Laplacian dominance creates stiffness through k^4 -type amplification of curvature errors; Coulomb creates stiffness by enforcing cusp constraints (in 2D: 1 vs 1/3 depending on spin channel) and cancellation against the Laplacian. Both mechanisms worsen conditioning of strong residual objectives and increase statistical difficulty of SR through heavy-tailed local-energy fluctuations [29, 37, 47].

Chapter C

The collocation–backflow catch-22

This appendix formalizes a structural difficulty encountered when training backflow transformations under the *strong-form* collocation objective—the one that minimises the variance of the pointwise local energy E_L —and presents extensive numerical evidence for the $N=6$, $\omega=0.5$ quantum dot. The core observation is a *catch-22*: backflow’s learning signal is concentrated near the nodal surface, but the local energy diverges there for any imperfect wavefunction, and any sampling strategy that includes these points produces gradients with unbounded variance. The result is an irreducible accuracy floor—roughly 0.3%, set by the Jastrow alone—that is absent under VMC + SR training of the same ansatz.

Two things about the scope of this result should be stated before the details, so the floor is not misread. First, it is a property of the *strong form* specifically: the weak-form (score-function / REINFORCE) collocation objective used for the results in the main text never differentiates through the Laplacian and does not inherit this pathology, which is why it trains the same ansatz to far below the 0.3% floor. The value here is therefore a diagnosis of one objective’s limits, not a measure of what MCMC-free training can achieve. Second, the analysis is what motivated the shift to the weak form in the first place; it is presented as a worked negative result—the kind that redirects a research programme—rather than as a headline number.

C.1 Setup and notation

We consider a Slater–Jastrow–backflow ansatz of the form

$$\Psi_\theta(\mathbf{R}) = \det \tilde{S}_\uparrow(\mathbf{R}; \theta_{\text{BF}}) \det \tilde{S}_\downarrow(\mathbf{R}; \theta_{\text{BF}}) \exp(J(\mathbf{R}; \theta_J)), \quad (\text{C.1})$$

where $\tilde{S}_\sigma$ is the Slater matrix evaluated at backflow-shifted coordinates $\tilde{\mathbf{r}}_i = \mathbf{r}_i + \Delta \mathbf{x}_i(\mathbf{R}; \theta_{\text{BF}})$, J is a neural Jastrow factor (PINN), and $\theta = (\theta_J, \theta_{\text{BF}})$ are trainable parameters. A CTNN message-passing network produces the backflow shifts $\Delta \mathbf{x}_i$.

We train by minimizing the *screened collocation* objective

$$\mathcal{L}(\theta) = \text{Var}_{\mathcal{S}}[E_L(\mathbf{R}; \theta)], \quad E_L = \frac{\hat{H}\Psi_\theta}{\Psi_\theta}, \quad (\text{C.2})$$

where the collocation set $\mathcal{S}$ is obtained by importance-sampling from a Gaussian proposal q and retaining the top- K points ranked by $|\Psi_\theta|^2/q$ (“screened collocation”). This top- K selection naturally suppresses near-node configurations where $|\Psi|^2 \approx 0$.

C.2 The divergence of E_L near nodes

The local energy decomposes as

$$E_L = -\frac{1}{2} \frac{\nabla^2 \Psi}{\Psi} + V = -\frac{1}{2} \left[\nabla^2 J + |\nabla J|^2 + 2 \nabla J \cdot \frac{\nabla D}{D} + \frac{\nabla^2 D}{D} \right] + V, \quad (\text{C.3})$$

where $D := \det \tilde{S}_\uparrow \det \tilde{S}_\downarrow$ and the last term $\nabla^2 D/D$ contains second derivatives of the backflow shifts $\nabla^2 \Delta \mathbf{x}_i$ amplified by the inverse determinant.

For the exact wavefunction, E_L is constant. If Ψ is an eigenstate, $E_L = E$ everywhere, and the apparent D^{-1} singularity is cancelled by the numerator vanishing at the same rate as the denominator. Concretely, for a smooth wavefunction with a simple node at surface $\mathcal{N}$, both D and $\nabla^2 D$ vanish linearly in the signed distance d from $\mathcal{N}$:

$$D(\mathbf{R}) = \mathcal{O}(d), \quad \nabla^2 D(\mathbf{R}) = \mathcal{O}(d), \quad \frac{\nabla^2 D}{D} \text{ is finite on } \mathcal{N}. \quad (\text{C.4})$$

For an approximate wavefunction, the cancellation generically fails. During training, the backflow parameters θ_{BF} are imperfect: the *learned* nodal surface $\mathcal{N}_\theta$ (where $D(\cdot; \theta) = 0$) does not coincide with the *true* nodal surface. Let d_θ denote the signed distance from the learned node and d^* that from the true node. Near a point where $d_\theta \approx 0$ but $d^* \neq 0$, the determinant $D \sim d_\theta \rightarrow 0$ while the kinetic contributions from the Jastrow and potential terms remain finite:

$$E_L(\mathbf{R}) \supset \frac{\nabla^2 D(\mathbf{R}; \theta)}{D(\mathbf{R}; \theta)} \sim \frac{c}{d_\theta} \rightarrow \pm\infty \quad \text{as } d_\theta \rightarrow 0, \ d^* \neq 0, \quad (\text{C.5})$$

for some configuration-dependent constant $c \neq 0$. Conversely, near a point on the true node ($d^* = 0$) but off the learned node ($d_\theta \neq 0$), $\Psi \sim d^* \rightarrow 0$ and the wavefunction assigns vanishing weight. The pathology (C.5) is therefore not an artifact of the nodal structure itself—it is a property of *mismatched* nodes, and is worst precisely when backflow still has something to learn.

C.3 The geometric mismatch argument

We now show that the variance of the local energy (and hence of the training gradients) generically diverges for imperfect backflow parameters.

Consider the contribution to $\text{Var}[E_L]$ from a neighborhood U_ε of the learned nodal surface:

$$\text{Var}[E_L] \geq \int_{U_\varepsilon} |\Psi(\mathbf{R}; \theta)|^2 (E_L(\mathbf{R}; \theta) - \bar{E})^2 d\mathbf{R}. \quad (\text{C.6})$$

When the node is “simple” (D vanishes linearly), we have

$$|\Psi|^2 \sim d_\theta^2, \quad E_L^2 \sim d_\theta^{-2\alpha}, \quad (\text{C.7})$$

where α characterises the severity of the mismatch. For the generic case where the numerator $\nabla^2 D$ does *not* vanish on $\mathcal{N}_\theta$ (because $\mathcal{N}_\theta \neq \mathcal{N}^*$), one has $\alpha = 1$ so that $E_L^2 \sim d_\theta^{-2}$.

In D -dimensional configuration space, the volume element in a tubular neighborhood of a codimension-1 surface scales as $dV \sim d_\theta^0 d\sigma dd_\theta$. Thus the integrand in (C.6) behaves as

$$|\Psi|^2 \cdot E_L^2 \cdot dV \sim d_\theta^2 \cdot d_\theta^{-2} \cdot dd_\theta = dd_\theta, \quad (\text{C.8})$$

which integrates to $\mathcal{O}(\varepsilon)$ and is therefore *finite*: a simple node does not, by itself, make $\text{Var}[E_L]$ diverge under the $|\Psi|^2$ measure. The instability enters one derivative later, in the *parameter gradient*. But the key observation is that this is the *best case*. In practice, the mismatch between $\mathcal{N}_\theta$ and $\mathcal{N}^*$ is not uniform: there exist regions where the surfaces cross at angles, creating pockets where $|d^*|/|d_\theta|$ is large (the true wavefunction is non-negligible while the learned determinant is near-zero), and there the E_L^2 spike is not compensated by $|\Psi|^2$ suppression.

More concretely, the gradient $\partial \mathcal{L} / \partial \theta_{\text{BF}}$ involves terms of the form

$$\frac{\partial E_L}{\partial \theta_{\text{BF}}} \supset \frac{1}{D} \frac{\partial (\nabla^2 D)}{\partial \theta_{\text{BF}}} - \frac{\nabla^2 D}{D^2} \frac{\partial D}{\partial \theta_{\text{BF}}}, \quad (\text{C.9})$$

both of which diverge as $D \rightarrow 0$. Even in the marginal case where $\text{Var}[E_L]$ is finite, the *gradient variance* (which involves these higher-order poles) generically diverges, making gradient-based optimisation impossible at points near the learned node.

C.4 The catch-22

The preceding analysis reveals a three-sided dilemma:

1. **Backflow learns at the node.** The purpose of backflow is to reshape the nodal surface of the Slater determinant. The gradient signal $\partial E_L / \partial \theta_{\text{BF}}$ is largest where the backflow shifts actually change which side of the node a configuration point lies on, i.e., where $D \approx 0$.
2. **E_L diverges at the node for imperfect θ .** As shown in §C.2–C.3, the local energy and its parameter gradient contain poles of the form D^{-1} and D^{-2} whenever the learned node does not coincide with the exact node. The resulting gradient variance is unbounded.
3. **Good collocation sampling must avoid the node.** Under screened collocation, the top- K selection ranks points by $|\Psi|^2/q \propto D^2 e^{2J}/q$. Since $D^2 \rightarrow 0$ at the node, near-node points are systematically excluded from the training batch. This is *protective*—it prevents gradient explosion—but it also removes the points where backflow’s learning signal lives.

A “catch-22” emerges:

Including near-node points gives divergent, noisy gradients and backflow shrinks defensively or diverges. Excluding them gives stable training but removes backflow’s learning signal. In either regime, collocation cannot effectively train backflow beyond an accuracy floor set by the Jastrow (PINN) quality alone.

C.5 Why VMC + SR escapes

Under VMC, the sampling distribution is $|\Psi_\theta|^2$, which assigns vanishing measure to the nodal surface. Near-node configurations are strongly *suppressed*—for a simple node the probability of lying within a distance ϵ scales as ϵ^3 —rather than strictly absent, which is enough to keep the D^{-1} singularity out of the gradient estimator in practice. However, VMC *still optimises the node position* through a subtle mechanism:

Indirect nodal information through bulk correlations. The variational energy $\langle E_L \rangle_{|\Psi|^2}$ is sensitive to nodal position because shifting the node changes the wavefunction *everywhere* (through normalisation and through the functional form of D). The SR force vector $g_i = \text{Cov}_{|\Psi|^2}[E_L, O_i]$ (where $O_i = \partial_{\theta_i} \log \Psi$) receives contributions from the *bulk*—regions where both $|\Psi|^2$ and E_L are well-behaved—that are *correlated with* nodal position but do not require evaluating E_L at the node.

SR amplifies the signal. The Fisher metric $S_{ij} = \text{Cov}[O_i, O_j]$ captures how sensitive $\log \Psi$ is to each parameter direction. By solving $(S + \epsilon I) \delta \theta = -g$, SR amplifies exactly those subtle bulk signals that encode nodal information, compensating for the fact that the node itself has zero weight under $|\Psi|^2$.

Core asymmetry. VMC obtains nodal information *indirectly* through a well-conditioned integral (the covariance g_i is finite because E_L is sampled away from the node); collocation tries to obtain it *directly* through a divergent pointwise quantity (E_L evaluated near the node). None of the sampling, loss, or preconditioning strategies we tested (§C.6) overcame this asymmetry within the strong-form collocation framework; we present it as a structural obstruction of the tested pipeline rather than a theorem that no collocation method can ever avoid it.

C.6 Experimental evidence

We present a systematic study for $N=6$ electrons in a 2D harmonic trap with $\omega=0.5$ (DMC reference: $E_{\text{DMC}} = 11.78484(6)$ [44]). The ansatz uses a PINN Jastrow ($\sim 15,000$ parameters) and a CTNN backflow network ($\sim 8,600$ parameters) with `bf_scale = 0.7`. All energies are evaluated via MCMC with 15 000 samples after burn-in.

C.6.1 Baseline: PINN alone vs. PINN + CTNN

Table C.1: Baseline collocation results for $N=6$, $\omega=0.5$. “PINN only” trains without backflow; “PINN + CTNN joint” trains both networks from scratch for 300 epochs with `bf_scale = 0.7`. The backflow provides only a marginal improvement of ~ 0.1 percentage points.

Configuration	E (Ha)	σ_E	err (%)
PINN only (300 ep)	11.834	0.003	0.42
PINN + CTNN joint (300 ep)	11.823	0.003	0.33
PINN + CTNN separate opt	11.822	0.003	0.32
DMC reference	11.78484	—	0.00

The backflow reduces the error from 0.42% to 0.32–0.33%, far short of the $< 0.01\%$ accuracy achievable with VMC + SR on the same ansatz (Table 6.2).

C.6.2 Sampling strategies

We tested multiple strategies to bring near-node information into the collocation batch, hoping to give the backflow a stronger learning signal.

Table C.2: Effect of sampling strategy on collocation accuracy. All experiments use 50-epoch fine-tuning from the best collocation checkpoint (0.35%), except “hard-smooth (scratch)” which trains 300 epochs from random initialisation.

Method	Description	err (%)	Outcome
Top- K baseline	standard screened collocation	0.35	stable
Hard mining + Huber	top- K of residuals	0.41	neutral
Hard mining + smoothness	+ $\ \nabla^2 \Delta x\ ^2$ penalty	0.38	slight help
SIR (single proposal)	$ \Psi ^2$ -reweighted resampling	0.47	worse
SIR (mixture)	Gaussian mixture proposal	0.44	worse
Gumbel top- K	stochastic top- K selection	5.00	catastrophic

Every strategy that includes more near-node points (SIR, Gumbel) produced *worse* results—consistent with the catch-22: the additional points carry divergent E_L values that corrupt the gradient. Gumbel top- K (which replaces deterministic selection with stochastic Gumbel-softmax) was catastrophic, with early stopping triggered by rapid energy divergence.

C.6.3 Loss modifications

Table C.3: Effect of loss function modifications. The hybrid loss adds a mean-energy (variational) term; the det filter discards points with small $|\det \tilde{S}|$.

Method	Description	err (%)	Outcome
Variance loss (baseline)	Huber on $E_L - \bar{E}$	0.33	best collocation
Hybrid anneal	$\beta \cdot \text{Var} + (1-\beta) \cdot \text{Mean}$, $\beta: 0.7 \rightarrow 0.15$	2.59	catastrophic
Gumbel + hybrid	stochastic top- K + hybrid	5.00	catastrophic
Hybrid + det filter	$\beta=0.2$ fixed + $ \det $ filter	24.75	broken

The mean-energy gradient $\nabla_\theta \langle E_L \rangle_S$ under biased collocation sampling does *not* equal the VMC gradient—the sampling distribution is not $|\Psi|^2$ —so the variational principle does not hold, and the backflow drifts to large, unphysical shifts ($|\Delta \mathbf{x}| \approx 0.83$, versus ~ 0.23 under variance-only training). The variance loss’s implicit constraint on backflow magnitude is *protective*, not pathological.

Table C.4: Regularisation of the E_L singularity (50-epoch fine-tune from 0.35% checkpoint). Each modification smooths or clips one of the divergent contributions to E_L .

Method	Modification	err (%)	Outcome
Soft Coulomb	$1/r \rightarrow 1/\sqrt{r^2 + \varepsilon^2}$, $\varepsilon=0.2$	1.41	much worse
Regularised det	$\log D \rightarrow \log \text{addexp}(\log D , \log \delta)$	0.49	worse
T clipping	$\max(T , T_{\max}), T_{\max}=30$	0.43	neutral

C.6.4 Gradient regularisation

We attacked the divergence sources in E_L directly:

Soft Coulomb trains on wrong physics (V is no longer the true potential). Regularising the determinant distorts the $\log \Psi$ landscape. Kinetic energy clipping is neutral because the clipped points are exactly those that carried the (noisy) backflow signal. None of the regularisations improve upon the baseline.

C.6.5 Gradient preconditioning

Table C.5: Gradient preconditioning experiments (300-epoch scratch training unless noted). K-FAC uses block-diagonal Kronecker-factored Fisher on the backflow network only.

Method	err (%)	Outcome
Vanilla Adam	0.33	baseline
K-FAC on CTNN	0.34	no improvement
K-FAC + smoothness	0.51	worse
Variance-minimising pretrain	0.54	worse
Adam fine-tune (30 ep, lr=1e-4)	0.34	no improvement
Adam burst (15 ep, lr=3e-4)	0.56	worse

K-FAC provides a block-diagonal approximation to the Fisher matrix and should, in principle, help with the ill-conditioned parameter-space geometry. In practice, it does not: the conditioning problem is *operator-level* (Appendices A–B), not merely parameter-space anisotropy, and K-FAC cannot address the fundamental divergence in E_L .

C.6.6 Summary of all interventions

The pattern is striking:

- Every intervention that *includes* more near-node points (SIR, Gumbel) *increases* the error—backflow shrinks defensively under gradient noise or diverges entirely.
- Every intervention that *changes* the loss to tolerate near-node contributions (hybrid mean-energy, det filter) produces catastrophic results because the collocation sampling is not $|\Psi|^2$ and the variational principle does not apply.
- The smoothness penalty $\lambda \|\nabla^2 \Delta \mathbf{x}\|^2$ helps modestly (from 0.59% to 0.38% without it vs. with it) by damping the $\nabla^2 D/D$ poles, but cannot eliminate the fundamental divergence.
- Ironically, the only method that trains backflow effectively is MCMC-based VMC + SR—which sidesteps the entire collocation framework.

C.7 Connection to Appendices A and B

The catch-22 is a specific manifestation of the general conditioning difficulties analysed in Appendices A–B:

Table C.6: Complete summary of all collocation-based backflow experiments for $N=6$, $\omega=0.5$. No intervention breaks the $\sim 0.32\%$ accuracy floor. For comparison, the same PINN + CTNN ansatz achieves $< 0.01\%$ under VMC + SR (Table 6.2).

Experiment	err (%)	Category
PINN + CTNN sep opt (best)	0.32	baseline
PINN + CTNN joint 300 ep	0.33	baseline
K-FAC on CTNN	0.34	preconditioner
Adam fine-tune 30 ep	0.34	optimiser
Top- K baseline (retrained)	0.35	sampling
Hard mining + smoothness	0.38	sampling
Hard mining + Huber	0.41	sampling
PINN only	0.42	no backflow
T clipping	0.43	regularisation
SIR (mixture)	0.44	sampling
SIR (single)	0.47	sampling
Regularised det	0.49	regularisation
K-FAC + smoothness	0.51	preconditioner
Var-min pretrain	0.54	pretrain
Adam burst lr=3e-4	0.56	optimiser
Soft Coulomb	1.41	regularisation
Hybrid loss (annealed)	2.59	loss
Gumbel top- K + hybrid	5.00	sampling + loss
Hybrid + det filter	24.75	loss + filter
Same ansatz, VMC + SR	<0.01	variational

Laplacian stiffness (Appendix A). The k^4 -type amplification of curvature errors (§A.3) is compounded by backflow, because $\nabla^2(\det \tilde{S})$ introduces second derivatives of $\Delta \mathbf{x}$ that are not present in a fixed-node ansatz. These additional high-frequency modes inflate the spectral spread of the training operator L^*L .

Coulomb singularity (Appendix B). The $1/r$ Coulomb potential (§B.1) adds a singular multiplication operator to $\hat{H}$. Near both the coalescence point ($r_{ij} \rightarrow 0$) and the node ($D \rightarrow 0$), E_L contains products of these two singularities, and the cusp cancellation (§B.2) must be maintained simultaneously with the nodal cancellation (C.4). In the strong residual squared $\|LE_L\|^2$, these compound singularities create an extremely stiff landscape.

The backflow-specific pathology. What distinguishes the backflow catch-22 from the general conditioning problem is the *self-referential* nature of the singularity: the backflow parameters θ_{BF} determine *where* the node is, so updating θ_{BF} moves the singularity in configuration space. Under the collocation loss, the gradient pushes θ_{BF} in a direction determined by E_L values at the current collocation points, but the resulting node shift invalidates those very points. VMC+SR avoids this because it samples from the *current* $|\Psi_\theta|^2$ at every step, so the sampling distribution co-adapts with the node position.

Chapter D

Post-catch-22 experimental history and philosophy

This appendix documents the full experimental program executed after the backflow catch-22 analysis in Appendix C. The goal here is not to present a single best number, but to make explicit the strategy, the failed branches, the partial recoveries, and the current status for difficult regimes ($\omega \in \{0.01, 0.001\}$ and larger N).

D.1 The experimental programme

After the catch-22 analysis the work shifted from searching for a single better architecture to engineering a training pipeline that makes a physically strong ansatz trainable across regimes. Four principles organised it: keep the Slater–Jastrow–backflow ansatz but treat sampling and preconditioning as first-class objects; replace one-shot “hero” runs with staged continuation (solve easy regimes, warm-start hard ones); separate claims by evidence tier (online probe, restored VMC-best, heavy exact VMC); and record negative results, because many theoretically plausible variants fail on gradient-noise geometry.

What was varied. We varied the *sampling* (Gaussian and Gaussian-mixture proposals scaled by $1/\sqrt{\omega}$; oversample-then-resample with ESS diagnostics; ESS-adaptive oversampling; weight tempering and log-weight clipping), the *objective* (finite-difference collocation residuals; a hybrid weak form; and pure RE-INFORCE, which became the reliable default, with the Laplacian kept in the reward path but never differentiated), the *preconditioner* (Adam; diagonal-Fisher natural gradient; and full SR in Woodbury/CG modes with damping anneals and trust-region caps), and the *training style* (single sweeps; wave-based cascades with checkpoint inheritance; asynchronous campaigns that relaunch from the best available checkpoint). Two empirical conclusions were robust. First, optimisation quality dominated architectural novelty: moderately sized BF+Jastrow models with stable training beat more ambitious variants with weaker optimisation geometry. Second, at low ω regularising the resampling distribution (ESS control, top-mass reduction) helped while hard cusp gating alone did not—a controlled $2 \times 2 \times 3$ matrix (hard gate on/off, resampling-regularisation on/off, three seeds) confirmed both, with no measurable gain from gating and a consistent gain from resampling regularisation.

Outcomes. SR-style preconditioning in CG mode, warm-started along the confinement cascade $\omega : 1.0 \rightarrow 0.5 \rightarrow 0.1 \rightarrow 0.01 \rightarrow 0.001$, was what carried performance from solved regimes into harder ones; cold-start SR at low ω did not converge. The full set of best energies is in §6.4.3; representative cascade successes are $N=6$ at $\omega=1.0$ ($E = 20.16167 \pm 0.00181$, +0.012%), $\omega=0.5$ ($E = 11.78466 \pm 0.00199$, −0.002%) and $\omega=0.1$ ($E = 3.56525 \pm 0.00046$, +0.32%), and $N=12$ at $\omega=1.0$ ($E = 65.70699 \pm 0.00595$, +0.010%). For $N=2$ the collocation route now covers the whole confinement range cleanly—+0.012%, +0.023%, −0.004%, +0.002% and +0.154% at $\omega = 1, 0.5, 0.1, 0.01, 0.001$ (zero-variance extrapolated)—so the paradigm is DMC-accurate wherever the importance sampling stays healthy and degrades only in the deep-Wigner limit.

D.2 Current frontier: low- ω and high- N stabilisation

The current unresolved front remains the combination of extreme correlation regimes and larger particle count:

- $\omega \in \{0.01, 0.001\}$: transfer helps, but reliability and reproducibility are not yet at the same confidence level as $\omega \in \{1.0, 0.5, 0.1\}$.
- $N \geq 20$: memory pressure, architecture-capacity constraints, and higher gradient noise make training significantly less stable than $N \in \{6, 12\}$.

Dedicated stabilisation runs introduced stricter SR trust constraints, heavier damping, ESS-adaptive sampling, rollback controls, and explicit process-level relaunch hygiene for GPU memory contention. These changes improved operational stability but have not yet closed the full accuracy gap for the hardest targets.

The $N=20$ collocation results. For completeness we record here the $N=20$ collocation energies referred to in §6.4.3; they are the first collocation-trained wavefunctions at this size, and they show the memory bottleneck in the clearest possible form. The backflow’s edge-feature tensor scales as $\mathcal{O}(N^2 \times h_{\text{bf}})$, and at $N=20$ fitting it in memory forces the hidden dimension down to $h_{\text{bf}}=48\text{--}64$. The best backflow run at $\omega=1.0$ then reached only $\sim+18\%$ error, while a Jastrow-only ansatz—freed of that tensor, and so able to afford a larger collocation budget and higher oversampling—reached $+1.32\%$ ($E = 157.934 \pm 0.018$). The architecture that should in principle be the more expressive is beaten by the simpler one, purely because the memory-forced capacity cut degrades its gradients more than its extra flexibility repays. Polishing the Jastrow checkpoints at reduced learning rate improved the $\omega=1.0$ error from $\sim+2.6\%$ to $+1.32\%$ over ~ 400 epochs and the $\omega=0.5$ error from $\sim+7.0\%$ to $+2.74\%$ ($E = 96.447 \pm 0.015$); the best $\omega=0.1$ result was $+5.53\%$ ($E = 31.637 \pm 0.008$). These are honest first numbers. The gap to the sub-percent accuracy reached at $N=6, 12$ follows from the memory-forced capacity reduction and the gradient noise it brings, not from the collocation objective; closing it is a question of a more memory-frugal backflow, not of a different method.

D.3 Interpretive summary

The post-catch-22 phase supports six robust conclusions.

1. The collocation program is not a single recipe but a coupled system: sampling, objective, and preconditioning must be tuned jointly.
2. Transfer/cascade is not optional for difficult regimes; it is the primary mechanism by which low- ω performance becomes attainable.
3. SR-style preconditioning is central for stable progress beyond early Adam-only basins.
4. Diagnostic metrics (ESS, top-mass concentration, rollback counts, probe-final gaps) are essential; scalar final energy alone is insufficient for method development.
5. Several intuitively attractive interventions (e.g., hard cusp gating in this tested range) did not deliver gains, and documenting these negatives was necessary to avoid circular search.
6. The method now demonstrates near-DMC quality in multiple nontrivial targets, but the strongest claims must still be bounded to regimes where repeated campaign evidence exists.

Chapter E

Declaration on the Use of Artificial Intelligence

The collocation approach in this thesis was developed over several months of failed experiments. Its central difficulty was not merely that a network did not converge, but that the parts of the Schrödinger residual that made the method physically meaningful also made its gradients unstable near the wavefunction's nodes. During that period, AI systems were used as sounding boards: to test interpretations of unexpected results, to formulate possible explanations, and to decide which experiment should come next. The exchange below is a short sample from one longer conversation at a point where the negative evidence had begun to make abandoning collocation seem plausible.

Author

I am trying to develop a PINN to find ground-state solutions of the Schrödinger equation. I have tried multiple approaches, but I am not succeeding. ...What do you think about this? What should we do from here?

AI response 1

...Anything that injects near-node points makes gradients explode or become uselessly noisy. ...

⋮

AI response 2

...This asymmetry is fundamental and cannot be resolved by sampling tricks, loss modifications, or gradient preconditioning within the collocation framework. ...

⋮

AI response 3

...Stick with SR + energy (or energy+variance) VMC. ...

Artificial-intelligence systems were therefore used for discussion, planning, and review, rather than as a source of results. They helped turn observations into hypotheses that could be checked, suggested alternative diagnostics when a run failed, and reviewed the final prose for clarity and mistakes. The code, the experiments, the scientific judgments, and the text of this thesis were written by the author. In particular, the recommendation shown above was not accepted as a methodological conclusion: it became one claim among several to investigate. The work that followed led to the collocation analysis reported in this thesis.

Bibliography

- [1] Giuseppe Carleo and Matthias Troyer. Solving the quantum many-body problem with artificial neural networks. *Science*, 355(6325):602–606, 2017. doi: 10.1126/science.aag2302.
- [2] David Pfau, James S. Spencer, Alexander G. D. G. Matthews, and W. Matthew C. Foulkes. Ab initio solution of the many-electron schrödinger equation with deep neural networks. *Physical Review Research*, 2:033429, 2020.
- [3] Jan Hermann, Zeno Schätzle, and Frank Noé. Deep-neural-network solution of the electronic schrödinger equation. *Nature Chemistry*, 12:891–897, 2020. doi: 10.1038/s41557-020-0544-y.
- [4] Di Luo and Bryan K. Clark. Backflow transformations via neural networks for quantum many-body wave functions. *Physical Review Letters*, 122:226401, 2019. doi: 10.1103/PhysRevLett.122.226401.
- [5] Gabriel Pescia, Jannes Nys, Jane Kim, Alessandro Lovato, and Giuseppe Carleo. Message-passing neural quantum states for the homogeneous electron gas. *Physical Review B*, 110(3):035108, 2024. doi: 10.1103/PhysRevB.110.035108.
- [6] Jane Kim, Gabriel Pescia, Bryce Fore, Jannes Nys, Giuseppe Carleo, Stefano Gandolfi, Morten Hjorth-Jensen, and Alessandro Lovato. Neural-network quantum states for ultracold fermi gases, 2023. Preprint; details as provided in source list.
- [7] R. Egger, W. Häusler, C. H. Mak, and H. Grabert. Crossover from fermi liquid to wigner molecule behavior in quantum dots. *Phys. Rev. Lett.*, 82:3320–3323, Apr 1999. doi: 10.1103/PhysRevLett.82.3320. URL <https://link.aps.org/doi/10.1103/PhysRevLett.82.3320>.
- [8] Minghui Kong, B. Partoens, and F. M. Peeters. Transition between ground state and metastable states in classical two-dimensional atoms. *Phys. Rev. E*, 65:046602, Mar 2002. doi: 10.1103/PhysRevE.65.046602. URL <https://link.aps.org/doi/10.1103/PhysRevE.65.046602>.
- [9] Vitaly A. Schweigert and François M. Peeters. Spectral properties of classical two-dimensional clusters. *Phys. Rev. B*, 51:7700–7713, Mar 1995. doi: 10.1103/PhysRevB.51.7700. URL <https://link.aps.org/doi/10.1103/PhysRevB.51.7700>.
- [10] M. Manninen and S.M. Reimann. Chapter 12 - metal clusters, quantum dots, and trapped atoms: From single-particle models to correlation. In Purusottam Jena and A. Welford Castleman, editors, *Nanoclusters*, volume 1 of *Science and Technology of Atomic, Molecular, Condensed Matter and Biological Systems*, pages 437–484. Elsevier, 2010. doi: <https://doi.org/10.1016/B978-0-444-53440-8.00012-4>. URL <https://www.sciencedirect.com/science/article/pii/B9780444534408000124>.
- [11] M. Mazars. Bond orientational order parameters in the crystalline phases of the classical yukawa-wigner bilayers. *Europhysics Letters*, 84(5):55002, dec 2008. doi: 10.1209/0295-5075/84/55002. URL <https://doi.org/10.1209/0295-5075/84/55002>.
- [12] A. V. Filinov, M. Bonitz, and Yu. E. Lozovik. Wigner crystallization in mesoscopic 2d electron systems. *Phys. Rev. Lett.*, 86:3851–3854, Apr 2001. doi: 10.1103/PhysRevLett.86.3851. URL <https://link.aps.org/doi/10.1103/PhysRevLett.86.3851>.
- [13] Walter Rudin. *Principles of Mathematical Analysis*. McGraw-Hill, 3 edition, 1976.
- [14] Arjun Berera and Luigi Del Debbio. *Quantum Mechanics*. Cambridge University Press, 2021.

- [15] David J. Griffiths and Darrell F. Schroeter. *Introduction to Quantum Mechanics*. Cambridge University Press, 2018.
- [16] Tosio Kato. On the eigenfunctions of many-particle systems in quantum mechanics. *Communications on Pure and Applied Mathematics*, 10:151, 1957.
- [17] Dirk P. Kroese, Tim Brereton, Thomas Taimre, and Zdravko I. Botev. Why the monte carlo method is so important today. *Wiley Interdisciplinary Reviews: Computational Statistics*, 6:386, 2014.
- [18] Ioan Kosztin, Byron Faber, and Klaus Schulten. Introduction to the diffusion monte carlo method. *American Journal of Physics*, 64:633, 1996.
- [19] Nicholas Metropolis, Arianna W. Rosenbluth, Marshall N. Rosenbluth, Augusta H. Teller, and Edward Teller. Equation of state calculations by fast computing machines. *The Journal of Chemical Physics*, 21:1087, 1953.
- [20] Siu A. Chin. Quadratic diffusion monte carlo algorithms for solving atomic many-body problems. *Physical Review A*, 1990.
- [21] Nicolaas Godfried van Kampen. *Stochastic Processes in Physics and Chemistry*, volume 1. Elsevier, 1992.
- [22] Julius Berner, Philipp Grohs, Gitta Kutyniok, and Philipp Petersen. *The Modern Mathematics of Deep Learning*, page 1–111. Cambridge University Press, 2022.
- [23] Trevor Hastie, Robert Tibshirani, and Jerome H. Friedman. *The Elements of Statistical Learning: Data Mining, Inference, and Prediction*, volume 2. Springer, 2009.
- [24] Lei Wu, Mingze Wang, and Weijie Su. The alignment property of sgd noise and how it helps select flat minima: A stability analysis, 2022. URL <https://arxiv.org/abs/2207.02628>.
- [25] Diederik Kingma and Jimmy Ba. Adam: A method for stochastic optimization. In *International Conference on Learning Representations (ICLR)*, San Diego, CA, USA, 2015.
- [26] Yushun Zhang, Congliang Chen, Naichen Shi, Ruoyu Sun, and Zhi-Quan Luo. Adam can converge without any modification on update rules. In *Advances in Neural Information Processing Systems*, page 35:28386, 2022.
- [27] Shun ichi Amari. Natural gradient works efficiently in learning. *Neural Computation*, 10:251, 1998.
- [28] S. Amari and S.C. Douglas. Why natural gradient? In *Proceedings of the 1998 IEEE International Conference on Acoustics, Speech and Signal Processing, ICASSP '98 (Cat. No.98CH36181)*, volume 2, pages 1213–1216 vol.2, 1998. doi: 10.1109/ICASSP.1998.675489.
- [29] Federico Becca and Sandro Sorella. *Quantum Monte Carlo approaches for correlated systems*. Cambridge University Press, 2017.
- [30] James Stokes, Josh Izaac, Nathan Killoran, and Giuseppe Carleo. Quantum natural gradient. *Quantum*, 4:269, 2020.
- [31] Arthur Jacot, Franck Gabriel, and Clément Hongler. Neural tangent kernel: Convergence and generalization in neural networks. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2018. URL <https://arxiv.org/abs/1806.07572>.
- [32] Manzil Zaheer, Satwik Kottur, Siamak Ravanbakhsh, Barnabas Poczos, Russ R. Salakhutdinov, and Alexander J. Smola. Deep sets. In *Advances in Neural Information Processing Systems*, volume 30, 2017.
- [33] Justin Gilmer, Samuel S. Schoenholz, Patrick F. Riley, Oriol Vinyals, and George E. Dahl. Neural message passing for quantum chemistry. In *Proceedings of the 34th International Conference on Machine Learning (ICML)*, volume 70 of PMLR, pages 1263–1272, 2017. URL <https://arxiv.org/abs/1704.01212>.
- [34] Peter W. Battaglia, Jessica B. Hamrick, Victor Bapst, et al. Relational inductive biases, deep learning, and graph networks. *arXiv preprint arXiv:1806.01261*, 2018. URL <https://arxiv.org/abs/1806.01261>.

- [35] George Cybenko. Approximation by superpositions of a sigmoidal function. *Mathematics of Control, Signals and Systems*, 2:303–314, 1989. doi: 10.1007/BF02551274. URL <https://doi.org/10.1007/BF02551274>.
- [36] Kurt Hornik. Approximation capabilities of multilayer feedforward networks. *Neural Networks*, 4(2):251–257, 1991. ISSN 0893-6080. doi: [https://doi.org/10.1016/0893-6080\(91\)90009-T](https://doi.org/10.1016/0893-6080(91)90009-T). URL <https://www.sciencedirect.com/science/article/pii/089360809190009T>.
- [37] Tim De Ryck and Siddhartha Mishra. Numerical analysis of physics-informed neural networks and related models in physics-informed machine learning. *Acta Numerica*, 33:633–713, 2024. doi: 10.1017/S0962492923000089.
- [38] Sandro Sorella. Green function monte carlo with stochastic reconfiguration. *Physical Review Letters*, 80(20):4558–4561, 1998. doi: 10.1103/PhysRevLett.80.4558.
- [39] Ronald J. Williams. Simple statistical gradient-following algorithms for connectionist reinforcement learning. *Machine Learning*, 8(3–4):229–256, 1992. doi: 10.1007/BF00992696.
- [40] Malvin H. Kalos, Dominique Levesque, and Loup Verlet. Helium at zero temperature with hard-sphere and other forces. *Physical Review A*, 9:2178, 1974.
- [41] Aki Vehtari, Daniel Simpson, Andrew Gelman, Yuling Yao, and Jonah Gabry. Pareto smoothed importance sampling. *Journal of Machine Learning Research*, 25, 2024. URL <https://arxiv.org/abs/1507.02646>.
- [42] H. Flyvbjerg and H. G. Petersen. Error estimates on averages of correlated data. *The Journal of Chemical Physics*, 91:461, 1989. American Institute of Physics.
- [43] Marius Jonsson. Standard error estimation by an automated blocking method. *Physical Review E*, 98:043304, 2018.
- [44] F. Pederiva, C. J. Umrigar, and E. Lipparini. Diffusion monte carlo study of circular quantum dots. *Phys. Rev. B*, 62:8120–8125, 2000. doi: 10.1103/PhysRevB.62.8120.
- [45] Daniel Haas. Deep learning methods for quantum many-body systems: A study on neural quantum states. Master’s thesis, University of Oslo, September 2024. Provides correlated-level (DMC-quality) reference energies for 2D parabolic quantum dots.
- [46] Chae-Yeun Park and Michael J. Kastoryano. Geometry of learning neural quantum states. *Physical Review Research*, 2:023232, 2020.
- [47] Sami Siljamäki. *Wave Function Methods for Quantum Dots in Magnetic Field*. PhD thesis, Helsinki University of Technology, 2003.