

Status of Vitamin D in Children with Pediatric Acute-Onset Neuropsychiatric Syndrome (PANS)

G. Delgado, R.C. Kaufmann, D.D. Siri.

Illinois Math and Science Academy / Midwest Allergy Sinus Asthma / SWIA Clinical Research, Normal, IL.

RATIONALE

- Numerous studies have indicated an association between vitamin D deficiency and the immune dysregulation and the pathogenesis of autoimmunity.
- An increased prevalence of vitamin D deficiency has been demonstrated in several pediatric autoimmune diseases, including autoimmune thyroid disease and Type-1 diabetes mellitus.
- Identification of children with Vitamin D deficiency may lead to the opportunity to influence the risk factors lead to the development of autoimmune diseases by vitamin D supplementation.

OBJECTIVE

- This study was undertaken to determine the prevalence of vitamin D deficiency in children with PANS/PANDAS, an autoimmune disorder characterized by abrupt-onset neuropsychiatric symptoms, often associated with infections such as Group A Streptococcus.

METHODS

- A chart review was performed retrospectively from a single PANS/PANDAS treatment center. Data from patients who presented to the specialty clinic for evaluation of PANS/PANDAS was collected. Additional information collected included 25-hydroxy vitamin D levels, age, gender, race, and age of diagnosis.
- Vitamin D status was stratified into three groups; deficiency (0 to < 20 ng/ml), insufficiency (20 to < 30 ng/ml), and sufficiency (30-100 ng/ml) (1). Patients were also stratified into two age groups; child (< 12 years old) and adolescent (12-17 years old).

RESULTS

- 91/129 patients with the diagnosis of PANS/PANDAS were able to be evaluated.

	Deficient	Insufficient	Sufficient	Total (n=91)
Child (<12 yo)	6 (8%)	18 (24%)	51 (68%)	75
Adolescent (12-17 yo)	1 (6.3%)	4 (25%)	11 (68.8%)	16
Male	4 (8.3%)	12 (25%)	32 (66.7%)	48
Female	3 (7%)	10 (23.3%)	30 (69.8%)	43
Total	7 (7.7%)	22 (24.2%)	62 (68.1%)	91

Table 1. Prevalence of vitamin D status by age and gender.




Figure 1. Age at PANS/ PANDAS diagnosis vs. 25-hydroxy vitamin D.

Figure 2. Vitamin D status of PANS/PANDAS pediatric population.

	PANS/PANDAS Cohort	US Child Cohort
Vitamin D Deficiency	7.7%	9%
Vitamin D Insufficiency	24.2%	61%

Odds Ratio: 0.843505 95% CI (-2.91 to 4.59)

Table 2. Prevalence of vitamin D status is shown in the PANS/PANDAS cohort compared with the prevalence in a recently studied in a general US pediatric population.(Kumar 2009 *Pediatrics*). The Odds Ratio comparing the two cohorts is noted, indicating a lower rate of vitamin D deficiency or insufficiency in the PANS/PANDAS cohort studied, versus the general US pediatric population.

CONCLUSIONS/DISCUSSION

- Results indicate that our patient population of children with PANS/PANDAS had a lower risk of vitamin D deficiency and insufficiency status than those of the general US pediatric population regardless of their age or gender.
- The results are inconsistent with the high prevalence of vitamin D deficiency/insufficiency status reported in children with other autoimmune diseases.
- Various factors that contribute to this negative study:
 - 97.7% of our patients are white (non-Hispanic). Children in our cohort were disproportionately skewed toward white race and a higher socioeconomic status than participants in the general US pediatric comparator study. Children with skin-of-color and lower socioeconomic status have a higher risk factor for vitamin D deficiency/insufficiency.
 - A higher rate of vitamin supplementation may have been present in our PANS/PANDAS cohort, due to earlier and more frequent medical evaluation and intervention, diagnostic delays and challenges contributing to higher nutritional interventions, and/or parental choice.
- More studies are needed to evaluate a larger, more diverse population of children with PANS/PANDAS and vitamin D status at earlier time points, prior to supplementation.
- Although our cohort population is small and from a single center, results also suggest that PANS/PANDAS may be underdiagnosed in children with skin-of-color and in those with lower socioeconomic status.

REFERENCES

- Endocrine Society of America 2. Yang et al. Clin Rev All Imm 2013 3. Dankers et al. Front Imm 2017 4. Kumar et al. Pediatrics 2009 5. Ranganathan et al. Pros Clin Res 2015 6. Szumilas, et al. J Can Acad Child Adolesc Psych 2010 7. Yeşiltepe-Mutlu et al. J Clin Res Ped Endo 2020 8. Taksler et al. Pub Health Nutr 2015



Acknowledgements: This study was self-funded. We thank the Illinois Math and Science Academy Business Mentorship program for supporting the research initiative, Bim Thomas, Christ Sengson, and Henry Anayabor for data entry assistance.