

Temă

Laborator 2.

a) $V = \{1, 2, 3, 4, 5, 6, 7\}$


$E = \{(1, 7), (7, 2), (2, 4), (4, 3), (4, 5), (5, 2), (2, 6)\}$

b) Modul 1: are o singură muchie incidentă, respectiv $(1, 7) \Rightarrow \text{Grad } 1$

Modul 2: are 3 muchii incidente, respectiv $(7, 2), (5, 2), (2, 6) \Rightarrow \text{Grad } 3$


Modul 5: are 2 muchii incidente, respectiv $(4, 5), (5, 2) \Rightarrow \text{Grad } 2$

c) Subgraf:



S-au eliminat nodurile $\{3, 1, 6\}$

Graf partial:



S-au eliminat muchiile $[7, 2]$, $[4, 3]$

d) Lanț:

$[3, 4, 5, 2, 6]$

Lanț elementar:


$[1, 7, 4, 5, 2]$

e) Ciclu:

$[7, 4, 5, 2, 7]$

Ciclu elementar:

$[7, 4, 5, 2, 7]$




1) Nodurile 1, 2, 3, 4, 5, 6 și 7 au grade impare
cea ce rezulta că grafel nu poate fi lanț earehem

Pavel Moravcs Floren Adrian

Grupa 2

1) Permutare 1, 7, 2, 4, 6, 5, 3



b)

	1	2	3	4	5	6	7
1	0	0	0	0	0	0	1
2	0	0	0	0	1	1	1
3	0	0	0	1	0	0	0
4	0	0	1	0	1	0	1
5	0	1	0	1	0	0	0
6	0	1	0	0	0	0	0
7	1	1	0	1	0	0	0

i)

	(1,7)	(2,7)	(7,4)	(4,3)	(4,5)	(5,2)	(2,6)
1	1	0	0	0	0	0	0
2	0	1	0	0	0	1	1
3	0	0	0	1	0	0	0
4	0	0	1	1	1	0	0
5	0	0	0	0	0	1	0
6	0	0	0	0	0	0	1
7	1	1	1	0	0	0	0

1: [1]
 2: [7, 5, 6]
 3: [4]
 4: [7, 3, 5]
 5: [2, 4]
 6: [2]