



Journal

Communications in Soil Science and Plant Analysis >

Volume 37, 2006 - Issue 15-20

54 0
Views CrossRef citations Altmetric

Poster Paper

Effect of Clinoptilolite Addition to Soil on Wheat Yield and Nitrogen Uptake

C. D. Tsadilas & G. Argyropoulos

Pages 2691-2699 | Received 14 Feb 2005, Accepted 14 Oct 2005, Published online: 31 Oct 2011

[Download citation](#) <http://dx.doi.org/10.1080/00103620600830088>

Select Language ▼

[Translator disclaimer](#)

Abstract

During the past years, appreciable amounts of zeolite-rich tuff that contains more than 70% clinoptilolite (Cp) have been discovered in Greece. The present study evaluates the ability of natural Greek Cp to increase the efficiency of nitrogen (N) fertilizer uptake in wheat. A pot experiment with winter wheat was conducted in a Typic Xerorthent that was fertilized with ammonium sulfate and amended with 0 to 60 ton/ha of Cp. Clinoptilolite application resulted in an increase of the cation exchange capacity of the soil from 9.5 to 13.6 meq/100 g (i.e., 43%). Soil ammoniacal N was greater in the samples amended with Cp at the boot stage, as was NO₃-N. Clinoptilolite addition increased total wheat yield (dry matter of hay plus seed) 52% (from 21.1 g/pot in the control to 32.0 g/pot) in the treatment with 60 ton/ha of Cp. The influence was greater for seed yield than hay yield. Clinoptilolite addition resulted in high increase in total N uptake, about 141% (from 156 mg/pot in the control to 376 mg/pot) in the treatment with the highest Cp rate. For hay, the increase was 133% (from 125 mg/pot to 291 mg/pot), whereas for seed it was 126% (from 31 mg/pot to 70 mg/pot) from control to the treatment with the highest Cp rate, resulting in a more efficient N fertilizer use. The optimum Cp addition rate was estimated as large as 15 ton/ha.

Keywords: [Clinoptilolite](#), [N efficiency](#), [nitrate leaching](#), [ammonium fixation](#)