

Mathematics

Imahn Shekhzadeh

July 25, 2025

Contents

1	Algebra	2
2	Measure Theory	5
3	Analysis	8
4	Inner Product and Normed Spaces	10
5	Metric Spaces	15
5.1	Complete metric spaces	23
5.2	Compactness	37
5.3	Baire Category Theorem	44
5.4	Continuity	45
6	Topological Spaces	49
6.1	Hausdorff Spaces	74
6.2	Continuity and Homeomorphisms	76
7	Linear Functional Analysis	80
7.1	Bounded Linear Operators	80
8	Set Theory	83
9	Differential Geometry	90
10	Normalizing Flows	91
11	Miscellaneous	95
A	Weierstrass Approximation Theorem	98
B	Construction of \mathbb{R}	105
	References	110

1 Algebra

Definition 1.1 (Action). Let G be a group (where e is the neutral element), and X a set. Then a (left) group action α of G on X is a function

$$\alpha : G \times X \rightarrow X \quad (1.1)$$

satisfying the following two axioms $(\forall g, h \in G) \wedge (\forall x \in X)$:

$$\begin{aligned} \alpha(e, x) &= x, \\ \alpha(g, \alpha(h, x)) &= \alpha(gh, x) \end{aligned}$$

The group action is commonly written as $g \cdot x$ s.t. the axioms become

$$e \cdot x = x, \quad (1.2)$$

$$g \cdot (h \cdot x) = (gh) \cdot x \quad (1.3)$$

A right group action of G on X is a function

$$\alpha : X \times G \rightarrow X \quad (1.4)$$

satisfying the following two axioms $(\forall g, h \in G) \wedge (\forall x \in X)$:

$$\alpha(x, e) = x, \quad (1.5)$$

$$\alpha(\alpha(x, g), h) = \alpha(x, gh) \quad (1.6)$$

Example 1.2. Consider $G = \text{SO}(2)$, i.e. the group of all rotations in \mathbb{R}^2 , with the group operation given by the composition of rotations, and let $X = \mathbb{S}^1$ be the unit circle in \mathbb{C} . Now consider the group action

$$\Phi : \text{SO}(2) \times \mathbb{S}^1 \rightarrow \mathbb{S}^1, (R_\theta, p) \mapsto R_\theta(p) \quad (1.7)$$

This is a group action on $X = \mathbb{S}^1$, since

$$\Phi(R_{\theta_1}, \Phi(R_{\theta_2}, p)) = \Phi(R_{\theta_1+\theta_2}, p).$$

In the short notation, one would write this as

$$R_{\theta_1} \cdot (R_{\theta_2} \cdot p) = R_{\theta_1+\theta_2} \cdot p.$$

Definition 1.3 (Signals¹). Let V be a vector space² and Ω a set, possibly with additional structure. Then the space of V -valued signals on Ω ,

$$\mathcal{X}(\Omega, V) = \{x : \Omega \rightarrow V\} \quad (1.8)$$

is a function space with a vector space structure. (Note that each $x \in \mathcal{X}(\Omega, V)$ is a signal.)

Addition and scalar multiplication of signals are defined as

$$(\alpha x + \beta y)(u) := \alpha x(u) + \beta y(u) \quad \forall u \in \Omega \quad \forall \alpha, \beta \in \mathbb{R}. \quad (1.9)$$

¹[1, p. 11]

²In computer vision, its dimensions would be called *channels*.

Lemma 1.4. If V is endowed with an inner product $\langle v, w \rangle_V$ and a measure μ on a σ -algebra defined on the set Ω (wrt which an integral can be defined), we have the following induced inner product on $\mathcal{X}(\Omega, V)$

$$\langle x, y \rangle_{\mathcal{X}(\Omega, V)} := \int_{\Omega} \underbrace{\langle x(u), y(u) \rangle_V}_{\in V} d\mu(u) \quad (1.10)$$

Proof. Trivial. ■

Remark 1.5. The domain (set) Ω can be discrete, in which case μ can be chosen to be the counting measure, in which case the integral becomes a sum.

Example 1.6. Let $\Omega = \mathbb{Z}_n \times \mathbb{Z}_n$, i.e. a two-dimensional grid, and x an RGB image, i.e. a signal $x : \Omega \rightarrow \mathbb{R}^3$, and f a function (such as a single-layer perceptron) operating on $3n^2$ -dimensional inputs.

Definition 1.7 (Homomorphism). Let $f : A \rightarrow B$ be a map between two sets A and B that are equipped with the same structure. Also assume that \cdot is an operation of the structure. Then f is said to be a homomorphism if $\forall x, y \in A$,

$$f(x \cdot y) = f(x) \cdot f(y). \quad (1.11)$$

Definition 1.8 (Group representation). An n -dimensional real **representation** of a group G is a map $\rho : G \rightarrow \text{GL}(n, \mathbb{R})$ (correspondingly, one defines an n -dimensional complex representation) [1, p. 15], where each $g \in G$ is mapped to an invertible matrix, under the *condition* that the map is a homomorphism, i.e. $\rho(gh) = \rho(g)\rho(h)$.

Example 1.9 (Group representation). Let $G = \mathbb{Z}_n^3$, where the group operation is addition. The group representation

$$\rho : \mathbb{Z}_n \rightarrow \text{GL}(1, \mathbb{C}), g \mapsto \exp\left(\frac{2\pi i g}{n}\right) \quad (1.12)$$

To see that this is indeed a group representation, note that

$$\rho(gh) = \rho(g + h) = \exp\left(\frac{2\pi i gh}{n}\right) = \exp\left(\frac{2\pi i (g + h)}{n}\right) = \exp\left(\frac{2\pi i g}{n}\right) \exp\left(\frac{2\pi i h}{n}\right), \quad (1.13)$$

i.e. the group representation is a homomorphism.

Definition 1.10 (Equivariance). Let ρ be a representation of a group G . A function $f : \mathcal{X}(\Omega, V) \rightarrow \mathcal{X}(\Omega, V)$ is G -equivariant if

$$f(\rho(g)x) = \rho(g)f(x) \quad \forall g \in G. \quad (1.14)$$

Remark 1.11. Note that $\rho(g) \in \text{GL}(n, \mathbb{K})$, $x \in \mathcal{X}(\Omega, V)$, and hence $\rho(g)x$ is a group action mediated by the representation ρ on the group element g , i.e. $\rho(g)x \in \mathcal{X}(\Omega, V)$. Also note that $f(x) \in \mathcal{X}(\Omega, V)$, i.e. $\rho(g)f(x)$ is a group action as well, where the group element g – mediated by the representation ρ – and hence $\rho(g)f(x) \in \mathcal{X}(\Omega, V)$.

The fact that we have a group action implies according to Def. 1.1 and Eq. (1.14),

$$f(\rho(g)(\rho(h)x)) \stackrel{(1.2)}{=} f((\rho(g)\rho(h))x) = (\rho(g)\rho(h))f(x). \quad (1.15)$$

³Cyclic group of order n , $\mathbb{Z}_n = \{0, 1, \dots, n-1\}$.

Definition 1.12 (Ring [2]). Let R be a set equipped with two binary operations, i.e. maps from $R \times R$ to R that are called *addition* and *multiplication* and that are denoted by $+$ and \cdot . Further, let there be two elements that are called *zero* and *unity* and that are denoted by 0 and 1 respectively. Then $(R, +, \cdot)$ is called a *ring* if

1. $(R, +, 0)$ is an abelian group, where 0 is the neutral element for the operation $+$,
2. $(R, \cdot, 1)$ is a monoid, where 1 is the neutral element for the operation \cdot ,
3. the *distributive laws* hold in R , i.e. for all $a, b, c \in R$, we have

$$((a + b) \cdot c = a \cdot c + b \cdot c) \wedge (a \cdot (b + c) = a \cdot b + a \cdot c),$$

4. we have $0 \cdot a = a \cdot 0 = 0$ for all $a \in R$.

Definition 1.13 (Field [2]). Let R be a commutative ring, then we say it is a *field* if $0 \neq 1$ in R and if every non-zero element has a multiplicative inverse.

2 Measure Theory

Definition 2.1 (Generated σ -algebra [3]). Let X be a set and $\mathcal{E} \subset \mathcal{P}(X)$ a non-empty collection of subsets of X . The *smallest* σ -algebra containing all the sets of \mathcal{E} is denoted by $\sigma(\mathcal{E})$.

Corollary 2.2. Let $\mathcal{E}_1, \mathcal{E}_2 \subset \mathcal{P}(X)$ be such that $\mathcal{E}_1 \subset \mathcal{E}_2$. Then $\sigma(\mathcal{E}_1) \subset \sigma(\mathcal{E}_2)$.

Definition 2.3 (Measurable function). Let (X, \mathcal{E}) and (Y, \mathcal{F}) be measurable spaces. A map $f : X \rightarrow Y$ is said to be \mathcal{E} -*measurable* if

$$f^{-1}(\mathcal{F}) := \{f^{-1}(A) | A \in \mathcal{F}\} := \{\{x \in X | f(x) \in A\} | A \in \mathcal{F}\} \subset \mathcal{E}. \quad (2.1)$$

Theorem 2.4 (Generator and measurable function [4]). Let (X, \mathcal{E}) and (Y, \mathcal{F}) be measurable spaces and $\mathcal{F} = \sigma(\mathcal{G})$, i.e. \mathcal{F} is the σ -algebra generated by a family $\mathcal{G} \subset \mathcal{P}(Y)$, where $\mathcal{P}(Y)$ denotes the power set of Y . Then $f : X \rightarrow Y$ is measurable if and only if

$$f^{-1}(G) \in \mathcal{E} \quad \forall G \in \mathcal{G}. \quad (2.2)$$

Proof. “ \Rightarrow ” Since $\mathcal{F} = \sigma(\mathcal{G})$, it obviously holds that $\mathcal{G} \subset \mathcal{F}$ and therefore $f^{-1}(G) \in \mathcal{E} \quad \forall G \in \mathcal{G}$ is ensured by f being measurable.

“ \Leftarrow ” Define the set $\mathcal{M} := \{B \subset Y | f^{-1}(B) \in \mathcal{A}\}$. First we want to convince ourselves that \mathcal{M} is a σ -algebra on Y .

1. $\emptyset \in \mathcal{M}$, since $f^{-1}(\emptyset) = \{x \in X | f(x) \in \emptyset\} = \emptyset$.
2. Let $B \in \mathcal{M}$, then also $Y \setminus B \in \mathcal{M}$, since $f^{-1}(Y \setminus B) = f^{-1}(Y) \setminus f^{-1}(B)$, as can be easily shown by using the definition of the complement of a set. Since $f^{-1}(Y) = X$, it follows that $f^{-1}(Y \setminus B) = X \setminus f^{-1}(B)$. Since by assumption $B \in \mathcal{M}$ (and therefore $f^{-1}(B) \in \mathcal{A}$) and \mathcal{A} itself is a σ -algebra, it follows that $X \setminus f^{-1}(B) \in \mathcal{A}$.
3. Let $B_i \in \mathcal{M}$ for $i \in \mathbb{N}$, then also $\cup_{i \in \mathbb{N}} B_i \in \mathcal{M}$, since

$$f^{-1}\left(\bigcup_{i \in \mathbb{N}} B_i\right) = \bigcup_{i \in \mathbb{N}} f^{-1}(B_i).$$

Since \mathcal{M} is a σ -algebra and since $\mathcal{G} \subset \mathcal{M} \Rightarrow \mathcal{F} = \sigma(\mathcal{G}) \subset \mathcal{M} = \sigma(\mathcal{M})$, it follows that f is measurable. ■

Lemma 2.5 (Push-forward measure). Let (X, \mathcal{E}) and (Y, \mathcal{F}) be measurable spaces. Given a measurable map $f : X \rightarrow Y$ and a measure μ on \mathcal{E} , let $f_{\#}\mu$ be defined by

$$f_{\#}\mu(A) := \mu(f^{-1}(A)) \quad \forall A \in \mathcal{F}. \quad (2.3)$$

$f_{\#}\mu$ is a measure on \mathcal{F} and called the *push-forward* of μ under f .

Proof. Obviously, $f_{\#}\mu(\emptyset) = \mu(f^{-1}(\emptyset)) \stackrel{(2.1)}{=} \mu(\emptyset) = 0$. Also, no matter what kind of set $A \in \mathcal{F}$ we take, since $\mu(A) \geq 0$, the same holds for $f_{\#}\mu(A)$. Finally, let $(A_n)_{n \in \mathbb{N}} \subset \mathcal{F}$ be

a sequence of mutually disjoint sets, then:

$$\begin{aligned}
f_{\#}\mu\left(\bigcup_{n\in\mathbb{N}} A_n\right) &= \mu\left(f^{-1}\left(\bigcup_{n\in\mathbb{N}} A_n\right)\right) \stackrel{(2.1)}{=} \mu\left(\left\{x \in X \mid f(x) \in \bigcup_{n\in\mathbb{N}} A_n\right\}\right) \\
&= \mu\left(\bigcup_{n\in\mathbb{N}} \{x \in X \mid f(x) \in A_n\}\right) = \mu\left(\bigcup_{n\in\mathbb{N}} f^{-1}(A_n)\right) \\
&= \sum_{n\in\mathbb{N}} \mu(f^{-1}(A_n)) = \sum_{n\in\mathbb{N}} f_{\#}\mu(A_n)
\end{aligned}$$

■

Corollary 2.6 (Push-forward of a probability measure). Let (X, \mathcal{E}) and (Y, \mathcal{F}) be measurable spaces. Given a measurable map $f : X \rightarrow Y$ and a probability measure on \mathcal{E} , the push-forward of μ under f , denoted by $f_{\#}\mu$, is also a probability measure.

Proof. Since μ is in particular a measure and thus $f_{\#}\mu$ is also a measure, we only need to show that

$$f_{\#}\mu(Y) = \mu(f^{-1}(Y)) = \mu(\{x \in X \mid f(x) \in Y\}) = \mu(X) = 1. \quad (2.4)$$

■

Definition 2.7 (σ -finite measure). Let (X, \mathcal{A}) be a measurable space and μ a measure on it. If there are sets $A_1, A_2, \dots \in \mathcal{A}$ with $\mu(A_n) < \infty \forall n \in \mathbb{N}$ that satisfy

$$\bigcup_{n\in\mathbb{N}} A_n = X \quad (2.5)$$

then we say μ is σ -finite.

Remark 2.8. Obviously, every finite measure is σ -finite; however, the converse does not necessarily hold [5].

Definition 2.9 (Absolutely continuous measures.). Let μ and ν be two measures on a σ -algebra \mathcal{A} . ν is called *absolutely continuous* w.r.t. μ , written as

$$\nu \ll \mu, \quad (2.6)$$

if for each $A \in \mathcal{A}$, $\mu(A) = 0$ implies $\nu(A) = 0$. If μ and ν are both absolutely continuous w.r.t each other μ and ν are called *equivalent*.

Theorem 2.10 (Radon-Nikodym Theorem [6]). Let μ be a σ -finite measure on a measurable space (S, \mathcal{A}) . Then it is equivalent:

1. $\nu \ll \mu$,
2. $d\nu = h d\mu$ for some measurable function $h : S \rightarrow \mathbb{R}_+$.

The density h then is μ -a.e. finite and μ -a.e. unique.

Lemma 2.11 (Frechét Inception Distance). For two multivariate Gaussian distributions $\mathcal{G}(\mu_x, \Sigma_x)$, $\mathcal{G}(\mu_y, \Sigma_y)$, the *Frechét Inception Distance* (FID) is defined as [https://arxiv.org/pdf/1706.08500.pdf]:

$$d(\mathcal{G}(\mu_x, \Sigma_x), \mathcal{G}(\mu_y, \Sigma_y)) := \sqrt{\|\mu_x - \mu_y\|_2^2 + \text{Tr}(\Sigma_x + \Sigma_y - 2(\Sigma_x \Sigma_y)^{\frac{1}{2}})}. \quad (2.7)$$

It is a metric.

Proof. Clearly, $d(\mathcal{G}(\mu_x, \Sigma_x), \mathcal{G}(\mu_x, \Sigma_x)) = 0$. Also,

$$d(\mathcal{G}(\mu_x, \Sigma_x), \mathcal{G}(\mu_y, \Sigma_y)) = d(\mathcal{G}(\mu_y, \Sigma_y), \mathcal{G}(\mu_x, \Sigma_x)),$$

which holds because $\text{Tr}(AB) = \text{Tr}(BA)$ for any matrices A and B . To see that

$$d(\mathcal{G}(\mu_x, \Sigma_x), \mathcal{G}(\mu_y, \Sigma_y)) \geq 0,$$

note that $\text{Tr}(\Sigma_x + \Sigma_y - 2(\Sigma_x \Sigma_y)^{1/2}) = \text{Tr}\left(\left(\Sigma_x^{1/2} - \Sigma_y^{1/2}\right)^2\right) \geq 0$, since the covariance matrices contain the variances on the diagonal, which are obviously non-negative. It now remains to be shown that also the triangle inequality is fulfilled. For this, note that

$$d(\mathcal{G}(\mu_x, \Sigma_x), \mathcal{G}(\mu_y, \Sigma_y)) = \sqrt{\|\mu_x - \mu_y\|_2^2 + \text{Tr}(\Sigma_x + \Sigma_y - 2(\Sigma_x \Sigma_y)^{\frac{1}{2}})} \quad (2.8)$$

$$= \sqrt{\|\mu_x - \mu_y\|_2^2 + \text{Tr}\left(\left(\Sigma_x^{1/2} - \Sigma_y^{1/2}\right)^2\right)} \quad (2.9)$$

$$= \sqrt{\|\mu_x - \mu_y\|_2^2 + \|\sigma_x - \sigma_y\|_2^2}, \quad (2.10)$$

where σ_x and σ_y denote vectors containing the standard deviations of the two Gaussian distributions. Clearly,

$$d(\mathcal{G}(\mu_x, \Sigma_x), \mathcal{G}(\mu_z, \Sigma_z)) = \sqrt{\|\mu_x - \mu_z\|_2^2 + \|\sigma_x - \sigma_z\|_2^2} \quad (2.11)$$

$$\leq \sqrt{\|\mu_x - \mu_y\|_2^2 + \|\mu_y - \mu_z\|_2^2 + \|\sigma_x - \sigma_y\|_2^2 + \|\sigma_y - \sigma_z\|_2^2} \quad (2.12)$$

$$\leq \sqrt{\|\mu_x - \mu_y\|_2^2 + \|\sigma_x - \sigma_y\|_2^2} + \sqrt{\|\mu_y - \mu_z\|_2^2 + \|\sigma_y - \sigma_z\|_2^2} \quad (2.13)$$

$$= d(\mathcal{G}(\mu_x, \Sigma_x), \mathcal{G}(\mu_y, \Sigma_y)) + d(\mathcal{G}(\mu_y, \Sigma_y), \mathcal{G}(\mu_z, \Sigma_z)), \quad (2.14)$$

since $\sqrt{x+y} \leq \sqrt{x} + \sqrt{y}$, as one can directly show by squaring for non-negative $x, y \in \mathbb{R}$. ■

Definition 2.12 (Convergence in probability). Assume we have a sequence of random variables $(X_n)_{n \in \mathbb{N}}$, defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. We say this sequence converges to another random variable X if

$$\forall \epsilon > 0 : \lim_{n \rightarrow \infty} \mathbb{P}\{|X_n - X| > \epsilon\} = 0. \quad (2.15)$$

3 Analysis

Theorem 3.1. (Bolzano⁴-Weierstrass⁵) Let $(x_n)_{n \in \mathbb{N}}$ be a bounded sequence of real numbers, then $(x_n)_{n \in \mathbb{N}}$ has a convergent subsequence.

Proof [7]. Let $(x_n)_{n \in \mathbb{N}}$ be a bounded sequence of real numbers, i.e. there exists an $L \in \mathbb{R}$ s.t. $|x_n| \leq L$ for all $n \in \mathbb{N}$. We now proceed in several steps to construct a convergent subsequence of $(x_n)_{n \in \mathbb{N}}$.

Step 1: Set $a_0 := -L$ and $b_0 := L$. Clearly, $|b_0 - a_0| = 2L$. We now divide the interval $[a_0, b_0]$ into two halves. At least one half contains infinitely many sequence elements. We now pick that half and denote it by $[a_1, b_1]$. Clearly, either $a_1 = a_0$ or $b_1 = b_0$. Also, $|b_1 - a_1| = |b_0 - a_0|/2 = L$ and $[a_1, b_1] \subset [a_0, b_0]$. Figure 1 visualizes this. Since there are




Figure 1: Construction of subsequence of $(x_n)_{n \in \mathbb{N}}$.

infinitely many sequence elements in $[a_1, b_1]$, we can choose a sequence element contained in $[a_1, b_1]$, which we denote by x_{n_1} .

We can now characterize the search procedure inductively: For $k \in \mathbb{N}$, let $[a_k, b_k]$ be s.t. $[a_k, b_k]$ contains infinitely many sequence elements of $(x_n)_{n \in \mathbb{N}}$, $[a_k, b_k] \subset [a_{k-1}, b_{k-1}]$ and $|b_k - a_k| = 2L/2^k = L/2^{k-1}$. Then we construct $[a_{k+1}, b_{k+1}]$ by dividing $[a_k, b_k]$ into two halves and choosing the half that contains infinitely many sequence elements, i.e. either $a_{k+1} = a_k$ or $b_{k+1} = b_k$. Clearly, $|b_{k+1} - a_{k+1}| = |b_k - a_k|/2 = L/2^k$. Since $[a_{k+1}, b_{k+1}]$ contains infinitely many sequence elements, we can choose an $x_{n_{k+1}} \in [a_{k+1}, b_{k+1}]$ s.t. $x_{n_{k+1}} > x_{n_k}$. In this way, we generate nested intervals of the form

$$\underbrace{[a_0, b_0]}_{\text{length } 2L} \supset \underbrace{[a_1, b_1]}_{\text{length } L} \supset \underbrace{[a_2, b_2]}_{\text{length } L/2} \supset \dots$$

Step 2: The sequence $(a_n)_{n \in \mathbb{N}}$ is monotonically increasing and bounded above by b_0 , so by Corollary 5.54, it converges to an $x \in \mathbb{R}$.

Step 3: We now show that $(x_{n_k})_{k \in \mathbb{N}}$ converges to x . For this, note that

$$|x_{n_k} - x| \leq |x_{n_k} - a_k| + |a_k - x| \leq \frac{L}{2^{k-1}} + |a_k - x| \xrightarrow{k \rightarrow \infty} 0.$$

■

Corollary 3.2 (Bolzano-Weierstrass). Let $(z_n)_{n \in \mathbb{N}}$ be a bounded sequence of complex numbers, then $(z_n)_{n \in \mathbb{N}}$ has a convergent subsequence.

Proof [8]. Since $(z_n)_{n \in \mathbb{N}}$ is bounded, the real sequence $(\operatorname{Re}(z_n))_{n \in \mathbb{N}}$ is also bounded, as $|\operatorname{Re}(z_n)| \leq |z_n|$. Thus, $(\operatorname{Re}(z_n))_{n \in \mathbb{N}}$ has a convergent subsequence, which implies that $(z_n)_{n \in \mathbb{N}}$ has a subsequence for which the real part converges, which we denote by $(z_{n_k})_{k \in \mathbb{N}}$. Since $|\operatorname{Im}(z_{n_k})| \leq |z_{n_k}|$, the sequence $(\operatorname{Im}(z_{n_k}))_{k \in \mathbb{N}}$ is also bounded and thus has a convergent subsequence. By Theorem 5.24, the real part of the subsequence will still converge. Therefore, there is a subsequence of $(z_n)_{n \in \mathbb{N}}$ for which both the real and imaginary parts converge. ■

⁴BERNARD BOLZANO (1781 – 1848), Bohemian mathematician and Catholic priest.

⁵KARL WEIERSTRASS (1815 – 1897), German mathematician, “father of modern analysis”.

Example 3.3. Consider the real sequence $(x_n)_{n \in \mathbb{N}}$ with $x_n := (-1)^n$, which is clearly bounded, but not convergent. However, $(x_{2n})_{n \in \mathbb{N}}$ and $(x_{2n+1})_{n \in \mathbb{N}}$ are convergent subsequences of $(x_n)_{n \in \mathbb{N}}$.

4 Inner Product and Normed Spaces

Definition 4.1 (Inner Product Space). Let \mathbb{K} be a field ($\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$). An inner product space is a vector space V over \mathbb{K} that allows for an **inner product**

$$\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{K} \quad (4.1)$$

satisfying the following properties $\forall \alpha, \beta \in \mathbb{K}; x, y, z \in V$:

1. **Conjugate symmetry**:

$$\langle x, y \rangle = \overline{\langle y, x \rangle}, \quad (4.2)$$

which implies that $\langle x, x \rangle \in \mathbb{R}$, even if $\mathbb{K} = \mathbb{C}$.⁶

For $\mathbb{K} = \mathbb{R}$, conjugate symmetry is exact symmetry.

2. **Linearity** (in the first argument):

$$\langle \alpha x + \beta y, z \rangle = \alpha \langle x, z \rangle + \beta \langle y, z \rangle \quad (4.3)$$

From the conjugate symmetry property, this means that we have semi-linearity in the second argument:

$$\langle x, \alpha y + \beta z \rangle = \overline{\langle \alpha y + \beta z, x \rangle} = \overline{\alpha \langle y, x \rangle + \beta \langle z, x \rangle} = \bar{\alpha} \langle x, y \rangle + \bar{\beta} \langle x, z \rangle. \quad (4.4)$$

3. **Positive-definiteness**:

$$\langle x, x \rangle \geq 0 \quad (4.5)$$

and

$$\langle x, x \rangle = 0 \Leftrightarrow x = 0. \quad (4.6)$$

Definition 4.2 (Normed Space). Let \mathbb{K} be a space ($\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$). Then a mapping $\|\cdot\| : X \rightarrow \mathbb{R}$ is called a **norm** if it satisfies the following properties for all $\varphi, \psi \in X$ and $\alpha \in \mathbb{K}$:

1. **Positivity**:

$$\|\varphi\| \geq 0 \quad (4.7)$$

2. **Definiteness**:

$$\|\varphi\| = 0 \Leftrightarrow \varphi = 0 \quad (4.8)$$

3. **Homogeneity**:

$$\|\alpha \varphi\| = |\alpha| \|\varphi\| \quad (4.9)$$

⁶Set $y = x$.

4. Triangle inequality:

$$\|\varphi + \psi\| \leq \|\varphi\| + \|\psi\| \quad (4.10)$$

A linear space X with a norm $\|\cdot\|$ is called a **normed linear space** or **normed space** for short. For a normed space, we shall use the notation $(X, \|\cdot\|)$.

Example 4.3. Let $X = \mathbb{R}^d$. Then for $1 \leq p < \infty$, the L^p norm of a vector $x \in X$ is defined as:

$$\|x\|_p := \left(\sum_{j=1}^d |x_j|^p \right)^{\frac{1}{p}} \quad (4.11)$$

In the limit $p \rightarrow \infty$, we obtain the so-called *supremum norm*:

$$\|x\|_\infty := \max_{1 \leq j \leq d} |x_j|. \quad (4.12)$$

In the special case of $p = 2$, we recover the Euclidean norm.

Proof. First, we show that Eq. (4.12) is indeed the limit of Eq. (4.11):

$$\|x\|_\infty = \max_{1 \leq j \leq d} |x_j| \leq \sum_{j=1}^d |x_j| \leq d \cdot \|x\|_\infty \quad (4.13)$$

$$\Rightarrow \|x\|_\infty^p = \left(\max_{1 \leq j \leq d} |x_j| \right)^p = \max_{1 \leq j \leq d} |x_j|^p \leq \sum_{j=1}^d |x_j|^p \leq d \cdot \max_{1 \leq j \leq d} |x_j|^p = d \|x\|_\infty^p \quad (4.14)$$

$$\Rightarrow \|x\|_\infty \leq \left(\sum_{j=1}^d |x_j|^p \right)^{\frac{1}{p}} \leq d^{\frac{1}{p}} \|x\|_\infty \quad (4.15)$$

$$\Rightarrow \lim_{p \rightarrow \infty} \|x\|_\infty = \|x\|_\infty \leq \lim_{p \rightarrow \infty} \left\{ \left(\sum_{j=1}^d |x_j|^p \right)^{\frac{1}{p}} \right\} \leq \lim_{p \rightarrow \infty} \left\{ d^{\frac{1}{p}} \|x\|_\infty \right\} = \|x\|_\infty \quad (4.16)$$

$$\Rightarrow \lim_{p \rightarrow \infty} \left\{ \left(\sum_{j=1}^d |x_j|^p \right)^{\frac{1}{p}} \right\} = \|x\|_\infty \quad (4.17)$$

To show the norm property of the L^p norm for $1 \leq p < \infty$, we will explicitly show the fulfilling properties of a norm, cf. Defn. (4.2), for all $x, y, z \in X$ and $\alpha \in \mathbb{R}$:

1. Positivity: $\|x\|_p \geq 0 \ \forall x \in X$,
2. Definiteness: $\|x\|_p = 0 \Leftrightarrow x = 0$,
3. Homogeneity:

$$\|\alpha x\|_p = \left(\sum_{j=1}^d |\alpha x_j|^p \right)^{\frac{1}{p}} = \left(\sum_{j=1}^d |\alpha|^p |x_j|^p \right)^{\frac{1}{p}} = |\alpha| \|x\|_p,$$

4. Triangle inequality:

$$\|x + y\|_p = \left(\sum_{j=1}^d |x_j + y_j|^p \right)^{\frac{1}{p}} \leq \left(\sum_{j=1}^d |x_j|^p + |y_j|^p \right)^{\frac{1}{p}} = \|x\|_p + \|y\|_p.$$

In case of $p = \infty$, we will only show the triangle inequality, since the other properties are trivial to prove:

$$\|x + y\|_\infty = \max_{1 \leq j \leq d} |x_j + y_j| \leq \max_{1 \leq j \leq d} \{|x_j| + |y_j|\} \leq \max_{1 \leq j \leq d} |x_j| + \max_{1 \leq j \leq d} |y_j| = \|x\|_\infty + \|y\|_\infty,$$

where the last inequality holds since for any $1 \leq j \leq d$, it holds that

$$\begin{aligned} & \left(|x_j| \leq \max_{1 \leq k \leq d} |x_k| \right) \wedge \left(|y_j| \leq \max_{1 \leq k \leq d} |y_k| \right) \\ & \Rightarrow |x_j| + |y_j| \leq \max_{1 \leq k \leq d} |x_k| + \max_{1 \leq k \leq d} |y_k| \\ & \Rightarrow \max_{1 \leq j \leq d} \{|x_j| + |y_j|\} \leq \max_{1 \leq k \leq d} |x_k| + \max_{1 \leq k \leq d} |y_k| \end{aligned}$$

■

Theorem 4.4. Let $(X, \|\cdot\|)$ be a normed space. Then the “second triangle inequality” holds:

$$||\varphi| - |\psi|| \leq \|\varphi - \psi\| \quad \forall \varphi, \psi \in X. \quad (4.18)$$

Proof. For $\varphi, \psi \in X$ we have

$$\|\varphi\| = \|\varphi - \psi + \psi\| \leq \|\varphi - \psi\| + \|\psi\| \Leftrightarrow \|\varphi\| - \|\psi\| \leq \|\varphi - \psi\|. \quad (4.19)$$

By exchanging the roles of φ and ψ we obtain

$$\|\psi\| - \|\varphi\| \leq \|\varphi - \psi\| \quad (4.20)$$

and thus

$$||\varphi| - |\psi|| \leq \|\varphi - \psi\|. \quad (4.21)$$

■

Theorem 4.5. Let $(X, \|\cdot\|)$ be a normed space. Then the addition, scalar multiplication and the norm itself are continuous.

Proof. • ad continuity of the addition: Let $(\varphi_n)_{n \in \mathbb{N}}$ and $(\psi_n)_{n \in \mathbb{N}}$ be convergent sequences in X with limit elements $\varphi, \psi \in X$, i.e. $\varphi_n \rightarrow \varphi$ and $\psi_n \rightarrow \psi$ for $n \rightarrow \infty$. Thus

$$0 \leq \|(\varphi_n + \psi_n) - (\varphi + \psi)\| \leq \|\varphi_n - \varphi\| + \|\psi_n - \psi\| \rightarrow 0 \quad \text{for } n \rightarrow \infty \quad (4.22)$$

and hence $\varphi_n + \psi_n \rightarrow \varphi + \psi$ for $n \rightarrow \infty$.

- ad continuity of the scalar multiplication: Let $(\alpha_n)_{n \in \mathbb{N}} \in \mathbb{K}$ converge to $\alpha \in \mathbb{K}$ and $(\varphi_n)_{n \in \mathbb{N}} \in X \rightarrow \varphi \in X$ for $n \rightarrow \infty$. Then

$$0 \leq \|\alpha_n \varphi_n - \alpha \varphi\| = \|\alpha_n (\varphi_n - \varphi) + (\alpha_n - \alpha) \varphi\| \leq \|\alpha_n (\varphi_n - \varphi)\| + \|(\alpha_n - \alpha) \varphi\| \quad (4.23)$$

$$\leq |\alpha_n| \|\varphi_n - \varphi\| + |\alpha_n - \alpha| \|\varphi\| \xrightarrow{n \rightarrow \infty} 0. \quad (4.24)$$

This implies $\alpha_n \varphi_n \rightarrow \alpha \varphi$ for $n \rightarrow \infty$.

- ad continuity of the norm: Let $\varphi_n \rightarrow \varphi$. With Theorem 4.4 we have:

$$0 \leq |\|\varphi_n\| - \|\varphi\|| \leq \|\varphi_n - \varphi\| \xrightarrow{n \rightarrow \infty} 0 \quad (4.25)$$

and hence $\|\varphi_n\| \rightarrow \|\varphi\|$ for $n \rightarrow \infty$. ■

Definition 4.6. Two norms $\|\cdot\|_a$ and $\|\cdot\|_b$ on a linear space X are said to be equivalent if and only if there exist positive constants $0 < c \leq C < \infty$ such that

$$c \|\varphi\|_b \leq \|\varphi\|_a \leq C \|\varphi\|_b \quad \forall \varphi \in X. \quad (4.26)$$

(It is also possible to write this as $\tilde{c} \|\varphi\|_a \leq \|\varphi\|_b \leq \tilde{C} \|\varphi\|_a$ with $\tilde{c} := C^{-1}$ and $\tilde{C} := c^{-1}$, where $0 < \tilde{c} \leq \tilde{C} < \infty$.)

Lemma 4.7. Let X be a linear space and the pairs $(\|\cdot\|_a, \|\cdot\|_c)$ and $(\|\cdot\|_b, \|\cdot\|_c)$ be equivalent. Then also the pair $(\|\cdot\|_a, \|\cdot\|_b)$ is equivalent.

Proof. By assumption, we know that

$$c \|\varphi\|_c \leq \|\varphi\|_a \leq C \|\varphi\|_c \quad \forall \varphi \in X \quad (4.27)$$

and

$$d \|\varphi\|_c \leq \|\varphi\|_b \leq D \|\varphi\|_c \quad \forall \varphi \in X \quad (4.28)$$

$$\Leftrightarrow \|\varphi\|_c \leq \frac{1}{d} \|\varphi\|_b \leq \frac{D}{d} \|\varphi\|_c. \quad (4.29)$$

$$\stackrel{(4.27)}{\Leftrightarrow} \frac{1}{C} \|\varphi\|_a \leq \|\varphi\|_c \leq \frac{1}{d} \|\varphi\|_b \leq \frac{D}{d} \|\varphi\|_c \leq \frac{D}{d \cdot c} \|\varphi\|_a \quad (4.30)$$

$$\Leftrightarrow \frac{1}{C} \|\varphi\|_a \leq \frac{1}{d} \|\varphi\|_b \leq \frac{D}{d \cdot c} \|\varphi\|_a \quad (4.31)$$

$$\Leftrightarrow \frac{d}{C} \|\varphi\|_a \leq \|\varphi\|_b \leq \frac{D}{c} \|\varphi\|_a. \quad (4.32)$$

It is clear that $0 < dC^{-1} \leq Dc^{-1} < \infty$ holds. ■

Theorem 4.8. On a *finite-dimensional* space X over a field \mathbb{K} all norms are equivalent.

Proof [9]. Let $\dim(X) = n$, $\{e_1, \dots, e_n\}$ be a basis of X and $\|\cdot\|$ a norm on X . We can now show that $\|\cdot\|$ is equivalent to the Euclidean norm $\|\sum_{i=1}^n \alpha_i e_i\|_2 = (\sum_{i=1}^n |\alpha_i|^2)^{1/2}$ as follows:

Set $K := \max\{\|e_1\|, \dots, \|e_n\|\} > 0$. Then from the triangle inequality for $\|\cdot\|$ we have:

$$\|x\| = \left\| \sum_{i=1}^n \alpha_i e_i \right\| \leq \sum_{i=1}^n \|\alpha_i e_i\| = \sum_{i=1}^n |\alpha_i| \|e_i\| \quad (4.33)$$

Since $(|\alpha_1|, \dots, |\alpha_n|)^T, (\|e_1\|, \dots, \|e_n\|)^T \in \mathbb{R}^n$ and

$$\left\langle (|\alpha_1|, \dots, |\alpha_n|), (\|e_1\|, \dots, \|e_n\|) \right\rangle = \sum_{i=1}^n |\alpha_i| \|e_i\| \quad (4.34)$$

we can use the Cauchy-Schwarz inequality:

$$\begin{aligned} \left\langle (|\alpha_1|, \dots, |\alpha_n|), (\|e_1\|, \dots, \|e_n\|) \right\rangle &\leq \left\| \sum_{i=1}^n \alpha_i e_i \right\|_2 \cdot \left\| \sum_{i=1}^n e_i \right\|_2 = \sqrt{\sum_{i=1}^n |\alpha_i|^2} \cdot \sqrt{\sum_{i=1}^n \|e_i\|^2} \\ &= \|x\|_2 \cdot \sqrt{\sum_{i=1}^n K^2} = K\sqrt{n} \|x\|_2 \quad \forall x \in X, \end{aligned} \quad (4.35)$$

where in the last line we used that $K = \max\{\|e_1\|, \dots, \|e_n\|\}$ and $x = \sum_{i=1}^n \alpha_i e_i$. Putting the last Eq. into Eq. (4.33), we have:

$$\|x\| \leq \sum_{i=1}^n |\alpha_i| \|e_i\| \leq K\sqrt{n} \|x\|_2 \quad \forall x \in X. \quad (4.37)$$

Now define the set

$$S := \{x \in X \mid \|x\|_2 = 1\}. \quad (4.38)$$

This set is closed since it is the preimage of the closed set $\{1\} \subset \mathbb{R}$ under the continuous function $\|\cdot\|_2$, cf. Theorem 4.5, 5.132. S is also closed since $S \subset B_r(0) = \{\psi \in X \mid \|\psi\|_2 < r\}$ for $r > 0$ (here, we take into account that every norm induces a metric). Thus, S is compact according to Heine-Borel (which applies to every finite-dimensional normed vector space). Since every continuous function takes its minimum on a compact set, we know that $\|\cdot\|$ has a minimum $m > 0$ on S . Since $x \cdot \|x\|_2^{-1} \in S$ for $x \neq 0$, we have (m is the minimum of the function $\|\cdot\|$):

$$m \|x\|_2 \leq \|x\| \quad \forall x \in X. \quad (4.39)$$

All in all, we proved:

$$m \|x\|_2 \leq \|x\| \leq K\sqrt{n} \|x\|_2 \quad \forall x \in X. \quad (4.40)$$

■

Definition 4.9 (Strongly equivalent metrics [10]). Let X be a linear space equipped with two metrics d and d' . Then the metrics are *strongly equivalent* if and only if there exist positive constants $0 < \alpha \leq \beta < \infty$ such that

$$\alpha d(x, y) \leq d'(x, y) \leq \beta d(x, y) \quad \forall x, y \in X. \quad (4.41)$$

Remark 4.10. Obviously, Eq. (4.26) can also be written as

$$c \|\varphi - \psi\|_b \leq \|\varphi - \psi\|_a \leq C \|\varphi - \psi\|_b \quad \forall \varphi, \psi \in X. \quad (4.42)$$

Thus, Theorem 4.8 holds for metric spaces as well if the metric is defined via $d(\varphi, \psi) := \|\varphi - \psi\|$ for all $\varphi, \psi \in X$.

5 Metric Spaces

Definition 5.1 (Metric Space [11]). Let X be a non-empty set. Then the map

$$d : X \times X \rightarrow \mathbb{R}$$

is called a *metric* on X if for all $\varphi, \psi, \chi \in X$ the following properties are given:

- **positivity:** $d(\varphi, \psi) \geq 0$,
- **symmetry:** $d(\varphi, \psi) = d(\psi, \varphi)$,
- **definiteness:** $d(\varphi, \psi) = 0$ iff $\varphi = \psi$,
- **triangle inequality:** $d(\varphi, \psi) \leq d(\varphi, \chi) + d(\chi, \psi)$.

Theorem 5.2. Let X be a linear space and d a metric on X . Then $\|x\| := d(x, 0)$ defines a norm on X if $d(tx, 0) = |t| d(x, 0)$ (“scaling property”) for all $t \in \mathbb{K}$, $x \in X$, and if $d(x + y, 0) = d(x, 0) + d(y, 0)$ for all $x, y \in X$ (“translation invariance”).

Proof. Positivity, definiteness and homogeneity of $\|\cdot\|$ are obvious, for the triangle inequality note that for all $x, y \in X$,

$$\begin{aligned} \|x + y\| &= d(x + y, 0) = d(x, -y) \leq d(x, 0) + d(0, -y) = d(x, 0) + d(-y, 0) \\ &= d(x, 0) + d(y, 0) = \|x\| + \|y\|. \end{aligned}$$

■

Theorem 5.3. Let (X, d) be a metric space, then (X, ρ) with

$$\rho(x, y) := \frac{d(x, y)}{1 + d(x, y)} \tag{5.1}$$

for all $x, y \in X$ is also a metric space.

Proof [12, 13]. Clearly, ρ is positive and symmetric. If $\rho(x, y) = 0$, then this is only possible iff $d(x, y) = 0$, which is equivalent to $x = y$, i.e. ρ is also definite. For the triangle inequality, first note that for all $a, b \in \mathbb{R}_{\geq 0}$ with $a \in [0, b]$, we have

$$\frac{a}{1 + a} \leq \frac{b}{1 + b}. \tag{5.2}$$

If $a = 0 = b$ or $a = 0$ and $b > 0$, then the inequality trivially holds. Thus, assume $a, b \neq 0$. Then

$$\frac{a}{1 + a} = \frac{1}{\frac{1}{a} + 1} \leq \frac{1}{\frac{1}{b} + 1} = \frac{b}{1 + b},$$

since $b \geq a$. Thus, for any $x, y, z \in X$, we have

$$\begin{aligned} \rho(x, y) &= \frac{d(x, y)}{1 + d(x, y)} \stackrel{(5.2)}{\leq} \frac{d(x, z) + d(z, y)}{1 + d(x, z) + d(z, y)} \\ &= \frac{d(x, z)}{1 + d(x, z) + d(z, y)} + \frac{d(z, y)}{1 + d(x, z) + d(z, y)} \\ &\leq \frac{d(x, z)}{1 + d(x, z)} + \frac{d(z, y)}{1 + d(z, y)} \\ &= \rho(x, z) + \rho(z, y). \end{aligned}$$

■

Remark 5.4. The metric ρ from Theorem 5.3 is bounded between 0 and 1, $\rho(x, y) \in [0, 1)$ for all $x, y \in X$.

Theorem 5.5. Let (X, d) be a metric space. Then we have the inequality

$$|d(\varphi, \psi) - d(\varphi', \psi')| \leq d(\varphi, \varphi') + d(\psi, \psi') \quad \forall \varphi, \varphi', \psi, \psi' \in X. \quad (5.3)$$

Proof. Let $\varphi, \varphi', \psi, \psi' \in X$ be arbitrary. Applying the triangle inequality twice, we have

$$d(\varphi, \psi) \leq d(\varphi, \varphi') + d(\varphi', \psi) \leq d(\varphi, \varphi') + d(\psi, \psi') + d(\psi', \varphi') \quad (5.4)$$

$$\Rightarrow d(\varphi, \psi) - d(\varphi', \psi') \leq d(\varphi, \varphi') + d(\psi, \psi') \quad (5.5)$$

Exchanging the roles of φ and φ' and ψ and ψ' , we get the stated inequality. \blacksquare

Remark 5.6 (Discrete Metric). Let X be an arbitrary set and $d(\varphi, \psi) = 1 \quad \forall \varphi, \psi \in X$ and $d(\varphi, \varphi) = 0$. Then d is a metric, called the *discrete metric* on X .

Definition 5.7 (Open Ball). Let (X, d) be a metric space, $\varphi \in X$ and $r > 0$. Then

$$B_r(\varphi) := \{\psi \in X \mid d(\varphi, \psi) < r\} \subset X \quad (5.6)$$

is called an *open ball* in X with middle point φ and radius r .

Example 5.8. For the discrete metric d according to Remark 5.6, the open balls can be characterized as follows:

$$B_r(\varphi) = \begin{cases} \{\varphi\}, & r \leq 1 \\ X, & r > 1 \end{cases} \quad (5.7)$$

Theorem 5.9. Let (X, d) be a metric space, and $B_{\tilde{r}}(\tilde{x})$ and $B_{r'}(x')$ be open balls with $\tilde{x}, x' \in X$ and $\tilde{r}, r' > 0$. Then for all $x \in X$ with $x \in B_{\tilde{r}}(\tilde{x}) \cap B_{r'}(x')$ there is an open ball containing x and that is contained in $B_{\tilde{r}}(\tilde{x}) \cap B_{r'}(x')$.

Proof [14]. The intuition of the proof is shown in Figure 2. If the intersection is empty, then the statement holds (vacuously). Let $x \in B_{\tilde{r}}(\tilde{x}) \cap B_{r'}(x')$, and thus $x \in B_{\tilde{r}}(\tilde{x})$ and $x \in B_{r'}(x')$. Define $r := \min\{\tilde{r} - d(x, \tilde{x}), r' - d(x, x')\}$, and note that $x \in B_r(x)$. For all $y \in B_r(x)$, we have $d(y, \tilde{x}) \leq d(y, x) + d(x, \tilde{x}) < r + d(x, \tilde{x}) \leq \tilde{r}$, i.e. $y \in B_{\tilde{r}}(\tilde{x})$. Similarly, we can show $y \in B_{r'}(x')$, and thus $y \in B_{\tilde{r}}(\tilde{x}) \cap B_{r'}(x')$, i.e. $B_r(x) \subset B_{\tilde{r}}(\tilde{x}) \cap B_{r'}(x')$. \blacksquare

Definition 5.10 (Closed ball). Let (X, d) be a metric space, $\varphi \in X$ and $r > 0$. Then

$$B_r[\varphi] := \{\psi \in X \mid d(\varphi, \psi) \leq r\} \subset X \quad (5.8)$$

is called a *closed ball* in X with middle point φ and radius r .

Definition 5.11 (Open Set). Let (X, d) be a metric space. Then a subset $U \subset X$ is called *open* in X if for every $\varphi \in U$ there is an open ball $B_r(\varphi)$ that is contained in U , i.e. $B_r(\varphi) \subset U$.

Definition 5.12 (Closed Set). Let (X, d) be a metric space. Then a subset $A \subset X$ is called *closed* in X if the complement $X \setminus A$ is open according to Definition 5.11.

Example 5.13. Let $X = \mathbb{R}$ with the standard metric. Then for each $x \in \mathbb{R}$, the singleton $\{x\} \subset \mathbb{R}$ is closed.




Figure 2: Any point in the intersection of two open balls lies in another open ball, which is contained in the intersection.




Figure 3: In a metric space, each open ball contains another open ball of smaller radius.

Proof. Note that $\mathbb{R} \setminus \{x\} = (-\infty, x) \cup (x, \infty)$ is open. ■

Theorem 5.14. Open balls are open.

Proof. An illustration of the idea of the proof is shown in Figure 3.

Let $f \in X$ and for $r > 0$ consider the open ball $U := B_r(f) \subset X$. By definition, for every $\varphi \in U$ it holds that $d(\varphi, f) < r$. Now we show that

$$B_{r-d(\varphi, f)}(\varphi) \subset U. \quad (5.9)$$

For this consider $\psi \in B_{r-d(\varphi,f)}(\varphi)$, i.e. $d(\psi, \varphi) < r - d(\varphi, f)$. With the triangle inequality we obtain:

$$d(f, \psi) \leq d(f, \varphi) + d(\varphi, \psi) < d(f, \varphi) + r - d(\varphi, f) = r, \quad (5.10)$$

i.e. $d(f, \psi) < r$ and thus $\psi \in U = B_r(f)$. ■

Theorem 5.15. Closed balls are closed.

Proof. Let $f \in X$ and for $r > 0$ consider the closed ball $B_r[f] \subset X$. By definition, for every $\varphi \in X \setminus B_r[f] \subset X$, it holds that $d(\varphi, f) > r$. We will now show that

$$B_{d(\varphi,f)-r}(\varphi) \subset X \setminus B_r[f], \quad (5.11)$$

which would prove that $B_r[f]$ is closed. Let $\psi \in B_{d(\varphi,f)-r}(\varphi)$, i.e. $d(\psi, \varphi) < d(\varphi, f) - r$. With the triangle inequality, we obtain:

$$d(f, \varphi) \leq d(f, \psi) + d(\psi, \varphi) \Leftrightarrow d(f, \psi) \geq d(f, \varphi) - d(\psi, \varphi) > d(f, \varphi) - d(\varphi, f) + r = r, \quad (5.12)$$

i.e. $d(f, \psi) > r$, and hence $\psi \notin B_r[f]$, implying that $\psi \in X \setminus B_r[f]$. ■

Theorem 5.16. Let (X, d) be a metric space. The collection of open sets as in Definition 5.11 gives a topology.

Proof. According to Defn. 5.1, we need to prove that X and \emptyset are open, that the union of an arbitrary number of open sets is open, and that the intersection of a finite number of open sets is open.

Clearly, X is open (since for any choice of $r > 0$ and $\varphi \in X$, $B_r(\varphi) \subset X$). \emptyset is also open, since nothing needs to be shown.

Now let $U_1, \dots, U_n \subset X$ be open, and consider the intersection

$$U := \bigcap_{i=1}^n U_i \subset X. \quad (5.13)$$

For any $\varphi \in U$, we know that $\varphi \in U_i$ for all $1 \leq i \leq n$. Since the U_i are open, there is a radius $r_i > 0$ such that $B_{r_i}(\varphi) \subset U_i$ for all $1 \leq i \leq n$. Defining $r := \min_{1 \leq i \leq n} \{r_i\} > 0$, we have that $B_r(\varphi) \subset U_i$ for all $1 \leq i \leq n$ and hence $B_r(\varphi) \subset U$.

Finally, let $U_i, i \in I$, be open, where I is an index set. Consider the union

$$U := \bigcup_{i \in I} U_i \subset X. \quad (5.14)$$

For any $\varphi \in U$ there is an index $i \in I$ such that $\varphi \in U_i$. Since U_i is open, there is an $r > 0$ such that $B_r(\varphi) \subset U_i \subset U$. ■

Definition 5.17 (Convergence of sequence [11]). Let (X, d) be a metric space. A sequence $(\varphi_n)_{n \in \mathbb{N}}$ with elements $\varphi_n \in X, n \in \mathbb{N}$, is called *convergent* if there is a $\varphi \in X$ s.t.

$$\lim_{n \rightarrow \infty} d(\varphi_n, \varphi) = 0, \quad (5.15)$$

i.e. for all $\epsilon > 0$ there exists an $N(\epsilon) \in \mathbb{N}$ s.t.

$$d(\varphi_n, \varphi) < \epsilon \quad \forall n \geq N(\epsilon). \quad (5.16)$$

φ is called the *limit* of the sequence $(\varphi_n)_{n \in \mathbb{N}}$. For this, we write

$$\lim_{n \rightarrow \infty} \varphi_n = \varphi \quad \text{or} \quad \varphi_n \xrightarrow{n \rightarrow \infty} \varphi. \quad (5.17)$$

Non-convergent sequence are called *divergent*.

Theorem 5.18. Let (X, d) be a metric space, and $(\varphi_n)_{n \in \mathbb{N}}$ a convergent sequence in X wrt d . Then the limit is unique.

Proof. Assume the limit is not unique, i.e. $(\varphi_n)_{n \in \mathbb{N}}$ converges to both $\varphi \in X$ and $\psi \in X$. with $\varphi \neq \psi$. Then for $\epsilon := d(\varphi, \psi)/2$ there exists an $N' \in \mathbb{N}$ s.t. for all $n \geq N'$, we have

$$d(\varphi_n, \varphi) < \frac{d(\varphi, \psi)}{2} = \epsilon$$

and an \tilde{N} s.t. for all $n \geq \tilde{N}$, we have

$$d(\varphi_n, \psi) < \frac{d(\varphi, \psi)}{2} = \epsilon.$$

Define $N := \max \{N', \tilde{N}\}$, then for all $n \geq N$, we have

$$d(\varphi, \psi) \leq d(\varphi, \varphi_n) + d(\varphi_n, \psi) < d(\varphi, \psi),$$

which is a contradiction. ■

Definition 5.17 defines what the convergence of a sequence $(x_n)_{n \in \mathbb{N}}$ in a metric space means. For functions, it turns out that there are different plausible notions of convergence.

Definition 5.19 (Pointwise convergence [15]). Let E be a set, (Y, d) a metric space and $(f_n)_{n \in \mathbb{N}}$ a sequence of functions mapping from E to Y . We say that $(f_n)_{n \in \mathbb{N}}$ converges *pointwisely* to an $f : E \rightarrow Y$ if for all $\epsilon > 0$ and all $x \in E$ there exists an $N = N(\epsilon, x)$ s.t.

$$d(f_n(x), f(x)) < \epsilon \quad \forall n \geq N.$$

Example 5.20. For $n \in \mathbb{N}$, let

$$f_n : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto \begin{cases} -1 & x \leq -\frac{1}{2^n}, \\ 2^n x & -\frac{1}{2^n} < x < \frac{1}{2^n}, \\ 1 & x \geq \frac{1}{2^n}, \end{cases}$$

where we consider \mathbb{R} with the standard metric [16]. The sequence $(f_n)_{n \in \mathbb{N}}$ converges pointwisely to the function

$$f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto \begin{cases} -1 & x < 0, \\ 0 & x = 0, \\ 1 & x > 0. \end{cases}$$

Proof. For $x < 0$, there exists an N s.t. $x \leq -1/2^n$ for all $n \geq N$, as $(-1/2^n)_{n \in \mathbb{N}}$ converges to 0. Then, $f_n(x) = f(x)$. Similarly, for $x > 0$, there exists an $N \in \mathbb{N}$ s.t. $x \geq 1/2^n$ for all $n \geq N$, as $(1/2^n)_{n \in \mathbb{N}}$ converges to 0. Then, $f_n(x) = f(x)$. Also, for $x = 0$, $f_n(x) = 0 = f(x)$ for all $n \in \mathbb{N}$. ■

Remark 5.21. Note that in the preceding example, every f_n is continuous, yet the pointwise limit f is not. This is one drawback of the notion of pointwise convergence. To mitigate this, we need a stronger notion of convergence.

Definition 5.22 (Uniform convergence [15]). Let E be a set, (Y, d) a metric space and $(f_n)_{n \in \mathbb{N}}$ a sequence of functions mapping from E to Y . We say that $(f_n)_{n \in \mathbb{N}}$ converges *uniformly* to an $f : E \rightarrow Y$ if for all $\epsilon > 0$ there exists an $N = N(\epsilon)$ such that for all $x \in E$,

$$d(f_n(x), f(x)) < \epsilon \quad \forall n \geq N.$$

Remark 5.23. Clearly, uniform convergence implies pointwise convergence.

Theorem 5.24. A sequence is convergent iff every subsequence converges to the same limit.

Proof [17]. “ \Leftarrow ” Every sequence has a trivial subsequence, namely itself.

“ \Rightarrow ” Let $(\varphi_{n_k})_{k \in \mathbb{N}}$ be a subsequence of $(\varphi_n)_{n \in \mathbb{N}}$. Note that $n_k \geq k$ for all $k \in \mathbb{N}$, which we will prove by induction: Clearly, $n_1 \geq 1$, and $n_k \geq k$ implies $n_{k+1} > n_k \geq k$, i.e. $n_{k+1} \geq k+1$.

We denote the limit of $(\varphi_n)_{n \in \mathbb{N}}$ by φ , and note that for all $\epsilon > 0$, there exists an $N \in \mathbb{N}$ s.t. for all $n \geq N$, we have $d(\varphi_n, \varphi) < \epsilon$. Thus, for all $k \geq N$ (and hence $n_k \geq N$), we have $d(\varphi_{n_k}, \varphi) < \epsilon$. ■

Definition 5.25 (Accumulation point). Let $U \subset X$. Then $\varphi \in X$ is called an *accumulation point* of U if there is a sequence $(\varphi_n)_{n \in \mathbb{N}}$ in U such that $\lim_{n \rightarrow \infty} \varphi_n = \varphi$.

Theorem 5.26. A subset $U \subset X$ is closed if and only if it contains all its accumulation points according to Defn. 5.25.

Proof. “ \Leftarrow ”: Let U contain all its accumulation points, and let $\varphi \in X \setminus U$. We will now show that there is an $n \in \mathbb{N}$ such that $B_{\frac{1}{n}}(\varphi) \subset X \setminus U$, proving that $X \setminus U$ is open. Assuming there is no such $n \in \mathbb{N}$, then $B_{\frac{1}{n}}(\varphi) \cap U \neq \emptyset$ for all $n \in \mathbb{N}$. This means that there is a sequence $(\varphi_n)_{n \in \mathbb{N}}$ in U such that $d(\varphi, \varphi_n) < \frac{1}{n}$ for all $n \in \mathbb{N}$. Hence, φ is an accumulation point of U , and since U contains all its accumulation points by assumption, it follows that $\varphi \in U$, contradicting the assumption that $\varphi \in X \setminus U$.

“ \Rightarrow ”: Let U be closed, i.e. $X \setminus U$ is open. Further, let $\varphi \in X$ be an accumulation point of U , i.e. there is a sequence $(\varphi_n)_{n \in \mathbb{N}}$ in U such that $\lim_{n \rightarrow \infty} \varphi_n = \varphi$. Hence, by definition of convergence, for all $\epsilon > 0$ there exists an $n \in \mathbb{N}$ such that $\varphi_n \in B(\varphi, \epsilon) \cap U$, implying that $\varphi \in U$. Thus, U contains all its accumulation points. ■

Corollary 5.27. Let $U \subset \mathbb{R}$ be closed and bounded, where on \mathbb{R} we consider the standard metric, then $\sup(U), \inf(U) \in U$, i.e. $\sup(U) = \max(U)$ and $\inf(U) = \min(U)$.

Proof [18]. Since U is bounded, $\inf(U)$ and $\sup(U)$ exist.

By Theorem 8.14, there exists a sequence $(x_n)_{n \in \mathbb{N}}$ in U s.t. $\sup(U) - \frac{1}{n} < x_n \leq \sup(U)$ for all $n \in \mathbb{N}$. Thus, $\lim_{n \rightarrow \infty} x_n = \sup(U)$, i.e. $\sup(U)$ is an accumulation point of U and since U is closed, $\sup(U) \in U$.

Similarly, one can show that $\inf(U) \in U$. ■

Definition 5.28 (Closure of set). The set

$$\bar{U} := \{\varphi \in X \mid \varphi \text{ is an accumulation point of } U\} \subset X \quad (5.18)$$

is called the closure of the set $U \subset X$.

Remark 5.29. Clearly, $U \subset \bar{U}$, since any $\varphi \in U$ is an accumulation point of U (consider the constant sequence, where for any $n \in \mathbb{N}$, $\varphi_n := \varphi$).

Remark 5.30. According to Theorem 5.26, a subset $U \subset X$ is closed if and only if it contains all its limit points, thus for a closed subset we have $\bar{U} \subset U$. With Remark 5.29, we get U is closed if and only if $U = \bar{U}$.

Theorem 5.31. Let $U \subset X$. Then its closure, \bar{U} , is closed.

Proof. According to Theorem 5.26, we need to prove that \bar{U} contains all its limit points. Let $\varphi \in X$ be an accumulation point of \bar{U} . Then for all $n \in \mathbb{N}$ there is a $\varphi_n \in \bar{U}$ such that $d(\varphi, \varphi_n) < 1/n$. By Def. 5.28, we also know that since $\varphi_n \in \bar{U}$, it is an accumulation point of U , i.e. for every $\varphi_n \in \bar{U}$ there is a $\psi_n \in U$ such that $d(\varphi_n, \psi_n) < 1/n$. Thus:

$$d(\psi_n, \varphi) \leq d(\psi_n, \varphi_n) + d(\varphi_n, \varphi) < \frac{1}{n} + \frac{1}{n} \xrightarrow{n \rightarrow \infty} 0, \quad (5.19)$$

i.e. $(\psi_n)_{n \in \mathbb{N}}$ is a sequence in U that converges to φ , i.e. φ is an accumulation point of U , hence $\varphi \in \bar{U}$. ■

Theorem 5.32. Let (X, d) be a metric space. Then $\varphi \in \bar{U}$ for $U \subset X$ if and only if for every open set $O \subset X$ containing φ , $U \cap O \neq \emptyset$.

Proof. “ \implies ” Since $\varphi \in \bar{U}$, there exists a sequence $(\varphi_n)_{n \in \mathbb{N}}$ converging to φ wrt d , i.e. for all $\epsilon > 0$ there exists an $N \in \mathbb{N}$ s.t. for all $n \geq N$, $d(\varphi_n, \varphi) < \epsilon$. Let $O \subset X$ be open, and let $\varphi \in O$. By definition, there exists an open ball $B_r(\varphi) \subset O$ with $r > 0$. Choose $\epsilon = r$ and $\psi = x_N \in U$, then $\psi \in B_r(\varphi) \subset O$.

“ \impliedby ” By assumption, for any non-empty open set $O \subset X$, $U \cap O \neq \emptyset$, where $U \subset X$. Let $\varphi \in X$. Since open balls are open, for any $\epsilon > 0$, $U \cap B_\epsilon(\varphi) \neq \emptyset$. Consider the sequence of open balls $(B_{1/n}(\varphi))_{n \in \mathbb{N}}$, for which $U \cap B_{\frac{1}{n}}(\varphi) \neq \emptyset$ holds. Thus for each $n \in \mathbb{N}$, there exists a point $\varphi_n \in X$ s.t. $\varphi_n \in B_{\frac{1}{n}}(\varphi)$ and $\varphi_n \in U$. Hence, by definition, $\varphi \in X$ is an accumulation point of U . ■

Theorem 5.33. Let $U \subset X$. Then its closure, \bar{U} , is the *smallest* closed set in X containing U , i.e.

$$\bar{U} = \bigcap \{G \mid G \text{ is closed in } X \text{ and } U \subset G\}. \quad (5.20)$$

Proof. From Remark 5.29, we know that $U \subset \bar{U}$, and from Theorem 5.31, we know that \bar{U} is closed. To prove that \bar{U} is the smallest *closed* set in X containing U , let $V \subset X$ be another closed set that contains U , i.e. $U \subset V$. Every sequence in U is also a sequence in V , i.e. every accumulation point of U is in V , i.e. $\bar{U} \subset V$. ■

Corollary 5.34. Let X be a metric space and $A, B \subset X$. If $A \subset B$, then $\bar{A} \subset \bar{B}$.

Proof [19]. \bar{B} is the smallest closed set containing B , and since $A \subset B$, this implies $A \subset \bar{B}$. Since \bar{A} is the smallest closed set containing A , we have $\bar{A} \subset \bar{B}$. ■

Definition 5.35 (Interior [20]). Let X be a metric space and $U \subset X$. Then

$$\text{int}(U) := \bigcup \{G \mid G \text{ is open in } X \text{ and } G \subset U\} \quad (5.21)$$

is the largest open set contained in U ; it is called the *interior* of U in X . A point $x \in \text{int}(U)$ is called an *interior point* of U .

Example 5.36. Equip \mathbb{R} with the standard metric, then $\text{int}(\mathbb{Q}) = \text{int}(\mathbb{R} \setminus \mathbb{Q}) = \emptyset$.

Example 5.37. Equip $[a, b]$ with the subspace metric, then $\text{int}([a, b]) = (a, b)$.

Example 5.38. Equip \mathbb{R}^2 with the Euclidean metric. Then $\text{int}(\mathbb{S}^1) = \emptyset$ and $\text{int}(\mathbb{D}^2) = B_1(0)$, where $\mathbb{D}^2 := \{x \in \mathbb{R}^2 \mid \|x\| \leq 1\} \subset \mathbb{R}^2$ is the unit disc.

Remark 5.39. Clearly, $U \subset X$ is open iff $\text{int}(U) = U$. Also, $x \in X$ is an interior point of U , i.e. $x \in U$, iff U is a neighborhood of x .

Theorem 5.40. Let X be a metric space and $U \subset X$, then

$$\text{int}(U) = X \setminus \overline{X \setminus U}. \quad (5.22)$$

Proof [21]. According to Definition 5.35, $\text{int}(U)$ is the largest open set contained in U . Thus, we need to show that a) $M := X \setminus \overline{X \setminus U}$ is open in X , b) $M \subset U$ and c) $O \subset M$ for any other open set $O \subset U$ in X .

- a) Since $\overline{X \setminus U}$ is closed in X , $M = X \setminus \overline{X \setminus U}$ is open in X .
- b) Since $X \setminus U \subset \overline{X \setminus U}$, this implies $M = X \setminus \overline{X \setminus U} \subset X \setminus (X \setminus U) = U$.
- c) Let $O \subset U$ be open, then $X \setminus O$ is closed and $X \setminus U \subset X \setminus O$. Corollary 5.34 now implies that $\overline{X \setminus U} \subset \overline{X \setminus O} = X \setminus O$. Thus, $X \setminus (X \setminus O) = O \subset X \setminus \overline{X \setminus U} = M$.

■

Theorem 5.41. Let X be a metric space and $U \subset X$. Then

$$X \setminus \text{int}(U) = \overline{X \setminus U}. \quad (5.23)$$

Proof [22]. We use the definitions and make use of de Morgan's law.

$$\begin{aligned} X \setminus \text{int}(U) &\stackrel{(5.22)}{=} X \setminus \bigcup \{G \mid G \text{ is open in } X \text{ and } G \subset U\} \\ &\stackrel{(8.13)}{=} \bigcap \{X \setminus G \mid X \setminus G \text{ is closed in } X \text{ and } X \setminus U \subset X \setminus G\} \\ &\stackrel{(5.20)}{=} \overline{X \setminus U}. \end{aligned}$$

■

Definition 5.42 (Bounded set). A subset $U \subset X$ is called *bounded* in X if it is contained in a closed ball in X , i.e. there exists $r > 0$ and $\varphi \in X$ s.t. $U \subset B_r[\varphi]$.

Theorem 5.43. Convergent sequences in X are fully contained in a bounded subset.

Proof. Let $(\varphi_n)_{n \in \mathbb{N}}$ be a convergent sequence in X with limit point $\varphi \in X$. Then there exists an $N \in \mathbb{N}$ s.t. $d(\varphi_n, \varphi) < \epsilon$ for all $n \geq N$ for a fixed $\epsilon > 0$. Defining

$$r := \max \left\{ \epsilon, \max_{n \leq N} \{d(\varphi, \varphi_n)\} \right\}, \quad (5.24)$$

we have $d(\varphi, \varphi_n) \leq r$ for all $n \in \mathbb{N}$. Hence, for any $n \in \mathbb{N}$, $\varphi_n \in B_r[\varphi]$. ■

Definition 5.44 (Diameter [20]). Let (X, d) be a metric space and $A \subset X$, then the *diameter* of A is defined as

$$\text{diam}(A) := \sup_{a, a' \in A} \{d(a, a')\}$$

Definition 5.45 (Isometry). Let (X, d) and (X', d') be metric spaces. Then $f : X \rightarrow Y$ is called a distance-preserving map (or *isometry*) if

$$d'(f(\varphi), f(\psi)) = d(\varphi, \psi) \quad \forall \varphi, \psi \in X \quad (5.25)$$

(X, d) and (X', d') are called *isometric* if there is a surjective isometry $f : X \rightarrow X'$.

Theorem 5.46. Let (X, d) and (X', d') be metric spaces and $f : X \rightarrow X'$ be an isometry. Then it is injective.

Proof. If $f(\varphi) = f(\psi)$ for any $\varphi, \psi \in X$, then

$$d'(f(\varphi), f(\psi)) = d'(f(\varphi), f(\varphi)) = d(\varphi, \psi) = 0, \quad (5.26)$$

i.e. $\varphi = \psi$. ■

Remark 5.47. Two isometric spaces (X, d) and (X', d') do not differ in their properties wrt their metrics, but instead in the properties of their individual elements. Any metric property holding in one metric space also holds in the other. Hence, in this sense, we say that (X, d) and (X', d') are *identical*.

Remark 5.48. Surjective isometries are hence bijective.

5.1 Complete metric spaces

Definition 5.49. A sequence $(\varphi_n)_{n \in \mathbb{N}}$ in a metric space (X, d) is a *Cauchy sequence* if

$$\lim_{n, m \rightarrow \infty} d(\varphi_n, \varphi_m) = 0, \quad (5.27)$$

i.e. for all $\epsilon > 0$ there exists an $N(\epsilon) \in \mathbb{N}$ s.t.

$$d(\varphi_n, \varphi_m) < \epsilon \quad \forall n, m \geq N. \quad (5.28)$$

Theorem 5.50. Every convergent sequence in X is a Cauchy sequence in X .

Proof. Let $(\varphi_n)_{n \in \mathbb{N}}$ be a convergent sequence in X with limit point $\varphi \in X$. Hence, for all $\epsilon > 0$, there is an $N(\epsilon) \in \mathbb{N}$ with $d(\varphi_n, \varphi) < \epsilon$ for all $n \geq N(\epsilon)$. Hence

$$d(\varphi_n, \varphi_m) \leq d(\varphi_n, \varphi) + d(\varphi, \varphi_m) < 2\epsilon \quad \forall n, m \geq N. \quad (5.29)$$

■

Theorem 5.51. A Cauchy sequence in a metric space (X, d) is convergent iff it contains a convergent subsequence.

Proof [23]. “ \implies ” Every sequence has a trivial subsequence, namely itself.

“ \impliedby ” Let $(\varphi_{n_k})_{k \in \mathbb{N}}$ be a subsequence of the Cauchy sequence $(\varphi_n)_{n \in \mathbb{N}}$, and assume that $\lim_{k \rightarrow \infty} \varphi_{n_k} = \varphi$ wrt d , where $\varphi \in X$, i.e. for all $\epsilon > 0$ there exists an $N' \in \mathbb{N}$ s.t. for all $k \geq N'$, $d(\varphi_{n_k}, \varphi) < \epsilon$ (note that $n_k \geq k$). Since $(\varphi_n)_{n \in \mathbb{N}}$ is Cauchy, there exists an \tilde{N} s.t. for all $m, n \geq \tilde{N}$, we have $d(\varphi_n, \varphi_m) < \epsilon$. Define $N := \max \{N', \tilde{N}\}$, then for all $k, n \geq N$,

$$d(\varphi_n, \varphi) \leq d(\varphi_n, \varphi_{n_k}) + d(\varphi_{n_k}, \varphi) < 2\epsilon,$$

and thus $(\varphi_n)_{n \in \mathbb{N}}$ converges to φ . ■

Remark 5.52. The converse of Theorem 5.50 is in general not true: Take $X = \mathbb{Q}$ and let $d(x, y) := |x - y|$, for any $x, y \in \mathbb{Q}$. Then we have a metric space (\mathbb{Q}, d) , but the sequence $(a_n)_{n \in \mathbb{N}}$ with

$$a_n := \left(1 + \frac{1}{n}\right)^n, \quad (5.30)$$

which is a Cauchy sequence and where all elements are in \mathbb{Q} , does not converge in \mathbb{Q} (the limit point is Euler’s number e). To prove this, we will make use of the following two Theorems.

Theorem 5.53. A real sequence $(a_n)_{n \in \mathbb{N}}$ that is monotonically increasing and upper bounded is a Cauchy sequence in (\mathbb{R}, d) with the standard metric.

Proof. ([24]) The completeness axiom states that any non-empty subset of \mathbb{R} that has an upper bound has a least upper bound in \mathbb{R} . Let M be the least upper bound for the sequence $(a_n)_{n \in \mathbb{N}}$, i.e. $a_n \leq M$ for all $n \in \mathbb{N}$. Since M is the least upper bound, for all $\epsilon > 0$ there is an $N \in \mathbb{N}$ s.t. $M - \epsilon < a_N \leq M$.

By assumption, the sequence is monotonically increasing, i.e. $a_m \geq a_n$ for any $m > n$. Without loss of generality, let $m > n$, then we have

$$M - \epsilon < a_N \leq a_n \leq a_m \leq M, \quad \forall (m, n \geq N) \wedge (m > n) \quad (5.31)$$

from which $a_m - a_n < \epsilon$ follows. Similarly, we can show that $a_n - a_m < \epsilon$ for $n > m$. Hence, $d(a_m, a_n) < \epsilon$ for any $m, n \geq N$. ■

Corollary 5.54. A real sequence $(a_n)_{n \in \mathbb{N}}$ that is monotonically increasing and upper bounded converges in (\mathbb{R}, d) wrt the standard metric.

Proof. This follows from the fact that (\mathbb{R}, d) is complete, cf. Proposition B.21, i.e. every Cauchy sequence in \mathbb{R} converges. ■

Theorem 5.55. Let $x_1, \dots, x_n \in \mathbb{R}_{\geq 0}$, $n \geq 1$, then the arithmetic mean is bigger than or equal to the geometric mean, i.e.

$$\frac{\sum_{i=1}^n x_i}{n} \geq \left(\prod_{i=1}^n x_i \right)^{1/n} \quad (5.32)$$

Proof. ([25]) Via induction. Let $n = 1$, then $x_1 \geq x_1$, which is true.

Assuming that Eq. (5.32) holds for any $n \in \mathbb{N}$, we will make the transition $n \mapsto n + 1$ and denote the arithmetic mean of x_1, \dots, x_{n+1} by α :

$$(n + 1)\alpha = \sum_{i=1}^{n+1} x_i \Leftrightarrow n\alpha = \sum_{i=1}^{n+1} x_i - \alpha \quad (5.33)$$

If $x_i = \alpha$ for all $1 \leq i \leq n + 1$, then we are done, since Eq. (5.32) would be an equality. If all the x_i are not equal to α , then WLOG, let there be exactly two x_i that are not equal to α . Place these two elements at the end, and noting that one needs to be bigger than α and the other smaller than α , WLOG we have

$$(x_n - \alpha > 0) \wedge (\alpha - x_{n+1} > 0) \Rightarrow (x_n - \alpha)(\alpha - x_{n+1}) > 0. \quad (5.34)$$

Now define

$$y := x_n + x_{n+1} - \alpha \geq x_n - \alpha \stackrel{(5.34)}{>} 0. \quad (5.35)$$

and substitute it into Eq. (5.33):

$$n\alpha = \sum_{i=1}^{n+1} x_i - \alpha = \sum_{i=1}^{n-1} x_i + x_n + x_{n+1} - \alpha = \sum_{i=1}^{n-1} x_i + y \quad (5.36)$$

Thus, α is also the arithmetic mean of the n numbers x_1, \dots, x_{n-1}, y . Making use of the induction hypothesis, we get

$$\alpha^{n+1} = \alpha\alpha^n \geq \alpha y \prod_{i=1}^{n-1} x_i. \quad (5.37)$$

From Eq.(5.34) we know that

$$y\alpha - x_n x_{n+1} = (x_n + x_{n+1} - \alpha)\alpha - x_n x_{n+1} = (x_n - \alpha)(\alpha - x_{n+1}) > 0 \Leftrightarrow y\alpha > x_n x_{n+1} \quad (5.38)$$

Substituting Eq. (5.38) into (5.37), we get

$$\alpha^{n+1} \geq \alpha y \prod_{i=1}^{n-1} x_i > x_n x_{n+1} \prod_{i=1}^{n-1} x_i = \prod_{i=1}^{n+1} x_i \Leftrightarrow \alpha \geq \left(\prod_{i=1}^{n+1} x_i \right)^{\frac{1}{n+1}}, \quad (5.39)$$

which proves the induction hypothesis. ■

Proof. (of Remark 5.52) Looking at the proof of Theorem 5.53, we will see that the statement of Theorem 5.53 also holds for sequences in \mathbb{Q} that are upper bounded and monotonically increasing, since any non-empty subset of $\mathbb{Q} \subset \mathbb{R}$ that is upper bounded has a least upper bound in \mathbb{R} (though not necessarily in \mathbb{Q}).

Hence, to show that the sequence $(a_n)_{n \in \mathbb{N}}$ with $a_n = (1 + n^{-1})^n$ is a Cauchy sequence in \mathbb{Q} , we only need to show that it is monotonically increasing and derive an upper bound.

For the monotonicity, we will make use of Theorem 5.55 ([26]). For this, define the x_i as follows:

$$x_i := \begin{cases} 1, & i = 1 \\ 1 + \frac{1}{n}, & 2 \leq i \leq n + 1 \end{cases} \quad (5.40)$$

Then we have

$$\frac{\sum_{i=1}^{n+1} x_i}{n+1} \geq \left(\prod_{i=1}^{n+1} x_i \right)^{\frac{1}{n+1}} \Leftrightarrow \left(\frac{\sum_{i=1}^{n+1} x_i}{n+1} \right)^{n+1} \geq \prod_{i=1}^{n+1} x_i \quad (5.41)$$

$$\Rightarrow \left(\frac{1 + n \left(1 + \frac{1}{n} \right)}{n+1} \right)^{n+1} = \left(1 + \frac{1}{n+1} \right)^{n+1} \geq \left(1 + \frac{1}{n} \right)^n \quad \forall n \in \mathbb{N}, \quad (5.42)$$

i.e. $a_{n+1} > a_n$.

To prove that the sequence $(a_n)_{n \in \mathbb{N}}$ is upper bounded ([27]), note that

$$a_n = \left(1 + \frac{1}{n} \right)^n = \sum_{k=0}^n \binom{n}{k} \frac{1}{n^k} = \sum_{k=0}^n \frac{n!}{k! (n-k)!} \frac{1}{n^k} = 1 + \sum_{k=1}^n \frac{n!}{k! (n-k)!} \frac{1}{n^k} \quad (5.43)$$

$$= 1 + \sum_{k=1}^n \frac{n(n-1) \dots (n-k+1)}{k!} \frac{1}{n^k} \leq 1 + \sum_{k=1}^n \frac{1}{k!} \leq 1 + \sum_{k=1}^n \frac{1}{2^{k-1}} \quad (5.44)$$

$$\leq 1 + \sum_{k=0}^{\infty} \frac{1}{2^k} = 3. \quad (5.45)$$

■

Example 5.56. We can instruct infinitely many counter-examples to Theorem 5.50 in \mathbb{Q} . Let

$$(a_n)_{n \in \mathbb{N}} := \lfloor 10^n \cdot a \rfloor \cdot 10^{-n}, \quad (5.46)$$

where $a \in \mathbb{R} \setminus \mathbb{Q}$. $(a_n)_{n \in \mathbb{N}}$ is a Cauchy sequence in \mathbb{Q} , which gives a to its n^{th} decimal point, converging to a . To see that it is a Cauchy sequence in \mathbb{Q} , note that all sequence elements are rational, that the sequence is bounded by a and that it is monotonically increasing. Thus, by the proof of Theorem 5.53, which also holds for \mathbb{Q} , $(a_n)_{n \in \mathbb{N}}$ is Cauchy in \mathbb{Q} .

Theorem 5.57. Any sequence $(a_n)_{n \in \mathbb{N}}$ with $a_n \in \mathbb{R}$ for any $n \in \mathbb{N}$ that is monotonically increasing and upper bounded for any $n \in \mathbb{N}$ converges in (\mathbb{R}, d) with $d(x, y) := |x - y|$ for any $x, y \in \mathbb{R}$.

Proof. The proof is identical to that of Theorem 5.53. With the same notation as there, we have, cf. Eq. (5.31):

$$M - \epsilon < a_n \leq M, \quad \forall n \geq N \quad (5.47)$$

$$\Rightarrow |a_n - M| < \epsilon, \quad (5.48)$$

i.e. the limit point of $(a_n)_{n \in \mathbb{N}}$ is M (the least upper bound). ■

Remark 5.58. Note that Theorem 5.57 does not hold in \mathbb{Q} , since the completeness axiom only guarantess the existence of a least upper bound in \mathbb{R} .

Theorem 5.59. Let (X, d) be a metric space. Then if $(\varphi_n)_{n \in \mathbb{N}}$ is a Cauchy sequence, it is bounded.

Proof. Fix $\epsilon > 0$ and $\psi \in X$. Since $(\varphi_n)_{n \in \mathbb{N}}$ is a Cauchy sequence, there exists an $N \in \mathbb{N}$ s.t. $d(\varphi_m, \varphi_n) < \epsilon$ for all $m, n \geq N$. Define

$$M := \max_{1 \leq i \leq N} \{d(\psi, \varphi_n)\}, \quad (5.49)$$

and note that the maximum exists since we take it over a set with finitely many elements. For $n \leq N$, we obviously have $d(\psi, \varphi_n) \leq M$. For $n > N$, we have

$$d(\psi, \varphi_n) \leq d(\psi, \varphi_N) + d(\varphi_N, \varphi_n) < M + \epsilon.$$

Thus, $(\varphi_n)_{n \in \mathbb{N}}$ is contained in $B_{M+\epsilon+1}[\psi]$. ■

Definition 5.60. Let (X, d) be a metric space. Then (X, d) is called *complete* if every Cauchy sequence in X converges to a $\varphi \in X$ wrt d .

Example 5.61. (\mathbb{C}, d) with $d(z_1, z_2) := |z_1 - z_2|$ is a complete metric space.

Proof. Let $(z_n)_{n \in \mathbb{N}}$ be a Cauchy sequence in \mathbb{C} , i.e. for all $\epsilon > 0$ there exists an $N \in \mathbb{N}$ s.t. $|z_n - z_m| < \epsilon$ for all $m, n \geq N$. Since $z_n = x_n + iy_n$ and $z_m = x_m + iy_m$ with $x_n, y_n, x_m, y_m \in \mathbb{R}$, we have

$$\epsilon > |z_n - z_m| = |x_n + iy_n - x_m - iy_m| = |x_n - x_m + i(y_n - y_m)| \geq |x_n - x_m|,$$

since for any $w \in \mathbb{C}$ we have $|w| \geq |\operatorname{Re}(w)|$. Also,

$$\epsilon > |z_n - z_m| = |x_n + iy_n - x_m - iy_m| = |x_n - x_m + i(y_n - y_m)| \geq |y_n - y_m|,$$

since for any $w \in \mathbb{C}$ we have $|w| \geq |\operatorname{Im}(w)|$. For all $n, m \geq N$, $\epsilon > |x_n - x_m|$ and $\epsilon > |y_n - y_m|$ implies that $(x_n)_{n \in \mathbb{N}}$ and $(y_n)_{n \in \mathbb{N}}$ are Cauchy sequences in \mathbb{R} , and since (\mathbb{R}, d) is complete, cf. Proposition B.21, $(x_n)_{n \in \mathbb{N}}$ converges to an $x \in \mathbb{R}$ and $(y_n)_{n \in \mathbb{N}}$ converges to a $y \in \mathbb{R}$. Thus, $(z_n)_{n \in \mathbb{N}}$ converges to $z = x + iy \in \mathbb{C}$. ■

Theorem 5.62. Let (X, d) be a metric space and consider the set Y . If $f : X \rightarrow Y$ is a bijection, then we can endow Y with the following metric d' :

$$d'(y_1, y_2) := d(f^{-1}(y_1), f^{-1}(y_2)) \quad \forall y_1, y_2 \in Y \quad (5.50)$$

(X, d) is complete iff (Y, d') is complete [28].

Proof. Let (X, d) be complete, i.e. every Cauchy sequence in X converges in X . Now let $(x_n)_{n \in \mathbb{N}}$ be a Cauchy sequence in X with limit point $\varphi \in X$. Consider the sequence $(y_n)_{n \in \mathbb{N}} := (f(x_n))_{n \in \mathbb{N}}$ in Y . Since X is Cauchy, we have $d(x_m, x_n) = d'(y_m, y_n) < \epsilon$ for all $\epsilon > 0$ and $m, n \geq N$. Thus, we have that $(y_n)_{n \in \mathbb{N}}$ is also a Cauchy sequence in Y . The limit point of $(y_n)_{n \in \mathbb{N}}$ is $f(\varphi)$, since $d'(y_n, f(\varphi)) = d(x_n, \varphi) < \epsilon'$ for all $\epsilon' > 0$ and $n \geq N'$. ■

Remark 5.63. To construct incomplete metric spaces, we can thus take an incomplete metric space (X, d) , construct a surjection $f : X \rightarrow Y$ to another set Y and endow Y with the metric d' from Eq. (5.50), then (Y, d') is also incomplete.

Example 5.64. Let $X = (-\frac{\pi}{2}, \frac{\pi}{2})$, and consider the metric $d(x_1, x_2) := |x_1 - x_2|$. Define $Y = \mathbb{R}$, and consider the map $f : X \rightarrow Y$, $x \mapsto \tan(x)$, which is surjective. On Y , consider the metric $d'(y_1, y_2) := d(\arctan(y_1), \arctan(y_2)) = |\arctan y_1 - \arctan y_2|$, cf. Eq. (5.50). Since (X, d) is incomplete, (Y, d') is incomplete as well. To see that (X, d) is incomplete, note that $(\frac{\pi}{2} - \frac{1}{n})_{n \in \mathbb{N}}$ or $(-\frac{\pi}{2} + \frac{1}{n})_{n \in \mathbb{N}}$ are Cauchy in X , yet do not converge in X .

Example 5.65 (Complete metric spaces). Any set X equipped with the discrete metric from Def. 5.6 is complete.

Example 5.66. The metric space (\mathbb{R}, d) with $d(x, y) := |x - y|$ is complete. Also, $(\mathbb{R}^d, \|x\|_p)$, cf. Example 4.3, for $1 \leq p \leq \infty$ is complete.

Example 5.67. Let $\mathcal{C}[a, b]$ be the space of all continuous functions $\chi : [a, b] \rightarrow \mathbb{R}$ with the metric

$$d_\infty(\varphi, \psi) = \|\varphi - \psi\|_\infty := \sup_{x \in [a, b]} \{|\varphi(x) - \psi(x)|\} \quad \forall \varphi, \psi \in \mathcal{C}[a, b]. \quad (5.51)$$

The metric space $(\mathcal{C}[a, b], d_\infty)$ is complete.

Proof. We first show that d_∞ is well-defined. Let $\varphi, \psi \in \mathcal{C}[a, b]$, then by Theorem 6.133, $\varphi([a, b]) \subset \mathbb{R}$ and $\psi([a, b]) \subset \mathbb{R}$ are compact. Thus, by Theorem 6.110, $\varphi([a, b]) \times \psi([a, b])$ is also compact. The function $|\varphi(x) - \psi(x)| : \varphi([a, b]) \times \psi([a, b]) \rightarrow \mathbb{R}$ is continuous, since the Euclidean metric is, cf. Theorem 6.93. Thus, by Corollary 6.134, the function $|\varphi(x) - \psi(x)|$ attains its maximum on $\varphi([a, b]) \times \psi([a, b])$, i.e. d_∞ is well-defined.

We now show that $(\mathcal{C}[a, b], d_\infty)$ is indeed a metric space:

1. positivity: $d_\infty(\varphi, \psi) = \sup_{x \in [a, b]} \{|\varphi(x) - \psi(x)|\} \geq 0$ for all $\varphi, \psi \in \mathcal{C}[a, b]$,
2. symmetry: $d_\infty(\varphi, \psi) = d_\infty(\psi, \varphi)$,
3. definiteness: $d_\infty(\varphi, \psi) = \sup_{x \in [a, b]} \{|\varphi(x) - \psi(x)|\} = 0 \Leftrightarrow |\varphi(x) - \psi(x)| = 0$ for all $x \in [a, b] \Leftrightarrow \varphi = \psi$,
4. triangle inequality: Let $\varphi, \psi, \chi \in \mathcal{C}[a, b]$, then

$$\begin{aligned} d_\infty(\varphi, \psi) &= \sup_{x \in [a, b]} \{|\varphi(x) - \psi(x)|\} \\ &= \sup_{x \in [a, b]} \{|\varphi(x) - \chi(x) + \chi(x) - \psi(x)|\} \\ &\stackrel{(11.18)}{\leq} \sup_{x \in [a, b]} \{|\varphi(x) - \chi(x)| + |\chi(x) - \psi(x)|\} \\ &\stackrel{(11.14)}{\leq} \sup_{x \in [a, b]} \{|\varphi(x) - \chi(x)|\} + \sup_{x \in [a, b]} \{|\chi(x) - \psi(x)|\} \\ &= d_\infty(\varphi, \chi) + d_\infty(\chi, \psi). \end{aligned}$$

Finally, we show that $(\mathcal{C}[a, b], d_\infty)$ is complete. Let $(\varphi_n)_{n \in \mathbb{N}}$ be a Cauchy sequence in $\mathcal{C}[a, b]$, i. e. for all $\epsilon > 0$ there exists an $N \in \mathbb{N}$ s.t. for all $m, n \geq N$, $d_\infty(\varphi_n, \varphi_m) < \epsilon$. Let $\epsilon > 0$ be arbitrary, then for any $x \in [a, b]$ we have

$$|\varphi_n(x) - \varphi_m(x)| \leq \sup_{x \in [a, b]} \{|\varphi_n(x) - \varphi_m(x)|\} = d_\infty(\varphi_n, \varphi_m) < \epsilon \quad \forall m, n \geq N. \quad (5.52)$$

Hence, $(\varphi_n(x))_{n \in \mathbb{N}}$ is a Cauchy sequence in \mathbb{R} for any $x \in [a, b]$ wrt the metric $d(x, y) := |x - y|$. Since (\mathbb{R}, d) is complete, we know that $\varphi(x) := \lim_{n \rightarrow \infty} \varphi_n(x)$ exists.

Finally, let us show that $\varphi \in \mathcal{C}[a, b]$. Since $(\varphi_n(x))_{n \in \mathbb{N}}$ converges in (\mathbb{R}, d) , we can choose an $N \in \mathbb{N}$ s.t. $|\varphi_n(x) - \varphi(x)| < \epsilon/3$ for all $n \geq N$ and all $x \in [a, b]$. And since $(\varphi_n)_{n \in \mathbb{N}}$ is a Cauchy sequence in $(\mathcal{C}[a, b], d_\infty)$ (by assumption), i.e. the φ_n are continuous, there exists a $\delta > 0$ s.t. $|\varphi_N(x) - \varphi_N(y)| < \epsilon/3$ with $|x - y| < \delta$. Let $x, y \in [a, b]$ be arbitrary, then if $|x - y| < \delta$, we get by applying the triangle inequality

$$|\varphi(x) - \varphi(y)| \leq |\varphi(x) - \varphi_N(x)| + |\varphi_N(x) - \varphi_N(y)| + |\varphi_N(y) - \varphi(y)| < \epsilon, \quad (5.53)$$

which shows the continuity of φ , cf. Def. 5.123, i.e. $\varphi \in \mathcal{C}[a, b]$. ■

Remark 5.68. Let (X, d) be a metric space. Then if $(\varphi_n)_{n \in \mathbb{N}}$ and $(\psi_n)_{n \in \mathbb{N}}$ are Cauchy sequences in (X, d) , their sum $(\varphi_n + \psi_n)_{n \in \mathbb{N}}$ is *not* necessarily a Cauchy sequence in X wrt d . (Note that we are assuming here that X is equipped with a binary operation $+$.)

Example 5.69. Consider the metric space (\mathbb{R}, d) with $d(x, y) := |\arctan x - \arctan y|$ for any $x, y \in \mathbb{R}$, cf. Example 5.64. Define the sequences $(a_n)_{n \in \mathbb{N}}$ as $a_n := n$ and $(b_n)_{n \in \mathbb{N}}$ as $b_n := (-1)^n - n$. Note that both sequences are Cauchy, but $(a_n + b_n)_{n \in \mathbb{N}} = ((-1)^n)_{n \in \mathbb{N}}$ is not.

Remark 5.70. Let (X, d) be a metric space. Then if $(\varphi_n)_{n \in \mathbb{N}}$ and $(\psi_n)_{n \in \mathbb{N}}$ are Cauchy sequences in (X, d) , their product $(\varphi_n \cdot \psi_n)_{n \in \mathbb{N}}$ is *not* necessarily a Cauchy sequence in X wrt d . (Note that we are assuming here that X is equipped with a binary operation \cdot .)

Example 5.71. Consider the metric space (\mathbb{R}, d) with $d(x, y) := |\arctan x - \arctan y|$ for any $x, y \in \mathbb{R}$, cf. Example 5.64. Define the sequences $(a_n)_{n \in \mathbb{N}}$ as $a_n := n$ and $(b_n)_{n \in \mathbb{N}}$ as $b_n := \frac{(-1)^n}{n}$. Note that both sequences are Cauchy, but $(a_n \cdot b_n)_{n \in \mathbb{N}} = ((-1)^n)_{n \in \mathbb{N}}$ is not.

Remark 5.72. In the special case of $X = \mathbb{Q}$ or $X = \mathbb{R}$ with $d(x, y) := |x - y|$ for any $x, y \in X$, then the sum and product of two Cauchy sequences is indeed Cauchy again, cf. Theorem B.3 and B.4.

Proof of Example 5.69. First, we will show that $(a_n)_{n \in \mathbb{N}} = (n)_{n \in \mathbb{N}}$ is Cauchy. For any $\epsilon > 0$, choose $N = N(\epsilon)$ s.t. $N > 1/\epsilon$, and note that for any $m, n \geq N$, we have

$$d(a_m, a_n) = |\arctan m - \arctan n| = \left| \arctan \left(\frac{m - n}{1 + mn} \right) \right| \stackrel{(*)}{\leq} \left| \frac{m - n}{1 + mn} \right| \leq \frac{|m - n|}{mn}, \quad (5.54)$$

where the inequality in $(*)$ can be obtained by using the integral definition of $\arctan x$. WLOG assume $n \leq m$, then Equation (5.54) can be further simplified to

$$d(a_m, a_n) \stackrel{(5.54)}{\leq} \frac{|m - n|}{mn} \leq \frac{m}{mn} \stackrel{n \geq N}{\leq} \frac{m}{mN} = \frac{1}{N} \stackrel{N > \epsilon^{-1}}{<} \epsilon.$$

To show that $(b_n)_{n \in \mathbb{N}} = ((-1)^n - n)_{n \in \mathbb{N}}$ is Cauchy, choose $N = N(\epsilon)$ s.t. $N > \frac{1}{\epsilon} + 2$, and

note that for $m, n \geq N$, we have

$$d(b_m, b_n) = |\arctan((-1)^m - m)| - |\arctan((-1)^n - n)| \quad (5.55)$$

$$= \left| \arctan \left(\frac{(-1)^m - m - (-1)^n + n}{1 + ((-1)^m - m) \cdot ((-1)^n - n)} \right) \right| \quad (5.56)$$

$$\leq \left| \frac{(-1)^m - m - (-1)^n + n}{1 + ((-1)^m - m) \cdot ((-1)^n - n)} \right| \quad (5.57)$$

$$= \left| \frac{1}{1 + ((-1)^m - m) \cdot ((-1)^n - n)} \right| \cdot |(-1)^m - m - (-1)^n + n| \quad (5.58)$$

$$\leq \left| \frac{1}{1 + ((-1)^m - m) \cdot ((-1)^n - n)} \right| |n - m| \quad (5.59)$$

$$= \left| \frac{m - n}{1 + (-1)^{m+n} - m \cdot (-1)^n - n \cdot (-1)^m + mn} \right| \quad (5.60)$$

$$\leq \left| \frac{m - n}{-m \cdot (-1)^n - n \cdot (-1)^m + mn} \right| \quad (5.61)$$

We will distinguish four cases:

1. If m and n are both uneven, then Eq. (5.61) becomes

$$\begin{aligned} d(b_m, b_n) &\stackrel{(5.61)}{\leq} \left| \frac{m - n}{-m \cdot (-1)^n - n \cdot (-1)^m + mn} \right| \\ &= \left| \frac{m - n}{m + n + mn} \right| \leq \left| \frac{m - n}{nm} \right|, \end{aligned}$$

which reduces to showing that $(a_n)_{n \in \mathbb{N}}$ is Cauchy, cf. Eq. (5.54).

2. If m and n are both even and WLOG we assume $n \leq m$, then Eq. (5.61) becomes

$$\begin{aligned} d(b_m, b_n) &\stackrel{(5.61)}{\leq} \left| \frac{m - n}{-m \cdot (-1)^n - n \cdot (-1)^m + mn} \right| \\ &= \left| \frac{m - n}{-m - n + mn} \right| \stackrel{n \leq m}{\leq} \frac{m}{|-m - n + mn|} = \frac{1}{|-1 - \frac{n}{m} + n|} \\ &\stackrel{(\star)}{=} \frac{1}{-1 - \frac{n}{m} + n} \stackrel{(\diamond)}{\leq} \frac{1}{n - 2} \stackrel{n \geq N}{\leq} \frac{1}{N - 2} \stackrel{N > \epsilon^{-1} + 2}{<} \epsilon, \end{aligned}$$

where the equality in (\star) and the inequality in (\star) follow from $n \geq N > 2$ and $n \leq m$, since

$$n \leq m \Rightarrow \frac{n}{m} \leq 1 \Rightarrow -1 - \frac{n}{m} + n \geq n - 2 > 0.$$

3. If m is uneven and n is even and WLOG we assume $n \leq m$, then we have

$$\begin{aligned} d(b_m, b_n) &\stackrel{(5.61)}{\leq} \left| \frac{m - n}{-m \cdot (-1)^n - n \cdot (-1)^m + mn} \right| = \left| \frac{m - n}{-m + n + mn} \right| \\ &\stackrel{n \leq m}{\leq} \frac{m}{|n - m + mn|} = \frac{1}{|\frac{n}{m} - 1 + n|} = \frac{1}{\frac{n}{m} - 1 + n} \leq \frac{1}{n - 1} \\ &\stackrel{n \geq N}{\leq} \frac{1}{N - 1} \stackrel{N > \epsilon^{-1} + 2}{<} \epsilon. \end{aligned}$$

4. If m is even and n is uneven, proceed similarly to 3.

Finally, note that it is obvious that $(a_n + b_n)_{n \in \mathbb{N}} = ((-1)^n)_{n \in \mathbb{N}}$ is not Cauchy. \blacksquare

Proof of Example 5.71. We have already shown in the proof of Example 5.69 that $(a_n)_{n \in \mathbb{N}} = (n)_{n \in \mathbb{N}}$ is Cauchy wrt $d(x, y) := |\arctan x - \arctan y|$ for any $x, y \in \mathbb{R}$. To show that $(b_n)_{n \in \mathbb{N}} = \left(\frac{(-1)^n}{n}\right)_{n \in \mathbb{N}}$ is Cauchy wrt d , choose $N = N(\epsilon) > \frac{2}{\epsilon}$, then we have for any $m, n \geq N$:

$$\begin{aligned} d(b_m, b_n) &= \left| \arctan \left(\frac{(-1)^m}{m} \right) - \arctan \left(\frac{(-1)^n}{n} \right) \right| \\ &\leq \left| \arctan \left(\frac{(-1)^m}{m} \right) \right| + \left| \arctan \left(\frac{(-1)^n}{n} \right) \right| \\ &\leq \left| \frac{(-1)^m}{m} \right| + \left| \frac{(-1)^n}{n} \right| = \frac{1}{m} + \frac{1}{n} \\ &\stackrel{\text{wlog}}{\leq} \frac{2}{n \leq m} \stackrel{n \geq N}{\leq} \frac{2}{N} \stackrel{N > 2\epsilon^{-1}}{<} \epsilon. \end{aligned}$$

\blacksquare

Theorem 5.73. Let (X, d) be a metric space. Then the following statements are true.

- (a) Complete subsets of (X, d) are closed.
- (b) If (X, d) is complete, then every closed subset of (X, d) is complete.

Proof. (a): Let $U \subset X$ be closed wrt d , and let $\varphi \in X$ be a limit point of \overline{U} , i.e. there exists a sequence $(\varphi_n)_{n \in \mathbb{N}}$ in U s.t. $\lim_{n \rightarrow \infty} \varphi_n = \varphi$. Hence, $(\varphi_n)_{n \in \mathbb{N}}$ converges in X , and is thus a Cauchy sequence in X . Since U is closed, the limit of the sequence lies in U , i.e. U is closed.

(b): Let (X, d) be complete, $U \subset X$ closed and $(\varphi_n)_{n \in \mathbb{N}}$ a Cauchy sequence in U , which converges to a $\varphi \in X$, since X is closed. Since φ is the limit of $(\varphi_n)_{n \in \mathbb{N}}$ and U is closed, it follows that $\varphi \in U$. \blacksquare

Definition 5.74. Let (X, d) be a metric space. Then $U \subset X$ is called *dense* in X if $\overline{U} = X$, i.e. to every $\varphi \in X$ there exists a sequence $(\varphi_n)_{n \in \mathbb{N}}$ in U that converges to φ wrt d . Alternatively, we can say that to every $\varphi \in X$ and $\epsilon > 0$ there is a $\psi \in U$ s.t. $d(\varphi, \psi) < \epsilon$ (set e.g. $\psi := \varphi_N$, where N is s.t. for all $n \geq N$, $d(\varphi_n, \varphi) < \epsilon$).

Example 5.75. \mathbb{Q} lies dense in \mathbb{R} wrt $d(x, y) := |x - y|$ for all $x, y \in \mathbb{R}$.

Theorem 5.76. If $\varphi \in \mathbb{R} \setminus \mathbb{Q}$ and $\psi \in \mathbb{Q}$, then $\varphi + \psi$ is irrational. This is because any $\psi \in \mathbb{Q}$ can be written as p/q , where $p, q \in \mathbb{Z}$. Thus: If $\varphi + \psi$ was rational, we would have

$$\varphi + \psi = \varphi + \frac{p}{q} = \frac{r}{s} \Rightarrow \varphi = \frac{r}{s} - \frac{p}{q} = \frac{rq - ps}{qs} \in \mathbb{Q}, \quad (5.62)$$

which is a contradiction to the assumption $\varphi \in \mathbb{R} \setminus \mathbb{Q}$. This proves that the sum of a rational and irrational number is always irrational.

Example 5.77 (Dense subsets). $\mathbb{R} \setminus \mathbb{Q}$ lies dense in \mathbb{R} wrt $d(x, y) := |x - y|$ for all $x, y \in \mathbb{R}$.

Proof. We need to show that to every $\varphi \in \mathbb{R}$ there exists a sequence $(\varphi_n)_{n \in \mathbb{N}} \in \mathbb{R} \setminus \mathbb{Q}$ s.t. $\varphi_n \xrightarrow{n \rightarrow \infty} \varphi$ wrt d . We already know that \mathbb{Q} lies dense in \mathbb{R} wrt d , i.e. there exists a sequence $(\psi_n)_{n \in \mathbb{N}}$ s.t. $\psi_n \xrightarrow{n \rightarrow \infty} \varphi$. Now define the sequence $(\varphi_n)_{n \in \mathbb{N}} := (\psi_n + \frac{a}{n})_{n \in \mathbb{N}}$, where $a \in \mathbb{R} \setminus \mathbb{Q}$. Each sequence element lies in $\mathbb{R} \setminus \mathbb{Q}$, since the sum of a rational and an irrational number is irrational, cf. Theorem 5.76, and converges to φ wrt d . ■

Example 5.78 (Weierstrass Approximation Theorem). The algebraic polynomials \mathcal{P} are dense on an interval $[a, b] \subset \mathbb{R}$ with $-\infty < a < b < \infty$ in $\mathcal{C}[a, b]$ wrt the metric d_∞ from Example 5.67. [29, Corollary 6.12]

Proof. We delegeate the proof of this celebrated Theorem to appendix A. ■

Example 5.79. Let (X, d) be a metric space. Then $U \subset X$ lies dense in $\overline{U} \subset X$ wrt d .

Theorem 5.80. Let (X, d) be a metric space. Then $U \subset X$ is dense iff every non-empty open set $O \subset X$ intersects U , i.e. $U \cap O \neq \emptyset$.

Proof [30]. “ \implies ” Use Theorem 5.32 and note that $U \subset X$ is dense, i.e. $\overline{U} = X$.

“ \impliedby ” We will prove the contraposition, i.e. assume $U \subset X$ is not dense, then $\overline{U} \neq X$, which implies $O := X \setminus \overline{U} \neq \emptyset$, i.e. $O \cap \overline{U} = \emptyset$. Since $U \subset \overline{U}$, $O \cap U = \emptyset$. Clearly, O is open, since it is the complement of a closed set. ■

Theorem 5.81. Let X be a metric space. Then $U \subset X$ is dense iff $\text{int}(X \setminus U) = \emptyset$.

Proof [31]. $U \subset X$ is dense $\Leftrightarrow X = \overline{U} \Leftrightarrow \text{int}(X \setminus U) \stackrel{(5.22)}{=} X \setminus \overline{U} = X \setminus X = \emptyset$ ■

Theorem 5.82. Let X be a metric space and let $U_1, U_2 \subset X$ be open and dense. Then $U_1 \cap U_2$ is also open and dense.

Proof [32]. Since U_1 is dense, by Theorem 5.80, for every non-empty open set $O \subset X$, we have $U_1 \cap O \neq \emptyset$. Since both U_1 and O are open, so is $U_1 \cap O$. Thus, since U_2 is also dense, $U_2 \cap U_1 \cap O \neq \emptyset$, which shows that $U_1 \cap U_2$ intersects every non-empty open set $O \subset X$. Finally, $U_1 \cap U_2$ is also open, as it is the finite intersection of two open sets. ■

Remark 5.83. Let $U_1, U_2 \subset X$ be dense. As the proof in the preceding Theorem shows, it is only necessary that either U_1 or U_2 is open for $U_1 \cap U_2$ to be dense. However, it is necessary that either U_1 or U_2 is open. If neither U_1 or U_2 is open, the resulting intersection may not be dense.

Example 5.84 (Proof [33]). Let $X = \mathbb{R}$ with the standard metric. Consider the rationals \mathbb{Q} and irrationals $\mathbb{R} \setminus \mathbb{Q}$. Neither set is open in \mathbb{R} . Their intersection is the empty set, which is clearly not dense in \mathbb{R} .

Remark 5.85. Let d and d' be two metrics on X . If $U \subset X$ is dense wrt d , U is not necessarily dense wrt d' .

Example 5.86. Consider $X = \mathbb{R}$ and let $d(x, y) := |x - y|$ for all $x, y \in \mathbb{R}$. While $U = \mathbb{Q} \subset X = \mathbb{R}$ is dense wrt d , it is not dense wrt the discrete metric from Remark 5.6, since for $r \leq 1$, the open balls are given by the singletons $\{\varphi\}$ for any $\varphi \in X$, cf. Example 5.8. Let $\varphi = \pi$, then $B_r(\pi) = \{\pi\}$ for any $r \leq 1$, but clearly, $\mathbb{Q} \cap \{\pi\} = \emptyset$, and thus by Theorem 5.80 \mathbb{Q} is not dense in \mathbb{R} wrt the discrete metric.

Remark 5.87. Let (X, d) be a metric space. Then there are examples of dense subsets $U \subset X$ wrt d that are not complete wrt d , and vice versa.

Example 5.88. Consider the cantor set \mathcal{C} on $[0, 1]$, which we consider as a subset of the metric space $([0, 1], d)$ with $d(x, y) := |x - y|$ for all $x, y \in [0, 1]$. Formally, it is defined as follows:

$$C_0 := [0, 1], \quad (5.63)$$

$$C_n := \frac{C_{n-1}}{3} \cup \left(\frac{2}{3} + \frac{C_{n-1}}{3} \right) = \frac{1}{3} (C_{n-1} \cup (2 + C_{n-1})), \quad (5.64)$$

$$\mathcal{C} := \bigcap_{n=0}^{\infty} C_n. \quad (5.65)$$

Intuitively, the Cantor set \mathcal{C} is created by iteratively removing the open middle third of each line segment.

Since for all $n \in \mathbb{N}$, C_n is closed (as the finite union of closed sets) – for a formal proof, use induction – and the arbitrary intersection of closed sets is closed, we have that $\mathcal{C} \subset [0, 1]$ is closed wrt d . Thus, by Theorem 5.73 b), \mathcal{C} is complete wrt d . Now consider the open set $O := (\frac{1}{2}, \frac{2}{3}) \subset [0, 1]$, which is open wrt d . Clearly, $O \cap \mathcal{C} = \emptyset$, and thus by Theorem 5.80 \mathcal{C} is not dense wrt d .

Example 5.89. \mathbb{Q} is a dense subset of $X = \mathbb{R}$, where (X, d) with $d(x, y) := |x - y|$ for all $x, y \in \mathbb{R}$. However, we have already established that \mathbb{Q} is not complete wrt d , e.g. cf. Remark 5.52.

Definition 5.90 (Completion of metric spaces). Let (X, d) be a metric space. Then a *completion* of X is a triplet $(\tilde{X}, \tilde{d}, f)$, where (\tilde{X}, \tilde{d}) is a complete metric space and $f : X \rightarrow \tilde{X}$ is an isometry with dense image, i.e. $f(X) \subset \tilde{X}$ is dense wrt \tilde{d} [34, Def. 2.2].

Remark 5.91. Let (X, d) be a metric space. If a completion $(\tilde{X}, \tilde{d}, f)$ exists, then X is isometric to the dense subset $f(X)$.

Proposition 5.92. To every metric space (X, d) , a completion $(\tilde{X}, \tilde{d}, f)$ exists and up to isometry, it is unique.

Remark 5.93. We can henceforth talk about *the* completion of a metric space.

Example 5.94. Consider the metric space (\mathbb{Q}, d) with $d(r_1, r_2) := |r_1 - r_2|$ for any $r_1, r_2 \in \mathbb{Q}$. In Appendix B, we will construct the completion of \mathbb{Q} (the real numbers \mathbb{R}) with the extended metric $d_{\text{ext}}(x, y) := |x - y|$ for any $x, y \in \mathbb{R}$ with an isometry $f : \mathbb{Q} \rightarrow \mathbb{R}$.

Remark 5.95. The completion of \mathbb{R} wrt $|\cdot|$ is needed for the proof of Proposition 5.92.

Proof of Proposition 5.92. Let (X, d) be a metric space. We will proceed in several steps.

Step 1: We will first construct \tilde{X} . Let $(\varphi_n)_{n \in \mathbb{N}}$ and $(\psi_n)_{n \in \mathbb{N}}$ be Cauchy sequences in X . Then by Theorem 5.5,

$$|d(\varphi_n, \psi_n) - d(\varphi_m, \psi_m)| \leq d(\varphi_n, \varphi_m) + d(\psi_n, \psi_m) \xrightarrow{n, m \rightarrow \infty} 0, \quad (5.66)$$

which implies that $(d(\varphi_n, \psi_n))_{n \in \mathbb{N}}$ is a Cauchy sequence in \mathbb{R} with the usual metric. By the completeness \mathbb{R} , every Cauchy sequence in \mathbb{R} converges, and thus the limit $\lim_{n \rightarrow \infty} d(\varphi, \psi_n)$ exists.

We will now construct an equivalence relation on the set of Cauchy sequences in X as follows: Let $(\varphi_n)_{n \in \mathbb{N}}$ and $(\psi_n)_{n \in \mathbb{N}}$ be Cauchy sequences in X , then we call them equivalent if $d(\varphi_n, \psi_n) \xrightarrow{n \rightarrow \infty} 0$. By Def. 8.2, we need to show reflexivity (which holds since $d(\varphi_n, \varphi_n) = 0$), symmetry (which holds since $d(\varphi_n, \psi_n) = d(\psi_n, \varphi_n)$) and transitivity (which holds because $d(\varphi_n, \chi_n) \leq d(\varphi_n, \psi_n) + d(\psi_n, \chi_n)$ for any Cauchy sequence $(\chi_n)_{n \in \mathbb{N}}$ in X).

We will denote the set of all equivalent Cauchy sequences in X by \tilde{X} ,

$$\tilde{X} = \{(\varphi_n)_{n \in \mathbb{N}} \mid (\varphi_n)_{n \in \mathbb{N}} \text{ is a Cauchy sequence in } X\} / \sim. \quad (5.67)$$

Step 2: We will now define a metric \tilde{d} on \tilde{X} . Let $\tilde{\varphi}, \tilde{\psi} \in \tilde{X}$ with representations $(\varphi_n)_{n \in \mathbb{N}}$ and $(\psi_n)_{n \in \mathbb{N}}$. We set

$$\tilde{d}(\tilde{\varphi}, \tilde{\psi}) := \lim_{n \rightarrow \infty} d(\varphi_n, \psi_n). \quad (5.68)$$

The map $\tilde{d} : \tilde{X} \times \tilde{X} \rightarrow \mathbb{R}$ is well-defined, since for other representations $(\varphi'_n)_{n \in \mathbb{N}} \sim (\varphi_n)_{n \in \mathbb{N}}$ and $(\psi'_n)_{n \in \mathbb{N}} \sim (\psi_n)_{n \in \mathbb{N}}$, we have by Theorem 5.5 the inequality

$$|d(\varphi'_n, \psi'_n) - d(\varphi_n, \psi_n)| \leq d(\varphi'_n, \varphi_n) + d(\psi'_n, \psi_n) \xrightarrow{n \rightarrow \infty} 0. \quad (5.69)$$

And we have already established in Step 1 that the Cauchy sequence $(d(\varphi_n, \psi_n))_{n \in \mathbb{N}} \in \mathbb{R}$ is convergent wrt the usual metric on \mathbb{R} .

To show that \tilde{d} is indeed a metric on \tilde{X} , note that for any $\tilde{\varphi}, \tilde{\psi}, \tilde{\chi} \in \tilde{X}$ (with representations $(\varphi_n)_{n \in \mathbb{N}}$, $(\psi_n)_{n \in \mathbb{N}}$ and $(\chi_n)_{n \in \mathbb{N}}$), we have:

1. Positivity: $\tilde{d}(\tilde{\varphi}, \tilde{\varphi}) = \lim_{n \rightarrow \infty} d(\varphi_n, \varphi_n) = 0$.
2. Symmetry: $\tilde{d}(\tilde{\varphi}, \tilde{\psi}) = \lim_{n \rightarrow \infty} d(\varphi_n, \psi_n) = \lim_{n \rightarrow \infty} d(\psi_n, \varphi_n) = \tilde{d}(\tilde{\psi}, \tilde{\varphi})$.
3. Definiteness: $\tilde{d}(\tilde{\varphi}, \tilde{\psi}) = \lim_{n \rightarrow \infty} d(\varphi_n, \psi_n) = 0$ iff $(\varphi_n)_{n \in \mathbb{N}} \sim (\psi_n)_{n \in \mathbb{N}}$, i.e. $\tilde{\varphi} = \tilde{\psi}$, cf. Theorem 8.4.
4. Triangle inequality:

$$\tilde{d}(\tilde{\varphi}, \tilde{\psi}) = \lim_{n \rightarrow \infty} d(\varphi_n, \psi_n) \leq \lim_{n \rightarrow \infty} \{d(\varphi_n, \chi_n) + d(\chi_n, \psi_n)\}$$

Since $(d(\varphi_n, \chi_n))_{n \in \mathbb{N}}$ and $(d(\chi_n, \psi_n))_{n \in \mathbb{N}}$ are Cauchy sequences in \mathbb{R} , also cf. Step 1, we have

$$\begin{aligned} \tilde{d}(\tilde{\varphi}, \tilde{\psi}) &\leq \lim_{n \rightarrow \infty} \{d(\varphi_n, \chi_n) + d(\chi_n, \psi_n)\} = \lim_{n \rightarrow \infty} d(\varphi_n, \chi_n) + \lim_{n \rightarrow \infty} d(\chi_n, \psi_n) \\ &= \tilde{d}(\tilde{\varphi}, \tilde{\chi}) + \tilde{d}(\tilde{\chi}, \tilde{\psi}), \end{aligned}$$

which shows the triangle inequality.

Step 3: We now embed X isometrically as a dense subset of \tilde{X} . For this, consider the map $f : X \rightarrow \tilde{X}, \varphi \mapsto [(\varphi_n)_{n \in \mathbb{N}}]$ with $\varphi_n = \varphi$ for any $n \in \mathbb{N}$. f is an isometry, since

$$\tilde{d}(f(\varphi), f(\psi)) = \lim_{n \rightarrow \infty} d(\varphi_n, \psi_n) = \lim_{n \rightarrow \infty} d(\varphi, \psi) = d(\varphi, \psi). \quad (5.70)$$

The image of f under X , i.e. $f(X) \subset \tilde{X}$, is dense in \tilde{X} wrt \tilde{d} , since for any $\tilde{\varphi} \in \tilde{X}$ with representation $(\varphi_n)_{n \in \mathbb{N}}$ the sequence $(f(\varphi_n))_{n \in \mathbb{N}} \in f(X)$ converges to $\tilde{\varphi}$ wrt \tilde{d} :

$$\lim_{n \rightarrow \infty} \tilde{d}(\tilde{\varphi}, f(\varphi_n)) = \lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} d(\varphi_m, \varphi_n) = 0, \quad (5.71)$$

since $(\varphi_n)_{n \in \mathbb{N}}$ is a Cauchy sequence in X wrt d .

Step 4: We will now show that (\tilde{X}, \tilde{d}) is a complete metric space.

Let $(\tilde{\varphi}_n)_{n \in \mathbb{N}}$ be a Cauchy sequence in \tilde{X} . Since $f(X) \subset \tilde{X}$ is dense wrt \tilde{d} , to every $\tilde{\varphi}_n$ there is a $\varphi_n \in X$ s.t. $\tilde{d}(\tilde{\varphi}_n, f(\varphi_n)) < 1/n$. Hence, by applying the triangle equality twice, we have

$$d(\varphi_n, \varphi_m) = \tilde{d}(f(\varphi_n), f(\varphi_m)) \leq \tilde{d}(f(\varphi_n), \tilde{\varphi}_n) + \tilde{d}(\tilde{\varphi}_n, \tilde{\varphi}_m) + \tilde{d}(\tilde{\varphi}_m, f(\varphi_m)) \quad (5.72)$$

$$< \frac{1}{n} + \tilde{d}(\tilde{\varphi}_n, \tilde{\varphi}_m) + \frac{1}{m} \xrightarrow{n, m \rightarrow \infty} 0, \quad (5.73)$$

since by assumption, $(\tilde{\varphi}_n)_{n \in \mathbb{N}}$ is a Cauchy sequence. Hence, $(\varphi_n)_{n \in \mathbb{N}}$ is a Cauchy sequence in X , and hence a representation for some $\tilde{\varphi} \in \tilde{X}$. We will now show that $(\tilde{\varphi}_n)_{n \in \mathbb{N}}$ converges to $\tilde{\varphi}$:

$$\tilde{d}(\tilde{\varphi}_n, \tilde{\varphi}) \leq \tilde{d}(\tilde{\varphi}, f(\varphi_n)) + \tilde{d}(f(\varphi_n), \tilde{\varphi}_n) < \tilde{d}(\tilde{\varphi}, f(\varphi_n)) + \frac{1}{n} \quad (5.74)$$

$$= \lim_{m \rightarrow \infty} d(f(\varphi_n)_m, \varphi_m) + \frac{1}{n} = \lim_{m \rightarrow \infty} d(\varphi_n, \varphi_m) + \frac{1}{n} \quad (5.75)$$

$$\Rightarrow \lim_{n \rightarrow \infty} \tilde{d}(\tilde{\varphi}_n, \tilde{\varphi}) \leq \lim_{n, m \rightarrow \infty} d(\varphi_n, \varphi_m) = 0. \quad (5.76)$$

Hence, (\tilde{X}, \tilde{d}) is complete.

Step 5: We will now show \tilde{X} , a completion of X , is unique up to isometry.

Let (Y, d_Y, f) and (Z, d_Z, g) be completions of the metric space (X, d) , i.e. (Y, d_Y) and (Z, d_Z) are complete metric spaces, and $f : X \rightarrow Y$ and $g : X \rightarrow Z$ are isometries with dense subsets $f(X) \subset Y$ wrt d_Y and $g(X) \subset Z$ wrt d_Z .

We will now show that (Y, d_Y) and (Z, d_Z) are isometric. For this, let $\psi \in Y$. Since $f(X) \subset Y$ is dense wrt d_Y , there is a sequence $(\varphi_n)_{n \in \mathbb{N}}$ in X s.t.

$$\lim_{n \rightarrow \infty} d_Y(\psi, f(\varphi_n)) = 0.$$

Now define $h : Y \rightarrow Z$ via

$$h(\psi) := \lim_{n \rightarrow \infty} g(\varphi_n). \quad (5.77)$$

Note the following properties of h :

- The limit in Eq. (5.77) exists, since

$$d_Z(g(\varphi_n), g(\varphi_m)) = d_X(\varphi_n, \varphi_m) = d_Y(f(\varphi_n), f(\varphi_m)) \quad (5.78)$$

$$\Rightarrow \lim_{n, m \rightarrow \infty} d_Z(g(\varphi_n), g(\varphi_m)) = \lim_{n, m \rightarrow \infty} d_Y(f(\varphi_n), f(\varphi_m)) = d_Y(\psi, \psi) = 0, \quad (5.79)$$

since by Theorem 5.5 we have that

$$|d_Y(f(\varphi_n), f(\varphi_m)) - d_Y(\psi, \psi)| \leq d_Y(f(\varphi_n), \psi) + d_Y(f(\varphi_m), f(\psi)).$$

Eq. (5.79) shows that $(g(\varphi_n))_{n \in \mathbb{N}}$ is a Cauchy sequence in Z , which also converges in Z , since by assumption, (Z, d_Z) is complete.

- Let $(\varphi'_n)_{n \in \mathbb{N}}$ be another sequence in X converging to ψ wrt d_Y . Then

$$\begin{aligned} d_Z(g(\varphi_n), g(\varphi'_n)) &= d_X(\varphi_n, \varphi'_n) = d_Y(f(\varphi_n), f(\varphi'_n)) \xrightarrow{n \rightarrow \infty} d_Y(\psi, \psi) = 0 \\ \Rightarrow \lim_{n \rightarrow \infty} g(\varphi_n) &= \lim_{n \rightarrow \infty} g(\varphi'_n), \end{aligned}$$

which makes the limit in Eq. (5.77) well-defined.

- The map h is an isometry, since from the triangle inequality and the inequality in Eq. (5.5), we have for arbitrary $\psi, \psi' \in Y$

$$\begin{aligned} &|d_Z(h(\psi), h(\psi')) - d_Y(\psi, \psi')| \\ &= \left| d_Z(h(\psi), h(\psi')) + \underbrace{d_Y(f(\varphi_n), f(\varphi'_n)) - d_Y(f(\varphi_n), f(\varphi'_n))}_{=0} - d_Y(\psi, \psi') \right| \\ &= \left| d_Z(h(\psi), h(\psi')) + d_Y(f(\varphi_n), f(\varphi'_n)) - \underbrace{d_Y(f(\varphi_n), f(\varphi'_n))}_{=d_X(\varphi_n, \varphi'_n)=d_Z(g(\varphi_n), g(\varphi'_n))} - d_Y(\psi, \psi') \right| \\ &\leq |d_Z(h(\psi), h(\psi')) - d_Z(g(\varphi_n), g(\varphi'_n))| + |d_Y(f(\varphi_n), f(\varphi'_n)) - d_Y(\psi, \psi')| \\ &\leq d_Z(h(\psi), g(\varphi_n)) + d_Z(h(\psi'), g(\varphi'_n)) + d_Y(f(\varphi_n), \psi) + d_Y(f(\varphi'_n), \psi') \\ &\xrightarrow{n \rightarrow \infty} 0, \end{aligned}$$

since we choose the sequences $(\varphi_n)_{n \in \mathbb{N}}$ and $(\varphi'_n)_{n \in \mathbb{N}}$ in X such that $f(\varphi_n) \xrightarrow{n \rightarrow \infty} \psi$ and $f(\varphi'_n) \xrightarrow{n \rightarrow \infty} \psi'$ (wrt d_Y each). By definition, cf. Eq. (5.77), we then have $h(\psi) = \lim_{n \rightarrow \infty} g(\varphi_n)$ and $h(\psi') = \lim_{n \rightarrow \infty} g(\varphi'_n)$.

The above calculation shows that $d_Z(h(\psi), h(\psi')) = d_Y(\psi, \psi')$ for arbitrary $\psi, \psi' \in Y$.

- Finally, h is surjective, since for every $\chi \in Z$ there is a sequence $(\varphi_n)_{n \in \mathbb{N}} \in X$ s.t. $\lim_{n \rightarrow \infty} g(\varphi_n) = \chi$, since $g(X)$ is dense in Z wrt d_Z . Now consider the limit $\psi := \lim_{n \rightarrow \infty} f(\varphi_n) \in Y$ and note that $\chi = h(\psi) = \lim_{n \rightarrow \infty} g(\varphi_n)$.

Note that the existence of the limit $\psi = \lim_{n \rightarrow \infty} f(\varphi_n) \in Y$ follows from

$$d_Y(f(\varphi_n), f(\varphi_m)) = d_X(\varphi_n, \varphi_m) = d_Z(g(\varphi_n), g(\varphi_m)) \xrightarrow{n \rightarrow \infty} 0, \quad (5.80)$$

i.e. $(f(\varphi_n))_{n \in \mathbb{N}}$ is a Cauchy sequence in Y , and hence by the completeness of (Y, d_Y) , the sequence also converges in Y .

This proof of the existence of the limit $\psi = \lim_{n \rightarrow \infty} f(\varphi_n)$ is very similar to how we proved the existence of the limit in Eq. (5.77) for the definition of h .

■

5.2 Compactness

Definition 5.96 (Covering [20]). Let X be a topological space, cf. Definition 6.1. A *covering* of X is a family \mathcal{G} of subsets of X s.t.

$$X = \bigcup \{G \mid G \in \mathcal{G}\}.$$

We also say that \mathcal{G} covers X . If every member of \mathcal{G} is open, then \mathcal{G} is called an *open covering* of X .

If $U \subset X$, then a family \mathcal{G} of subsets of X is called a covering of U if

$$U \subset \bigcup \{G \mid G \in \mathcal{G}\}.$$

Definition 5.97 (Compact [20]). Let X be a topological space and $U \subset X$. Then U is called

- *compact* if every open covering of U contains a finite sub-covering, i.e. if to every family $\{V_j\}_{j \in J}$ of open sets with $U \subset \bigcup_{j \in J} V_j$ there are finitely many V_{j_1}, \dots, V_{j_n} , $n \in \mathbb{N}$, s.t.

$$U \subset \bigcup_{k=1}^n V_{j_k},$$

- *sequentially compact* if every sequence in U contains a convergent subsequence in U .

Example 5.98. Consider $U := (0, 1]$, which is a subset of the topological space \mathbb{R} , where we equip \mathbb{R} with the standard topology. Then $V_j := (1/j, 1 + 1/j)$, $j \in \mathbb{N}$, are an open covering of U , but U is not compact, since there is no *finite* covering of U with open sets from from $\{V_j\}_{j \in \mathbb{N}}$.

U is also not sequentially compact, since the sequence $(1/n)_{n \in \mathbb{N}}$ does not contain a convergent subsequence in U ($0 \notin U$).

Example 5.99. $(\mathbb{R}, |\cdot|)$ is obviously not compact. Also, it is not sequentially compact, since $(n)_{n \in \mathbb{N}}$ does not contain a convergent subsequence in \mathbb{R} .

Definition 5.100 (Totally bounded set). A subset U of a metric space is called *totally bounded* if for every $\epsilon > 0$ there are finitely many elements $\varphi_1, \dots, \varphi_n \in U$ s.t.

$$U \subset \bigcup_{i=1}^n B_\epsilon(\varphi_i),$$

i.e. to every $\varphi \in U$ there is a $\varphi_k \in U$, $1 \leq k \leq n$, s.t. $d(\varphi, \varphi_k) < \epsilon$.

Theorem 5.101. Every totally bounded set is bounded.

Proof. Let $\varphi_1, \dots, \varphi_n \in U$ be s.t.

$$U \subset \bigcup_{i=1}^n B_\epsilon(\varphi_i)$$

for all $\epsilon > 0$. Now note that

$$U \subset \bigcup_{i=1}^n B_\epsilon(\varphi_i) \subset \bigcup_{i=1}^n B_\epsilon[\varphi_i],$$

and that by Theorem 5.16, the finite union of closed sets is also closed. Thus, U is bounded. ■

Remark 5.102. The converse of Theorem 5.101 is not true in general: There are bounded sets that are not totally bounded.

Example 5.103. Consider \mathbb{R} with the metric $d(x, y) := \min\{1, |x - y|\}$. Then \mathbb{R} (as a subset of \mathbb{R}) is bounded, but not totally bounded.

Proof. \mathbb{R} is bounded wrt d , since $\mathbb{R} \subset B_r[0]$, where $r \geq 1$. However, \mathbb{R} is not totally bounded. For this, let $\epsilon \leq 1$ be arbitrary, then it is clear that we need infinitely many open balls to cover \mathbb{Z} , and thus also \mathbb{R} . ■

Theorem 5.104. Every sequentially compact set is totally bounded.

Proof. We will prove this by contradiction. So let U be sequentially compact, but not totally bounded. The fact that U is totally bounded can be written with quantifiers:

$$\forall \epsilon > 0 \exists \psi_1, \dots, \psi_n \in U : U \subset \bigcup_{i=1}^n B_\epsilon(\psi_i)$$

So if U is not totally bounded, we can write this as

$$\exists \epsilon > 0 \forall \psi_1, \dots, \psi_n \in U : \exists \psi \in U, \psi \notin \bigcup_{i=1}^n B_\epsilon(\psi_i)$$

So there is an $\epsilon > 0$ s.t. for arbitrary $\psi_1, \dots, \psi_n \in U$, $n \in \mathbb{N}$, there is a $\psi_{n+1} \in U$ s.t. $d(\psi_{n+1}, \psi_j) \geq \epsilon$ for all $j \in \{1, \dots, n\}$. Thus, we can also find a $\psi_{n+2} \in U$ s.t. $d(\psi_{n+2}, \psi_j) \geq \epsilon$ for all $j \in \{1, \dots, n+1\}$. Hence, we can recursively construct a sequence $(\psi_n)_{n \in \mathbb{N}}$ in U s.t.

$$d(\psi_l, \psi_k) \geq \epsilon \quad \text{for } n < k < l.$$

However, such a sequence cannot contain a convergent subsequence, which is in contradiction to U being sequentially compact. ■

Definition 5.105. A set U of a metric space is called *separable* if U contains a countable dense subset.

Example 5.106. Consider the metric space (\mathbb{R}, d) , where d is the usual metric. Then $\mathbb{Q} \subset \mathbb{R}$ is separable wrt d .

Theorem 5.107. Every totally bounded set is separable.

Proof. Let U be totally bounded. There for every $\epsilon_m = 1/m > 0$ there is a finite sequence of elements $\varphi_{1,m}, \dots, \varphi_{n_m,m} \in U$ s.t.

$$U \subset \bigcup_{j=1}^{n_m} B_{\epsilon_m}(\varphi_{j,m}).$$

Consider the set

$$V := \{\varphi_{1,1}, \dots, \varphi_{n_1,1}, \varphi_{1,2}, \dots, \varphi_{n_2,2}, \varphi_{1,3}, \dots, \varphi_{n_3,3}, \dots\},$$

which is the set of elements of the sequence $(\varphi_{1,m}, \dots, \varphi_{n_m,m})_{m \in \mathbb{N}}$. Obviously, V is countable. Also, for every $\varphi \in U$ and $\epsilon > 0$, there is an $\epsilon_m = 1/m < \epsilon$ and a $\varphi_{j,m} \in V$ s.t. $d(\varphi_{j,m}, \varphi) < \epsilon_m = 1/m < \epsilon$. Thus, V is dense in U . ■

Theorem 5.108. Let U be a subset of a metric space, then the following statements are equivalent:

- (a) U is compact,
- (b) U is sequentially compact,
- (c) U is totally bounded and complete.

Proof. “(a) \implies (b)”: Proof by contradiction: Let U be compact, but not sequentially compact. Then there is a sequence $(\varphi_n)_{n \in \mathbb{N}}$ in U that does not have a convergent subsequence. Thus, to every $\varphi \in U$ there exists an open ball $B_\varphi = B_{r_\varphi}(\varphi)$, where B_φ only contains finitely many elements of $(\varphi_n)_{n \in \mathbb{N}}$.

However, since U is compact, there are finitely many elements $\psi_1, \dots, \psi_n \in U$ s.t.

$$U \subset \bigcup_{k=1}^n B_{\psi_k},$$

where B_{ψ_k} only contains finitely many points of $(\varphi_n)_{n \in \mathbb{N}}$. But then, U contains only finitely many elements of $(\varphi_n)_{n \in \mathbb{N}}$, which is a contradiction.

“(b) \implies (c)”: Let U be sequentially compact and thus totally bounded by Theorem (5.104). It now remains to be shown that U is complete. Let $(\varphi_n)_{n \in \mathbb{N}}$ be a Cauchy sequence in U , then it contains a convergent subsequence in U , since U is sequentially compact, and thus by Theorem 5.51, $(\varphi_n)_{n \in \mathbb{N}}$ converges.

“(c) \implies (a)”: Let U be complete and totally bounded. We will show that U is compact via contradiction.

Let $\{V_j\}_{j \in J}$ be an open cover of U s.t.

$$U \subset \bigcup_{j \in J} V_j. \tag{5.81}$$

We now define

$$\mathcal{B} := \left\{ B \subset U \mid B \subset \bigcup_{i \in I} V_i \Rightarrow I \text{ infinite, where } I \subset J \right\}$$

and we will show that $U \notin \mathcal{B}$.

Since U is totally bounded, for all $\epsilon > 0$ there is an open cover

$$U \subset \bigcup_{i=1}^{n_\epsilon} B_\epsilon(\varphi_i) \tag{5.82}$$

with points $\varphi_1, \dots, \varphi_n \in U$. Further, to *each* $B \in \mathcal{B}$, there is an $i(\epsilon)$ with $B_\epsilon(\varphi_i) \cap B \in \mathcal{B}$. This holds because according to Eq. (5.82),

$$B \cap U = B \subset \left(\bigcup_{i=1}^{n_\epsilon} B_\epsilon(\varphi_i) \right) \cap B = \bigcup_{i=1}^{n_\epsilon} (B_\epsilon(\varphi_i) \cap B),$$

where the last equality follows from the distributive law for sets. Since $B_\epsilon(\varphi_i) \cap B \subset B$, we have equality, i.e.

$$B = \bigcup_{i=1}^{n_\epsilon} (B_\epsilon(\varphi_i) \cap B).$$

Since we assumed $B \in \mathcal{B}$, this implies that there is an $i(\epsilon)$ s.t. $B_\epsilon(\varphi_i) \cap B \in \mathcal{B}$, since otherwise, B could be covered by a finite collection of $\{V_i\}_{i \in I}$.

If $U \in \mathcal{B}$, define $B_1 := U \in \mathcal{B}$, and then recursively construct (for $\epsilon_k = 1/k$)

$$B_k := B_{\epsilon_k}(\varphi_k) \cap B_{k-1} \in \mathcal{B} \quad \text{for } k \geq 2. \quad (5.83)$$

By construction, we have

$$B_k \subset B_{k-1} \subset \dots \subset B_2 \subset B_1 = U.$$

Now, for any n , choose $\psi_n \in B_n$, then for all $l, k \in \mathbb{N}$ we have (WLOG, let $l \geq k$),

$$d(\psi_k, \psi_l) \leq d(\psi_k, \varphi_k) + d(\varphi_k, \psi_l) < 2\epsilon_k, \quad (5.84)$$

and hence $(\psi_n)_{n \in \mathbb{N}}$ is a Cauchy sequence in U . Since U is complete, $(\psi_n)_{n \in \mathbb{N}}$ converges to a $\psi \in U$ wrt d . Since $\psi \in V_{j_0}$ for a $j_0 \in J$, cf. Eq. (5.81), for sufficiently large k , we have

$$B_k \subset B_{\epsilon_k}(\varphi_k) \subset B_{2\epsilon_k}(\psi_k) \subset B_{2\epsilon_k + d(\psi_k, \psi)}(\psi) \subset V_{j_0}.$$

For the second inclusion, note that for any $\varphi \in B_{\epsilon_k}(\varphi_k)$,

$$d(\varphi, \psi_k) \leq d(\psi_k, \varphi_k) + d(\varphi_k, \varphi) < 2\epsilon_k.$$

For the third inclusion, note that for any $\chi \in B_{2\epsilon_k}(\psi_k)$,

$$d(\psi, \chi) \leq d(\psi, \psi_k) + d(\psi_k, \chi) < d(\psi, \psi_k) + 2\epsilon_k.$$

For the final inclusion, note that V_{j_0} is open, i.e. for $\psi \in U$ there is an $r > 0$ s.t. $B_r(\psi) \subset V_{j_0}$. Since $2\epsilon_k + d(\psi_k, \psi) = 2/k + d(\psi_k, \psi)$ becomes arbitrarily small, for sufficiently large k , the last inclusion holds.

However, $B_k \in V_{j_0}$ is a contradiction to $B_k \in \mathcal{B}$, cf. the construction of the B_k in Eq. (5.83). ■

Corollary 5.109. Compact sets in metric spaces are closed and bounded.

Proof. Let (X, d) be a metric space, and let $U \subset X$ be compact.

Let $\varphi \in \overline{U}$, i.e. φ is an accumulation point of U , and thus there exists a sequence $(\varphi_n)_{n \in \mathbb{N}}$ in U converging to φ wrt d . Thus, by Theorem 5.24, every subsequence of $(\varphi_n)_{n \in \mathbb{N}}$ also converges to φ in \overline{U} . Since U is sequentially compact, cf. Theorem 5.108, there is a subsequence $(\varphi_{n_k})_{k \in \mathbb{N}}$ of $(\varphi_n)_{n \in \mathbb{N}}$ converging to $\psi \in U \subset \overline{U}$, and since limits of convergent

sequences are unique, cf. Theorem 5.18, it follows that $\varphi = \psi$, i.e. $\varphi \in U$, which shows that U is closed.

To see that U is bounded, note that by Theorem 5.108, U is totally bounded. Hence, for any fixed $\epsilon > 0$, there exist points $\varphi_1, \dots, \varphi_{n_\epsilon} \in U$ with

$$U \subset \bigcup_{i=1}^{n_\epsilon} B_\epsilon(\varphi_i),$$

i.e. for $\varphi \in U$ we have $\varphi \in B_\epsilon(\varphi_j)$ for $1 \leq j \leq n$. Now define

$$r := \epsilon + \max_{1 \leq j \leq n} \{d(\varphi_1, \varphi_j)\},$$

then we have

$$d(\varphi, \varphi_1) \leq d(\varphi, \varphi_j) + d(\varphi_j, \varphi_1) < \epsilon + r - \epsilon = r.$$

Thus, $U \subset B_r[\varphi_1]$. ■

Remark 5.110. For topological spaces, this does not hold in general.

Definition 5.111 (Relatively compact). A subset U of a metric space (X, d) is called *relatively compact* in X if its closure $\overline{U} \subset X$ is compact.

Theorem 5.112. $U \subset X$ is sequentially compact iff U is relatively compact and closed.

Proof. “ \implies ” Since U is sequentially compact, it is compact, and thus closed, cf. Theorem 6.122. Since it is closed, we have that $U = \overline{U}$, and thus \overline{U} is compact.

“ \impliedby ” Since U is closed, we have that $U = \overline{U}$, and since U is relatively compact in X , $U = \overline{U}$ is compact, and hence also sequentially compact. ■

Theorem 5.113. A subset $U \subset X$ of a metric space (X, d) is relatively compact iff every sequence in U contains a convergent subsequence in X .

Proof. “ \implies ” Let U be relatively compact, i.e. \overline{U} is compact, and thus also sequentially compact. Since every sequence $(\varphi_n)_{n \in \mathbb{N}}$ in U also lies in \overline{U} , $U \subset \overline{U}$, $(\varphi_n)_{n \in \mathbb{N}}$ contains a convergent subsequence in $\overline{U} \subset X$.

“ \impliedby ” Let every sequence in U contain a convergent subsequence in X . We now show that \overline{U} is sequentially compact, which implies its compactness.

Let $(\varphi_n)_{n \in \mathbb{N}}$ be a sequence in \overline{U} . Since U is dense in \overline{U} , for every φ_n , $n \in \mathbb{N}$, there is a $\psi_n \in U$ s.t. $d(\varphi_n, \psi_n) < 1/n$. That way we can construct a sequence $(\psi_n)_{n \in \mathbb{N}}$ in U , and by assumption it contains a convergent subsequence $(\psi_{n_k})_{k \in \mathbb{N}}$ in X ; denote the limit by $\psi \in X$. Now note that

$$d(\varphi_{n_k}, \psi) \leq d(\varphi_{n_k}, \psi_{n_k}) + d(\psi_{n_k}, \psi) < \frac{2}{k} \xrightarrow{k \rightarrow \infty} 0,$$

i.e. every sequence $(\varphi_n)_{n \in \mathbb{N}}$ in \overline{U} contains a convergent subsequence $(\varphi_{n_k})_{k \in \mathbb{N}}$ in X . ■

Theorem 5.114. Relationship between totally bounded and relatively compact subsets [35, Theorem 4.27]:

- (a) Any relatively compact subset of a metric space is totally bounded.
- (b) Any totally bounded subset of a complete metric space is relatively compact.

Proof. (a): Let U be relatively compact, i.e. \overline{U} is compact, and thus by Theorem 5.108 also totally bounded. Since $U \subset \overline{U}$, U is also totally bounded.

(b): Let $U \subset X$ be totally bounded, where (X, d) is a complete metric space. This means that for any fixed $\epsilon > 0$, there exists an $n_\epsilon \in \mathbb{N}$ s.t.

$$U \subset \bigcup_{i=1}^{n_\epsilon} B_\epsilon(\varphi_i),$$

where $\varphi_1, \dots, \varphi_{n_\epsilon} \in U$. Since U lies dense in \overline{U} , for any $\psi \in \overline{U}$ there is a $\varphi \in U$ s.t. $d(\varphi, \psi) < \epsilon$. Since $\varphi \in U$, there is a $j \in \{1, \dots, n_\epsilon\}$ s.t. $\varphi \in B_\epsilon(\varphi_j)$. Then

$$d(\psi, \varphi_j) \leq d(\psi, \varphi) + d(\varphi, \varphi_j) < 2\epsilon,$$

i.e. \overline{U} can be covered as follows:

$$\overline{U} \subset \bigcup_{i=1}^{n_\epsilon} B_{2\epsilon}(\varphi_i),$$

implying that \overline{U} is totally bounded. Since \overline{U} is closed and by Theorem 5.73, closed subsets of complete metric spaces are also complete, by Theorem 5.108, \overline{U} is compact. ■

We now generalize Example 5.67 by considering functions that are defined on arbitrary compact metric spaces X .

Theorem 5.115. Let X be a compact metric space and denote by $\mathcal{C}[X]$ the space of all continuous functions $\chi : X \rightarrow \mathbb{K}$, where $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$. Then $(\mathcal{C}[X], d_\infty)$ with

$$d_\infty(\varphi, \psi) = \|\varphi - \psi\|_\infty := \sup_{x \in X} \{|\varphi(x) - \psi(x)|\} \quad \forall \varphi, \psi \in \mathcal{C}[X]$$

is a complete metric space.

Proof. Replace $[a, b]$ in the proof of Example 5.67 by X and \mathbb{R} by \mathbb{K} , and note that (\mathbb{C}, d) with $d(z_1, z_2) := |z_1 - z_2|$ is also a complete metric space, cf. Example 5.61. ■

We generalize Example 5.67 once more.

Theorem 5.116. Let X be a compact metric space and (Y, d) a complete metric space. Denote the space of all continuous functions $\chi : X \rightarrow Y$ by $\mathcal{C}[X, Y]$, then $(\mathcal{C}[X, Y], d_\infty)$ with

$$d_\infty(\varphi, \psi) := \sup_{x \in X} \{d(\varphi(x), \psi(x))\} \quad \forall \varphi, \psi \in \mathcal{C}[X, Y] \quad (5.85)$$

is a complete metric space.

Proof. Replace \mathbb{K} by Y in the proof of the preceding Theorem, $|\cdot - \cdot|$ by $d(\cdot, \cdot)$ and note that (Y, d) is complete by assumption. ■

Theorem 5.117 (Arzelà⁷-Ascoli⁸). Let (X, d) be a compact metric space. Then $U \subset \mathcal{C}[X]$ is relatively compact iff U is bounded in $(\mathcal{C}[X], d_\infty)$ and uniformly equicontinuous.

⁷CESARE ARZELÀ (1847 – 1912), Italian mathematician.

⁸GIULIO ASCOLI (1843 – 1896), Jewish-Italian mathematician.

Proof [36]. “ \Leftarrow ” Let $U \subset \mathcal{C}[X]$ be bounded and uniformly equicontinuous.

Since X is compact, by Theorems 5.104, 5.107 and 5.108, X is separable, i.e. there is a set $S \subset X$ that is dense and countable. We denote the (countably many) points of S by $(x_n)_{n \in \mathbb{N}}$.

Let $(\varphi_n)_{n \in \mathbb{N}}$ be a sequence in U . We now show that $(\varphi_n)_{n \in \mathbb{N}}$ contains a uniformly convergent subsequence, which we do via a standard diagonal argument. Since $U \subset \mathcal{C}[X]$ is bounded, there exists a $C > 0$ s.t. for all $f \in U$ and $x \in X$, $|f(x)| \leq C$. Thus, the sequence $(\varphi_n(x_1))_{n \in \mathbb{N}}$ (of real or complex values) is bounded. The Bolzano-Weierstrass Theorem 3.1 now implies that $(\varphi_n(x_1))_{n \in \mathbb{N}}$ has a convergent subsequence $(\varphi_{n_{1,l}}(x_1))_{l \in \mathbb{N}}$ in the real or complex numbers. Further, the sequence $(\varphi_{n_{1,l}}(x_2))_{l \in \mathbb{N}}$ is bounded and has a convergent subsequence, which we denote by $(\varphi_{n_{2,l}}(x_2))_{l \in \mathbb{N}}$. Note that the sequence of functions $(\varphi_{n_{2,l}})_{l \in \mathbb{N}}$ converges at both x_1 and x_2 , as $(\varphi_{n_{2,l}})_{l \in \mathbb{N}}$ is a subsequence of $(\varphi_{n_{1,l}})_{l \in \mathbb{N}}$. This way, through repeated selection, we inductively obtain a countable collection of subsequences $(\varphi_{n_{k,l}})_{l \in \mathbb{N}}$ of the original sequence $(\varphi_n)_{n \in \mathbb{N}}$

$$\begin{array}{cccc} \varphi_{n_{1,1}} & \varphi_{n_{1,2}} & \varphi_{n_{1,3}} & \cdots \\ \varphi_{n_{2,1}} & \varphi_{n_{2,2}} & \varphi_{n_{2,3}} & \cdots \\ \varphi_{n_{3,1}} & \varphi_{n_{3,2}} & \varphi_{n_{3,3}} & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{array}$$

where the sequence in the n -th row converges at x_1, \dots, x_n , and each row is a subsequence of the one above it.

For every l , the sequence $(\varphi_{n_{k,k}})_{k \geq l}$ is a subsequence of $(\varphi_{n_{l,k}})_{k \in \mathbb{N}}$. Thus, for any l , the sequence $(\varphi_{n_{k,k}}(x_l))_{k \in \mathbb{N}}$ converges, i.e. the sequence $(\varphi_{n_{k,k}})_{k \in \mathbb{N}}$ converges pointwisely on S .

We will now show that $(\varphi_{n_{k,k}})_{k \in \mathbb{N}}$ converges uniformly on X and that it is uniformly Cauchy. Let $\delta > 0$ (we choose the same δ as the δ for which U is uniformly equicontinuous). Since S is dense in X , i.e. for every $x \in X$ there exists an $x_k \in S$ s.t. $d(x, x_k) < \delta$, we have that the collection of open balls $\{B_\delta(x_k)\}_{k \in \mathbb{N}}$ forms an open cover of X , i.e.

$$X = \bigcup_{k \in \mathbb{N}} B_\delta(x_k).$$

Since X is compact, there exists a $j \in \mathbb{N}$ s.t.

$$X = \bigcup_{1 \leq k \leq j} B_\delta(x_k).$$

As $(\varphi_{n_{k,k}})_{k \in \mathbb{N}}$ converges pointwisely on S , for all $\epsilon > 0$ and all $x_k \in \{x_1, \dots, x_j\}$, there exists an $N_k = N_k(\epsilon)$ s.t. $|\varphi_{n_{l,l}}(x_k) - \varphi_{n_{m,m}}(x_k)| < \epsilon$ for all $n_{l,l}, n_{m,m} \geq N_k$, cf. Theorem 5.50. Setting $N := \max_{1 \leq k \leq j} \{N_k\}$, we have that for all $\epsilon > 0$ there exists an $N \in \mathbb{N}$ s.t. for all $x_k \in \{x_1, \dots, x_j\}$,

$$|\varphi_{n_{l,l}}(x_k) - \varphi_{n_{m,m}}(x_k)| < \epsilon \quad \forall n_{l,l}, n_{m,m} \geq N,$$

which implies that $(\varphi_{n_{k,k}})_{k \in \mathbb{N}}$ converges uniformly on S . Since S is dense, i.e. for all $x \in X$ there is an x_k with $1 \leq k \leq j$ s.t. $d(x, x_k) < \delta$, we have

$$\begin{aligned} & |\varphi_{n_{l,l}}(x) - \varphi_{n_{m,m}}(x)| \\ & \leq |\varphi_{n_{l,l}}(x) - \varphi_{n_{l,l}}(x_k)| + |\varphi_{n_{l,l}}(x_k) - \varphi_{n_{m,m}}(x_k)| + |\varphi_{n_{m,m}}(x_k) - \varphi_{n_{m,m}}(x)| \\ & < 3\epsilon \end{aligned}$$

for all $n_{l,l}, n_{m,m} \geq N$, where we used that U is uniformly equicontinuous twice. Our estimate shows that $(\varphi_{n_{k,k}})_{k \in \mathbb{N}}$ is uniformly Cauchy in $\mathcal{C}[X]$, and thus uniformly convergent on X . As \mathbb{R} and \mathbb{C} are complete, this implies that $(\varphi_{n_{k,k}})_{k \in \mathbb{N}}$ converges to a $\varphi \in \mathcal{C}[X]$, i.e. φ is an accumulation point of U . Thus, \overline{U} is sequentially compact, and by Theorem 5.108 also compact, i.e. U is relatively compact.

" \implies " Let $U \subset \mathcal{C}[X]$ be relatively compact, i.e. \overline{U} is compact and by Theorem 5.108 also totally bounded. Thus, for every $\epsilon > 0$, there exist $\varphi_1, \dots, \varphi_n \in \overline{U}$ s.t.

$$U \subset \overline{U} \subset \bigcup_{i=1}^n B_\epsilon(\varphi_i),$$

i.e. for every $\varphi \in U$ there exists a $\varphi_i \in \{\varphi_1, \dots, \varphi_n\}$ s.t. $d_\infty(\varphi, \varphi_i) < \epsilon$. The inclusion shows that U is bounded.

Since X is compact, the functions $\varphi_1, \dots, \varphi_n \in \overline{U} \subset \mathcal{C}[X]$ are uniformly continuous on X , i.e. for every $\epsilon > 0$ there is a $\delta_i = \delta_i(\epsilon)$ s.t. for all $x, y \in X$ with $d(x, y) < \delta_i$,

$$|\varphi_i(x) - \varphi_i(y)| < \epsilon.$$

Now define $\delta := \min_{1 \leq i \leq n} \{\delta_i\}$. Then for all $\epsilon > 0$, δ satisfies the condition that for all $\varphi \in U$ and all $x, y \in X$ with $d(x, y) < \delta$,

$$\begin{aligned} |\varphi(x) - \varphi(y)| &\leq |\varphi(x) - \varphi_i(x)| + |\varphi_i(x) - \varphi_i(y)| + |\varphi_i(y) - \varphi(y)| \\ &< d_\infty(\varphi, \varphi_i) + \epsilon + d_\infty(\varphi_i, \varphi) \\ &< 3\epsilon, \end{aligned}$$

where we chose φ_i s.t. $d_\infty(\varphi, \varphi_i) < \epsilon$. This shows that U is also uniformly equicontinuous. ■

Sometimes, the following version is called the Arzelà-Ascoli Theorem in the literature.

Corollary 5.118. Let (X, d) be a compact metric space. Then $U \subset \mathcal{C}[X]$ is compact iff U is bounded in $(\mathcal{C}[X], d_\infty)$, uniformly equicontinuous and closed.

Proof. The statement directly follows from Theorem 5.112. ■

An application of the Arzelà-Ascoli Theorem is given in Example 7.7.

5.3 Baire Category Theorem

In Theorem 5.82, we have already established that the finite intersection of open dense sets yields an open dense set again. While in the case of infinite intersections, it is inevitable that we lose the openness property of the resulting intersection, in case of countable intersections, it turns out that we still have a dense set, as long as we assume that the underlying metric space is complete.

Theorem 5.119 (Baire Category Theorem). Let

There are many applications of the Baire Category Theorem, cf. Ref. [37] for some. One application is to show that the irrationals $\mathbb{R} \setminus \mathbb{Q}$ are dense. In Example 5.77, another proof of this statement can be found.

Example 5.120. Let $X = \mathbb{R}$ with the standard metric. According to Example 5.13, for any $q \in \mathbb{Q}$, the singleton $\{q\}$ is closed in \mathbb{R} , i.e. $\mathbb{R} \setminus \{q\}$ is open. $\mathbb{R} \setminus \{q\}$ is also dense. Thus, by Eq. (8.13), $\bigcap_{q \in \mathbb{Q}} \mathbb{R} \setminus \{q\} = \mathbb{R} \setminus \bigcup_{q \in \mathbb{Q}} \{q\} = \mathbb{R} \setminus \mathbb{Q}$. By the Baire Category Theorem, we know that $\mathbb{R} \setminus \mathbb{Q}$ is dense.

For uncountable intersections of open and dense sets of a complete metric space, the resulting set is not dense, as the following example shows [38].

Example 5.121. Let $X = \mathbb{R}$ with the standard metric. For any $x \in \mathbb{R}$, the singleton $\{x\}$ is closed, i.e. $\mathbb{R} \setminus \{x\}$ is open. $\mathbb{R} \setminus \{x\}$ is also dense. However, by Eq. (8.13), $\bigcap_{x \in \mathbb{R}} \mathbb{R} \setminus \{x\} = \mathbb{R} \setminus \bigcup_{x \in \mathbb{R}} \{x\} = \emptyset$, which is not dense in \mathbb{R} .

Note that in the formulation of the Baire Category Theorem, it is vital that the underlying metric space is complete. Otherwise, the Baire Category Theorem may not hold, as the following example shows [39].

Example 5.122. Let $X = \mathbb{Q}$ with its metric inherited from \mathbb{R} . Then for any $q \in \mathbb{Q}$, $\mathbb{Q} \setminus \{q\}$ is open and dense, since $\{q\}$ is closed and by Theorem 5.41, $\overline{\mathbb{Q} \setminus \{q\}} = \mathbb{Q} \setminus \text{int}(\{q\}) = \mathbb{Q} \setminus \emptyset = \mathbb{Q}$. However,

$$\bigcap_{q \in \mathbb{Q}} \mathbb{Q} \setminus \{q\} = \mathbb{Q} \setminus \bigcup_{q \in \mathbb{Q}} \{q\} = \emptyset,$$

which is clearly not dense in \mathbb{Q} .

5.4 Continuity

Definition 5.123 (ϵ - δ definition of continuity [40]). Let (X, d) and (Y, d') be metric spaces. A function $f : X \rightarrow Y$ is called *continuous* at $p \in X$ if

$$\forall \epsilon > 0 \exists \delta(\epsilon, p) > 0 \forall z \in X : d(z, p) < \delta \Rightarrow d'(f(z), f(p)) < \epsilon. \quad (5.86)$$

f is called continuous on X if it is continuous for every $p \in X$, i.e.

$$\forall \epsilon > 0 \forall p \in X \exists \delta(\epsilon, p) > 0 \forall z \in X : d(z, p) < \delta \Rightarrow d'(f(z), f(p)) < \epsilon. \quad (5.87)$$

Remark 5.124. The notion of continuity depends on the metrics. For example [41], let $X = \mathbb{R}$, and let d be the usual metric on \mathbb{R} , and d' the discrete metric. Then the function $\text{id} : (\mathbb{R}, d') \rightarrow (\mathbb{R}, d), x \mapsto x$ is continuous for all $p \in \mathbb{R}$, since for $\delta = 1$, if we have $d'(z, p) < \delta = 1$ for all $z \in \mathbb{R}$, then $z = p$, and thus $d(z, p) = 0 < \epsilon$ for any $\epsilon > 0$.

Conversely, $\text{id} : (\mathbb{R}, d) \rightarrow (\mathbb{R}, d'), x \mapsto x$ is not continuous on \mathbb{R} . To see this, let $\epsilon = 1$, consider arbitrary $\delta > 0$ and define $p := \min\{1, \delta\} \in \mathbb{R}$ and $z := p/2 \neq p$. Then $d(z, p) = |p|/2 < \delta$, but $d'(z, p) = 1 \geq \epsilon$.

Definition 5.125 (Uniform continuity [42]). Let (X, d) and (Y, d') be metric spaces. A function $f : X \rightarrow Y$ is called *uniformly continuous* on X

$$\forall \epsilon > 0 \exists \delta(\epsilon) > 0 \forall p, z \in X : d(z, p) < \delta \Rightarrow d'(f(z), f(p)) < \epsilon. \quad (5.88)$$

Remark 5.126. While uniform continuity implies continuity, the opposite is in general not true.

Example 5.127. Consider the metric space $(\mathbb{R}_{>0}, d)$, where $d(x, y) := |x - y|$ for any $x, y \in \mathbb{R}_{>0}$. The function $f : \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$ is continuous on $\mathbb{R}_{>0}$, but not uniformly continuous on $\mathbb{R}_{>0}$.

Proof. To see that f is not uniformly continuous on $\mathbb{R}_{>0}$, choose $\epsilon = 1$, then for all $\delta > 0$ take $x := \min\{\delta, 1\} \in \mathbb{R}_{>0}$ and $y := x/2 \in \mathbb{R}_{>0}$, then $|x - y| = \frac{x}{2} < \delta$, but

$$|f(x) - f(y)| = \left| \frac{1}{x} - \frac{1}{y} \right| = \frac{1}{x} \geq 1 = \epsilon.$$

■

Definition 5.128. (Uniform equicontinuity [43]) Let (X, d) and (Y, d') be metric spaces and \mathcal{F} a family of functions from X to Y . Then we say that \mathcal{F} is *uniformly equicontinuous* if

$$\forall \epsilon > 0 \exists \delta(\epsilon) > 0 \forall f \in \mathcal{F} \forall p, z \in X : d(z, p) < \delta \Rightarrow d'(f(z), f(p)) < \epsilon. \quad (5.89)$$

Remark 5.129. If \mathcal{F} is uniformly equicontinuous, then clearly, every $f \in \mathcal{F}$ is uniformly continuous on X . However, conversely, if \mathcal{F} is a family of uniformly continuous functions on X , then \mathcal{F} is not necessarily uniformly equicontinuous [44]. To see this, consider the family of functions

$$\mathcal{F} := \left\{ f_n : [0, 1] \rightarrow \mathbb{R}, x \mapsto \frac{x^2}{x^2 + (1 - nx)^2} \mid n \in \mathbb{N} \right\}.$$

Clearly, each $f_n \in \mathcal{F}$ is uniformly continuous, since the domain $[0, 1]$ is compact and f_n is continuous on $[0, 1]$, as it is the product of two continuous functions on $[0, 1]$ and the denominator is bigger than 0 for all $n \in \mathbb{N}$ and $x \in [0, 1]$. However, \mathcal{F} is not uniformly equicontinuous. Let $\epsilon = 1$, then for all $\delta > 0$, take $p = 0$ and $z = 1/n$. There exists an $n \in \mathbb{N}$ s.t. $|p - z| = \frac{1}{n} < \delta$, yet

$$|f_n(z) - f_n(p)| = |1 - 0| = 1 \geq \epsilon.$$

Theorem 5.130. Let (X, d) and (Y, d') be metric spaces. A function $f : X \rightarrow Y$ is continuous at $p \in X$ iff for every sequence $(p_n)_{n \in \mathbb{N}}$ in X that converges to p wrt d , the sequence $(f(p_n))_{n \in \mathbb{N}}$ converges to $f(p)$ wrt d' .

Proof. “ \implies ” Let f be continuous at $p \in X$, and $(p_n)_{n \in \mathbb{N}}$ a sequence converging to p . Then for all $\epsilon > 0$ there exists a $\delta(\epsilon, p)$ s.t. for all $z \in X$, $d(z, p) < \delta$ implies $d'(f(z), f(p)) < \epsilon$. Since $(p_n)_{n \in \mathbb{N}}$ converges to p wrt d , for the same δ , there exists an $N(\delta)$ s.t. for all $n \geq N$, $d(p_n, p) < \delta$, and thus $d'(f(p_n), f(p)) < \epsilon$.

“ \impliedby ” Let $(p_n)_{n \in \mathbb{N}}$ be a sequence in X converging to $p \in X$ wrt d , and $(f(p_n))_{n \in \mathbb{N}}$ converging to $f(p)$ wrt d' . Assume that f is not continuous at p , i.e. there exists an $\epsilon > 0$ s.t. for all $\delta > 0$ there exists a $z \in X$ s.t. $d(z, p) < \delta$, but $d'(f(z), f(p)) \geq \epsilon$. Fix this $\epsilon > 0$, and consider an arbitrary $\delta > 0$. Since $(p_n)_{n \in \mathbb{N}}$ converges to p , there exists an $N(\delta) \in \mathbb{N}$ s.t. for all $n \geq N$, $d(p_n, p) < \delta$, but $d'(f(p_n), f(p)) \geq \epsilon$. But this is a contradiction to $(f(p_n))_{n \in \mathbb{N}}$ converging to $f(p)$ wrt d' . ■

Lemma 5.131. Consider the map $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$. Then f is continuous if all the $f_i : \mathbb{R}^m \rightarrow \mathbb{R}$, $i \in \{1, \dots, n\}$ are continuous.

Proof. Since all metrics are strongly equivalent, it does not matter which metric we equip \mathbb{R} , \mathbb{R}^m and \mathbb{R}^n with, cf. Remark 4.10; for the following, consider the Euclidean metric. Suppose each f_i is continuous for all $i \in \{1, \dots, n\}$, i.e.:

$$\forall p \in \mathbb{R}^m : \forall \epsilon_i > 0 \exists \delta_i > 0 : d_{\mathbb{R}}(f_i(p), f_i(z)) < \frac{\epsilon_i}{\sqrt{n}} \quad \forall z \in D \quad \text{with } d_{\mathbb{R}^m}(p, z) < \delta_i. \quad (5.90)$$

Now define $\epsilon := \max\{\epsilon_1, \dots, \epsilon_n\}$. Thus:

$$d_{\mathbb{R}^n}(f(p), f(z)) = \sqrt{\sum_{i=1}^n (f_i(p) - f_i(z))^2} = \sqrt{\sum_{i=1}^n d_{\mathbb{R}}^2(f_i(p), f_i(z))} < \sqrt{\sum_{i=1}^n \left(\frac{\epsilon}{\sqrt{n}}\right)^2} = \epsilon. \quad (5.91)$$

■

Theorem 5.132. For a map $f : X \rightarrow Y$ between two metric spaces (X, d_X) and (Y, d_Y) the following statements are equivalent:

- (i) f is continuous,
- (ii) preimages $f^{-1}(V) := \{x \in X \mid f(x) \in V\}$ of all open sets $V \subset Y$ are open,
- (iii) preimages $f^{-1}(A)$ of all closed sets $A \subset Y$ are closed.

Proof. (i) \Rightarrow (ii) Assume that f is continuous and that $V \subset Y$ is open and let $a \in f^{-1}(V)$. Since V is an open set, $\exists \epsilon > 0$ s.t. $B_{\epsilon}(f(a)) \subset V$, cf. Def. 5.11. By assumption, f is continuous at $a \in f^{-1}(V) \subset X$ and therefore $\forall \epsilon > 0 \exists \delta > 0$ s.t.

$$d_X(x, a) < \delta \Rightarrow d_Y(f(x), f(a)) < \epsilon \quad \forall x \in X.$$

Put differently, $x \in B_{\delta}(a)$ implies $f(x) \in B_{\epsilon}(f(a)) \subset V$. Thus, $B_{\delta}(a) \subset f^{-1}(V) \Rightarrow f^{-1}(V) \subset X$ is an open set, cf. Def. 5.11.

(ii) \Rightarrow (i) Assume that $f^{-1}(V) \subset X$ is open for $V \subset Y$ open and let $a \in X$, $\epsilon > 0$. From Theorem 5.14 we know that $B_{\epsilon}(f(a)) \subset Y$ is open. Thus, by assumption, $f^{-1}(B_{\epsilon}(f(a))) = \{x \in X \mid f(x) \in B_{\epsilon}(f(a))\} = \{x \in X \mid f(x) \in \{y \in Y \mid d_Y(y, f(a)) < \epsilon\}\}$ is open as well. Clearly, $a \in f^{-1}(B_{\epsilon}(f(a)))$. Therefore, it follows from Def. 5.11 that $\exists \delta > 0$ s.t. $B_{\delta}(a) \subset f^{-1}(B_{\epsilon}(f(a)))$. Thus, $\forall x \in B_{\delta}(a) : d_X(x, a) < \delta \Rightarrow d_Y(f(x), f(a)) < \epsilon$. This proves that $f : X \rightarrow Y$ is continuous in every point $a \in X$. [45]

(ii) \Rightarrow (iii) Assume that the preimages $f^{-1}(V)$ of all open sets $V \subset Y$ are open. Since $f^{-1}(Y \setminus V) = f^{-1}(Y) \setminus f^{-1}(V)$, as can be easily shown by using the definition of the complement of a set, we have for all open sets $V \subset Y$:

$$f^{-1}(Y \setminus V) = f^{-1}(Y) \setminus f^{-1}(V) = X \setminus f^{-1}(V). \quad (5.92)$$

Since $f^{-1}(V)$ is open by assumption, $f^{-1}(Y \setminus V) = X \setminus f^{-1}(V)$ is closed.

(iii) \Rightarrow (ii) Assume that the preimages $f^{-1}(A)$ of all closed sets $A \subset Y$ are closed. Then:

$$f^{-1}(Y \setminus A) = f^{-1}(Y) \setminus f^{-1}(A) = X \setminus f^{-1}(A). \quad (5.93)$$

Since $f^{-1}(A)$ is closed by assumption, $f^{-1}(Y \setminus A) = X \setminus f^{-1}(A)$ is open. [46]

■

Theorem 5.133. Let (X, d) be a compact metric space and (Y, ρ) a metric space. A continuous function $f : X \rightarrow Y$ is uniformly continuous.

Proof [47]. Since f is continuous, for all $\epsilon > 0$ and all $x \in X$, there exists a $\delta_x > 0$ s.t. $\{f(y) \mid y \in X, d(x, y) < \delta_x\} = f(B_{\delta_x}^d(x)) \subset B_{\epsilon/2}^\rho(f(x)) = \{z \in Y \mid \rho(z, f(x)) < \epsilon/2\}$. Note that the family $\{B_{\delta_{x_i/2}}^d(x_i)\}_{x_i \in X}$ is an open cover of X , and since X is compact, there exists a finite subcover of X , which we denote by $\{B_{\delta_{x_i/2}}^d(x_i)\}_{i=1}^n$, where $n \in \mathbb{N}$. Now define $\delta := \min_{1 \leq i \leq n} \delta_{x_i}/2$.

Now suppose that for arbitrary $x, y \in X$, $d(x, y) < \delta$. Note that there is a $j \in \{1, \dots, n\}$ s.t. $x \in B_{\delta_{x_j/2}}^d(x_j)$. Thus,

$$d(y, x_j) \leq d(y, x) + d(x, x_j) < \delta + \frac{\delta_{x_j}}{2} \leq \delta_{x_j},$$

i.e. $y \in B_{\delta_{x_j}}^d(x_j)$, which implies that $f(y) \in f(B_{\delta_{x_j}}^d(x_j)) \subset B_{\epsilon/2}^\rho(f(x_j))$. Also, $x \in B_{\delta_{x_j/2}}^d(x_j)$ implies that $f(x) \in f(B_{\delta_{x_j/2}}^d(x_j)) \subset B_{\epsilon/2}^\rho(f(x_j))$. Thus,

$$\rho(f(x), f(y)) \leq \rho(f(x), f(x_j)) + \rho(f(x_j), f(y)) < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

Since δ only depends on ϵ , this shows that f is uniformly continuous on X . ■

Remark 5.134. Note that for the preceding theorem, it is vital that X is compact.

Example 5.135. Take the function $f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto x^2$, which is not uniformly continuous, where on \mathbb{R} , we consider the usual metric.

Proof [48]. Choose $\epsilon = 1$, and let $\delta > 0$ be arbitrary. Choose $x := \frac{1}{\delta} \in \mathbb{R}$ and let $y = x + \delta/2$; clearly, $d(x, y) < \delta$. However,

$$d(f(x), f(y)) = \left| x^2 - \left(x + \frac{\delta}{2} \right)^2 \right| = \left| x\delta + \frac{\delta^2}{4} \right| = 1 + \frac{\delta^2}{4} \geq 1 = \epsilon.$$

■

6 Topological Spaces

Definition 6.1 (Topological Space [20]). A topology on a set X is a collection τ of subsets of X such that

- the intersection of two members of τ is in τ ,
- the union of any collection of members of τ is in τ ,
- $\emptyset, X \in \tau$.

A set X endowed with a topological structure τ on it is called a *topological space*. The elements of X are called *points*, and the members of τ are called the *open sets*. We also say that members of τ are *open in X* .

Remark 6.2. Instead of defining a topology via open sets, it is alternatively possible to use closed sets. Then the above axioms become:

- the union of two closed sets in τ is closed,
- the intersection of any collection of closed sets in τ is closed,
- $\emptyset, X \in \tau$.

Proof. Let I be an index set and $\{U_i\}_{i \in I}$ a family of open subsets of X , i.e. $\{X \setminus U_i\}_{i \in I}$ is a family of closed subsets of X . Then

$$\bigcap_{i \in I} X \setminus U_i \stackrel{(8.13)}{=} X \setminus \bigcup_{i \in I} U_i,$$

is closed, as $\bigcup_{i \in I} U_i$ is open.

Let U_1, U_2 be open sets, then

$$\bigcup_{i \in \{1,2\}} X \setminus U_i \stackrel{(8.14)}{=} X \setminus \bigcap_{i \in \{1,2\}} U_i$$

is closed, as $\bigcap_{i \in \{1,2\}} U_i$ is open.

Since \emptyset and X are open, $X \setminus \emptyset = X$ and $X \setminus X = \emptyset$ are closed. ■

Example 6.3 (Trivial topology). For every set X , the trivial topology is defined as $\tau := \{\emptyset, X\}$.

Example 6.4 (Discrete topology). For every set X , the discrete topology is defined as $\tau := \mathcal{P}(X)$, i.e. every subset of X is an open set.

Remark 6.5. Consider the discrete topological space $(X, \mathcal{P}(X))$, and note that for every member $U \in \mathcal{P}(X)$, $X \setminus U \subset X \in \mathcal{P}(X)$, and thus X is both an open and a closed set.

Example 6.6 (Co-finite topology). For every set X , the collection of all subsets of X whose complements are finite forms a topology τ on X :

$$\tau := \{U \subset X \mid U = \emptyset \vee X \setminus U \text{ finite}\}$$

Proof. Clearly, $\emptyset \in \tau$. Also, $X \in \tau$, since $X \setminus X = \emptyset$. Let $U_1, U_2 \in \tau$, then if $U_1 = \emptyset$ or $U_2 = \emptyset$, we have $U_1 \cap U_2 = \emptyset \in \tau$. Otherwise, assume that $U_1 \neq \emptyset$ and $U_2 \neq \emptyset$, then by de Morgan's law, cf. Theorem 8.23, $X \setminus (U_1 \cap U_2) = (X \setminus U_1) \cup (X \setminus U_2)$, and since the union of two finite sets is finite again, $U_1 \cap U_2 \in \tau$ follows. Now let $\{U_i\}_{i \in I} \subset \tau$, where I is an arbitrary index set. Then $X \setminus \bigcup_{i \in I} U_i = \bigcap_{i \in I} X \setminus U_i \in \tau$, cf. Theorem 8.23, since the arbitrary union of countable sets is finite again. Thus, the arbitrary union of members of τ is a member of τ again. ■

Remark 6.7. Note that if X is already finite, then $X \setminus U$ for any $U \subset X$ is always finite, and thus the co-finite topology is the discrete topology in that case.

Example 6.8 (Co-countable topology). For every set X , the collection of all subsets of X whose complements are countable forms a topology τ on X :

$$\tau := \{U \subset X \mid U = \emptyset \vee X \setminus U \text{ countable}\}$$

Proof. Clearly, $\emptyset \in \tau$. Also, $X \in \tau$, since $X \setminus X = \emptyset$ is countable. Let $U_1, U_2 \in \tau$, then if $U_1 = \emptyset$ or $U_2 = \emptyset$, we have $U_1 \cap U_2 = \emptyset \in \tau$. Otherwise, assume that $U_1 \neq \emptyset$ and $U_2 \neq \emptyset$, then by de Morgan's law, cf. Theorem 8.23, $X \setminus (U_1 \cap U_2) = (X \setminus U_1) \cup (X \setminus U_2)$, and since the union of two countable sets is countable again, $U_1 \cap U_2 \in \tau$ follows. Now let $\{U_i\}_{i \in I} \subset \tau$, where I is an arbitrary index set. Then $X \setminus \bigcup_{i \in I} U_i = \bigcap_{i \in I} X \setminus U_i \in \tau$, cf. Theorem 8.23, since the arbitrary union of countable sets is countable again. Thus, the arbitrary union of members of τ is a member of τ again. ■

Remark 6.9. Note that if X is already finite, then $X \setminus U$ for any $U \subset X$ is always countable, and thus the co-countable topology is the discrete topology in that case.

Definition 6.10 (Quotient topology). Let X be a topological space, and let \sim be an equivalence relation on X . Denote the set of all equivalence classes by X/\sim . We denote the canonical surjection, cf. Definition 8.3, by $\pi : X \rightarrow X/\sim$, which maps each $x \in X$ to its equivalence class $[x]$. The set $\{U \subset X/\sim \mid \pi^{-1}(U) \text{ is open in } X\}$ is called the *quotient topology* on X/\sim .

Remark 6.11. The quotient topology is indeed a topology.

Proof. Clearly, $X = \pi^{-1}(X/\sim)$, and X is open, thus X/\sim is a member of the quotient topology. Also, $\emptyset = \pi^{-1}(\emptyset)$, and \emptyset is open in X , thus \emptyset is a member of the quotient topology. With Eq. (8.9) and (8.10), one can easily show that the finite intersection and arbitrary union of members of the quotient topology is in the quotient topology again. Thus, the quotient topology forms a topology on X/\sim . ■

Definition 6.12 (Open map). Let (X, τ_X) and (Y, τ_Y) be topological spaces, then $f : X \rightarrow Y$ is called an *open map* if f maps open sets in τ_X to open sets in τ_Y .

Definition 6.13 (Neighborhood [20]). Let (X, τ) be a topological space and $x \in X$, then a set $N \subset X$ is called a *neighborhood* of x in X if there is an open set $U \in \tau$ with $x \in U \subset N$. If N is open, then we say that N is an *open neighborhood* of x in X .

Theorem 6.14. $U \subset X$ is open iff U is a neighborhood of each of its points.

Proof. “ \Rightarrow ” By definition.

“ \Leftarrow ” If U is a neighborhood of each of its points, then there is an open set U_x in X s.t. $x \in U_x \subset U$. Note that [49]

$$U = \bigcup_{x \in U} \{x\} \subset \bigcup_{U_x \subset U} U_x \subset U,$$

which shows that $U = \bigcup_{U_x \subset U} U_x$, completing the proof. \blacksquare

Definition 6.15 (Neighborhood basis [20]). Let X be a topological space and $x \in X$. A *neighborhood basis* (also called *local basis*) at x is a collection \mathcal{B}_x of neighborhoods of x s.t. each neighborhood of x contains some member of \mathcal{B}_x .

Example 6.16. The family of all neighborhoods of a point is a neighborhood basis at that point.

Example 6.17. The family of all open neighborhoods of a point is a neighborhood basis at that point.

Example 6.18. If X is a topological space, where the topology is the discrete topology, then a neighborhood basis to $x \in X$ is $\mathcal{B}_x = \{\{x\}\}$.

Example 6.19. If X is a metric space, then the open r -balls at x for all $r > 0$ form a neighborhood basis at x .

Example 6.20. If X is a metric space, then $\mathcal{B}_x = \{B_{1/n}(x) \mid n \in \mathbb{N}\}$ forms a neighborhood basis at x .

Definition 6.21 ([20]). A topological space is *first-countable* if it has a countable neighborhood basis at each point.

Example 6.22. Every metric space is first-countable.

Proof. For any point x , the local basis at x given by the collection $\mathcal{B}_x = \{B_{1/n}^d(x) \mid n \in \mathbb{N}\}$ is countable. \blacksquare

Definition 6.23. Let τ and τ' be topologies on X . Then we call τ' *finer* than τ if $\tau \subset \tau'$, and we say that τ is *coarser* than τ' . In either case, we say that τ and τ' are *comparable*, otherwise we say they are non-comparable.

Remark 6.24. The trivial topology on X is the coarsest topology, and the discrete topology the finest on X . Thus, for a finite set X , the number of topologies τ on X that we can define is finite, since $\mathcal{P}(X)$ is the “biggest” topology on X that we can define, and $|\mathcal{P}(X)| = 2^{|X|}$.

Remark 6.25. An example of non-comparable topologies can be easily constructed: Let $X = \{a, b\}$, and consider $\tau_1 = \{\emptyset, \{a\}, X\}$ and $\tau_2 = \{\emptyset, \{b\}, X\}$, then these topologies are non-comparable.

Given a topological space, let us now look at constructing new topological spaces.

Theorem 6.26 (Subspace topology). Let (X, τ) be a topological space, then for any $S \subset X$ the set $\tau_S := \{S \cap U \mid U \in \tau\} \subset \mathcal{P}(S)$ defines a topology on S , which we call the *subspace topology*.

Proof. Let $S_1, S_2 \in \tau_S$, i.e. there exist $U_1, U_2 \in \tau$ s.t. $S_1 = S \cap U_1$ and $S_2 = S \cap U_2$. Then for the intersection, we have

$$S_1 \cap S_2 = S \cap U_1 \cap S \cap U_2 = S \cap (U_1 \cap U_2) \in \tau_S,$$

since $U_1 \cap U_2 \in \tau$.

Let I be an index set, and $\{S_i\}_{i \in I} \subset \tau_S$, where for each $S_j \in \{S_i\}_{i \in I}$, there is a $U_j \in \tau$, then by Eq. (8.4) we have

$$\bigcup_{i \in I} S_i = \bigcup_{i \in I} (S \cap U_i) = S \cap \left(\bigcup_{i \in I} U_i \right) \in \tau_S,$$

since the arbitrary union of members of τ is in τ again.

Finally, note that clearly, $\emptyset, S \in \tau_S$. ■

Theorem 6.27. Let X be a first-countable topological space, and consider $S \subset X$ with the subspace topology. Then S is also first-countable.

Proof [50]. Let $p \in S$, and let $(V_n)_{n \in \mathbb{N}}$ be a collection of neighborhoods of p in X s.t. each neighborhood of p contains some V_n , $n \in \mathbb{N}$.

Note that $N \subset X$ is a neighborhood of p if there is an open U in X s.t. $p \in U \subset N$. This implies that $p \in U \cap S \subset N \cap S$, i.e. $N \cap S$ is a neighborhood of p in S . Thus, $(V_n \cap S)_{n \in \mathbb{N}}$ is a collection of neighborhoods of p in S s.t. each neighborhood of p in S contains some $V_n \cap S$, $n \in \mathbb{N}$. ■

Theorem 6.28. Let I be a (finite, countable or uncountable) index set, then if (X, τ_i) , $i \in I$, is a topological space, so is the intersection of the topologies, i.e. $(X, \bigcap_{i \in I} \tau_i)$ is a topological space.

Proof. Let $U, V \in \bigcap_{i \in I} \tau_i \subset \mathcal{P}(X)$, i.e. for any $i \in I$, $U \in \tau_i$ and $V \in \tau_i$, and since τ_i is a topology, $U \cap V \in \tau_i$ for all i , implying $U \cap V \in \bigcap_{i \in I} \tau_i$.

Now let J be another index set, and for any $j \in J$, let $U_j \in \bigcap_{i \in I} \tau_i$, i.e. for any $i \in I$, $U_j \in \tau_i$, and since τ_i is a topology, $\bigcup_{j \in J} U_j \in \tau_i$ for all i , implying $\bigcup_{j \in J} U_j \in \bigcap_{i \in I} \tau_i$.

Finally, note that $\emptyset, X \in \tau_i$ for all $i \in I$, and thus $\emptyset, X \in \bigcap_{i \in I} \tau_i$. ■

Corollary 6.29. Let X be a set, $\mathcal{S} \subset \mathcal{P}(X)$ and I an index set. Then if (X, τ_i) are topologic spaces, $i \in I$, containing \mathcal{S} , i.e. $\mathcal{S} \subset \tau_i$ for each $i \in I$, their intersection $\bigcap_{i \in I} \tau_i$ is a topology on X , denoted by $\tau(\mathcal{S})$, which is the smallest topology on X containing \mathcal{S} . Note that this holds even for $\mathcal{S} = \emptyset$, since \emptyset is the subset of every set. Note that the intersection is guaranteed to be non-empty, since the discrete topology contains any \mathcal{S} .

This corollary motivates the following definition:

Definition 6.30 (Subbasis [20]). Let (X, τ) be a topological space, then a *subbasis* for τ is a collection of subsets $\mathcal{S} \subset \mathcal{P}(X)$ s.t. $\tau = \tau(\mathcal{S})$, and we call $\tau(\mathcal{S})$ the topology generated by \mathcal{S} .

We now seek to characterize $\tau(\mathcal{S})$.

Theorem 6.31. Let X be a set, $\mathcal{S} \subset \mathcal{P}(X)$, then the collection τ of all unions of finite intersections of members of \mathcal{S} is a topology on X .

Proof. Since the union of an empty collection of sets is \emptyset , and the intersection of an empty collection of sets is X [51], we have $\emptyset, X \in \tau$.

Let I be an index set, and consider the collection $\{A_i\}_{i \in I}$, where $A_i \in \tau$ for each $i \in I$. Since each A_i is the union of a finite intersection of members of \mathcal{S} , there exists an index set J_i s.t. $A_i = \bigcup_{j \in J_i} S_j$, where S_j is the finite intersection of members of \mathcal{S} for each $j \in J_i$. Thus, by Theorem 8.16,

$$\bigcup_{i \in I} A_i = \bigcup_{i \in I} \left(\bigcup_{j \in J_i} S_j \right) = \bigcup_{j \in \bigcup_{i \in I} J_i} S_j \in \tau.$$

Now let $I = \{1, \dots, n\}$ for some $n \in \mathbb{N}$, i.e. I is a finite index set, and consider the collection $\{A_i\}_{i \in I}$, where $A_i \in \tau$ for each $i \in I$. Now note that by Theorem 8.17,

$$\bigcap_{i \in I} A_i = \bigcap_{i \in I} \left(\bigcup_{j \in J_i} S_j \right) = \bigcup_{f \in \prod_{i \in I} J_i} \left(\bigcap_{i \in I} S_{f_i} \right) \in \tau, \quad (6.1)$$

where $f \in \prod_{i \in I} J_i$, i.e. $f = (f_1, \dots, f_n)$ with $f_i \in J_i$ for each $i \in I$. ■

Remark 6.32. Let X be a set, $\mathcal{S} \subset \mathcal{P}(X)$, then $\tau(\mathcal{S})$ is the topology τ from Theorem 6.31, i.e. it consists of all unions of finite intersections of members of \mathcal{S} . To see this, note that clearly, τ contains all members of \mathcal{S} (since each member is a finite intersection consisting of itself), and thus $\tau(\mathcal{S}) \subset \tau$. And since $\tau(\mathcal{S})$ is a topology on X , it also contains all unions of finite intersections of members of \mathcal{S} , i.e. $\tau \subset \tau(\mathcal{S})$, and thus $\tau = \tau(\mathcal{S})$, i.e. $\tau(\mathcal{S})$ is completely determined by \mathcal{S} .

Example 6.33. Let $X = \{a, b, c, d\}$ and $\mathcal{S} = \{\{a, b, c\}, \{b, c, d\}\}$, then the topology generated by \mathcal{S} is $\tau(\mathcal{S}) = \{\emptyset, \{b, c\}, \{a, b, c\}, \{b, c, d\}, X\}$.

Example 6.34. Let X be a set, and $\tau \subset \mathcal{P}(X)$ a topology on X . Then τ is a subbasis for itself.

Definition 6.35 (Order topology [20]). Let (X, \prec) be an ordered set, and for each pair of elements $a, b \in X$, define

$$\begin{aligned} (a, b) &:= \{x \in X \mid a \prec x \prec b\} && \text{open interval,} \\ [a, b] &:= \{x \in X \mid a \preceq x \preceq b\} && \text{closed interval,} \\ [a, b) &:= \{x \in X \mid a \preceq x \prec b\} && \text{right half-open or left half-closed interval,} \\ (a, b] &:= \{x \in X \mid a \prec x \preceq b\} && \text{left half-open or right half-closed interval.} \end{aligned}$$

For each $a \in X$, we define

$$\begin{aligned} (-\infty, a) &:= \{x \in X \mid x \prec a\} && \text{open ray or one-sided open interval,} \\ (a, \infty) &:= \{x \in X \mid a \prec x\} && \text{open ray or one-sided open interval,} \\ (-\infty, a] &:= \{x \in X \mid x \preceq a\} && \text{closed ray or one-sided closed interval,} \\ [a, \infty) &:= \{x \in X \mid a \preceq x\} && \text{closed ray or one-sided closed interval.} \end{aligned}$$

The *order topology* on X is the topology generated by the subbasis consisting of all open rays $(-\infty, a)$ and $(a, +\infty)$, where a ranges over X .

Remark 6.36. Notice that if $a_0 \in X$ is the smallest element, then $(-\infty, a) = [a_0, a)$, and $(a, +\infty) = (a, b_0]$ if $b_0 \in X$ is the largest element.

Remark 6.37. Note that for all $a, b \in X$, $(a, b) = (a, \infty) \cap (-\infty, b)$, i.e. every open interval is the finite intersection of open rays. Also, any closed interval $[a, b]$ is the complement of two open rays, namely $[a, b] = X \setminus ((-\infty, a) \cap (b, \infty))$.

Remark 6.38. Theorem 6.28 motivates the following question: Is the union of topologies a topology again? The answer is: The union of topologies does not necessarily yield a topology again.

Example 6.39. Consider $X = \{1, 2, 3\}$, and $\tau_1 = \{\emptyset, \{1\}, \{2\}, \{1, 2\}, X\}$ and $\tau_2 = \{\emptyset, \{3\}, X\}$, then both (X, τ_1) and (X, τ_2) are topological spaces, but $(X, \tau_1 \cup \tau_2)$ is not, since $\tau_1 \cup \tau_2 = \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, X\}$. Note that $\{1\}, \{3\} \in \tau_1 \cup \tau_2$, but $\{1\} \cup \{3\} \notin \tau_1 \cup \tau_2$.

Since the operations of union and intersection are both involved in the construction of a topology from a subbasis, an important simplification occurs if the open sets are constructed only by taking unions of members of \mathcal{S} ; this situation motivates the following.

Definition 6.40 (Basis). A *basis* of a topology (M, τ) is a collection of open sets \mathcal{B} such that for all $U \in \tau$ there exists an index set I and corresponding $B_i \in \mathcal{B}$ s. t.

$$\bigcup_{i \in I} B_i = U. \quad (6.2)$$

Example 6.41. Every topology is a basis for itself.

Remark 6.42. The empty set \emptyset does not need to be contained in \mathcal{B} , since it is the union of an empty collection of sets in \mathcal{B} .

Theorem 6.43. Let (X, τ) be a topological space, and fix $S \subset X$, then $\tau_S = \{S \cap U \mid U \in \tau\}$ is the subspace topology on S , cf. Theorem 6.26. If \mathcal{B} is a basis for τ , then

$$\mathcal{B}_S := \{S \cap B \mid B \in \mathcal{B}\} \quad (6.3)$$

is a basis for τ_S .

Proof. Let $U_S \in \tau_S$, then there exists a $U \in \tau$ s.t. $U_S = S \cap U$. We know there exists a collection of members of \mathcal{B} s.t. $U = \bigcup_{i \in I} B_i$, where I is an index set and $B_i \in \mathcal{B}$ for all $i \in I$, and thus

$$U_S = S \cap U = S \cap \left(\bigcup_{i \in I} B_i \right) \stackrel{(8.4)}{=} \bigcup_{i \in I} (S \cap B_i),$$

and thus \mathcal{B}_S is a basis for τ_S . ■

Theorem 6.44. A collection \mathcal{B} of open subsets of a space X is a basis iff for each open $U \subset X$ and each point $x \in U$, there is a $B \in \mathcal{B}$ s.t. $x \in B \subset U$.

Proof. “ \implies ” Let \mathcal{B} be a basis, then there exists an index set I s.t. an open $U \subset X$ can be written as $U = \bigcup_{i \in I} B_i$. Thus, for $x \in U$, there is an $i_0 \in I$ s.t. $x \in B_{i_0} \subset U$.

“ \impliedby ” Let $U \subset X$ be open, and fix $x \in U$, then there is a $B_x \in \mathcal{B}$ s.t. $B_x \subset U$. We claim that

$$U = \bigcup_{x' \in U} B_{x'}.$$

For this, note that $B_x \subset \bigcup_{x' \in U} B_{x'}$, and $B_x \subset U$ implies $\bigcup_{x' \in U} B_{x'} \subset U$. Thus, U can be written as a union of a collection of members of \mathcal{B} . ■

Corollary 6.45. Let (X, τ) be a topological space, and \mathcal{B} a basis for τ . Then a subset $U \subset X$ is open iff for all $x \in U$, there is a $B \in \mathcal{B}$ s.t. $x \in B \subset U$.

Proof. “ \implies ” Use Theorem 6.44.

“ \impliedby ” Identical to the proof in Theorem 6.44: U can be written as the union of open sets (basis members), and is thus open itself. ■

Remark 6.46. Let (X, τ) be a topological space, then a family \mathcal{S} of subsets of X is a subbasis for τ iff the family of all finite intersections of members of \mathcal{S} , denoted by \mathcal{B} , forms a basis for τ . This basis \mathcal{B} is said to be generated by \mathcal{S} , and it holds that $\mathcal{S} \subset \mathcal{B}$.

Example 6.47. Let X be an ordered set, and consider the order topology on X , i.e. according to Definition 6.35, the subbasis consists of all open rays $(-\infty, a)$ and $(a, +\infty)$, where $a \in X$ is arbitrary. This implies that the generated basis by this subbasis consists of all open rays, all open intervals (a, b) , cf. Remark 6.37, the empty set, since $(a, b) = \emptyset$ for $b \preceq a$, and X .

Let us look at when a given collection \mathcal{B} of subsets of X can serve as a basis for a topology on X .

Remark 6.48. Let \mathcal{B} be a basis for some topology on X . Since X is in the topology, there is an index set I s.t. $X = \bigcup_{i \in I} B_i$ with $B_i \in \mathcal{B}$ for all $i \in I$. And, for any $B_1, B_2 \in \mathcal{B}$, $B_1 \cap B_2$ is open (since B_1, B_2 are open), and thus there exists an index set $J \subset I$ s.t. $x \in B_1 \cap B_2$ implies $x \in \bigcup_{j \in J} B_j$, which in turn implies there exists a $j_0 \in J$ (WLOG, let $j_0 = 3$) s.t. $x \in B_3 \subset B_1 \cap B_2$. In fact, these conditions are also sufficient, as the next Theorem shows.

Theorem 6.49. Let \mathcal{B} be a collection of subsets of X s.t. there is an index set I with $X = \bigcup_{i \in I} B_i$, where $B_i \in \mathcal{B}$ for all $i \in I$, and for every two members B_1, B_2 of \mathcal{B} and for each point $x \in B_1 \cap B_2$ there is a $B_3 \in \mathcal{B}$ with $x \in B_3 \subset B_1 \cap B_2$. Then

$$\tau(\mathcal{B}) := \{U \subset X \mid \forall x \in U \exists B_x \in \mathcal{B} : x \in B_x \subset U\} = \left\{ \bigcup_{C \in \mathcal{C}} C \mid \mathcal{C} \subset \mathcal{B} \right\} \quad (6.4)$$

is the unique topology generated on X for which \mathcal{B} is a basis, and we say that $\tau(\mathcal{B})$ is generated by \mathcal{B} .

Proof. First, we will show that $\tau(\mathcal{B})$ is indeed a topology on X .

Let $U_1, U_2 \in \tau(\mathcal{B})$, then for all $x \in U_k$ there exist $B_{x,k} \in \mathcal{B}$ s.t. $x \in B_{x,k} \subset U_k$ with $k \in \{1, 2\}$. By hypothesis, for all $x \in B_{x,1} \cap B_{x,2}$, there exists a $B_{x,3} \in \mathcal{B}$ s.t. $x \in B_{x,3} \subset B_{x,1} \cap B_{x,2} \subset U_1 \cap U_2$, and thus $U_1 \cap U_2 \in \tau(\mathcal{B})$.

Now let J be an index set, and let $U_j \in \tau(\mathcal{B})$ for all $j \in J$. Then $x \in \bigcup_{j \in J} U_j$ implies there exists a $j_0 \in J$ s.t. $x \in U_{j_0}$. There is a $B_x \in \mathcal{B}$ s.t. $x \in B_x \subset U_{j_0} \subset \bigcup_{j \in J} U_j$, and thus $\bigcup_{j \in J} U_j \in \tau(\mathcal{B})$.

Also, note that $\emptyset \in \tau(\mathcal{B})$ (vacuously) and $X \in \tau(\mathcal{B})$, since $x \in X = \bigcup_{i \in I} B_i$ implies there exists a $i_0 \in I$ s.t. $x \in B_{x,i_0} \subset X$.

Now let us show that \mathcal{B} is basis for $\tau(\mathcal{B})$. For this, note that $\mathcal{B} \subset \tau(\mathcal{B})$, i.e. \mathcal{B} is a collection of open sets. And for any $U \in \tau(\mathcal{B})$, $x \in U$ implies there exists a $B_x \in \mathcal{B}$ with $x \in B_x \subset U$. Thus, $\bigcup_{x \in U} B_x \subset U$. For fixed $x \in U$, we also have $x \in B_x \subset \bigcup_{x \in U} B_x$, and thus $U \subset \bigcup_{x \in U} B_x$, i.e. $U = \bigcup_{x \in U} B_x = \bigcup_{B \in \mathcal{B}, B \subset U} B$.

Finally, we will show that $\tau(\mathcal{B})$ is the unique topology on X for which \mathcal{B} is a basis. For this, define $\tilde{\tau}(\mathcal{B}) := \{\bigcup_{C \in \mathcal{C}} C \mid \mathcal{C} \subset \mathcal{B}\} \subset \mathcal{P}(X)$, and note that $\tau(\mathcal{B}) = \tilde{\tau}(\mathcal{B})$. To see this, let $U \in \tilde{\tau}(\mathcal{B})$, i.e. $U = \bigcup_{C \in \mathcal{C}} C$ for some $\mathcal{C} \subset \mathcal{B}$. Thus, $x \in U$ implies there is a $C' \in \mathcal{C} \subset \mathcal{B}$ s.t. $x \in C' \subset U$. Conversely, if $U \in \tau(\mathcal{B})$, then we have already shown that $U = \bigcup_{x \in U} B_x$, where $x \in B_x \subset U$ and $B_x \in \mathcal{B}$. Thus, take $\mathcal{C} := \{B_x \subset U \mid B_x \in \mathcal{B}\} \subset \mathcal{B}$, then $U = \bigcup_{C \in \mathcal{C}} C$, i.e. $U \in \tilde{\tau}(\mathcal{B})$, and thus $\tau(\mathcal{B}) = \tilde{\tau}(\mathcal{B})$.

Now, if $\tau'(\mathcal{B})$ is another topology on X that has \mathcal{B} as a base, then $U \in \tau'(\mathcal{B})$ means that we can write U as a union of members of \mathcal{B} , since we assume \mathcal{B} is a basis for $\tau'(\mathcal{B})$, and thus $U \in \tau(\mathcal{B})$, i.e. $\tau'(\mathcal{B}) \subset \tau(\mathcal{B})$. Conversely, let $U \in \tau(\mathcal{B})$. Note that all members of \mathcal{B} are in τ' , since a basis is a subset of the topology. Since U is the union of members of \mathcal{B} , and a topology is closed under unions, we have that $U \in \tau'(\mathcal{B})$, i.e. $\tau(\mathcal{B}) \subset \tau'(\mathcal{B})$, which in total shows $\tau(\mathcal{B}) = \tau'(\mathcal{B})$ [52]. ■

Theorem 6.50. Let \mathcal{B} be a collection of subsets of X s.t. it fulfills the requirements of Theorem 6.49. Then $\tau(\mathcal{B})$, the topology generated by \mathcal{B} , is the intersection of all topologies containing \mathcal{B} .

Proof. Let I be an index set, then if (X, τ_i) are topological spaces, $i \in I$, containing \mathcal{B} , we want to show that $\tau(\mathcal{B}) = \bigcap_{i \in I} \tau_i$. From Corollary 6.29, we know that $\bigcap_{i \in I} \tau_i$ is a topology on X .

“ \subset ” Let $U \in \tau(\mathcal{B})$, then for all $x \in U$, there is a $B_x \in \mathcal{B}$ s.t. $x \in B_x \subset U$. We have $U = \bigcup_{x \in U} B_x$, and thus $U \in \bigcap_{i \in I} \tau_i$, since all $B_x \in \mathcal{B}$ are in $\bigcap_{i \in I} \tau_i$.

“ \supset ” We have $\tau(\mathcal{B}) = \tau_{i_0}$ for some $i_0 \in I$, since $\mathcal{B} \subset \tau(\mathcal{B})$, and hence $\bigcap_{i \in I} \tau_i \subset \tau(\mathcal{B})$. ■

Remark 6.51. A subbasis is not necessarily a basis, cf. Example 6.33, but by Theorem 6.50, every basis is a subbasis.

Remark 6.52. The notions of a (sub)basis provide shortcuts for defining topologies. It is easier to specify a basis of a topology than to explicitly define the whole topology, i.e. describe all open sets. Specifying a subbasis is even simpler. However, the price for this convenience is that it becomes more difficult to identify which sets are open. Figure 4 illustrates this.

Remark 6.53. Let (X, τ) be a topological space, and since τ is a basis for itself, the topology generated by τ , denoted by $\mathcal{T}(\tau)$, is τ , since by Theorem 6.49,

$$\mathcal{T}(\tau) = \left\{ \bigcup_{C \in \mathcal{C}} C \mid \mathcal{C} \subset \tau \right\} = \tau,$$

since for any $\mathcal{C} \subset \tau$, the union of open sets is an open set again, and for every open set $U \in \tau$, consider $\mathcal{C} = \{U\} \subset \tau$, and thus $\bigcup_{C \in \mathcal{C}} C = U$.

Theorem 6.54. Let (X, τ_X) and (Y, τ_Y) be topological spaces. Then the following properties of a function $f : X \rightarrow Y$ are equivalent [20]:

- (a) f is an open map.
- (b) For every $x \in X$ and each neighborhood U of $x \in X$, there is a neighborhood W of $f(x)$ in Y with $W \subset f(U)$.




Figure 4: Figure taken from [53].

- (c) For every $x \in X$ and each basic neighborhood B of $x \in X$, $f(B)$ is a neighborhood of $f(x)$ in Y .
- (d) The image of each member of a basis for X is open in Y .

Proof. “(a) \Rightarrow (b)” Let U be a neighborhood of an $x \in X$, then there exists an open set $O \in \tau_X$ s.t. $x \in O \subset U$. Define $W := f(O)$, then $W \in \tau_Y$, since f is an open map, and $W \subset f(U)$, since $O \subset U$.

“(b) \Rightarrow (c)” Let B be a neighborhood of $x \in X$, then there is a neighborhood W of $f(x)$ in Y s.t. $f(x) \in V \subset W \subset f(B)$, where $V \in \tau_Y$, and thus $f(B)$ is a neighborhood of $f(x)$ in Y .

“(c) \Rightarrow (d)” Use Theorem 6.14.

“(d) \Rightarrow (a)” Let \mathcal{B} be a basis for the topology on X , and $f(B) \in \tau_Y$ for all $B \in \mathcal{B}$. An open $U \subset X$ can be written as the union of a collection $\{B_i\}_{i \in I} \subset \mathcal{B}$, where I is an index set, and thus:

$$f(U) = f\left(\bigcup_{i \in I} B_i\right) = \bigcup_{i \in I} f(B_i) \in \tau_Y.$$

■

Example 6.55 (Metric topology). In a metric space X , the collection of all open balls

$$\mathcal{B} := \{B_r(x) \mid x \in X, r > 0\} \tag{6.5}$$

generates the topology

$$\tau(\mathcal{B}) = \{U \subset X \mid \forall x \in U \exists B_r(x) \in \mathcal{B} : x \in B_r(x) \subset U\},$$

called the *metric topology*, cf. Eq. (6.4) and Theorem 6.49, and \mathcal{B} is a basis for $\tau(\mathcal{B})$. This holds because for any fixed $r > 0$, $X = \bigcup_{x \in X} B_r(x)$, and for any point in the intersection of two open balls, there is an open ball containing that point, and the open ball is contained in the intersection, cf. Theorem 5.9.

Remark 6.56. Note that it is important to consider the collection of *all* open balls in Example 6.55. For if we considered the collection of all open balls with *fixed* radius $r > 0$, it is not guaranteed that for all $B_r(x)$ and $B_r(x')$ and each point $y \in B_r(x) \cap B_r(x')$, we can find a $B_r(z)$ with $z \in X$ s.t. $y \in B_r(z) \subset B_r(x) \cap B_r(x')$, also cf. Fig. 2.

Example 6.57. Let (X, d) be a metric space, where d is the discrete metric from Remark 5.6. Then the collection $\mathcal{B} := \{\varphi \in X\} \cup \{X\}$, which is the collection of all open balls, cf. Example 5.8, generates a topology $\tau(\mathcal{B})$ on X by Example 6.55, for which \mathcal{B} is a basis. The generated topology on X is the discrete topology from Example 6.4. Note that $\mathcal{B}' := \{\varphi \in X\}$ generates the same topology on X .

Definition 6.58 (Metrizable topological space [20]). Let (X, τ) be a topological space, then it is called *metrizable* if there is a metric d on X , whose collection of all open balls generates the topology τ .

Definition 6.59 (Standard topology of \mathbb{R}). Let $X = \mathbb{R}$, and d the usual metric on X . Then for $\varphi \in X$ and $r > 0$, the open ball is

$$B_r(\varphi) = \{\psi \in X \mid |\varphi - \psi| < r\} = (\varphi - r, \varphi + r),$$

and the collection of all such open balls is

$$\mathcal{B} = \{(\varphi - r, \varphi + r) \mid \varphi \in X, r > 0\} = \{(a, b) \mid a, b \in X, a < b\},$$

and the topology generated by this basis is called the *standard topology* of \mathbb{R} .

Definition 6.60 (Lower limit topology of \mathbb{R}). Let $X = \mathbb{R}$, and \mathcal{B} the collection of all right half-open intervals, i.e. $\mathcal{B} = \{[a, b) \mid a, b \in \mathbb{R}, a < b\}$. Clearly, there is a collection of right half-open intervals s.t. \mathbb{R} equals to its union, and the intersection of two right half-open intervals is either empty or itself a right half-open interval. Thus, for all x in the intersection there is a right half-open interval that contains x and is contained in the intersection. Thus, by Theorem 6.49, \mathcal{B} generates a topology, which is called the *lower limit topology* of \mathbb{R} .

Definition 6.61 (Upper limit topology of \mathbb{R}). Let $X = \mathbb{R}$, and \mathcal{B} the collection of all left half-open intervals, i.e. $\mathcal{B} = \{(a, b] \mid a, b \in \mathbb{R}, a < b\}$. Clearly, there is a collection of left half-open intervals s.t. \mathbb{R} equals to its union, and the intersection of two left half-open intervals is either empty or itself a left half-open interval. Thus, for all x in the intersection there is a left half-open interval that contains x and is contained in the intersection. Thus, by Theorem 6.49, \mathcal{B} generates a topology, which is called the *upper limit topology* of \mathbb{R} .

Remark 6.62. Let $X = \mathbb{R}$, and \mathcal{B} the collection of all closed intervals, i.e. $\mathcal{B} = \{[a, b] \mid a, b \in \mathbb{R}, a \leq b\}$. Note that there is a collection of closed intervals s.t. \mathbb{R} is its union, and the intersection of two closed intervals is either empty or closed. Thus, for all x in the intersection, there is a closed interval that contains x and that is contained in the intersection. Thus, by Theorem 6.49, \mathcal{B} generates a topology, which is the discrete topology on \mathbb{R} , i.e. $\tau(\mathcal{B}) = \mathcal{P}(\mathbb{R})$, since for any $x \in \mathbb{R}$, we have $x \in [x, x] \in \mathcal{B}$.

Remark 6.63. The lower and upper limit topologies of \mathbb{R} are not metrizable [54].

Example 6.64. The order topology on \mathbb{R} (with the usual order relation) coincides with the standard topology of \mathbb{R} , since all open rays $(-\infty, a)$ and (a, ∞) can be written as a union of open intervals. Conversely, every open interval, which is an open set in the standard topology, is contained in the order topology.

Definition 6.65. The *Euclidean topology* of \mathbb{R}^n is generated by the basis of open balls, where the metric is the Euclidean metric.

Theorem 6.66. Let $\tau(\mathcal{B})$ and $\tau(\mathcal{B}')$ be topologies on X generated by the bases \mathcal{B} and \mathcal{B}' respectively, cf. Theorem 6.49. Then we have:

$$\tau(\mathcal{B}) \subset \tau(\mathcal{B}') \text{ (}\tau(\mathcal{B}) \text{ is coarser than } \tau(\mathcal{B}')\text{)} \Leftrightarrow \forall B \in \mathcal{B} \forall x \in B \exists B' \in \mathcal{B}' : x \in B' \subset B.$$

Proof. “ \Rightarrow ” Let $\tau(\mathcal{B}) \subset \tau(\mathcal{B}')$, then for all $B \in \mathcal{B}$, we have $B \in \tau(\mathcal{B})$, and thus $B \in \tau(\mathcal{B}')$, i.e. by construction, for all $x \in B$ there exists a $B'_x \in \mathcal{B}'$ s.t. $x \in B'_x \subset B$.

“ \Leftarrow ” Let $U \in \tau(\mathcal{B})$, i.e. for all $x \in U$ there exists a $B_x \in \mathcal{B}$ s.t. $x \in B_x \subset U$. By assumption, to this $B_x \in \mathcal{B}$, there exists a $B'_x \in \mathcal{B}'$ s.t. $x \in B'_x \subset B_x \subset U$, which shows that $U \in \tau(\mathcal{B}')$. ■

Corollary 6.67. Let $\tau(\mathcal{B})$ and $\tau(\mathcal{B}')$ be topologies on X generated by the bases \mathcal{B} and \mathcal{B}' respectively, then $\tau(\mathcal{B}) = \tau(\mathcal{B}')$ iff for each $B \in \mathcal{B}$ and $x \in B$, there exists a $B' \in \mathcal{B}'$ s.t. $x \in B' \subset B$, and for each $B' \in \mathcal{B}'$ and $x \in B'$ there exists a $B \in \mathcal{B}$ s.t. $x \in B \subset B'$.

Remark 6.68. Let $X = \mathbb{R}$, and denote by τ its standard topology, by τ_l the lower limit topology, and by τ_u the upper limit topology. Then $\tau \subset \tau_l$ and $\tau \subset \tau_u$, i.e. the standard topology is coarser than the lower and upper limit topologies, since for all open intervals in τ of the form (a, b) with $a, b \in \mathbb{R}$ and $a < b$, and all points $x \in (a, b)$, the right half-open interval $[x, b)$ contains x and is a subset of (a, b) , and the left half-open interval $(a, x]$ contains x and is a subset of (a, b) . Thus, τ is coarser than τ_l and τ_u , cf. Theorem 6.66. However, note that the standard topology is not equal to either the lower or upper limit topology, since there is no open interval that would be contained in intervals of the form $[a, b)$ or $(a, b]$ and contain each point of the intervals.

Remark 6.69. Let $\tau(\mathcal{B})$ be the topology on X generated by the basis \mathcal{B} , then Corollary 6.67 implies that there might be another, distinct basis \mathcal{B}' s.t. $\tau(\mathcal{B}) = \tau(\mathcal{B}')$.

Example 6.70. Let $X = \{a, b, c\}$, then $\mathcal{B} = \{\{a, b\}, \{b, c\}, \{c, a\}, \{a\}, \{b\}, \{c\}\}$ and $\mathcal{B}' = \{\{a\}, \{b\}, \{c\}\}$ are collection of subsets of X s.t. the requirements of Theorem 6.49 are fulfilled [55], and we have $\tau(\mathcal{B}) = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{c, a\}, X\} = \tau(\mathcal{B}')$.

Example 6.71. Let $X = \mathbb{R}$, and \mathcal{B} the collection of all open intervals, then $\tau(\mathcal{B})$ is the standard topology, cf. Example 6.59. Now fix $c, d \in \mathbb{R}$, and consider

$$\mathcal{B}' := \{(a, b) \mid a, b \in \mathbb{R}, a < b, a \neq c \wedge b \neq d\},$$

i.e. the collection of all open intervals, where one specific interval was removed. Since \mathcal{B}' fulfills the requirements of Theorem 6.49, it generates the topology $\tau(\mathcal{B}')$. Clearly, by Corollary 6.67, $\tau(\mathcal{B}) = \tau(\mathcal{B}')$, and since c, d were arbitrary, it means that there infinitely many bases generating the standard topology of \mathbb{R} .

Remark 6.72. Let $X = \mathbb{R}$, and consider the collection of all open intervals with rational endpoints, i.e.

$$\mathcal{B} = \{(a, b) \mid a, b \in \mathbb{Q}, a < b\}.$$

Clearly, \mathbb{R} can be written as the union of all such intervals, and the intersection of two open intervals is either empty or an open interval again. Thus, \mathcal{B} fulfills the requirements of Theorem 6.49, and the generated topology is the standard topology of \mathbb{R} . To see this, consider an open interval of the form (a, b) with $a, b \in \mathbb{R}$ and $a \neq b$. Then for all $x \in (a, b)$, by the Archimedean property of \mathbb{R} , cf. Theorem B.23, we know that there is $p \in \mathbb{Q}$ s.t. $a < p < x$ and a $q \in \mathbb{Q}$ s.t. $x < q < b$, thus (p, q) contains x and $(p, q) \subset (a, b)$. Conversely, if we choose $(p, q) \in \mathcal{B}$, note that $p, q \in \mathbb{Q} \subset \mathbb{R}$, and thus by Corollary 6.67, $\tau(\mathcal{B})$ also generates the standard topology of \mathbb{R} .

Definition 6.73 (Topological equivalence of metrics [20]). Let (X, d) and (X, d') be metric spaces. We denote by \mathcal{B} the collection of all d -open balls, generating the metric topology $\tau(\mathcal{B})$, cf. Example 6.55, and by \mathcal{B}' the collection of all d' -open balls generating the metric topology $\tau(\mathcal{B}')$. If $\tau(\mathcal{B}) = \tau(\mathcal{B}')$, we say that d and d' are (*topologically*) *equivalent*.

Theorem 6.74. Two metrics d and d' on a set X are equivalent if and only if for each $x \in X$ and for each $\epsilon > 0$, there exist real numbers $\delta_1, \delta_2 > 0$ (depending upon x and ϵ) such that $B_{\delta_1}^d(x) \subset B_\epsilon^{d'}(x)$ and $B_{\delta_2}^{d'}(x) \subset B_\epsilon^d(x)$. [20].

Proof. Let the notation be as in Definition 6.73.

“ \implies ” For each $x \in X$ and for each $\epsilon > 0$, the open ball $B_\epsilon^d(x) \in \mathcal{B}$ in the metric space (X, d) is open in the topology $\tau(\mathcal{B})$, and the open ball $B_\epsilon^{d'}(x) \in \mathcal{B}'$ in the metric space (X, d') is open in the topology $\tau(\mathcal{B}')$. Hence, by Corollary 6.67, there exist δ_1, δ_2 such that $B_{\delta_1}^{d'}(x) \subset B_\epsilon^d(x)$ and $B_{\delta_2}^d(x) \subset B_\epsilon^{d'}(x)$.

“ \impliedby ” Suppose that $B_{\delta_1}^{d'}(x) \subset B_\epsilon^d(x)$ and $B_{\delta_2}^d(x) \subset B_\epsilon^{d'}(x)$ hold for each $x \in X$ and $\epsilon > 0$. Given an open set $U \subset X$ in the topology $\tau(\mathcal{B})$ and a point $x \in U$, there exists an open ball $B_r^d(x) \subset U$ with $r > 0$. By assumption, there exists an open ball $B_{\delta_1}^{d'}(x) \subset B_r^d(x) \subset U$, proving that U is an open set in the topology $\tau(\mathcal{B}')$. Similarly, one can show that every open set in the topology $\tau(\mathcal{B}')$ is also open in $\tau(\mathcal{B})$. Hence, by Corollary 6.67, $\tau(\mathcal{B}) = \tau(\mathcal{B}')$. \blacksquare

Theorem 6.75. If two metrics d and d' on a set X are strongly equivalent, cf. Def. 4.9, then they are also (topologically) equivalent.

Proof. [56] By definition of strong equivalence, we know that

$$\alpha d(x, y) \leq d'(x, y) \leq \beta d(x, y) \quad \forall x, y \in X, \alpha, \beta > 0. \quad (6.6)$$

According to Theorem 6.74, we need to show that for all $x \in X$ and $\epsilon > 0$, there exist $\delta_1, \delta_2 > 0$ such that

$$\left(B_{\delta_1}^d(x) \subset B_\epsilon^{d'}(x) \right) \wedge \left(B_{\delta_2}^{d'}(x) \subset B_\epsilon^d(x) \right) \quad (6.7)$$

To show the first inclusion, let $\delta_1 \leq \epsilon/\beta$, then we have for all $x, y \in X$:

$$\frac{d'(x, y)}{\beta} \leq d(x, y) < \delta_1 \leq \frac{\epsilon}{\beta} \Rightarrow d'(x, y) < \epsilon. \quad (6.8)$$

To show the second inclusion, let $\delta_2 \leq \epsilon\alpha$, then we have for all $x, y \in X$:

$$\alpha d(x, y) \leq d'(x, y) < \delta_2 \leq \epsilon\alpha \Rightarrow d(x, y) < \epsilon. \quad (6.9)$$

■

Theorem 6.76. Let (X, d) be a metric space, and consider the metric

$$\rho(x, y) := \frac{d(x, y)}{1 + d(x, y)}$$

for all $x, y \in X$, cf. Theorem 5.3. Then d and ρ are topologically equivalent.

Proof. Choose arbitrary $x \in X$ and $\epsilon > 0$. Choose $\delta_1 := \epsilon/(1 - \epsilon)$, then $B_{\delta_1}^d(x) \subset B_\epsilon^\rho(x)$, since for any $y \in B_{\delta_1}^d(x)$, we have

$$\begin{aligned} d(x, y) < \delta_1 = \frac{\epsilon}{1 - \epsilon} &\Leftrightarrow d(x, y) - \epsilon d(x, y) < \epsilon \Leftrightarrow d(x, y) < \epsilon + \epsilon d(x, y) = \epsilon(1 + d(x, y)) \\ &\Leftrightarrow \frac{d(x, y)}{1 + d(x, y)} = \rho(x, y) \leq \epsilon, \end{aligned}$$

i.e. $y \in B_\epsilon^\rho(x)$. Conversely, choose $\delta_2 := \epsilon$, then $B_{\delta_2}^\rho(x) \subset B_\epsilon^d(x)$, since for any $y \in B_{\delta_2}^\rho(x)$, $\rho(x, y) \leq d(x, y)$.

Thus, by Theorem 6.74, d and ρ are topologically equivalent. ■

Remark 6.77. There exist metrics d and d' on a set X that are (topologically) equivalent, but not strongly equivalent. [57]

Example 6.78. Let $X := (0, 1]$, and for any $x, y \in X$, define $d(x, y) := |x - y|$ and $d'(x, y) := |x^{-1} - y^{-1}|$. Clearly, d and d' define metrics on X , and they are not strongly equivalent, since there is no $\beta > 0$ s.t. $d'(x, y) \leq \beta d(x, y)$ for all $x, y \in X$.

However, d and d' are (topologically) equivalent. To see this, remember that acc. to Theorem 6.74, we need to prove that for any $x \in X$ and for any $\epsilon > 0$, there exist $\delta_1(x, \epsilon)$ and $\delta_2(x, \epsilon)$ s.t. $B_{\delta_1}^d(x) \subset B_\epsilon^{d'}(x)$ and $B_{\delta_2}^{d'}(x) \subset B_\epsilon^d(x)$.

To show the first inclusion, choose any

$$\delta_1 < \frac{\epsilon x^2}{\epsilon x + 1}. \quad (6.10)$$

Now we have

$$y \in B_{\delta_1}^d(x) \Rightarrow |y - x| < \delta_1 \Rightarrow \frac{1}{xy} |x - y| = \left| \frac{1}{x} - \frac{1}{y} \right| < \frac{\delta_1}{xy} \quad (6.11)$$

Since $z \leq |z|$ for any $z \in \mathbb{R}$,

$$y \in B_{\delta_1}^d(x) \Rightarrow x - y \leq |x - y| < \delta_1 \Rightarrow x - \delta_1 < y \Rightarrow \frac{1}{y} < \frac{1}{x - \delta_1}, \quad (6.12)$$

and hence

$$y \in B_{\delta_1}^d(x) \stackrel{(6.11)}{\Rightarrow} \left| \frac{1}{x} - \frac{1}{y} \right| < \frac{\delta_1}{xy} \stackrel{(6.12)}{<} \frac{\delta_1}{x(x - \delta_1)} = \frac{\delta_1}{x^2 - x\delta_1} \stackrel{(6.10)}{<} \frac{\epsilon x^2}{(\epsilon x + 1)(x^2 - \frac{\epsilon x^3}{\epsilon x + 1})} \quad (6.13)$$

$$= \frac{\epsilon x^2}{\epsilon x^3 - \frac{\epsilon^2 x^4}{\epsilon x + 1} + x^2 - \frac{\epsilon x^3}{\epsilon x + 1}} = \frac{\epsilon x^2}{\frac{\epsilon^2 x^4 + \epsilon x^3}{\epsilon x + 1} - \frac{\epsilon^2 x^4}{\epsilon x + 1} + x^2 - \frac{\epsilon x^3}{\epsilon x + 1}} = \epsilon. \quad (6.14)$$

To show the second inclusion, i.e. $B_{\delta_2}^{d'}(x) \subset B_\epsilon^d(x)$, choose $\delta_2 := \epsilon$, then we have:

$$y \in B_{\delta_2}^{d'}(x) \Rightarrow \left| \frac{1}{x} - \frac{1}{y} \right| < \delta_2 \Rightarrow xy \left| \frac{1}{x} - \frac{1}{y} \right| = |y - x| < \delta_2 xy \stackrel{x, y \in X = (0, 1]}{\leq} \delta_2 = \epsilon. \quad (6.15)$$

Theorem 6.79. Two metrics d and d' on a set X are (topologically) equivalent iff each convergent sequence in X wrt d is also convergent in X wrt d' . In that case, their limits coincide.

Proof. “ \implies ” Let d and d' be equivalent metrics on X , and let $(\varphi_n)_{n \in \mathbb{N}}$ be a sequence in X , which converges to $\varphi \in X$ wrt d , i.e.

$$d(\varphi, \varphi_n) \xrightarrow{n \rightarrow \infty} 0. \quad (6.16)$$

According to Theorem 6.74, for any $\epsilon > 0$, there is a $\delta > 0$ s.t. $B_\delta^d(\varphi) \subset B_\epsilon^{d'}(\varphi)$. To this δ , there exists an $N \in \mathbb{N}$ s.t. $d(\varphi, \varphi_n) < \delta$ for all $n \geq N$, cf. Def. 5.17. Hence:

$$\varphi_n \in B_\delta^d(\varphi) \subset B_\epsilon^{d'}(\varphi), \quad \forall n \geq N \quad (6.17)$$

i.e. $d'(\varphi, \varphi_n) < \epsilon$ for any $\epsilon > 0$.

“ \impliedby ” Let $U \subset X$ be d -open in X , i.e. open wrt d . Then $X \setminus U$ is d -closed in X . Let $\varphi \in X$ be a d' -accumulation point of $X \setminus U$, i.e. there exists a sequence $(\varphi_n)_{n \in \mathbb{N}}$ in $X \setminus U$ s.t. $\varphi_n \xrightarrow{d'} \varphi$ for $n \rightarrow \infty$. By assumption, $(\varphi_n)_{n \in \mathbb{N}}$ converges to a $\psi \in X$ wrt d . Since $X \setminus U$ is closed, by Theorem 5.26, $\psi \in X \setminus U$ holds.

Let us now show that $\varphi = \psi$. For this, consider the extended sequence

$$(\varphi_1, \varphi, \varphi_2, \varphi, \varphi_3, \varphi, \dots). \quad (6.18)$$

The extended sequence converges to φ wrt d' , and hence it also converges wrt d . The subsequence $(\varphi_n)_{n \in \mathbb{N}}$ of the extended sequence converges to ψ wrt d . Hence, $\varphi = \psi$, and the d' -accumulation point $\varphi = \psi$ of $X \setminus U$ lies in $X \setminus U$, i.e. $X \setminus U$ is also d' -closed (by Theorem 5.26). This in turn implies that U is d' -open.

We have shown that d and d' induce the same topology on X , and are hence equivalent. ■

Theorem 6.80. Let (X, τ) be a metrizable topological space, and d a metric on X inducing τ . Then for each $\lambda > 0$ there exists a metric d_λ on X s.t. d_λ is (topologically) equivalent to d and $\text{diam}(X) \leq \lambda$ wrt d_λ [20].

Proof. For all $x, y \in X$, define $d_\lambda(x, y) := \min\{\lambda, d(x, y)\}$. Clearly, d_λ is positive, and

$$d_\lambda(x, y) = \min\{\lambda, d(x, y)\} = 0 \Leftrightarrow d(x, y) = 0 \Leftrightarrow x = y,$$

i.e. d_λ is also definite. Clearly, d_λ is symmetric, to prove the triangle inequality, let $z \in X$, and note that

$$\begin{aligned} d_\lambda(x, z) &= \min\{\lambda, d(x, z)\} \leq \min\{\lambda, d(x, y) + d(y, z)\} \\ &\leq \min\{\lambda, d(x, y)\} + \min\{\lambda, d(y, z)\} = d_\lambda(x, y) + d_\lambda(y, z). \end{aligned}$$

Thus, d_λ is a metric on X , and clearly $\text{diam}(X) \leq \lambda$ wrt d_λ .

For all $x \in X$ and $\epsilon > 0$, we have

$$B_\epsilon^d(x) = \{y \in X \mid d(x, y) < \epsilon\} \subset \{y \in X \mid \min\{\lambda, d(x, y)\} < \epsilon\} = B_\epsilon^{d_\lambda}(x)$$

and for $\delta := \min\{\lambda, \epsilon\}$, we have

$$B_\delta^{d_\lambda}(x) = \{y \in X \mid \min\{\lambda, d(x, y)\} < \min\{\lambda, \epsilon\}\} \subset B_\epsilon^d(x),$$


since if $d_\lambda(x, y) = d(x, y)$, then $d(x, y) < \epsilon$, otherwise $d_\lambda(x, y) = \lambda$. In that case, if $\epsilon \leq \lambda$, we have $\min\{\lambda, \epsilon\} = \epsilon$, and thus there is no $y \in X$ fulfilling $\lambda < \epsilon$, i.e. $B_\delta^{d_\lambda}(x) = \{x\} \subset B_\epsilon^d(x)$. Otherwise, $\lambda < \epsilon$, but there is no $y \in X$ fulfilling $\lambda < \lambda$, i.e. $B_\delta^{d_\lambda}(x) = \{x\} \subset B_\epsilon^d(x)$. Thus, by Theorem 6.74, d_λ is topologically equivalent to d . ■

Let us now look at what topological structure we can endow $X \times Y := \{(x, y) \mid x \in X, y \in Y\}$ with if (X, τ_X) and (Y, τ_Y) are topological spaces.

Remark 6.81. The naive way would be to define the collection of open sets as

$$\mathcal{B} := \{U \times V \mid U \in \tau_X, V \in \tau_Y\} \subset \mathcal{P}(X \times Y). \quad (6.19)$$

While $\emptyset \in \mathcal{B}$ (since $\emptyset \in \tau_X$, $Y \in \tau_Y$, and $\emptyset \times Y = \emptyset$), $X \times Y \in \mathcal{B}$, and if $U_1 \times V_1 \in \mathcal{B}$ and $U_2 \times V_2 \in \mathcal{B}$, then $(U_1 \times V_1) \cap (U_2 \times V_2) \in \mathcal{B}$, i.e. the intersection of two members of \mathcal{B} is also in \mathcal{B} again⁹. However, note that the union of two members of \mathcal{B} is not necessarily in \mathcal{B} again. For example, consider these two rectangles, then their union is clearly not a rectangle again:



Mathematically, we can describe these two rectangles as $(1, 2) \times (1, 2)$ and $(1.5, 2.5) \times (1.5, 2.5)$, and we consider $X = Y = \mathbb{R}$ with the standard topology.

⁹This holds because

$$\begin{aligned} (U_1 \times V_1) \cap (U_2 \times V_2) &= \{(u, v) \mid (u, v) \in U_1 \times V_1 \wedge (u, v) \in U_2 \times V_2\} \\ &= \{(u, v) \mid u \in U_1 \wedge v \in V_1 \wedge u \in U_2 \wedge v \in V_2\} \\ &= \{(u, v) \mid u \in U_1 \cap U_2 \wedge v \in V_1 \cap V_2\} \\ &= (U_1 \cap U_2) \times (V_1 \cap V_2), \end{aligned}$$

and we have $U_1, U_2 \in \tau_X$ and $V_1, V_2 \in \tau_Y$, i.e. $U_1 \cap U_2 \in \tau_X$ and $V_1 \cap V_2 \in \tau_Y$.

Definition 6.82 (Product topology [20]). Let \mathcal{B} be defined as in Eq. 6.19, then \mathcal{B} fulfills the requirements of Theorem 6.49, and thus \mathcal{B} generates a topology $\tau(\mathcal{B})$ on $X \times Y$, called the *product topology*. When $X \times Y$ is equipped with the product topology, it is also called the *product space*; X and Y are called the *coordinate spaces*. The maps

$$\begin{aligned} p_X : X \times Y &\rightarrow X, (x, y) \mapsto x, \\ p_Y : X \times Y &\rightarrow Y, (x, y) \mapsto y \end{aligned} \tag{6.20}$$

are projections of the product space $X \times Y$ onto the coordinates spaces.

Theorem 6.83. Let (X, τ_X) and (Y, τ_Y) be topological spaces. The projections p_X and p_Y from the product space onto the coordinates spaces, cf. Eq. (6.20), are continuous, the product topology is the smallest topology for which this is true, and p_X and p_Y are open maps.

Proof. Let $U \in \tau_X$, then

$$p_X^{-1}(U) = \{(x, y) \in X \times Y \mid p_X((x, y)) \in U\} = \{(x, y) \in X \times Y \mid x \in U\} = U \times Y,$$

and thus p_X is continuous in the product topology on $X \times Y$, cf. Definition 6.126. Similarly, we can show the continuity of p_Y .

Now suppose there is a topology \mathcal{T} on $X \times Y$ s.t. both p_X and p_Y are continuous. Let $U \in \tau_X$ and $V \in \tau_Y$, then $p_X^{-1}(U) = U \times Y \in \mathcal{T}$, $p_Y^{-1}(V) = X \times V \in \mathcal{T}$, and thus

$$p_X^{-1}(U) \cap p_Y^{-1}(V) = (U \times Y) \cap (X \times V) = (U \cap X) \times (Y \cap V) = U \times V, \tag{6.21}$$

i.e. all basis members of the product topology on $X \times Y$ are contained in \mathcal{T} (note that \mathcal{T} is a basis for itself, generating \mathcal{T} , cf. Remark 6.53), and thus by Theorem 6.66, \mathcal{T} is finer than the product topology on $X \times Y$, i.e. the product topology is the smallest topology on which p_X and p_Y are continuous.

Finally, note that p_X and p_Y are open maps. To see this, note that if $U \in \tau_X$ and $V \in \tau_Y$, then $U \times V$ is open in the product topology, and $p_X(U \times V) = U \in \tau_X$ and $p_Y(U \times V) = V \in \tau_Y$. ■

Remark 6.84. Define $\mathcal{S} := \{p_X^{-1}(U) \mid U \in \tau_X\} \cup \{p_Y^{-1}(V) \mid V \in \tau_Y\}$, then \mathcal{S} is a sub-basis for the product topology on $X \times Y$, since $p_X^{-1}(U) \cap p_Y^{-1}(V) = U \times V \in \mathcal{B}$, cf. Eq. (6.21) and (6.19).

If bases of the coordinate spaces are known, then we can construct a smaller basis for the product topology than in Eq. (6.19).

Theorem 6.85. Let \mathcal{C} be a basis for τ_X , and \mathcal{D} be a basis for τ_Y , then

$$\mathcal{B}' := \{C \times D \mid C \in \mathcal{C}, D \in \mathcal{D}\}$$

is a basis for the product topology on $X \times Y$.

Proof. By Definition 6.82, $\mathcal{B} = \{U \times V \mid U \in \tau_X, V \in \tau_Y\}$ is a basis for the product topology on $X \times Y$. Let $U \times V \in \mathcal{B}$, i.e. $U \in \tau_X$ and $V \in \tau_Y$. Then by Theorem 6.44, for all $x \in U$ there is a $C_x \in \mathcal{C}$ s.t. $x \in C_x \subset U$, and for all $y \in V$ there is a $D_y \in \mathcal{D}$ s.t. $y \in D_y \subset V$. Thus, for all $U \times V \in \mathcal{B}$ and all $(x, y) \in U \times V$, there is $C_x \times D_y \in \mathcal{B}'$ s.t. $(x, y) \in C_x \times D_y \subset U \times V$.

Conversely, if $C \times D \in \mathcal{B}'$, i.e. $C \in \mathcal{C}$ and $D \in \mathcal{D}$, then $C \times D \in \mathcal{B}$, and thus by Corollary 6.67, \mathcal{B}' generates the product topology on $X \times Y$. ■

Remark 6.86. Let (X, τ_X) and (Y, τ_Y) be topological spaces, $A \subset X$ and $B \subset Y$. There are two ways in which we can define a topology on $A \times B$. First, notice that $A \times B \subset X \times Y$, and thus we can endow $A \times B$ with the subspace topology, cf. Theorem 6.26. According to Theorem 6.43, sets of the form $(A \times B) \cap (U \times V)$, where $U \in \tau_X$ and $V \in \tau_Y$, are members of the basis for this topology.

Since $A \subset X$, A is endowed with the subspace topology from X , similarly for $B \subset Y$, and thus $A \times B$ is the product topology of the subspace topologies on A and B , where the basis members are of form $(A \cap U) \times (B \cap V)$ for $U \in \tau_X$ and $V \in \tau_Y$. Note that $(A \cap U) \times (B \cap V) = (A \times B) \cap (U \times V)$, and thus both topologies coincide.

Theorem 6.87. Let X, Y, Z be topological spaces. A function $f : Z \rightarrow X \times Y$ is continuous iff the compositions $p_X \circ f : Z \rightarrow X$ and $p_Y \circ f : Z \rightarrow Y$ are continuous.

Proof. “ \implies ” If f is continuous, then $p_X \circ f$ and $p_Y \circ f$ are continuous, since they are the composition of continuous functions.

“ \impliedby ” Let $p_X \circ f$ and $p_Y \circ f$ be continuous. Let $U \subset X$ and $V \subset Y$ be open, then we want to show that $f^{-1}(U \times V) = \{z \in Z \mid f(z) \in U \times V\}$ is open in Z . For this, note that

$$(p_X \circ f)^{-1}(U) \cap (p_Y \circ f)^{-1}(V) = \{z \in Z \mid (p_X \circ f)(z) \in U\} \cap \{z \in Z \mid (p_Y \circ f)(z) \in V\}$$

and

$$\begin{aligned} z \in f^{-1}(U \times V) &\Leftrightarrow f(z) \in U \times V \Leftrightarrow (p_X \circ f)(z) \in U \wedge (p_Y \circ f)(z) \in V \\ &\Leftrightarrow z \in (p_X \circ f)^{-1}(U) \cap (p_Y \circ f)^{-1}(V), \end{aligned}$$

i.e. $f^{-1}(U \times V) = (p_X \circ f)^{-1}(U) \cap (p_Y \circ f)^{-1}(V)$, and since $p_X \circ f$ and $p_Y \circ f$ are continuous by assumption, $(p_X \circ f)^{-1}(U)$ and $(p_Y \circ f)^{-1}(V)$ are open in Z , and so is their intersection. \blacksquare

Remark 6.88. Let $(X, \tau_X), (Y, \tau_Y)$ and (Z, τ_Z) be topological spaces. The operation of forming the product of topological spaces is commutative, in the sense that there is an obvious canonical homeomorphism

$$f : X \times Y \rightarrow Y \times X, (x, y) \mapsto (y, x),$$

where $X \times Y$ is equipped with the product topology on $X \times Y$, and $Y \times X$ is equipped with the product topology on $Y \times X$.

The operation of forming the product of topological spaces is also associative, in the sense that there is an obvious canonical homeomorphism

$$g : (X \times Y) \times Z \rightarrow X \times (Y \times Z), ((x, y), z) \mapsto (x, (y, z)),$$

where $(X \times Y) \times Z$ is equipped with the product topology on $(X \times Y) \times Z$, and $X \times (Y \times Z)$ is equipped with the product topology on $X \times (Y \times Z)$.

Proof. We use Theorem 6.130 for the proof.

Clearly, f and g are bijective. f is continuous, since a basis for the product topology on $Y \times X$ is the set $\{V \times U \mid V \in \tau_Y, U \in \tau_X\}$, and

$$\begin{aligned} f^{-1}(V \times U) &= \{(x, y) \in X \times Y \mid f((x, y)) \in V \times U\} \\ &= \{(x, y) \in X \times Y \mid (y, x) \in V \times U\} = U \times V, \end{aligned}$$

and $U \times V$ is a member of the basis for the product topology on $X \times Y$.

$f^{-1} : Y \times X \rightarrow X \times Y$ is continuous, since a basis for the product topology on $X \times Y$ is $\{U \times V \mid U \in \tau_X, V \in \tau_Y\}$, and

$$(f^{-1})^{-1}(U \times V) = f(U \times V) = \{f((u, v)) \mid (u, v) \in U \times V\} = V \times U,$$

and $V \times U$ is a member of the basis for the product topology on $Y \times X$. Thus, f is a homeomorphism.

g is continuous, since a basis for the product topology on $X \times (Y \times Z)$ is $\{U \times (V \times W) \mid U \in \tau_X, V \in \tau_Y, W \in \tau_Z\}$, and

$$\begin{aligned} g^{-1}(U \times (V \times W)) &= \{((x, y), z) \in (X \times Y) \times Z \mid g((x, y), z) \in U \times (V \times W)\} \\ &= \{((x, y), z) \in (X \times Y) \times Z \mid (x, (y, z)) \in U \times (V \times W)\} \\ &= (U \times V) \times W, \end{aligned}$$

and $(U \times V) \times W$ is a member of the basis for the product topology on $(X \times Y) \times Z$.

Similarly, we can show that g^{-1} is continuous, and thus g is a homeomorphism. ■

Definition 6.89 (Product of topological spaces [20]). Define $I := \{1, \dots, n\}$, for some $n \in \mathbb{N}$, and let (X, τ_i) be topological spaces for all $i \in I$, then based on Remark 6.88, we can inductively define the product of topological spaces to be

$$\prod_{i=1}^n X_i = X_1 \times \dots \times X_n := (X_1 \times \dots \times X_{n-1}) \times X_n. \quad (6.22)$$

Remark 6.90. The results for the finite product of topological spaces follow by induction from the corresponding results for the product of two topological spaces. For example, if (X, τ_i) are topological spaces for all $i \in \{1, \dots, n\}$ for some $n \in \mathbb{N}$, and \mathcal{B}_i is a basis for τ_i , then Theorem 6.85 implies that $\mathcal{B} := \{\prod_{i=1}^n U_i \mid U_i \in \mathcal{B}_i\}$ is a basis for the product topology on $\prod_{i=1}^n X_i$.

Theorem 6.91. For all $i \in \{1, \dots, n\}$, with $n \in \mathbb{N}$, let (X_i, τ_i) be metrizable topological spaces. Then $\prod_{i=1}^n X_i$ with the product topology on it is metrizable.

Proof [58]. First, we construct a metric on $X := \prod_{i=1}^n X_i$. Since each X_i is a metrizable topological space, there is a metric d_i on X_i inducing the topology on X_i . Define

$$d(x, y) := \max_{1 \leq i \leq n} \{d_i(x_i, y_i)\}, \quad x, y \in X. \quad (6.23)$$

Note that d is a metric on X : d is positive, since $d_i(x_i, y_i) \geq 0$, and thus $d(x, y) \geq 0$ for all $x, y \in X$. Further,

$$d(x, y) = 0 \Leftrightarrow \forall i \in \{1, \dots, n\} : d_i(x_i, y_i) = 0 \Leftrightarrow \forall i \in \{1, \dots, n\} : x_i = y_i \Leftrightarrow x = y,$$

i.e. d is definite. Clearly, d is also symmetric. Finally, for all $x, y, z \in X$, we have

$$\begin{aligned} d(x, z) &= \max_{1 \leq i \leq n} \{d_i(x_i, z_i)\} \leq \max_{1 \leq i \leq n} \{d_i(x_i, y_i) + d_i(y_i, z_i)\} \\ &\stackrel{(11.5)}{\leq} \max_{1 \leq i \leq n} \{d_i(x_i, y_i)\} + \max_{1 \leq i \leq n} \{d_i(y_i, z_i)\} = d(x, y) + d(y, z). \end{aligned}$$

We will now prove that the topology generated (induced) by d is the product topology on X .

For all $i \in \{1, \dots, n\}$, choose open $U_i \subset X_i$, then $U := \prod_{i=1}^n U_i$ is a basis member of the product topology on X . Since the U_i are open, by Theorem 6.44, for all $u_i \in U_i$ there is an open ball $B_{\epsilon_i}(u_i)$ that is contained in U_i . Define $\epsilon := \min_{1 \leq i \leq n} \epsilon_i > 0$, then for all $u \in U$ we have

$$B_\epsilon(u) = \{x \in X \mid d(x, u) < \epsilon\} = \left\{x \in X \mid \max_{1 \leq i \leq n} \{d_i(x_i, u_i)\} < \min_{1 \leq i \leq n} \{\epsilon_i\}\right\}$$

$$\stackrel{(\star)}{\subset} \prod_{i=1}^n B_{\epsilon_i}(u_i) \subset \prod_{i=1}^n U_i = U,$$

where (\star) holds, since $x \in B_\epsilon(u)$ implies that for all $i \in \{1, \dots, n\}$,

$$d_i(x_i, u_i) \leq \max_{1 \leq i \leq n} \{d_i(x_i, u_i)\} < \epsilon \leq \epsilon_i,$$

and thus for all i , $x_i \in B_{\epsilon_i}(u_i)$.

Conversely, consider a basis member $B_\epsilon(x)$ of the induced topology by d for some $x \in X$ and $\epsilon > 0$. Let $y \in \prod_{i=1}^n B_\epsilon(x_i)$, then for all i ,

$$d_i(x_i, y_i) < \epsilon \Rightarrow \max_{1 \leq i \leq n} \{d_i(x_i, y_i)\} < \epsilon \Rightarrow y \in B_\epsilon(x),$$

i.e. $x \in \prod_{i=1}^n B_\epsilon(x_i) \subset B_\epsilon(x)$, where $\prod_{i=1}^n B_\epsilon(x_i)$ is a basis member of the product topology on X , since each $B_\epsilon(x_i)$ is a basis member of X_i .

By Corollary 6.67, the induced topology by d on X and the product topology on X coincide. ■

Remark 6.92. In Theorem 6.91, we constructed the metric $d(x, y) := \max_{1 \leq i \leq n} \{d_i(x_i, y_i)\}$ on $X := \prod_{i=1}^n X_i$, where X_i is a metrizable topological space, and d_i a metric on X_i inducing the topology on X_i , and showed that it induces the product topology on X . Note that there are other metrics on X that are strongly – and thus topologically – equivalent to d [58], for example the L^p metrics for $p \in \mathbb{N} \cup \{\infty\}$:

$$d_p(x, y) := \left(\sum_{i=1}^n d_i(x_i, y_i)^p \right)^{1/p}, \quad x, y \in X,$$

where d_i is a metric on X_i , also cf. Example 4.3. For $p = \infty$, we obtain the maximum metric defined previously.

Theorem 6.93. Equip \mathbb{R} with its standard topology, let X be a metrizable topological space, i.e. there is a metric d inducing its topology, then the metric $d : X \times X \rightarrow \mathbb{R}$ is continuous, when considering the product topology on $X \times X$.

Proof [59, 60]. A basis for the standard topology of \mathbb{R} is given by open intervals (a, b) , where $a < b$. By Theorem 6.130, we need to show that the preimage of (a, b) is open in the product topology; note that

$$d^{-1}((a, b)) = \{(x, y) \in X \times X \mid d(x, y) \in (a, b)\} = \{(x, y) \in X \times X \mid a < d(x, y) < b\}.$$

By Corollary 6.45, we only need to show that there is a basis member B of the product topology of $X \times X$ s.t. for all $(x, y) \in d^{-1}((a, b))$, $(x, y) \in B \subset d^{-1}((a, b))$.

Let $(x, y) \in d^{-1}((a, b))$ be arbitrary, and define $\epsilon := \min\{d(x, y) - a, b - d(x, y)\}$. Now consider $z \in B_\epsilon(d(x, y))$, i.e. $|d(x, y) - z| < \epsilon$. Since for all $r \in \mathbb{R}$, $r \leq |r|$ and $-r \leq |r|$, we have $z \in (a, b)$, i.e. $B_\epsilon(d(x, y)) \subset (a, b)$.

Now consider $B_{\epsilon/2}(x) \times B_{\epsilon/2}(y)$, which is a basis member for the product topology on $X \times X$, and let $(x', y') \in B_{\epsilon/2}(x) \times B_{\epsilon/2}(y)$, then we have

$$d(x', y') \leq d(x', y) + d(y, y') \leq d(x, y) + d(x', x) + d(y, y') < d(x, y) + \epsilon$$

and

$$d(x, y) \leq d(x, y') + d(y', y) \leq d(x', y') + d(x', x) + d(y', y) < d(x', y') + \epsilon.$$

Thus, $|d(x, y) - d(x', y')| < \epsilon$, which implies $d(x', y') \in B_\epsilon(d(x, y)) \subset (a, b)$, i.e. $(x', y') \in d^{-1}((a, b))$, which shows that $B_{\epsilon/2}(x) \times B_{\epsilon/2}(y) \subset d^{-1}((a, b))$, and we also have $(x, y) \in B_{\epsilon/2}(x) \times B_{\epsilon/2}(y)$. ■

Let us now look at how to define a topology on the infinite Cartesian product of sets.

Theorem 6.94. Let A be an index set, and consider the collection $\{X_\alpha\}_{\alpha \in A}$, where X_α is a topological space for each $\alpha \in A$. Consider the set

$$\mathcal{B} := \left\{ \prod_{\alpha \in A} U_\alpha \mid U_\alpha \subset X_\alpha \text{ open} \right\}, \quad (6.24)$$

then it fulfills the requirements of Theorem 6.49.

Proof. Choose $U_\alpha = X_\alpha$ for every $\alpha \in A$, then $\prod_{\alpha \in A} X_\alpha = \prod_{\alpha \in A} U_\alpha$.

Now let $U_\alpha \subset X_\alpha$ and $V_\alpha \subset X_\alpha$ for all $\alpha \in A$, then $\prod_{\alpha \in A} U_\alpha \in \mathcal{B}$ and $\prod_{\alpha \in A} V_\alpha \in \mathcal{B}$. Since $(\prod_{\alpha \in A} U_\alpha) \cap (\prod_{\alpha \in A} V_\alpha) = \prod_{\alpha \in A} (U_\alpha \cap V_\alpha) \in \mathcal{B}$, cf. Theorem 8.22, we are done. ■

Definition 6.95 (Box topology [20]). Since \mathcal{B} from Eq. (6.24) fulfills the requirements of Theorem 6.49, there is a topology for which \mathcal{B} is a basis. This topology on $\prod_{\alpha \in A} X_\alpha$ is called the *box topology*. For $\beta \in A$, the map

$$p_\beta : \prod_{\alpha \in A} X_\alpha \rightarrow X_\beta, (x_\alpha)_{\alpha \in A} \mapsto x_\beta \quad (6.25)$$

is the *projection* onto the β -th factor, also cf. Definition 8.20 for the notation.

Remark 6.96. The projections p_β are continuous.

Proof. Let $U_\beta \subset X_\beta$ be open, then

$$\begin{aligned} p_\beta^{-1}(U_\beta) &= \left\{ (x_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} X_\alpha \mid p_\beta((x_\alpha)_{\alpha \in A}) \in U_\beta \right\} \\ &= \left\{ (x_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} X_\alpha \mid x_\beta \in U_\beta \right\} \\ &= \prod_{\alpha \in A} U_\alpha \in \mathcal{B}, \end{aligned}$$

where $U_\alpha = X_\alpha$ for $\alpha \neq \beta$, and $U_\alpha = U_\beta$ for $\alpha = \beta$. ■

Remark 6.97. If A is a finite index set, then the box topology is the product topology.

Remark 6.98. Note that if A is an infinite index set and we equip $\prod_{\alpha \in A} X_\alpha$ with the box topology, then if $p_\beta \circ f$ is continuous for some $\beta \in A$, where f maps from a topological space to $\prod_{\alpha \in A} X_\alpha$, this does not imply that f is continuous, unlike where A is finite, cf. Theorem 6.87.

Example 6.99 ([20]). Let $A = \mathbb{N}$ and $X_\alpha = \mathbb{R}$ for all $\alpha \in \mathbb{N}$. Now consider the box topology on $\prod_{\alpha \in \mathbb{N}} X_\alpha = \mathbb{R}^\mathbb{N}$. Now define $f : \mathbb{R} \rightarrow \mathbb{R}^\mathbb{N}, t \mapsto (t)_{n \in \mathbb{N}}$, where on \mathbb{R} we consider the standard topology. If for any $n \in \mathbb{N}$, $p_n : \mathbb{R}^\mathbb{N} \rightarrow \mathbb{R}$ is the projection map, then $p_n \circ f : \mathbb{R} \rightarrow \mathbb{R}$ is the identity map on \mathbb{R} , which is continuous. However, f is not continuous. To see this, define $B := \prod_{n \in \mathbb{N}} U_n$, where $U_n := (-1/n, +1/n)$. Clearly, B is open in $\mathbb{R}^\mathbb{N}$, yet

$$\begin{aligned} f^{-1}(B) &= \left\{ y \in \mathbb{R} \mid f(y) \in \prod_{n \in \mathbb{N}} U_n \right\} \\ &= \left\{ y \in \mathbb{R} \mid (y)_{n \in \mathbb{N}} \in \left\{ (u_n)_{n \in \mathbb{N}} \mid u_n \in \left(-\frac{1}{n}, \frac{1}{n} \right) \quad \forall n \in \mathbb{N} \right\} \right\} \\ &= \bigcap_{n \in \mathbb{N}} \left(-\frac{1}{n}, \frac{1}{n} \right) = \{0\}, \end{aligned}$$

which is not an open set in the standard topology of \mathbb{R} .

Remark 6.100. By Theorem 6.83, the product topology on a finite set is the smallest topology that makes the projections onto the coordinates spaces continuous. This forms the basis for the new definition of the product topology on $\prod_{\alpha \in A} X_\alpha$. Note that the continuity of the projection $p_\beta : \prod_{\alpha \in A} X_\alpha \rightarrow X_\beta$ requires $p_\beta^{-1}(U_\beta)$ to be open in $\prod_{\alpha \in A} X_\alpha$, where U_β is open in X_β , also cf. Remark 6.84.

Definition 6.101 (Tychonoff topology[20]). The topology on $\prod_{\alpha \in A} X_\alpha$ generated by the subbasis $\{p_\alpha^{-1}(U_\alpha) \mid U_\alpha \in \tau_\alpha, \alpha \in A\}$ is called the *product topology* (or *Tychonoff*¹⁰ *topology*). With this topology, $\prod_{\alpha \in A} X_\alpha$ is referred to as the product space.

Remark 6.102. The basis generated by the subbasis for the Tychonoff topology on $\prod_{\alpha \in A} X_\alpha$ consists of sets $\prod_{\alpha \in A} U_\alpha$, cf. Remark 6.46, where $U_\alpha = X_\alpha$ for all but a finite number of α 's, and $U_\alpha \in \tau_\alpha$ for all α . This holds because for $n \in \mathbb{N}$, we have

$$\bigcap_{i=1}^n p_{\alpha_i}^{-1}(U_{\alpha_i}) = \prod_{\alpha \in A} U_\alpha,$$

where $U_\alpha = X_\alpha$ for $\alpha \notin \{\alpha_1, \dots, \alpha_n\}$.

Remark 6.103. By Theorem 6.66, the box topology is finer than the Tychonoff topology.

Theorem 6.104. Given a family $X_\alpha, \alpha \in A$, the Tychonoff topology on $\prod_{\alpha \in A} X_\alpha$ is the smallest topology for which all projections $p_\beta : \prod_{\alpha \in A} X_\alpha \rightarrow X_\beta$ are continuous. Also, the projections are open maps.

¹⁰ANDREY NIKOLAYEVICH TIKHONOV

Proof. Let U_β be open in X_β , then $p_\beta^{-1}(U_\beta) = \prod_{\alpha \in A} U_\alpha$, where $U_\alpha = X_\alpha$ for $\alpha \neq \beta$, which is a basis member, and thus the p_β are continuous.

Suppose \mathcal{T} is a topology on $\prod_{\alpha \in A} X_\alpha$ s.t. the projections p_β are continuous. For $n \in \mathbb{N}$, let $U_{\alpha_1}, \dots, U_{\alpha_n}$ be open in X_{α_i} , and consider their intersection

$$\bigcap_{i=1}^n p_{\alpha_i}^{-1}(U_{\alpha_i}) = \prod_{\alpha \in A} U_\alpha,$$

where $U_\alpha = X_\alpha$ for $\alpha \notin \{\alpha_1, \dots, \alpha_n\}$. This implies that all basis members of the Tychonoff topology are contained in \mathcal{T} , and thus \mathcal{T} is finer than the Tychonoff topology, i.e. the Tychonoff topology is the smallest topology on $\prod_{\alpha \in A} X_\alpha$ s.t. the projections are continuous, cf. Theorem 6.66.

Finally, let U_α , $\alpha \in A$, be open in X_α , then $p_\beta(\prod_{\alpha \in A} U_\alpha) = U_\beta$, which is open, and thus the p_β are open maps. ■

Theorem 6.105. Given a family X_α , $\alpha \in A$, of topological spaces, if \mathcal{B}_α is a basis for the topology of X_α , then the collection \mathcal{B}' of sets of the form $\prod_{\alpha \in A} B_\alpha$, where $B_\alpha = X_\alpha$ for all but finitely many α 's and $B_\alpha \in \mathcal{B}_\alpha$ for the remaining indices, is a basis for the product topology on $\prod_{\alpha \in A} X_\alpha$.

Proof. From Remark 6.102, we know that the collection \mathcal{B} of sets of the form $\prod_{\alpha \in A} U_\alpha$, where $U_\alpha = X_\alpha$ for all but finitely many α 's, and U_α being open in X_α for all α , is a basis for the product topology on $\prod_{\alpha \in A} X_\alpha$. Since \mathcal{B}_α is a basis for X_α , by Theorem 6.44, for all $x_\alpha \in U_\alpha$, there exists a $B_\alpha \in \mathcal{B}_\alpha$ s.t. $x_\alpha \in B_\alpha \subset U_\alpha$. Thus, for all $x = (x_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} U_\alpha$, there exists a $\prod_{\alpha \in A} B_\alpha \in \mathcal{B}'$ s.t. $x \in \prod_{\alpha \in A} B_\alpha \subset \prod_{\alpha \in A} U_\alpha$.

Conversely, it is obvious that members of \mathcal{B}' are open in the product topology on $\prod_{\alpha \in A} X_\alpha$. Thus, by Corollary 6.67, \mathcal{B}' is also a basis for the product topology on $\prod_{\alpha \in A} X_\alpha$. ■

Theorem 6.106. Let X_α , $\alpha \in A$, be a family of topological spaces, and let Y be a topological space. A function $f : Y \rightarrow \prod_{\alpha \in A} X_\alpha$, where $\prod_{\alpha \in A} X_\alpha$ is equipped with the product topology, is continuous iff $p_\alpha \circ f : Y \rightarrow X_\alpha$ is continuous, where p_α is the projection onto the α -th factor, cf. Eq. (6.25). This property characterizes the product topology for $\prod_{\alpha \in A} X_\alpha$.

Proof. If f is continuous, so is $p_\alpha \circ f$, since p_α is continuous, cf. Remark 6.96.

Conversely, assume that $p_\alpha \circ f$ is continuous, then for any open $U_\alpha \subset X_\alpha$, $(p_\alpha \circ f)^{-1}(U_\alpha)$ is open in Y . Since $(p_\alpha \circ f)^{-1}(U_\alpha) = f^{-1}(p_\alpha^{-1}(U_\alpha))$, and p_α is continuous, i.e. $p_\alpha^{-1}(U_\alpha)$ is open in $\prod_{\alpha \in A} X_\alpha$, this implies that f is continuous, as $p_\alpha^{-1}(U_\alpha)$ is a subbasis for the product topology on $\prod_{\alpha \in A} X_\alpha$, also cf. Theorem 8.19.

To see that $p_\alpha \circ f$ continuous iff f is continuous characterizes the product topology for $\prod_{\alpha \in A} X_\alpha$, define $X := \prod_{\alpha \in A} X_\alpha$ and denote the product topology on X by \mathcal{T} . Let \mathcal{T}' be another topology on X s.t. $f : Y \rightarrow (X, \mathcal{T}')$ is continuous iff $p_\alpha \circ f : Y \rightarrow X_\alpha$ is continuous. Since the identity $\iota : (X, \mathcal{T}') \rightarrow (X, \mathcal{T})$ is continuous, $p_\alpha \circ \iota = p_\alpha : (X, \mathcal{T}') \rightarrow X_\alpha$ is continuous.

Let $i : (X, \mathcal{T}) \rightarrow (X, \mathcal{T}')$ be the identity map, which is bijective, then the compositions

$$(X, \mathcal{T}) \xrightarrow{i} (X, \mathcal{T}') \xrightarrow{p_\alpha} X_\alpha \quad \text{and} \quad (X, \mathcal{T}') \xrightarrow{i^{-1}} (X, \mathcal{T}) \xrightarrow{p_\alpha} X_\alpha$$

are continuous. To see this, let U_α be open in X_α , then

$$(p_\alpha \circ i)^{-1}(U_\alpha) = i^{-1}(p_\alpha^{-1}(U_\alpha)) = p_\alpha^{-1}(U_\alpha) \in \mathcal{T}.$$

Also,

$$(p_\alpha \circ i^{-1})^{-1}(U_\alpha) = i(p_\alpha(U_\alpha)) = p_\alpha(U_\alpha) \in \mathcal{T}'.$$

Thus, i and i^{-1} are continuous, i.e. i is a homeomorphism. By Theorem 6.139, this implies that $\mathcal{T} = \mathcal{T}'$. \blacksquare

Theorem 6.107. For all $n \in \mathbb{N}$, let (X_n, τ_n) be metrizable topological spaces. Then $\prod_{i \in \mathbb{N}} X_i$, endowed with the product topology, is metrizable.

Proof. [61, 62] First, we construct a metric on $X := \prod_{i \in \mathbb{N}} X_i$. Since each X_i is a metrizable topological space, there is a metric d_i on X_i inducing the topology on X_i . For all $x, y \in X$, define

$$d(x, y) := \sum_{n \in \mathbb{N}} \frac{d_n(x_n, y_n)}{2^n (1 + d_n(x_n, y_n))}.$$

From Theorem 5.3, we know that $\rho_n(x_n, y_n) := d_n(x_n, y_n)/(1 + d_n(x_n, y_n))$ is a metric bounded above by 1, which is topologically equivalent to d_n , cf. Theorem 6.76.

Thus, instead of considering $d(x, y)$ as defined above, we can consider

$$\rho(x, y) := \sum_{n \in \mathbb{N}} \frac{\rho_n(x_n, y_n)}{2^n}. \quad (6.26)$$

Clearly, this definition is well-defined, since the series converges absolutely, as $\rho_n < 1$ for all n .

To see that ρ is a metric, note that ρ is clearly positive, definite and symmetric. For the triangle inequality, note that for all $x, y, z \in X$, we have

$$\begin{aligned} \rho(x, z) &= \sum_{n=1}^{\infty} \frac{\rho_n(x_n, z_n)}{2^n} \leq \sum_{n=1}^{\infty} \frac{\rho_n(x_n, y_n) + \rho_n(y_n, z_n)}{2^n} \\ &= \sum_{n=1}^{\infty} \frac{\rho_n(x_n, y_n)}{2^n} + \sum_{n=1}^{\infty} \frac{\rho_n(y_n, z_n)}{2^n} = \rho(x, y) + \rho(y, z) \end{aligned}$$

cf. Ref. [63], which shows the triangle inequality, and thus ρ is a metric.

We will now prove that the topology generated (induced) by ρ is the product topology on X .

Let U be a member of the basis of the product topology, i.e. $U = \prod_{n \in \mathbb{N}} U_n$, where all $U_n \subset X_n$ are open in X_n and where we have a finite set $F \subset \mathbb{N}$ s.t. $n \notin F$ if $U_n = X_n$. We want to show that for any $x \in U$, there is an open ball $B_r^\rho(x)$ s.t. $B_r^\rho(x) \subset U$. For every $n \in F$, we have $x_n \in U_n$, and so there is $r_n > 0$ s.t. $B_{r_n}^{\rho_n}(x_n) \subset U_n$, since ρ_n induces the topology on X_n . As F is finite, we can find an $r \in (0, 1)$ s.t. $r \leq r_n/2^n$ for all $n \in F$.

Now take any $y \in B_r^\rho(x)$, i.e. $\rho(x, y) < r$. For $n \in F$, we know that

$$\frac{\rho_n(x_n, y_n)}{2^n} \stackrel{(6.26)}{\leq} \rho(x, y) < r \leq \frac{r_n}{2^n} \Rightarrow \rho_n(x_n, y_n) < r_n \Rightarrow y_n \in B_{r_n}^{\rho_n}(x_n) \subset U_n.$$

Thus, for all $n \in F$, $y_n \in U_n$, and for all $n \notin F$, $y_n \in X_n$, implying that $y \in U$, i.e. $B_r^\rho(x) \subset U$.

We also want to show that if we consider $B_r^\rho(x)$ for an arbitrary $r > 0$ and $x \in X$, then there is a basis member U of the product topology on X s.t. $x \in U \subset B_r^\rho(x)$. For this, choose an $n \in \mathbb{N}$ s.t. $1/2^N < r/2$. For $1 \leq k \leq N$, define $U_k := B_{r/(2N)}^{\rho_k}(x_k) \subset X_k$, and for

$k > N$, define $U_k := X_k$. Setting $U := \prod_{k \in \mathbb{N}} U_k$, we will now show that $x \in U \subset B_r^\rho(x)$. Clearly, $x \in U$. Now, if $y \in U$, then for $1 \leq k \leq N$, we have

$$\rho_k(x_k, y_k) < \frac{r}{2N} \Rightarrow \sum_{k=1}^N \frac{\rho_k(x_k, y_k)}{2^k} \leq \sum_{k=1}^N \rho_k(x_k, y_k) < \frac{Nr}{2N} = \frac{r}{2}, \quad (6.27)$$

and for $k > N$, we have

$$\sum_{k=N+1}^{\infty} \frac{\rho_k(x_k, y_k)}{2^k} < \sum_{k=N+1}^{\infty} \frac{1}{2^k} = \sum_{k=1}^{\infty} \frac{1}{2^k} - \sum_{k=1}^N \frac{1}{2^k} = 1 - \left(1 - \frac{1}{2^N}\right) = \frac{1}{2^N} < \frac{r}{2}, \quad (6.28)$$

and thus

$$\rho(x, y) = \sum_{k=1}^{\infty} \frac{\rho_k(x_k, y_k)}{2^k} = \sum_{k=1}^N \frac{\rho_k(x_k, y_k)}{2^k} + \sum_{k=N+1}^{\infty} \frac{\rho_k(x_k, y_k)}{2^k} \stackrel{(6.27), (6.28)}{<} \frac{r}{2} + \frac{r}{2} < r,$$

i.e. $y \in B_r^\rho(x)$.

Thus, by Corollary 6.67, ρ induces the product topology on X . ■

Remark 6.108. The uncountable product of metrizable topological spaces is not necessarily metrizable, since every metrizable topological space is first-countable, yet $\mathbb{R}^{\mathbb{R}}$, equipped with the product topology, and where we consider \mathbb{R} with the standard topology, is not first-countable, since $\{0, 1\}^{\mathbb{R}}$, where we consider $\{0, 1\} \subset \mathbb{R}$ with the subspace topology, is not first-countable.

Theorem 6.109 (Heine-Borel Theorem). Let \mathbb{R} be equipped with the standard topology. Then a closed interval $J = [a, b] \subset \mathbb{R}$ with $a \leq b$ is compact.

Proof [64–66]. Let \mathcal{G} be a family of open subsets of \mathbb{R} covering J , and consider the set

$$X := \{x \in J \mid [a, x] \text{ is covered by a finite subfamily of } \mathcal{G}\}.$$

We want to show that $b \in X$.

First, note that $a \in X$, as $[a, a] = \{a\}$, which can be covered by a single member of \mathcal{G} . Thus, X is non-empty. X is bounded above by b , hence $c := \sup X$ exists. As $a \leq c \leq b$, there exists a set $G_0 \in \mathcal{G}$ s.t. $c \in G_0$. Since G_0 is open in \mathbb{R} , there is a $\delta > 0$ s.t. $(c - \delta, c + \delta) \subset G_0$.

Since $c = \sup X$, by Theorem 8.14, there exists an $x \in X$ s.t. $c \geq x > c - \delta$. Thus, $[a, c + \delta) = [a, x] \cup (c - \delta, c + \delta)$ can be covered by a finite subfamily of \mathcal{G} , as $[a, x]$ can and $(c - \delta, c + \delta) \subset G_0 \in \mathcal{G}$. Thus, for any $x' \in [a, c + \delta) \cap [a, b]$, we have $x' \in X$.

It holds that $c = b$, which we will prove via contradiction. It is clear that the possibility $c > b$ is excluded, as b is an upper bound of X and c is the lowest upper bound of X . Assume that $a \leq c < b$. Then we can find a $t \in \mathbb{R}$ s.t. $c < t < b$ and $t < c + \delta$, i.e. t is one of the x' values, implying that $t \in X$. This is a contradiction, as $t > \sup X = c$. This shows that $c = b$, i.e. $c = b$ is one of the x' values, hence $b \in X$. ■

Theorem 6.110. Let X and Y be compact topological spaces, then so is $X \times Y$.

Proof [20]. Let \mathcal{G} be an open cover of $X \times Y$. Fix an arbitrary $x \in X$, then for any $y \in Y$, there is an open set $G \in \mathcal{G}$ s.t. $(x, y) \in G$. Since G is open, there are open sets

$U_y \subset X$ and $V_y \subset Y$ s.t. $(x, y) \in U_y \times V_y \subset G$, cf. Corollary 6.45. Note that U_y and V_y also depend on x , yet since x is fixed, we only denote the dependence on y .

Since Y is compact, the open covering $\{V_y \mid y \in Y\}$ has a finite subcovering of Y , i.e. there are finitely many points $y_1, \dots, y_{n(x)}$ in Y s.t. $Y = \bigcup_{i=1}^{n(x)} V_{y_i}$, where $n(x) \in \mathbb{N}$. Define $U_x := \bigcap_{i=1}^{n(x)} U_{y_i}$ and note that U_x is an open neighborhood of x , since it is the intersection of finitely many open sets, and each U_{y_i} contains x .

For $(x, y_i) \in U_{y_i} \times V_{y_i}$, there exists a $G_x^i \in \mathcal{G}$ s.t. $(x, y_i) \in U_{y_i} \times V_{y_i} \subset G_x^i$. Note that

$$U_x \times V_{y_i} \subset U_{y_i} \times V_{y_i} \subset G_x^i \Rightarrow \bigcup_{i=1}^{n(x)} (U_x \times V_{y_i}) \stackrel{(8.17)}{=} U_x \times \bigcup_{i=1}^{n(x)} V_{y_i} = U_x \times Y \subset \bigcup_{i=1}^{n(x)} G_x^i. \quad (6.29)$$

Now consider the open covering $\{U_x \mid x \in X\}$ of X . Since X is compact, there are finitely many points x_1, \dots, x_m in X s.t. $X = \bigcup_{j=1}^m U_{x_j}$, where $m \in \mathbb{N}$. Thus,

$$X \times Y = \left(\bigcup_{j=1}^m U_{x_j} \right) \times Y \stackrel{(8.17)}{=} \bigcup_{j=1}^m (U_{x_j} \times Y) \stackrel{(6.29)}{=} \bigcup_{j=1}^m \bigcup_{i=1}^{n(x)} G_{x_j}^i,$$

where equality holds, since $G_{x_j}^i \subset X \times Y$ and their union can thus not be a superset of $X \times Y$. We have thus shown that each open cover of $X \times Y$ has a finite sub-covering. ■

Corollary 6.111. By induction, it follows that a finite product of compact topological spaces is also a compact topological space.

Remark 6.112. A visual animation for the proof of the preceding Theorem is given in Ref. [67].

Example 6.113. From the Heine-Borel Theorem 6.109, we already know that $I^1 = [0, 1]$ is compact. The preceding corollary implies that for any $n \in \mathbb{N}$, the unit n -cube $I^n := \prod_{i=1}^n I^1$ is also compact.

Theorem 6.114. Let X be a compact topological space and $U \subset X$ closed. Then U is also compact.

Proof [68]. Let $\mathcal{G} := \{U_\alpha\}_{\alpha \in A}$ be an open cover of U , where A is an index set and $U_\alpha \subset X$ for every α . Since \mathcal{G} is an open cover of U , $\mathcal{G} \cup \{X \setminus U\}$ is an open cover of X (note that $X \setminus U$ is open, since U is closed by assumption). Since X is compact, there exist finitely many indices $\alpha_1, \dots, \alpha_n \in A$ with $n \in \mathbb{N}$ s.t. $\{U_{\alpha_1}, \dots, U_{\alpha_n}, X \setminus U\}$ covers X , which implies that $\{U_{\alpha_1}, \dots, U_{\alpha_n}\}$ is a cover of U . ■

Theorem 6.115 (Generalized Heine-Borel Theorem). A subset $A \subset \mathbb{R}^n$ is compact iff it is closed and bounded.

Proof [20]. “ \Leftarrow ” Let $A \subset \mathbb{R}^n$ be closed and bounded. For each $i \in \{1, \dots, n\}$, let $p_i : \mathbb{R}^n \rightarrow \mathbb{R}$ denote the projection onto the i th factor. Then $p_i(A) \subset \mathbb{R}$ is bounded. To see this, note that since A is bounded, there is an $r > 0$ and a $\varphi = (\varphi_1, \dots, \varphi_n) \in \mathbb{R}^n$ s.t. $A \subset B_r[\varphi] = \left\{ \psi = (\psi_1, \dots, \psi_n) \in \mathbb{R}^n \mid \sqrt{\sum_{i=1}^n (\varphi_i - \psi_i)^2} \leq r \right\}$, where we used the

fact that the product topology on \mathbb{R}^n is induced by the Euclidean metric. Also, for each $i \in \{1, \dots, n\}$,

$$\begin{aligned} p_i(A) &= \{p_i(x) \mid x = (x_1, \dots, x_n) \in A\} \\ &= \{x_i \mid x \in A\} \\ &\subset \{x_i \mid x \in B_r[\varphi]\} \\ &= \left\{ x_i \mid \sqrt{\sum_{i=1}^n (\varphi_i - x_i)^2} \leq r \right\} \\ &\subset B_r[\varphi_i] = \{y \in \mathbb{R} \mid |y - \varphi_i| \leq r\}, \end{aligned}$$

since $|x - \varphi_i| = (x - \varphi_i)^2 \leq \sqrt{\sum_{i=1}^n (\varphi_i - x)^2} \leq r$. Thus, $p_i(A)$ is bounded as well. Defining $a_i := \varphi_i - r$ and $b_i := \varphi_i + r$, we have $p_i(A) \subset [a_i, b_i]$. It now holds that $A \subset \prod_{i=1}^n p_i(A)$, since for any $x = (x_1, \dots, x_n) \in A$, we have $x_i \in p_i(A)$ for each i in $\{1, \dots, n\}$. Thus, $A \subset \prod_{i=1}^n p_i(A) \subset \prod_{i=1}^n [a_i, b_i] =: X$. By Theorem 6.109 and 6.110, X is compact.

We now show that A is also closed in X , equipped with the subspace topology. Since $A \subset \mathbb{R}^n$ is closed, $\mathbb{R}^n \setminus A$ is open. Thus, $X \cap (\mathbb{R}^n \setminus A)$ is open in the subspace topology of X . Since $X \cap (\mathbb{R}^n \setminus A) = X \setminus A$, $A \subset X$ is closed. To see why $X \cap (\mathbb{R}^n \setminus A) = X \setminus A$, let $x \in X \cap (\mathbb{R}^n \setminus A)$, i.e. $x \in X$ and $x \in \mathbb{R}^n$ and $x \notin A$, which implies $x \in X \setminus A$.

Since $A \subset X$ is closed and X is compact, by Theorem 6.114, A is compact.

“ \implies ” Let $A \subset \mathbb{R}^n$ be compact. Since \mathbb{R}^n is a Hausdorff space, A is closed, cf. Theorem 6.122. Since A is compact, every open cover of A contains a finite sub-covering. Fix an $a \in A$ and consider the open covering $\{B_m(a) \mid m \in \mathbb{N}\}$ of A , i.e. the open covering consists of open balls around a with varying radii $m \in \mathbb{N}$. Since A is compact, there is a finite sub-covering, and since $B_m(a) \cup B_n(a) = B_n(a)$ if $n \geq m$, there is an $N \in \mathbb{N}$ s.t. $A \subset B_N(a) \subset B_N[a]$, showing that A is bounded. ■

Definition 6.116. Let (X, τ) be a topological space, then a sequence $(\varphi_n)_{n \in \mathbb{N}}$ is said to converge to a point $\varphi \in X$ if for each neighborhood U of x , there exists an $N \in \mathbb{N}$ s.t. for all $n \geq N$, we have $\varphi_n \in U$ [69].

Remark 6.117. Unlike metric spaces, the limit point of a converging sequence in topological spaces is not guaranteed to be unique.

Example 6.118. Consider $X = \mathbb{R}$ and the topology $\tau = \{\mathbb{R}, \emptyset\}$. Then a sequence $(\varphi_n)_{n \in \mathbb{N}}$ in \mathbb{R} converges to any point in \mathbb{R} , since the neighborhood of any $x \in \mathbb{R}$ is \mathbb{R} .

6.1 Hausdorff Spaces

Definition 6.119 (Hausdorff). A topological space (X, τ) is called *Hausdorff* if for all $p, q \in X$ there exist $U, V \in \tau$ s.t. $p \in U$, $q \in V$ and $U \cap V = \emptyset$.

Remark 6.120. Note that U is an open neighborhood of $p \in X$ and V an open neighborhood of $q \in X$.

Theorem 6.121. All metric spaces are Hausdorff spaces.




Figure 5: Visualization for why a metric space is a Hausdorff space.

Proof. A visualization is shown in Fig. 5. To prove this more rigorously, define $U := B_r(p)$ and $V := B_r(q)$ with radius $r := \frac{d(p,q)}{2}$, where from Theorem 5.14 we know that U and V are open. Suppose $U \cap V = \emptyset$ would not hold. Then there exists a $z \in U \cap V$ with

$$d(p, z) < \frac{d(p, q)}{2} \quad (6.30)$$

and

$$d(q, z) < \frac{d(p, q)}{2}. \quad (6.31)$$

Therefore, by the triangle inequality for metric spaces, we have:

$$d(p, q) \leq d(p, z) + d(q, z) < d(p, q), \quad (6.32)$$

which is clearly a contradiction. ■

Theorem 6.122. Let X be a Hausdorff space, and $K \subset X$ compact, then K is closed.

Proof [20, 70]. Fix $x \in X \setminus K$, then for each $y \in K$, there exist open sets $U_y, V_y \subset X$ s.t. $x \in U_y$ and $y \in V_y$ with $U_y \cap V_y = \emptyset$, since $x \neq y$. Clearly, the family $\{V_y \mid y \in K\}$ of subsets of X is an open covering of K , and since K is compact, there exist finitely many points y_1, \dots, y_n with $n \in \mathbb{N}$ and $y_k \in K$ for all $k \in \{1, \dots, n\}$ s.t. $K \subset \bigcup_{k=1}^n V_{y_k} =: V$. Define $U := \bigcap_{k=1}^n U_{y_k}$ and note that $x \in U$.

We now show that $U \cap V \neq \emptyset$ via contradiction. Assume there is a $z \in U \cap V$, i.e. $z \in U$ and $z \in V$. This implies that $z \in U_{y_k}$ for all $k \in \{1, \dots, n\}$ and that there is a $k' \in \{1, \dots, n\}$ s.t. $z \in V_{y_{k'}}$. However, this is a contradiction to $U_{y_{k'}} \cap V_{y_{k'}} = \emptyset$.

Since $U \cap V \neq \emptyset$, $x \in U \subset X \setminus V \subset X \setminus K$, and since $x \in X \setminus K$ was arbitrary, by Theorem 6.14, $X \setminus K$ is open, implying that K is closed. ■

Definition 6.123. Let (X, τ) be a topological space. Then it is said to be *second countable* if τ has a countable basis.

Example 6.124. For any $n > 0$, the topological spaces \mathbb{R}^n are second countable.

Remark 6.125. Metric spaces are not automatically second countable. For example, take an uncountable set X , endow it with the discrete metric and because of Example 6.57, 6.55, the topological space X is not second countable, since X is uncountable.

6.2 Continuity and Homeomorphisms

In topological spaces (which are not necessarily metrizable), we define continuity as follows [20]:

Definition 6.126. Let (X, τ_X) and (Y, τ_Y) be topological spaces. Then a function $f : X \rightarrow Y$ is called continuous on X if for every $U \in \tau_Y$, $f^{-1}(U) := \{x \in X \mid f(x) \in U\} \in \tau_X$.

Example 6.127. Let (X, τ_X) be a topological space, then the identity $\text{id} : X \rightarrow X$ is continuous.

Proof. Let $U \in \tau_X$, then $f^{-1}(U) = U \in \tau_X$. ■

Remark 6.128. In other words, the preimage of an open set is always open. For topological spaces that are metrizable, the ϵ - δ definition of continuity as in Def. 5.123 is equivalent to the preceding definition, cf. Theorem 5.132.

Theorem 6.129. Let (X, τ_X) and (Y, τ_Y) be topological spaces and let $A \subset X$ be equipped with the subspace topology. If $f : X \rightarrow Y$ is continuous, then so is $f|_A : A \rightarrow Y$.

Proof [71]. Let $U \in \tau_Y$, then

$$f|_A^{-1}(U) = \{a \in A \mid f(a) \in U\} = \{x \in X \mid f(x) \in U\} \cap A = f^{-1}(U) \cap A.$$

Since $f : X \rightarrow Y$ is continuous, this implies that $f^{-1}(U)$ is open in X and thus $f^{-1}(U) \cap A$ is open in the subspace topology of A . ■

Theorem 6.130. Let (X, τ_X) and (Y, τ_Y) be topological spaces, and \mathcal{B} a basis or a subbasis for τ_Y , then $f : X \rightarrow Y$ is continuous iff the inverse image under f of every member of \mathcal{B} is open in τ_X [20].

Proof. Let \mathcal{B} be a basis, then if f is continuous, $f^{-1}(B) \in \tau_X$ for all $B \in \mathcal{B}$. Conversely, every open set in τ_Y can be written as the union of basis members, i.e. $V = \bigcup_{i \in I} B_i$, where I is an index set, and by Theorem 8.19,

$$f^{-1}(V) = f^{-1}\left(\bigcup_{i \in I} B_i\right) = \bigcup_{i \in I} f^{-1}(B_i) \in \tau_X,$$

since $f^{-1}(B_i) \in \tau_X$ for all $B_i \in \mathcal{B}$.

If \mathcal{B} is a subbasis for τ_Y , and f is continuous, then $f^{-1}(B) \in \tau_X$ for all $B \in \mathcal{B}$, since $\mathcal{B} \subset \tau_Y$. Conversely, let $f^{-1}(B) \in \tau_X$ for all $B \in \mathcal{B}$. We know that each $V \in \tau_Y$ can be written as the union of a finite intersection of members of \mathcal{B} , i.e. $V = \bigcup_{i \in I} \left(\bigcap_{j \in J} B_{i,j}\right)$, where J is a finite index set and I an index set, and by Theorem 8.19,

$$f^{-1}(V) = f^{-1}\left(\bigcup_{i \in I} \left(\bigcap_{j \in J} B_{i,j}\right)\right) = \bigcup_{i \in I} \left(\bigcap_{j \in J} f^{-1}(B_{i,j})\right) \in \tau_X,$$

since $f^{-1}(B_{i,j}) \in \tau_X$ for all $B_{i,j} \in \mathcal{B}$. ■

Example 6.131. Consider the function $f : [0, 2\pi) \rightarrow \mathbb{S}^1, t \mapsto e^{it}$, where

$$\mathbb{S}^1 := \{e^{it} \mid t \in [0, 2\pi)\} = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}$$

is the unit circle in \mathbb{R}^2 . Let $[0, 2\pi)$ be equipped with the order topology, cf. Definition 6.35, and $\mathbb{S}^1 \subset \mathbb{R}^2$ be equipped with the subspace topology, cf. Theorem 6.26, where \mathbb{R}^2 is equipped with the Euclidean topology.

Proof. By Theorem 6.43, the collection of all open arcs of \mathbb{S}^1 is a basis for the subspace topology on \mathbb{S}^1 , since any open arc is the intersection of the full circle with an open ball. Figure 6 visualizes this.




Figure 6: Every open arc is the intersection of \mathbb{S}^1 with an open ball.

Let $G \subset \mathbb{S}^1$ be an open arc that does not contain $1 \in \mathbb{S}^1$, then $f^{-1}(G) = (a, b)$, where $0 < a < b < 2\pi$.

If G does contain $1 \in \mathbb{S}^1$, then $f^{-1}(G) = [0, a) \cup (b, 2\pi)$, where $0 < a < b < 2\pi$. Thus, regardless of whether G contains 1, $f^{-1}(G)$ is open in $[0, 2\pi)$ – note that $f^{-1}(G)$ is open in \mathbb{R} – and thus f is continuous by Theorem 6.130. ■

Theorem 6.132. Let (X, τ_X) , (Y, τ_Y) and (Z, τ_Z) be topological spaces, and $f : X \rightarrow Y$, $g : Y \rightarrow Z$ be continuous, then $g \circ f : X \rightarrow Z$ is continuous as well.

Proof. Let $W \in \tau_Z$, then $(g \circ f)^{-1}(W) = (f^{-1} \circ g^{-1})(W)$, and since $g^{-1}(W) \in \tau_Y$, this implies that $(f^{-1} \circ g^{-1})(W) \in \tau_X$. ■

Theorem 6.133. Let X be a compact topological space, Y a topological space and $f : X \rightarrow Y$ a continuous function, then $f(X)$ is a compact subspace of Y .

Proof [72]. Let $\{U_i\}_{i \in I}$ be an open cover of $f(X)$. Since f is continuous, $f^{-1}(U_i)$ is open in X for every $i \in I$. Note that $\{f^{-1}(U_i)\}_{i \in I}$ is an open cover of X , since

$$f(X) \subset \bigcup_{i \in I} U_i \Rightarrow X \subset f^{-1}(f(X)) \subset f^{-1}\left(\bigcup_{i \in I} U_i\right) = \bigcup_{i \in I} f^{-1}(U_i),$$

cf. Eq. (8.16). Since X is compact, there is a finite subcover $\{f^{-1}(U_j)\}_{j \in J}$ of X , where $J \subset I$ is finite. Thus,

$$X = \bigcup_{j \in J} f^{-1}(U_j) \Rightarrow f(X) = f\left(\bigcup_{j \in J} f^{-1}(U_j)\right) = \bigcup_{j \in J} f(f^{-1}(U_j)) \subset \bigcup_{j \in J} U_j,$$

cf. Eq. (8.15), which shows that every open cover of $f(X)$ has a finite subcover. ■

Corollary 6.134. Let X be a compact topological space, equip \mathbb{R} with the standard topology and let $f : X \rightarrow \mathbb{R}$ be continuous. Then f attains its minimum and maximum on X .

Proof [73]. Since f is continuous and X compact, $f(X) \subset \mathbb{R}$ is compact, and thus by Theorem 6.115 closed and bounded. Corollary 5.27 now implies that $\sup(f(X)), \inf(f(x)) \in f(X)$ and $\sup(f(X)) = \max(f(X))$ as well as $\inf(f(X)) = \min(f(X))$. Thus, there exist $x_0, x_1 \in X$ s.t. $\min(f(X)) = f(x_0)$ and $\max(f(X)) = f(x_1)$. ■

Definition 6.135 (Homeomorphism [20]). A *homeomorphism* between two topological spaces X and Y is an invertible function $f : X \rightarrow Y$ such that f and f^{-1} are continuous.

Example 6.136. The Euclidean space \mathbb{R}^n , equipped with the usual topology, is homeomorphic to the open ball $B_r(\varphi) = \{x \in \mathbb{R}^n \mid \|x - \varphi\| < r\} \subset \mathbb{R}^n$ (consider the open ball as a metric space with the *induced metric* from the whole space of \mathbb{R}^n and then equip it with the usual topology on a metric space given by the open sets).

Proof. Consider the map

$$f : \mathbb{R}^n \rightarrow B_r(\varphi), \quad x \mapsto \frac{r \cdot (x - \varphi)}{1 + \|x - \varphi\|}.$$

Obviously, f is continuous with inverse

$$f^{-1} : B_r(\varphi) \rightarrow \mathbb{R}^n, \quad x \mapsto \frac{x}{r - \|x\|} + \varphi,$$

which is also continuous. One can easily show that f and f^{-1} are inverses to each other. Thus, f describes a homeomorphism. ■

Example 6.137 (Stereographic projection). The unit sphere S^n , embedded in \mathbb{R}^{n+1} , without the north pole, i.e. $S^n \setminus \{p\} \subset \mathbb{R}^{n+1}$, where $S^n := \{x \in \mathbb{R}^{n+1} \mid \|x\|_2 = 1\}$ and $p := \{x \in \mathbb{R}^{n+1} \mid x_i = 0 \ \forall i \in [1, n], x_{n+1} = 1\}$, is homeomorphic to \mathbb{R}^n (consider the unit sphere as a metric space with the *induced metric* from the whole space of \mathbb{R}^{n+1} and then equip it with the usual topology on a metric space given by the open sets).

Proof. Consider the map

$$f : S^n \setminus \{p\} \rightarrow \mathbb{R}^n, \quad x \mapsto \frac{1}{1 - x_{n+1}} (x_1, \dots, x_n)^T. \quad (6.33)$$

Obviously, f is continuous. Its inverse is given by

$$f^{-1} : \mathbb{R}^n \rightarrow S^n \setminus \{p\}, \quad x \mapsto \begin{pmatrix} 2x_1/(1 + \|x\|_2^2) \\ \vdots \\ 2x_n/(1 + \|x\|_2^2) \\ 1 - 2/(1 + \|x\|_2^2) \end{pmatrix}. \quad (6.34)$$

It is trivial to show that f and f^{-1} are inverses to each other. One can also easily show that $\|f^{-1}(x)\|_2 = 1$; thus, f^{-1} does indeed bring us to the unit ball S^n . To see that f^{-1} is continuous, note that all components are continuous and thus, according to Lemma 5.131, the function itself is continuous. ■

Theorem 6.138. Let (X, τ_X) , (Y, τ_Y) and (Z, τ_Z) be topological spaces, and $f : X \rightarrow Y$, $g : Y \rightarrow Z$ be homeomorphisms, then $g \circ f : X \rightarrow Z$ is a homeomorphism.

Proof. The composition of bijective functions is bijective again. From Theorem 6.132, we know that $g \circ f$ is continuous. To show the continuity of $(g \circ f)^{-1} : Z \rightarrow X$, note that if $U \in \tau_X$, then $(g \circ f)(U) = g(f(U))$, and since f and g are homeomorphisms, we have $f(U) \in \tau_Y$, and thus $g(f(U)) \in \tau_Z$. ■

Theorem 6.139. Let (X, τ_1) and (X, τ_2) be topological spaces. The identity map $\iota : (X, \tau_1) \rightarrow (X, \tau_2), x \mapsto x$ is a homeomorphism iff $\tau_1 = \tau_2$.

Proof [74]. Let $\iota : (X, \tau_1) \rightarrow (X, \tau_2), x \mapsto x$ be a homeomorphism. We know that τ_1 is a base generating itself, as is τ_2 a base generating itself. Consider $U \in \tau_2$, then for all $x \in U$, $x \in \iota^{-1}(U) = U \in \tau_1$, since ι is continuous. Also, for any $U \in \tau_1$ and $x \in U$, $x \in \iota(U) = U \in \tau_2$, since ι^{-1} is continuous. Thus, by Corollary 6.67, $\tau_1 = \tau_2$.

Conversely, let $\tau_1 = \tau_2$, and consider $\iota : (X, \tau_1) \rightarrow (X, \tau_2), x \mapsto x$. Let $U \in \tau_2$, then $\iota^{-1}(U) = U \in \tau_1 = \tau_2$, and thus ι is continuous. Let $U \in \tau_1$, then $\iota(U) = U \in \tau_2 = \tau_1$, and thus also ι^{-1} is continuous. Further, ι is bijective, implying that ι is a homeomorphism. ■

7 Linear Functional Analysis

7.1 Bounded Linear Operators

Let us start by defining the notions of *linearity* and *boundedness* of operators (functionals).

Definition 7.1. Let X, Y be normed linear spaces. Then an operator $A : X \rightarrow Y$ is called *linear* if

$$A(\alpha\varphi + \beta\psi) = \alpha A(\varphi) + \beta A(\psi) \quad \forall \varphi, \psi \in X, \forall \alpha, \beta \in \mathbb{K}. \quad (7.1)$$

Theorem 7.2. A linear operator $A : X \rightarrow Y$ from a normed linear space $(X, \|\cdot\|_X)$ into another normed linear space $(Y, \|\cdot\|_Y)$ is continuous on X iff it is continuous at a single point in X .

Proof. “ \implies ” By definition.

“ \impliedby ” Let A be continuous at $\varphi_0 \in X$, and let $(\varphi_n)_{n \in \mathbb{N}} \subset X$ converge to $\varphi \in X$ wrt $\|\cdot\|_X$, then

$$A(\varphi_n) = A(\varphi_n + \varphi_0 - \varphi) = A(\varphi_0 - \varphi) \xrightarrow{n \rightarrow \infty} A(\varphi) \quad (7.2)$$

wrt $\|\cdot\|_Y$. ■

Definition 7.3. Let $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ be normed linear spaces. Then an operator $A : X \rightarrow Y$ is called *bounded* if there is a constant $C(A) > 0$ s.t.

$$\|A\varphi\|_Y \leq C \|\varphi\|_X \quad \forall \varphi \in X. \quad (7.3)$$

Any such constant C is called an *upper bound* for A .

Theorem 7.4. A linear operator $A : X \rightarrow Y$ between normed spaces $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ is bounded iff the operator norm

$$\|A\| := \sup_{\|\varphi\|_X=1} \|A\varphi\|_Y = \sup_{\varphi \in X \setminus \{0\}} \frac{\|A\varphi\|_Y}{\|\varphi\|_X} \quad (7.4)$$

on the linear space $\mathcal{L}(X, Y)$ of all bounded linear operators between the two normed spaces $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ is finite, i.e.

$$\|A\| < \infty. \quad (7.5)$$

The operator norm is the smallest upper bound for A .

Proof. First, let us prove that the operator norm does indeed define a norm on $\mathcal{L}(X, Y)$. Note that the sum of two operators $A, B \in \mathcal{L}(X, Y)$ is defined pointwise, i.e. for $\varphi \in X$: $(A + B)(\varphi) := A\varphi + B\varphi$. Similarly, scalar multiplication is defined for $\alpha \in \mathbb{K}$ as $(\alpha A)(\varphi) := \alpha A\varphi$.

The positivity, definiteness and homogeneity of $\|A\|$ are trivial to show. For the triangle equality, note that for any $A, B \in \mathcal{L}(X, Y)$, we have

$$\|A + B\| = \sup_{\|\varphi\|_X=1} \|(A + B)(\varphi)\|_Y = \sup_{\|\varphi\|_X=1} \|A\varphi + B\varphi\|_Y \quad (7.6)$$

$$\leq \sup_{\|\varphi\|_X=1} \{\|A\varphi\|_Y + \|B\varphi\|_Y\} \quad (7.7)$$

$$\leq \sup_{\|\varphi\|_X=1} \{\|A\varphi\|_Y\} + \sup_{\|\varphi\|_X=1} \{\|B\varphi\|_Y\} \quad (7.8)$$

$$= \|A\| + \|B\|. \quad (7.9)$$

Now to the statement A bounded iff $\|A\| < \infty$.

“ \implies ” Let A be upper bounded with bound $C > 0$, then

$$\|A\| = \sup_{\|\varphi\|_X=1} \|A\varphi\|_Y \leq C < \infty. \quad (7.10)$$

This also shows that $\|A\|$ is the smallest upper bound for A .

“ \impliedby ” We have

$$\|A\| \|\varphi\|_X = \sup_{\varphi \in X \setminus \{0\}} \frac{\|A\varphi\|_Y}{\|\varphi\|_X} \|\varphi\|_X = \sup_{\varphi \in X \setminus \{0\}} \|A\varphi\|_Y \geq \|A\varphi\|_Y, \quad (7.11)$$

which shows that A is bounded with upper bound $C = \|A\| < \infty$ (note that for $v = 0$, there is nothing to show). \blacksquare

Theorem 7.5. A linear operator $A : X \rightarrow Y$ for normed linear spaces $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ is continuous iff it is bounded.

Proof. “ \impliedby ” Since A is bounded, by definition, we know that for all $\varphi \in X$, there is an upper bound $C > 0$ s.t.

$$\|A\varphi\|_Y \leq C \cdot \|\varphi\|_X. \quad (7.12)$$

For an arbitrary $\epsilon > 0$, choose $\delta := \epsilon/C$, then for $\|\varphi - \psi\|_X < \delta = \epsilon/C$ (where $\psi \in X$ is arbitrary) we have because of the linearity of A

$$\|A\varphi - A\psi\|_Y = \|A(\varphi - \psi)\|_Y \leq C \|\varphi - \psi\|_X < \epsilon, \quad (7.13)$$

which proves the continuity of A , cf. Def. 5.123 [75].

“ \implies ” Let A be continuous on X . Then by Theorem 5.132, we know that $A^{-1}(B_1^Y(0))$ is open, since $B_1^Y(0)$ is open. By the linearity of A , $0 \in A^{-1}(B_1^Y(0))$, since $A(0) = 0 \in B_1^Y(0)$. This means that there is an $r > 0$ s.t. $B_r^Y(0) \subset A^{-1}(B_1^Y(0))$, i.e. the image of $B_r^Y(0)$ is contained in $B_1^Y(0)$.

Now, for any $a > 1$, choose $C := a/r$, and we will show that C is an upper bound for A . For $\varphi \in X \setminus \{0\}$ (for $\varphi = 0$, there is nothing to show), note that

$$\left\| \frac{r}{a\|\varphi\|_X} \varphi \right\|_X = \frac{r}{a} < r,$$

i.e.

$$\frac{r}{a\|\varphi\|_X} \varphi \in B_r^Y(0) \Rightarrow A\left(\frac{r}{a\|\varphi\|_X} \varphi\right) = \frac{r}{a\|\varphi\|_X} A\varphi \in B_1^Y(0) \quad (7.14)$$

$$\Rightarrow \left\| \frac{r}{a\|\varphi\|_X} A\varphi \right\|_Y < 1 \Rightarrow \|A\varphi\|_Y \leq \frac{a}{r} \|\varphi\|_X = C \|\varphi\|_X, \quad (7.15)$$

which completes the proof [76, p. 2]. \blacksquare

Definition 7.6 ([9]). Let X and Y be normed linear spaces and $T : X \rightarrow Y$ a linear operator between them. Then T is called *compact* if $T(B_1^{\|\cdot\|_X}[0]) \subset Y$ is relatively compact, where $B_1^{\|\cdot\|_X}[0]$ is the closed unit ball in X , i.e. the image of the unit ball must be relatively compact.

Example 7.7. Let $k : [0, 1] \times [0, 1] \rightarrow \mathbb{K}$, where $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$, be continuous and let $f \in \mathcal{C}[0, 1]$. Now define the operator

$$T_k : \mathcal{C}[0, 1] \rightarrow \mathcal{C}[0, 1], (T_k f)(x) := \int_0^1 k(x, t) f(t) dt,$$

which is known as the *Fredholm integral equation of first kind*. The operator T_k is compact.

Proof. We need to show that

$$M := T_k (B_1^{d_\infty}[0]) = \{T_k f \mid f \in \mathcal{C}[0, 1], \|f\|_\infty \leq 1\} \subset \mathcal{C}[0, 1]$$

is relatively compact.

We will first show that M is bounded. For all $f \in M$ and all $x \in [0, 1]$, we have

$$|(T_k f)(x)| = \left| \int_0^1 k(x, t) f(t) dt \right| \leq \int_0^1 |k(x, t)| \cdot |f(t)| dt \leq \|k\|_\infty \|f\|_\infty \int_0^1 dt \leq \|k\|_\infty,$$

which shows that M is bounded.

We will now show that M is uniformly equicontinuous. Since the continuous function k is defined on a compact set, k is uniformly continuous on $[0, 1] \times [0, 1]$, i.e. for all $\epsilon > 0$ there exists a $\delta > 0$ s.t. for all $(x, t) \in [0, 1] \times [0, 1]$ and $(x', t') \in [0, 1] \times [0, 1]$, $d_{\text{Euclid}}((x, t), (x', t')) < \delta$ implies $|k(x, t) - k(x', t')| < \epsilon$. For all $\epsilon > 0$, we take this same δ . Then for all $T_k f \in M$ and $x, x' \in [0, 1]$, with $|x - x'| = d_{\text{Euclid}}((x, t), (x', t)) < \delta$, we have

$$\begin{aligned} |(T_k f)(x) - (T_k f)(x')| &= \left| \int_0^1 k(x, t) f(t) dt - \int_0^1 k(x', t) f(t) dt \right| \\ &= \left| \int_0^1 (k(x, t) - k(x', t)) f(t) dt \right| \\ &\leq \int_0^1 |k(x, t) - k(x', t)| \cdot |f(t)| dt \\ &\stackrel{f \in B_1^{d_\infty}[0]}{\leq} \int_0^1 |k(x, t) - k(x', t)| dt \\ &< \epsilon, \end{aligned}$$

since k is uniformly continuous. This shows that M is uniformly equicontinuous.

Thus, by the Arzelà-Ascoli Theorem 5.117, M is relatively compact, and thus T_k is a compact operator. ■

8 Set Theory

Definition 8.1 (Binary Relation [77]). A *binary relation* over a set X is some relation R where for all $x, y \in X$ the statement xRy is either true or false.

Definition 8.2 (Equivalence Relation [78]). An *equivalence relation* on a set X is a binary relation \sim with the following properties $\forall x, y, z \in X$:

- **reflexivity:** $x \sim x$,
- **symmetry:** $x \sim y \Leftrightarrow y \sim x$,
- **transitivity:** $(x \sim y) \wedge (y \sim z) \Rightarrow x \sim z$.

Definition 8.3 (Equivalence Class). Let \sim be an equivalence relation on X . Then the *equivalence class* of an element $x \in X$ is defined as

$$[x] := \{y \in X \mid x \sim y\} = \{y \in X \mid y \sim x\} \subset X. \quad (8.1)$$

The set of all equivalence classes in X wrt an equivalence relation \sim is denoted by X/\sim . Note that $X/\sim \subset P(X)$, where $P(X)$ denotes the power set of X [79]. The surjective map $f : X \rightarrow X/\sim, x \mapsto [x]$ is called the *canonical surjection*.

Theorem 8.4. Let \sim be an equivalence relation on X . For any $x, y \in X$, it holds that $[x] = [y]$ iff $x \sim y$.

Proof. “ \Rightarrow ” Since $x \in [x] = [y]$ we have $x \sim y$.

“ \Leftarrow ” Let $x \sim y$ for arbitrary $x, y \in X$ hold, and consider $x' \in [x]$, i.e. $x' \sim x$. By transitivity, we have $x' \sim y$, i.e. $[x] \subset [y]$. Similarly, we show $[y] \subset [x]$, and hence $[x] = [y]$. ■

Theorem 8.5. Let \sim be an equivalence relation on X . Then for $x, y \in X$, we either have $[x] \cap [y] = \emptyset$ or $[x] = [y]$.

Proof. We know that the statement $x \sim y$ is either true or false. If it is true, then by Theorem 8.4, $[x] = [y]$. If $x \not\sim y$, then no $x' \in [x] = \{z \in X \mid x \sim z\}$ can lie in $[y] = \{z' \in X \mid x \sim z'\}$ (and vice versa), because otherwise, by transitivity, $x \sim y$, which we said is false. Hence, $[x] \cap [y] = \emptyset$. ■

Definition 8.6. Every element of an equivalence class characterizes it, and can be used to *represent* it. Such a chosen element is called a *representative*.

Definition 8.7 (Partially Ordered Set [80]). A *partially ordered set* (X, \leq) is a set X , equipped with a binary relation \leq , that satisfies the following properties $\forall x, y, z \in X$:

- **reflexivity:** $x \leq x$,
- **antisymmetry:** $(x \leq y) \wedge (y \leq x) \Rightarrow x = y$,
- **transitivity:** $(x \leq y) \wedge (y \leq z) \Rightarrow x \leq z$.

Definition 8.8 (Incomparability). In Definition 8.7, the phrasing “partially ordered” is used to emphasize that there might exist elements $x, y \in X$ s.t. both $x \leq y$ and $y \leq x$ are wrong. These pairs are called *incomparable*. If either $x \leq y$ or $y \leq x$ is true, then we say that the pair is *comparable*.

Example 8.9. Consider $X := \{\{1\}, \{2\}, \{1, 2\}\}$ with \subset as partial ordering. Obviously, the elements $\{1\}$ and $\{2\}$ are incomparable.

Definition 8.10 (Chain, Upper Bound, Maximal Element). For preparing the Kuratowski-Zorn lemma, the following definitions come in handy:

- a) A *chain* C is a partially ordered set where every pair of elements in C is comparable. One might also say that C is a *totally ordered set*.
- b) An *upper bound* (if existent) of a subset $S \subset X$, where X is a partially ordered set, is an element $u \in X$ such that

$$s \leq u \quad \forall s \in S. \quad (8.2)$$

Since $S \subset X$, S itself is a partially ordered set.

- c) A *maximal element* (if existent) of a partially ordered set X is an element $m \in X$ such that

$$\text{if } m \leq x \text{ for some } x \in X, \text{ then } x = m. \quad (8.3)$$

This is equivalent to saying that there is no $x \in X$ such that $m \leq x$ and $x \neq m$.

Remark 8.11. For an arbitrary partially ordered set X , a maximal element (if existent) does not have to be unique. For example, consider $X := \{\{1\}, \{2\}, \{3\}, \{1, 2\}\}$ with \subset as partial ordering. Both $\{3\}$ and $\{1, 2\}$ are maximal elements. However, if we consider chains, then maximal elements are indeed unique by definition.

Theorem 8.12 (Kuratowski-Zorn Lemma). Let (M, \leq) be a non-empty partially ordered set. If every chain $C \subset M$ has an upper bound, then M has a maximal element.

Remark 8.13. The upper bound of every chain $C \subset M$ need not be in C , by definition of a chain, but it must be in M .

Theorem 8.14. If $A \subset \mathbb{R}$, then $M = \sup A$ iff M is an upper bound of A and for every $M' < M$ there exists an $x \in A$ s.t. $x > M'$.

Proof [81]. “ \Leftarrow ” M is an upper bound and for any $M' < M$ there is a bigger element in A , hence M is the lowest upper bound of A .

“ \Rightarrow ” Since $M = \sup A$, M is an upper bound of A and if $M' < M$, then M' is not an upper bound of A , since M is the lowest upper bound of A , and thus there exists an $x \in A$ s.t. $x > M'$. ■

Theorem 8.15 (Distributive law for set operations [82]). Let B be a set, and $\{A_i\}$ be a collection of sets, where I is a (finite, countable or uncountable) index set. Then we have

$$\left(\bigcup_{i \in I} A_i \right) \cap B = \bigcup_{i \in I} (A_i \cap B) \quad (8.4)$$

and

$$\left(\bigcap_{i \in I} A_i \right) \cup B = \bigcap_{i \in I} (A_i \cup B). \quad (8.5)$$

Proof. We will only show Eq. (8.4), since the proof of (8.5) is similar.

If

$$\begin{aligned} x \in \left(\bigcup_{i \in I} A_i \right) \cap B &\Rightarrow \left(x \in \bigcup_{i \in I} A_i \right) \wedge (x \in B) \Rightarrow (\exists j \in I : x \in A_j) \wedge (x \in B) \\ &\Rightarrow x \in A_j \cap B \subset \bigcup_{i \in I} (A_i \cap B), \end{aligned}$$

i.e.

$$\left(\bigcup_{i \in I} A_i \right) \cap B \subset \bigcup_{i \in I} (A_i \cap B).$$

Conversely, if

$$\begin{aligned} x \in \bigcup_{i \in I} (A_i \cap B) &\Rightarrow \exists j \in I : x \in A_j \cap B \Rightarrow \left(x \in A_j \subset \bigcup_{i \in I} A_i \right) \wedge (x \in B) \\ &\Rightarrow x \in \left(\bigcup_{i \in I} A_i \right) \cap B, \end{aligned}$$

i.e.

$$\bigcup_{i \in I} (A_i \cap B) \subset \left(\bigcup_{i \in I} A_i \right) \cap B,$$

which completes the proof. ■

Theorem 8.16. Let I be an index set, and for each $i \in I$, J_i shall be an index set as well. Then we have [83]

$$\bigcup_{i \in I} \left(\bigcup_{j \in J_i} A_j \right) = \bigcup_{j \in \bigcup_{i \in I} J_i} A_j. \quad (8.6)$$

Proof. “ \subset ” Let $a \in \bigcup_{i \in I} \left(\bigcup_{j \in J_i} A_j \right)$, i.e. there exists a $i_0 \in I$ s.t.

$$a \in \bigcup_{j \in J_{i_0}} A_j \subset \bigcup_{j \in \bigcup_{i \in I} J_i} A_j.$$

“ \supset ”

$$\begin{aligned} a &\in \bigcup_{j \in \bigcup_{i \in I} J_i} A_j \\ &\Rightarrow \exists j_0 \in \bigcup_{i \in I} J_i : a \in A_{j_0} \\ &\Rightarrow \exists i_0 \in I : (j_0 \in J_{i_0}) \wedge (a \in A_{j_0}) \\ &\Rightarrow a \in A_{j_0} \subset \bigcup_{i \in I} \left(\bigcup_{j \in J_i} A_j \right) \end{aligned}$$

■

Theorem 8.17. Let I be a *finite* index set, i.e. $I = \{1, \dots, n\}$ for some $n \in \mathbb{N}$, and for each $i \in I$, J_i shall be an index set as well. Then we have

$$\bigcap_{i \in I} \left(\bigcup_{j \in J_i} A_j \right) = \bigcup_{f \in J} \left(\bigcap_{i \in I} A_{f_i} \right), \quad (8.7)$$

where $J := \prod_{i \in I} J_i = J_1 \times \dots \times J_n$, i.e. $f \in J$ is an n -tuple, and hence $f = (f_1, \dots, f_n)$ with $f_i \in J_i$ for each $i \in I$.

Proof. Via induction. For $n = 1$, we have $I = \{1\}$ and $f = f_1 \in J_1$, and thus

$$\bigcap_{i \in I} \left(\bigcup_{j \in J_i} A_j \right) = \bigcup_{j \in J_1} A_j = \bigcup_{f \in J_1} \left(\bigcap_{i \in I} A_{f_i} \right).$$

For the induction step, note that if Eq. (8.7) holds for an $n \in \mathbb{N}$, and we define $I' := \{1, \dots, n, n+1\}$ and $J' := J \times \prod J_{n+1} = \prod_{i \in I'} J_i$, then

$$\begin{aligned} \bigcap_{i \in I'} \left(\bigcup_{j \in J_i} A_j \right) &= \left(\bigcup_{j \in J_1} A_j \right) \cap \dots \cap \left(\bigcup_{j \in J_n} A_j \right) \cap \left(\bigcup_{j \in J_{n+1}} A_j \right) \\ &\stackrel{\text{IA}}{=} \left(\bigcup_{f \in J} \left(\bigcap_{i \in I} A_{f_i} \right) \right) \cap \left(\bigcup_{j \in J_{n+1}} A_j \right) \\ &\stackrel{(8.4)}{=} \bigcup_{f \in J} \left(\left(\bigcap_{i \in I} A_{f_i} \right) \cap \left(\bigcup_{j \in J_{n+1}} A_j \right) \right) \\ &\stackrel{(8.4)}{=} \bigcup_{f \in J} \bigcup_{f_{n+1} \in J_{n+1}} \left(\bigcap_{i \in I'} A_{f_i} \right) \\ &= \bigcup_{f \in J'} \left(\bigcap_{i \in I'} A_{f_i} \right). \end{aligned}$$

■

Theorem 8.18. Let X, Y be sets and $f : X \rightarrow Y$ a map between them. Further, let I be an index set and $\{S_i\}_{i \in I}$ be a family of subsets of X , then [84]

$$f \left(\bigcup_{i \in I} S_i \right) = \bigcup_{i \in I} f(S_i). \quad (8.8)$$

Proof. We have

$$\begin{aligned} f \left(\bigcup_{i \in I} S_i \right) &= f(\{x \mid \exists i_0 \in I : x \in S_{i_0}\}) \\ &= \{f(x) \mid \exists i_0 \in I : x \in S_{i_0}\} \\ &= \bigcup_{i \in I} f(S_i). \end{aligned}$$

■

Theorem 8.19. Let X, Y be sets and $f : X \rightarrow Y$ a map between them. Further, let I be an index set and $U_i \subset Y$ for all $i \in I$. Then

$$\bigcup_{i \in I} f^{-1}(U_i) = f^{-1} \left(\bigcup_{i \in I} U_i \right) \quad (8.9)$$

and

$$\bigcap_{i \in I} f^{-1}(U_i) = f^{-1} \left(\bigcap_{i \in I} U_i \right), \quad (8.10)$$

where the preimage $f^{-1}(V) := \{x \in X \mid f(x) \in V\}$ for $V \subset Y$ [85, 86].

Proof. 1. We first prove Eq. (8.9).

“ \subset ” Let $x \in \bigcup_{i \in I} f^{-1}(U_i)$, i.e. there exists an $i_0 \in I$ s.t.

$$x \in f^{-1}(U_{i_0}) = \{z \in X \mid f(z) \in U_{i_0}\} \subset \left\{ z \in X \mid f(z) \in \bigcup_{i \in I} U_i \right\},$$

i.e. $x \in f^{-1} \left(\bigcup_{i \in I} U_i \right)$.

“ \supset ” Let $x \in f^{-1} \left(\bigcup_{i \in I} U_i \right) = \{z \in X \mid f(z) \in \bigcup_{i \in I} U_i\}$, i.e. there exists an $i_0 \in I$ s.t. $f(x) \in U_{i_0}$, which implies $x \in f^{-1}(U_{i_0}) \subset \bigcup_{i \in I} f^{-1}(U_i)$.

2. We now prove Eq. (8.10):

$$\begin{aligned} x \in \bigcap_{i \in I} f^{-1}(U_i) &\Leftrightarrow \forall i \in I : x \in f^{-1}(U_i) = \{z \in X \mid f(z) \in U_i\} \\ &\Leftrightarrow x \in f^{-1} \left(\bigcap_{i \in I} U_i \right) = \left\{ z \in X \mid f(z) \in \bigcap_{i \in I} U_i \right\}. \end{aligned}$$

■

Definition 8.20 (Infinite cartesian product [20]). Let A be a countable or uncountable index set, i.e. A is not finite. Let $\{X_\alpha\}_{\alpha \in A}$ be a collection of sets. Then we define the cartesian product

$$\prod_{\alpha \in A} X_\alpha := \left\{ x : A \rightarrow \bigcup_{\alpha \in A} X_\alpha \mid x(\alpha) \in X_\alpha \ \forall \alpha \in A \right\}, \quad (8.11)$$

i.e. the cartesian product is the set of all functions mapping from A to the union of X_α . We write $x_\alpha = x(\alpha)$ and denote the function x by $x = (x_\alpha)_{\alpha \in A}$. If $X_\alpha = X$ for all $\alpha \in A$, then $\prod_{\alpha \in A} X_\alpha$ is the set of all functions $A \rightarrow X$, which we denote by X^A .

Remark 8.21. If any $X_\alpha = \emptyset$, then $\prod_{\alpha \in A} X_\alpha = \emptyset$, since there is no function x s.t. $x(\alpha) \in X_\alpha$.

Theorem 8.22. Let $\{U_\alpha\}_{\alpha \in A}$ and $\{V_\alpha\}_{\alpha \in A}$ be families of sets, where A is an index set, then

$$\left(\prod_{\alpha \in A} U_\alpha \right) \cap \left(\prod_{\alpha \in A} V_\alpha \right) = \prod_{\alpha \in A} (U_\alpha \cap V_\alpha). \quad (8.12)$$

Proof. When looking at the definition of the infinite Cartesian product, we notice that the codomain of the functions are only relevant to indicate that $x(\alpha) \in X_\alpha$, i.e. two functions $x : A \rightarrow B$ and $x' : A \rightarrow B'$ are identical as long as $\text{ran}(x) = \text{ran}(x') \subset B$ and $\text{ran}(x) = \text{ran}(x') \subset B'$. Thus, defining

$$\Omega := \left(\bigcup_{\alpha \in A} U_\alpha \right) \cup \left(\bigcup_{\alpha \in A} V_\alpha \right),$$

we have

$$\begin{aligned} & \left(\prod_{\alpha \in A} U_\alpha \right) \cap \left(\prod_{\alpha \in A} V_\alpha \right) \\ &= \left\{ u : A \rightarrow \bigcup_{\alpha \in A} U_\alpha \mid u(\alpha) \in U_\alpha \ \forall \alpha \in A \right\} \cap \left\{ v : A \rightarrow \bigcup_{\alpha \in A} V_\alpha \mid v(\alpha) \in V_\alpha \ \forall \alpha \in A \right\} \\ &= \{ u : A \rightarrow \Omega \mid u(\alpha) \in U_\alpha \ \forall \alpha \in A \} \cap \{ v : A \rightarrow \Omega \mid v(\alpha) \in V_\alpha \ \forall \alpha \in A \} \\ &= \{ w : A \rightarrow \Omega \mid w(\alpha) \in U_\alpha \cap V_\alpha \ \forall \alpha \in A \} \\ &= \prod_{\alpha \in A} (U_\alpha \cap V_\alpha). \end{aligned}$$

■

Theorem 8.23. Let X be a set, I an arbitrary index set, and $U_i \subset X$ for all $i \in I$. Then

$$X \setminus \bigcup_{i \in I} U_i = \bigcap_{i \in I} (X \setminus U_i). \quad (8.13)$$

Similarly,

$$X \setminus \bigcap_{i \in I} U_i = \bigcup_{i \in I} (X \setminus U_i). \quad (8.14)$$

Proof. Eq. (8.13):

“ \subset ”

$$x \in X \setminus \bigcup_{i \in I} U_i \Rightarrow x \in X \wedge x \notin \bigcup_{i \in I} U_i \Rightarrow x \in X \wedge (x \notin U_i \ \forall i \in I) \Rightarrow x \in \bigcap_{i \in I} (X \setminus U_i).$$

“ \supset ”

$$x \in \bigcap_{i \in I} (X \setminus U_i) \Rightarrow (x \in X \wedge x \notin U_i) \ \forall i \in I \Rightarrow x \in X \wedge x \notin \bigcup_{i \in I} U_i \Rightarrow x \in X \setminus \bigcup_{i \in I} U_i.$$

Eq. (8.14):

“ \subset ”

$$\begin{aligned} x \in X \setminus \bigcap_{i \in I} U_i &\Rightarrow x \in X \wedge x \notin \bigcap_{i \in I} U_i \Rightarrow x \in X \wedge (\exists i \in I : x \notin U_i) \Rightarrow \exists i \in I : x \in X \wedge x \notin U_i \\ &\Rightarrow \exists i \in I : x \in X \setminus U_i \Rightarrow x \in \bigcup_{i \in I} (X \setminus U_i) \end{aligned}$$

“ \supset ”

$$x \in \bigcup_{i \in I} (X \setminus U_i) \Rightarrow \exists i \in I : x \in X \wedge x \notin U_i \Rightarrow x \in X \wedge x \notin \bigcap_{i \in I} U_i \Rightarrow x \in X \setminus \bigcap_{i \in I} U_i.$$

■

Theorem 8.24. Let X, Y be sets and $f : X \rightarrow Y$ a map between them. Then

$$\forall E \subset Y : f(f^{-1}(E)) \subset E. \quad (8.15)$$

and

$$\forall A \subset X : A \subset f^{-1}(f(A)). \quad (8.16)$$

Proof.

$$\begin{aligned} f(f^{-1}(E)) &= f(\{x \in X \mid f(x) \in E\}) \\ &= \{f(x) \in Y \mid x \in X \wedge f(x) \in E\} \\ &= \{f(x) \in E \mid x \in X\} \subset E. \end{aligned}$$

Also,

$$\begin{aligned} x \in A &\Rightarrow f(x) \in f(A) = \{f(x) \mid x \in A\} \Leftrightarrow x \in f^{-1}(f(A)) = \{x \in X \mid f(x) \in f(A)\} \\ &\Rightarrow A \subset f^{-1}(f(A)). \end{aligned}$$

■

Theorem 8.25. Let $\{B_i\}_{i \in I}$ be a family of sets, where I is an arbitrary index set. Also, let A be a set. Then

$$\bigcup_{i \in I} (A \times B_i) = A \times \bigcup_{i \in I} B_i. \quad (8.17)$$

Proof.

$$\begin{aligned} \bigcup_{i \in I} (A \times B_i) &= \bigcup_{i \in I} \{(a, x) \mid a \in A \wedge x \in B_i\} \\ &= \left\{ (a, x) \mid a \in A \wedge x \in \bigcup_{i \in I} B_i \right\} \\ &= A \times \bigcup_{i \in I} B_i \end{aligned}$$

■

9 Differential Geometry

Definition 9.1 (Smooth Atlas [87]). Let M be a second countable Hausdorff topological space. An n -dimensional smooth atlas on M is a collection of maps

$$\mathcal{A} = \{(\varphi_i, U_i) \mid i \in I\}, \quad \varphi_i : U_i \rightarrow \varphi_i(U_i) \subset \mathbb{R}^n$$

such that all $U_i \subset M$ are open, all φ_i are homeomorphisms, I is an index set and

- $\{U_i \mid i \in I\}$ is an open covering of M ,
- $\varphi_i \circ \varphi_j^{-1} : \varphi_j(U_i \cap U_j) \rightarrow \varphi_i(U_i \cap U_j)$ are smooth $\forall i, j \in I$.

The tuples (φ_i, U_i) , $i \in I$, are so-called *charts* on M , the maps $\varphi_i \circ \varphi_j^{-1}$ are called *transition maps* or *changes of coordinates* and n is the *dimension* of M .

Remark 9.2. To see why the domain of the transition maps $\varphi_i \circ \varphi_j^{-1}$ is $\varphi_j(U_i \cap U_j)$, note that the expression $\varphi_i(\varphi_j^{-1}(x))$ only makes sense if

$$(x \in \varphi_j(U_j)) \wedge (\varphi_j^{-1}(x) \in U_i) \Rightarrow (x \in \varphi_j(U_j)) \wedge (x \in \varphi_j(U_i)) \Rightarrow x \in \varphi_j(U_j \cap U_i).$$

Similarly, we can convince ourselves that the codomain of the transition maps $\varphi_i \circ \varphi_j^{-1}$ is given by $\varphi_i(U_i \cap U_j)$. Since $x \in \varphi_j(U_j)$, it follows that $\varphi_j^{-1}(x) \in U_j$. In addition, due to the domain of the homeomorphism φ_i , it must hold that $\varphi_j^{-1}(x) \in U_i$. Thus:

$$\begin{aligned} & (\varphi_j^{-1}(x) \in U_j) \wedge (\varphi_j^{-1}(x) \in U_i) \\ \Rightarrow & (\varphi_i(\varphi_j^{-1}(x)) \in \varphi_i(U_j)) \wedge (\varphi_i(\varphi_j^{-1}(x)) \in \varphi_i(U_i)) \\ \Rightarrow & \varphi_i(\varphi_j^{-1}(x)) \in \varphi_i(U_i \cap U_j). \end{aligned}$$

Definition 9.3 (Equivalence of Atlases). Let M be a second countable Hausdorff topological space. Two atlases \mathcal{A} and \mathcal{B} on M are called *equivalent* if $\mathcal{A} \cup \mathcal{B}$ is an atlas on M .

Remark 9.4. To see that not all atlases are equivalent to each other, consider $M = \mathbb{R}$ (which is a second countable Hausdorff topological space). Consider the atlases $\mathcal{A} = \{(\varphi, M)\}$ with $\varphi : M \rightarrow M$, $x \mapsto x$ and $\mathcal{B} = \{(\psi, M)\}$ with $\psi : M \rightarrow M$, $x \mapsto x^3$. The atlases are not equivalent, since $\varphi \circ \psi^{-1} : M \rightarrow M$, $x \mapsto \sqrt[3]{x}$ is not smooth (the derivative is not continuous).

10 Normalizing Flows

Definition 10.1. Let M, N be two manifolds, $g : M \rightarrow N$ a differentiable map. If g is a bijection and its inverse $g^{-1} : N \rightarrow M$ is differentiable as well, then we call f a *diffeomorphism*. We talk of a C^k *diffeomorphism* if both g and g^{-1} are k -times continuously differentiable.

Theorem 10.2 (Change of variables [88]). Let $U, V \subset \mathbb{R}^n$ be open subsets and $T : U \rightarrow V$ a diffeomorphism, cf. Def. 10.1. Then the function $f : V \rightarrow \mathbb{C} \cup \{\infty\}$ is integrable over V if and only if the function

$$(f \circ T) \cdot \left| \det \left(\frac{\partial T_\mu}{\partial x_\nu} \right)_{\mu\nu} \right| \quad (10.1)$$

is integrable over U . In this case, it holds that

$$\int_U (f \circ T)(x) \cdot \left| \det \left(\frac{\partial T_\mu}{\partial x_\nu}(x) \right)_{\mu,\nu} \right| dx = \int_V f(y) dy. \quad (10.2)$$

Remark 10.3. If T is a diffeomorphism, then also T^{-1} is a diffeomorphism, thus we could also have chosen T^{-1} in the formulation of Theorem 10.2.

Theorem 10.4 (Inverse function theorem [89]). Let $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be continuously differentiable on some open set $V \subset \mathbb{R}^n$ containing \mathbf{a} and suppose $\det Jg(\mathbf{a}) \neq 0$, where J shall be the Jacobi matrix of g . Then there is some open set containing \mathbf{a} and an open set $W \subset \mathbb{R}^n$ containing $g(\mathbf{a})$ such that $g : V \rightarrow W$ has a continuous inverse $g^{-1} : W \rightarrow V$ which is differentiable for all $\mathbf{y} \in W$.

As matrices, we can write this as

$$J(g^{-1})(\mathbf{y}) = [Jg(g^{-1}(\mathbf{y}))]^{-1} \quad (10.3)$$

Remark 10.5. An example for a function that is invertible and continuously differentiable but not a diffeomorphism is $g : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto x^3$. Its inverse is obviously $g^{-1} : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto \sqrt[3]{x}$, cf. [90] for a nice plot. However, $\frac{dg^{-1}}{dx}|_{x=0}$ does not exist. The reason is that $\det Jg(0) = 0$ and hence the inverse function theorem does not apply.

- Let \mathbf{U} be a random variable and let $p(\mathbf{U})$ describe the probability distribution of it, e.g. a uniform distribution between 0 and 1. We now make a simple transformation and obtain a new random variable \mathbf{X} , where we again denote by $p(\mathbf{X})$ the probability distribution of \mathbf{X} . We obtain \mathbf{X} in the following way:

$$p(\mathbf{X}) = p(\mathbf{U}) \left| \det \left(\frac{\partial \mathbf{f}}{\partial \mathbf{U}} \right) \right|^{-1}, \quad (10.4)$$

where \mathbf{f} denotes an invertible (and hence bijective) mapping.

- Without proof, it holds that

$$\left| \det \left(\frac{\partial \mathbf{f}}{\partial \mathbf{U}} \right) \right|^{-1} = \left| \det \left(\frac{\partial \mathbf{f}^{-1}}{\partial \mathbf{U}} \right) \right| \quad (10.5)$$

and thus we can rewrite Eq. (10.4) as

$$p(\mathbf{U}) = p(\mathbf{X}) \left| \det \left(\frac{\partial \mathbf{f}^{-1}}{\partial \mathbf{U}} \right) \right|^{-1}. \quad (10.6)$$

Since we assumed \mathbf{f} to be invertible, \mathbf{f}^{-1} is well-defined.

- In practice, we will want \mathbf{f} to be both invertible and to have a **tractable** Jacobian, i.e. a Jacobian that we can easily calculate. For \mathbf{f} to have a Jacobian at all, each of its first-order partial derivatives must exist [91]. So-called *autoregressive flows* have the property that their Jacobian is an upper triangular matrix. For an upper triangular matrix, it holds that its determinant is given by the product of its diagonal elements [92].

Definition 10.6 (Determinant). Let D be an $n \times n$ matrix and let S_n denote the symmetric group over n . Then the determinant of D is defined as:

$$\det(D) := \sum_{\sigma \in S_n} \left(\text{sgn}(\sigma) \prod_{i=1}^n a_{i, \sigma(i)} \right), \quad (10.7)$$

cf. [93].

Lemma 10.7. Let A be a $k \times k$, 0 an $k \times n$, C an $n \times k$ and D an $n \times n$ matrix; then

$$\det \left(\begin{pmatrix} A & 0 \\ C & D \end{pmatrix} \right) = \det(A) \det(D). \quad (10.8)$$

Proof. Define

$$B := \begin{pmatrix} A & 0 \\ C & D \end{pmatrix}. \quad (10.9)$$

Clearly,

$$b_{i,j} = \begin{cases} a_{i,j} & i, j \leq k, \\ 0 & i \leq k, j \geq k+1, \\ c_{i-k,j} & i \geq k+1, j \leq k, \\ d_{i-k,j-k} & i, j \geq k+1. \end{cases} \quad (10.10)$$

We can write the determinant of B as

$$\det(B) = \sum_{\sigma \in S_{n+k}} \text{sgn}(\sigma) \prod_{i=1}^{n+k} b_{i, \sigma(i)}. \quad (10.11)$$

From Eq. (10.10) we know that all summands of the form $\sigma(i) = j$ with $i \leq k, j \geq k+1$ are 0. Therefore, we can consider all permutations of the form $\sigma(i) = j$ with $i, j \leq k$ or $\sigma(i) = j$ with $i \geq k+1, j \leq k$. We can also write this in the form $\sigma(i) = \pi(i)$ for $i \leq k$

and $\sigma(k+i) = k + \tau(i)$ for $1 \leq i \leq n$, where $\pi \in S_k$ and $\tau \in S_n$. Denote the set of all such permutations by \tilde{S}_{k+n} . Thus:

$$\det(B) = \sum_{\sigma \in \tilde{S}_{k+n}} \text{sgn}(\sigma) \prod_{i=1}^{n+k} b_{i,\sigma(i)} \quad (10.12)$$

$$= \sum_{\sigma \in \tilde{S}_{k+n}} \text{sgn}(\sigma) \prod_{i=1}^k b_{i,\sigma(i)} \prod_{i=k+1}^{n+k} b_{i,\sigma(i)} \quad (10.13)$$

$$\stackrel{(10.10)}{=} \sum_{\sigma \in \tilde{S}_{k+n}} \text{sgn}(\sigma) \prod_{i=1}^k a_{i,\sigma(i)} \prod_{i=k+1}^{n+k} d_{i-k,\sigma(i)-k} \quad (10.14)$$

$$= \sum_{\sigma \in \tilde{S}_{k+n}} \text{sgn}(\sigma) \prod_{i=1}^k a_{i,\sigma(i)} \prod_{i=1}^n d_{i,\sigma(i+k)-k} \quad (10.15)$$

$$= \sum_{\pi \in S_k, \tau \in S_n} \text{sgn}(\pi) \text{sgn}(\tau) \prod_{i=1}^k a_{i,\pi(i)} \prod_{i=1}^n d_{i,\tau(i)} \quad (10.16)$$

$$= \sum_{\pi \in S_k} \text{sgn}(\pi) \prod_{i=1}^k a_{i,\pi(i)} \sum_{\tau \in S_n} \text{sgn}(\tau) \prod_{i=1}^n d_{i,\tau(i)} \quad (10.17)$$

$$= \det(A) \det(D) \quad (10.18)$$

[94] ■

Definition 10.8. Let $h(\cdot; \theta) : \mathbb{R} \rightarrow \mathbb{R}$ be a bijection parametrized by θ . Then an *autoregressive model* is a function

$$g : \mathbb{R}^D \rightarrow \mathbb{R}^D, \begin{pmatrix} x_1 \\ \dots \\ x_D \end{pmatrix} \mapsto \begin{pmatrix} h(x_1; \Theta_1) \\ h(x_2; \Theta_2(x_1)) \\ \dots \\ h(x_D; \Theta_D(x_1, \dots, x_{D-1})) \end{pmatrix} \quad (10.19)$$

The functions Θ_t for $t = 2, \dots, D$ are arbitrary functions whose domain is \mathbb{R}^{t-1} , Θ_1 is a constant.

Remark 10.9. The Jacobian matrix of an autoregressive flow is given as follows:

$$Dg = \begin{pmatrix} \partial g_1 / \partial x_1 & \partial g_1 / \partial x_2 & \dots & \partial g_1 / \partial x_D \\ \partial g_2 / \partial x_1 & \partial g_2 / \partial x_2 & \dots & \partial g_2 / \partial x_D \\ \vdots & \vdots & \ddots & \vdots \\ \partial g_D / \partial x_1 & \partial g_D / \partial x_2 & \dots & \partial g_D / \partial x_D \end{pmatrix} \quad (10.20)$$

One can easily convince oneself that Dg is a lower triangular matrix.

Theorem 10.10. Normalizing flows in \mathbb{R}^D come from the push-forward of a measure.

Proof. Let $\mathcal{Y}, \mathcal{Z} \subset \mathbb{R}^D$ be open, $\Sigma_{\mathcal{Y}} = \mathcal{B}(\mathcal{Y})$, $\Sigma_{\mathcal{Z}} = \mathcal{B}(\mathcal{Z})$ and $g : \mathcal{Z} \rightarrow \mathcal{Y}$ be a diffeomorphism. The function $g : \mathcal{Z} \rightarrow \mathcal{Y}$ is measurable if and only if $g^{-1}(G) \in \Sigma_{\mathcal{Z}}$ for every set G that is open in \mathcal{Y} , cf. Theorem 2.4 (Borel σ -algebras are generated by the open sets). Since for functions between two topological spaces it holds that they are continuous if and

only if the inverse image of an open set is again open, we have that g is indeed measurable.

Now define the probability measure μ on the measurable space $(\mathcal{Z}, \Sigma_{\mathcal{Z}})$ as

$$\mu(I) := \int_I p_{\mathbf{Z}}(z) d\lambda(z) \quad \forall I \in \Sigma_{\mathcal{Z}}, \quad (10.21)$$

where $p_{\mathbf{Z}} : \mathcal{Z} \rightarrow \mathbb{R}$ shall be a PDF and λ the Lebesgue measure. (The Lebesgue measure is defined on the completion of $\mathcal{B}(\mathbb{R}^D)$, which is “larger” than $\mathcal{B}(\mathbb{R}^D)$.) By considering the push-forward of μ under the measurable map g , we have $\forall J \in \Sigma_{\mathcal{Y}}$:

$$g_{\star}\mu(J) = \mu(g^{-1}(J)) \stackrel{(10.21)}{=} \int_{g^{-1}(J)} p_{\mathbf{Z}}(z) d\lambda(z) \quad (10.22)$$

Since by assumption $g : \mathcal{Z} \rightarrow \mathcal{Y}$ and therefore also the inverse $g^{-1} : \mathcal{Y} \rightarrow \mathcal{Z}$ are a diffeomorphism, we can use the change of variables formula from Theorem 10.2, since we assume $p_{\mathbf{Z}}$ to be integrable over \mathcal{Z} . We apply Theorem 10.2 to g^{-1} instead of g , cf. Remark 10.3:

$$g_{\star}\mu(J) = \int_{g^{-1}(J)} p_{\mathbf{Z}}(z) d\lambda(z) = \int_J \underbrace{(p_{\mathbf{Z}} \circ g^{-1})(y) \cdot |\det Dg^{-1}(y)|}_{=p_{\mathbf{Y}}(y)} d\lambda(y) \quad (10.23)$$

■

Definition 10.11. A rational-quadratic function takes the form of a quotient of two quadratic polynomials.

$$\frac{\alpha^{(k)}(\xi)}{\beta^{(k)}(\xi)} = \frac{a_0\xi^2 + a_1\xi + a_2}{b_0\xi^2 + b_1\xi + b_2} \quad (10.24)$$

11 Miscellaneous

Lemma 11.1 (Chain rule for KL-divergences). Let $p(x, y)$ and $q(x, y)$ be two arbitrary PDF's. Then the following holds:

$$D_{\text{KL}}(p(x, y) || q(x, y)) = D_{\text{KL}}(p(x) || q(x)) + D_{\text{KL}}(p(y|x) || q(y|x)) \quad (11.1)$$

Proof. Brute-force calculation yields:

$$D_{\text{KL}}(p(x, y) || q(x, y)) = \int \int p(x, y) \log \frac{p(x, y)}{q(x, y)} d\lambda(x) d\lambda(y) \quad (11.2)$$

$$= \int \int p(x, y) \log \frac{p(x)p(y|x)}{q(x)q(y|x)} d\lambda(x) d\lambda(y) \quad (11.3)$$

$$= \int \int p(x, y) \log \frac{p(x)}{q(x)} d\lambda(x) d\lambda(y) + \int \int p(x, y) \log \frac{p(y|x)}{q(y|x)} d\lambda(x) d\lambda(y) \quad (11.4)$$

$$= D_{\text{KL}}(p(x) || q(x)) + \int p(x) d\lambda(x) \int p(y|x) \log \frac{p(y|x)}{q(y|x)} d\lambda(y) \quad (11.5)$$

$$= D_{\text{KL}}(p(x) || q(x)) + D_{\text{KL}}(p(y|x) || q(y|x)) \quad (11.6)$$

■

Definition 11.2 (Mutual information). Let $(X, Y) \sim P_{(X, Y)} \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$, where $\mathcal{P}(\mathcal{X} \times \mathcal{Y})$ is the space of all probability measures over the space $\mathcal{X} \times \mathcal{Y}$. The mutual information between the random variables X and Y is now defined as [95, Def. 10.1]:

$$I(X; Y) := D_{\text{KL}}(P_{(X, Y)} || P_X \otimes P_Y), \quad (11.7)$$

where P_X and P_Y are the marginal measures of the coupling measure $P_{(X, Y)}$ and $P_X \otimes P_Y$ is the induced **product measure**.

Definition 11.3 (Divergence). Let p and q be two probability distributions, then a divergence D must satisfy: $D(p || q) \geq 0 \forall p, q$ and $D(p || q) = 0 \Leftrightarrow p = q$ a.e. Note that the triangle inequality and the symmetry property need not be satisfied in general.

Example 11.4 (f -Divergence). Let P and Q be two probability measures defined on the σ -algebra over a space Ω such that $P \ll Q$, i.e. P is absolutely continuous wrt Q . Then, for a convex function $f : [0, \infty) \rightarrow \mathbb{R}$ s.t.

$$(i) \ f(1) = 0,$$

$$(ii) \ f \text{ is strictly convex at } x = 1,$$

the f -divergence is now defined as [96],

$$D_f(P || Q) := \int_{\Omega} f\left(\frac{dP}{dQ}\right) dQ, \quad (11.8)$$

where dP/dQ is the Radon-Nikodym derivative.

Proof. 1. **Non-negativity:** We know that

$$\mathbb{E}_Q \left[\frac{dP}{dQ} \right] = \int_{\Omega} \frac{dP}{dQ} dQ = 1. \quad (11.9)$$

Since f is a convex function, by Jensen's inequality,

$$0 = f(1) = f \left(\mathbb{E}_Q \left[\frac{dP}{dQ} \right] \right) \leq \mathbb{E}_Q \left[f \left(\frac{dP}{dQ} \right) \right] = \int_{\Omega} f \left(\frac{dP}{dQ} \right) dQ \stackrel{(11.8)}{=} D_f(P||Q). \quad (11.10)$$

2. **Zero iff equal:** If $P = Q$ a.e., then $dP/dQ = 1$ a.e., and hence

$$D_f(P||Q) = \int_{\Omega} f(1) dQ = \int_{\Omega} 0 dQ = 0. \quad (11.11)$$

Conversely, if $D_f(P||Q) = 0$, since f is strictly convex at 1 and convex everywhere else, this implies that the only minimum point of f is at $f(1) = 0$. Therefore, the integrand of Eq. (11.8), i.e. $f \left(\frac{dP}{dQ} \right)$, must be 0 a.e., implying that

$$\frac{dP}{dQ} = 1 \Rightarrow P = Q. \quad (11.12)$$

■

Theorem 11.5. For all $a, b, c, d \in \mathbb{R}$, we have

$$\max\{a + b, c + d\} \leq \max\{a, c\} + \max\{b, d\}. \quad (11.13)$$

Proof [97]. We have $a + b \leq \max\{a, c\} + \max\{b, d\}$ and $c + d \leq \max\{a, c\} + \max\{b, d\}$, from which Eq. (11.13) follows. ■

Theorem 11.6. Let D be a set and $f, g : D \rightarrow \mathbb{R}$ real-valued, bounded functions, then

$$\sup_{x \in D} \{(f + g)(x)\} \leq \sup_{x \in D} \{f(x)\} + \sup_{x \in D} \{g(x)\}. \quad (11.14)$$

and

$$\inf_{x \in D} \{(f + g)(x)\} \geq \inf_{x \in D} \{f(x)\} + \inf_{x \in D} \{g(x)\}. \quad (11.15)$$

Proof [98, 99]. Clearly, for all $x \in D$, $f(x) \leq \sup_{x \in D} \{f(x)\}$ and $g(x) \leq \sup_{x \in D} \{g(x)\}$, implying that $(f + g)(x) \leq \sup_{x \in D} \{f(x)\} + \sup_{x \in D} \{g(x)\}$, which shows that $\sup_{x \in D} \{f(x)\} + \sup_{x \in D} \{g(x)\}$ is an upper bound for $f + g$, yet the smallest upper bound of $f + g$ may be smaller, so $\sup_{x \in D} \{(f + g)(x)\} \leq \sup_{x \in D} \{f(x)\} + \sup_{x \in D} \{g(x)\}$.

Also, for all $x \in D$, $f(x) \geq \inf_{x \in D} \{f(x)\}$ and $g(x) \geq \inf_{x \in D} \{g(x)\}$, and thus $(f + g)(x) \geq \inf_{x \in D} \{f(x)\} + \inf_{x \in D} \{g(x)\}$, which shows that $\inf_{x \in D} \{f(x)\} + \inf_{x \in D} \{g(x)\}$ is a lower bound for $f + g$. Since the biggest lower bound for $f + g$ may be bigger, we have $\inf_{x \in D} \{(f + g)(x)\} \geq \inf_{x \in D} \{f(x)\} + \inf_{x \in D} \{g(x)\}$. ■

Remark 11.7. Note that there are cases where

$$\sup_{x \in D} \{(f + g)(x)\} < \sup_{x \in D} \{f(x)\} + \sup_{x \in D} \{g(x)\}$$

and

$$\inf_{x \in D} \{(f + g)(x)\} > \inf_{x \in D} \{f(x)\} + \inf_{x \in D} \{g(x)\}.$$

Example 11.8. Define $f : [a, b] \rightarrow \mathbb{R}, x \mapsto x$ and $g : [a, b] \rightarrow \mathbb{R}, x \mapsto -x$ with $b > a \geq 0$. Clearly,

$$\sup_{x \in [a, b]} \{(f + g)(x)\} = 0 < \sup_{x \in [a, b]} \{f(x)\} + \sup_{x \in [a, b]} \{g(x)\} = b - a$$

and

$$\inf_{x \in [a, b]} \{(f + g)(x)\} = 0 > \inf_{x \in [a, b]} \{f(x)\} + \inf_{x \in [a, b]} \{g(x)\} = a - b.$$

Corollary 11.9. Let D be a set and $f, g : D \rightarrow \mathbb{C}$ complex-valued functions s.t. $|f|$ and $|g|$ are bounded, then

$$\sup_{x \in D} \{(|f| + |g|)(x)\} \leq \sup_{x \in D} \{|f|(x)\} + \sup_{x \in D} \{|g|(x)\} \quad (11.16)$$

and

$$\inf_{x \in D} \{(|f| + |g|)(x)\} \geq \inf_{x \in D} \{|f|(x)\} + \inf_{x \in D} \{|g|(x)\}. \quad (11.17)$$

Theorem 11.10. Let D be a set and $f, g : D \rightarrow \mathbb{R}$, where $f \leq g$. If g is bounded from above, then

$$\sup_{x \in D} f(x) \leq \sup_{x \in D} g(x) \quad (11.18)$$

and if f is bounded from below, then

$$\inf_{x \in D} f(x) \leq \inf_{x \in D} g(x). \quad (11.19)$$

Proof [81]. For all $x \in [a, b]$, if $f \leq g$ and g is bounded from above, then

$$f(x) \leq g(x) \leq \sup_{x \in D} g(x),$$

i.e. $\sup_{x \in D} g(x)$ is an upper bound for $f(x)$. Since there may be a smaller upper bound, we have

$$\sup_{x \in D} f(x) \leq \sup_{x \in D} g(x).$$

If g is bounded from below, then

$$\inf_{x \in D} f(x) \leq f(x) \leq g(x),$$

i.e. $\inf_{x \in D} f(x)$ is a lower bound for $g(x)$. Since there may be a bigger lower bound, we have

$$\inf_{x \in D} f(x) \leq \inf_{x \in D} g(x).$$

■

Corollary 11.11. Let D be a set and $f, g : D \rightarrow \mathbb{C}$ s.t. $|f| \leq |g|$. If $|g|$ is bounded from above, then

$$\sup_{x \in D} |f|(x) \leq \sup_{x \in D} |g|(x) \quad (11.20)$$

and if $|f|$ is bounded from below, then

$$\inf_{x \in D} |f|(x) \leq \inf_{x \in D} |g|(x). \quad (11.21)$$

A Weierstrass Approximation Theorem

In the following, we will prove the Weierstrass approximation theorem, as stated in Example 5.78. The proof will use *Korovkin*¹¹ sequences. The approach is taken from Chapter 6 of [29].

Definition A.1 (Positivity of Operator). A linear operator $K : \mathcal{C}[a, b] \rightarrow \mathcal{C}[a, b]$ is *positive* on $\mathcal{C}[a, b]$ if $Kf \geq 0$ for all $f \in \mathcal{C}[a, b]$ satisfying $f \geq 0$, where all inequalities are taken pointwise on $[a, b]$.

Definition A.2 (Monotonicity of Operator). A linear operator $K : \mathcal{C}[a, b] \rightarrow \mathcal{C}[a, b]$ is *monotone* on $\mathcal{C}[a, b]$ if for any $f, g \in \mathcal{C}[a, b]$ satisfying $f \leq g$, we have $Kf \leq Kg$, where all inequalities are taken pointwise on $[a, b]$.

Remark A.3. A linear operator is positive iff it is monotone.

Proof. “ \implies ” Consider $h := g - f \in \mathcal{C}[a, b]$, where $g, f \in \mathcal{C}[a, b]$, with $h = g - f \geq 0$, i.e. $f \leq g$, then by the positivity and linearity of K we have $Kh = Kg - Kf \geq 0$, or equivalently $Kf \leq Kg$.

“ \impliedby ” Assume that for any $f, g \in \mathcal{C}[a, b]$ satisfying $f \leq g$, we have $Kf \leq Kg$. Take $f = 0$, then $g \geq 0$ implies $Kg \geq 0$ for any $g \in \mathcal{C}[a, b]$. ■

Definition A.4 (Korovkin sequence). A sequence $(K_n)_{n \in \mathbb{N}}$ of linear and positive operators $K_n : \mathcal{C}[a, b] \rightarrow \mathcal{C}[a, b]$ is called a *Korovkin sequence* on $\mathcal{C}[a, b]$ if $K_n p \xrightarrow{n \rightarrow \infty} p$ wrt d_∞ for all $p \in \mathcal{P}_2$, where \mathcal{P}_n is the linear space of all univariate polynomials of degree at most n defined on $[a, b]$.

Theorem A.5 (Korovkin, 1953). For a compact interval $[a, b] \subset \mathbb{R}$, let $(K_n)_{n \in \mathbb{N}}$ be a Korovkin sequence on $\mathcal{C}[a, b]$. Then for any $f \in \mathcal{C}[a, b]$, we have

$$K_n f \xrightarrow{n \rightarrow \infty} f \quad \text{wrt } d_\infty. \quad (\text{A.1})$$

Proof. Let $f \in \mathcal{C}[a, b]$ be arbitrary. Then, f is bounded on $[a, b]$, cf. Theorem 6.133, i.e. there is an $M > 0$ s.t. $\|f\|_\infty \leq M$. Also, f is uniformly continuous on $[a, b]$, cf. Theorem 5.133, i.e.

$$\forall \epsilon > 0 \exists \delta > 0 \forall x, y \in [a, b] : |x - y| < \delta \Rightarrow |f(x) - f(y)| < \epsilon. \quad (\text{A.2})$$

Now let $t \in [a, b]$ be fixed, $x \in [a, b]$ and $\epsilon > 0$ be arbitrary, then if $|x - t| < \delta$, we have $|f(x) - f(t)| < \epsilon$. Applying the linear and monotone operator K_n , for $n \in \mathbb{N}$, to both sides of the inequality (wrt x), we have

$$(K_n f)(x) - f(t)(K_n 1)(x) < \epsilon (K_n 1)(x) \quad (\text{A.3})$$

$$\Rightarrow |(K_n f)(x) - f(t)(K_n 1)(x)| < \epsilon |(K_n 1)(x)|. \quad (\text{A.4})$$

By assumption, for any $\tilde{\epsilon} > 0$ there is an $N(\tilde{\epsilon}) \in \mathbb{N}$ satisfying

$$d_\infty(K_n x^k, x^k) = \|K_n x^k - x^k\|_\infty < \tilde{\epsilon} \quad \text{for } k \in \{0, 1, 2\} \forall n \geq N. \quad (\text{A.5})$$

This in particular implies

$$|(K_n 1)(x)| \leq \|K_n 1\|_\infty = \|K_n 1 - 1 + 1\|_\infty \leq \|K_n 1 - 1\|_\infty + 1 < \tilde{\epsilon} + 1 \quad (\text{A.6})$$

¹¹PAVEL PETROVICH KOROVKIN (1913 – 1985), Russian mathematician.

for all $n \geq N$. Using the triangle inequality and putting Eq. (A.6) into (A.4), we get the following inequality:

$$|(K_n f)(x) - f(t)| \leq |(K_n f)(x) - f(t)(K_n 1)(x)| + |f(t)(K_n 1)(x) - f(t)| \quad (\text{A.7})$$

$$\stackrel{(\text{A.4})}{<} \epsilon |(K_n 1)(x)| + |f(t)| |(K_n 1)(x) - 1| \quad (\text{A.8})$$

$$\stackrel{(\text{A.6})}{<} \epsilon(\tilde{\epsilon} + 1) + M \|K_n 1 - 1\|_\infty \quad \forall n \geq N \quad (\text{A.9})$$

$$< \epsilon(\epsilon + 1) + M\tilde{\epsilon} \quad \forall n \geq N. \quad (\text{A.10})$$

For $x = t$, we have the inequality

$$|(K_n f)(t) - f(t)| < \epsilon(\tilde{\epsilon} + 1) \quad \forall n \geq N. \quad (\text{A.11})$$

The RHS can be bounded from above by an arbitrarily small $\hat{\epsilon} > 0$, so that for some $N' = N(\hat{\epsilon}) \in \mathbb{N}$ we have

$$d_\infty(K_n f, f) < \hat{\epsilon} \quad \forall n \geq N'. \quad (\text{A.12})$$

■

In the following, we will explicitly construct a Korovkin sequence on $\mathcal{C}[a, b]$.

Remark A.6. In the following, we might restrict ourselves to the continuous functions $\mathcal{C}[0, 1]$ on $[0, 1]$. This is WLOG, since for any $x \in [a, b]$, we can transform it into $[0, 1]$ via a simple affine-linear transformation.

Definition A.7. Let $x \in [0, 1]$, then the *Bernstein¹² polynomials* are defined as

$$\beta_j^{(n)}(x) := \binom{n}{j} x^j (1-x)^{n-j} \in \mathcal{P}_n \quad \text{with } 0 \leq j \leq n, n \in \mathbb{N}_0 \quad (\text{A.13})$$

Example A.8. Here are some Bernstein polynomials:

$$\mathbf{n} = \mathbf{0} : \quad \beta_0^{(0)}(x) = 1$$

$$\mathbf{n} = \mathbf{1} : \quad \beta_0^{(1)}(x) = 1 - x \quad \beta_1^{(1)}(x) = x$$

$$\mathbf{n} = \mathbf{2} : \quad \beta_0^{(2)}(x) = (1-x)^2 \quad \beta_1^{(2)}(x) = 2x(1-x) \quad \beta_2^{(2)}(x) = x^2$$

$$\mathbf{n} = \mathbf{3} : \quad \beta_0^{(3)}(x) = (1-x)^3 \quad \beta_1^{(3)}(x) = 3x(1-x)^2 \quad \beta_2^{(3)}(x) = 3x^2(1-x) \quad \beta_3^{(3)}(x) = x^3$$

Remark A.9. The Bernstein polynomials $\beta_0^{(n)}, \dots, \beta_n^{(n)} \in \mathcal{P}_n$, for $n \in \mathbb{N}_0, \dots$

- (a) ... form a basis for the polynomial space \mathcal{P}_n ;
- (b) ... are positive on $[0, 1]$, i.e. $\beta_j^{(n)}(x) \geq 0$ for all $x \in [0, 1]$;
- (c) ... are symmetrical around $x = 1/2$, i.e. $\beta_j^{(n)}(x) = \beta_{n-j}^{(n)}(1-x)$;
- (d) ... are a *partition of unity* on $[0, 1]$, i.e.

$$\sum_{j=0}^n \beta_j^{(n)}(x) = 1 \quad \forall x \in [0, 1]. \quad (\text{A.14})$$

¹²SERGEI NATANOVICH BERNSTEIN (1880 – 1968)

Proof. (a) From linear algebra, we know that a set is a basis of a finite-dimensional vector space iff it has the same dimension and all vectors in the set are linearly independent. Since $\dim(\mathcal{P}_n) = n + 1$ and $\{\beta_0^{(n)}(x), \dots, \beta_n^{(n)}(x)\}$ contains $n + 1$ vectors, it only remains to show that the $\beta_0^{(n)}(x), \dots, \beta_n^{(n)}(x)$ are linearly independent. For this, note that [100, proof Lemma 2.5 ix)]

$$\beta_j^{(n)}(x) = \binom{n}{j} x^j (1-x)^{n-j} \in \mathcal{P}_n \quad \text{with } 0 \leq j \leq n \quad (\text{A.15})$$

$$= \binom{n}{j} x^j \sum_{i=0}^{n-j} \binom{n-j}{i} (-x)^i \quad (\text{A.16})$$

$$= \sum_{i=0}^{n-j} \binom{n}{j} \binom{n-j}{i} (-1)^i x^{j+i} \quad (\text{A.17})$$

$$= \sum_{i=j}^n \binom{n}{j} \binom{n-j}{i-j} (-1)^{i-j} x^i \quad (\text{A.18})$$

$$= \sum_{i=j}^n \binom{n}{i} \binom{i}{j} (-1)^{i-j} x^i \quad (\text{A.19})$$

Now consider the equation (with $\lambda_k \in \mathbb{R}$) for all $0 \leq k \leq n$:

$$0 = \sum_{k=0}^n \lambda_k \beta_k^{(n)} \stackrel{(\text{A.19})}{=} \sum_{k=0}^n \lambda_k \sum_{i=k}^n \binom{n}{i} \binom{i}{k} (-1)^{i-k} x^i \quad (\text{A.20})$$

For the double sum, we will use the following identity with $a_{ik} \in \mathbb{R}$ [101, p. 2]:

$$\sum_{k=0}^n \sum_{i=k}^n a_{ik} = \sum_{0 \leq k \leq i \leq n} a_{ik} = \sum_{i=0}^n \sum_{k=0}^i a_{ik} = \sum_{k=0}^n \sum_{i=0}^k a_{ki}, \quad (\text{A.21})$$

since $0 \leq i \leq n$, and for a fixed i , $0 \leq k \leq i$, and where we exchanged the indices ($i \mapsto k$, $k \mapsto i$). Thus, Eq. (A.20) becomes

$$0 = \sum_{k=0}^n \lambda_k \beta_k^{(n)} = \sum_{k=0}^n x^k \sum_{i=0}^k \lambda_i \binom{n}{k} \binom{k}{i} (-1)^{k-i} \quad (\text{A.22})$$

$$= \lambda_0 + (-n\lambda_0 + n\lambda_1)x + \left(\lambda_0 n(n-1) - \lambda_1 n(n-1) + \frac{\lambda_2 n(n-1)}{2} \right) x^2 + \dots \quad (\text{A.23})$$

Thus, every polynomial $p \in \mathcal{P}_n$ can be written as the linear combination of $\{\beta_0^{(n)}(x), \dots, \beta_n^{(n)}(x)\}$. Also, note that we already know that the monomials $\{x^0, x^1, \dots, x^n\}$ form a basis for \mathcal{P}_n , thus

$$\lambda_0 = 0 \quad (\text{A.24})$$

$$-n\lambda_0 + n\lambda_1 = 0 \quad (\text{A.25})$$

$$\left(\lambda_0 n(n-1) - \lambda_1 n(n-1) + \frac{\lambda_2 n(n-1)}{2} \right) = 0 \quad (\text{A.26})$$

\vdots

By putting $\lambda_0 = 0$ into Eq. (A.25), we get $\lambda_1 = 0$. Putting $\lambda_0 = \lambda_1 = 0$ into Eq. (A.26), we have $\lambda_2 = 0$. Thus, we can recursively show that $\lambda_k = 0$ for all $0 \leq k \leq n$, and hence the set $\left\{ \beta_j^{(n)}(x) \right\}_{0 \leq j \leq n}$ is linearly independent.

(b) This is straightforward, since $x^j \geq 0$, $(1-x)^{n-j} \geq 0$ and $\binom{n}{j} \geq 0$ for all $x \in [0, 1]$, hence $\beta_j^{(n)}(x) \geq 0$.

(c) We have

$$\beta_{n-j}^{(n)}(1-x) = \binom{n}{n-j} (1-x)^{n-j} x^j = \binom{n}{j} (1-x)^{n-j} x^j = \beta_j^{(n)}(x). \quad (\text{A.27})$$

(d) By the binomial theorem we have

$$\sum_{j=0}^n \beta_j^{(n)}(x) = \sum_{j=0}^n \binom{n}{j} x^j (1-x)^{n-j} = (x + (1-x))^n = 1. \quad (\text{A.28})$$

■

Definition A.10. For $n \in \mathbb{N}$, define the *Bernstein operator* $B_n : \mathcal{C}[0, 1] \rightarrow \mathcal{P}_n$ as

$$(B_n f)(x) := \sum_{j=0}^n f\left(\frac{j}{n}\right) \beta_j^{(n)}(x) \quad \forall f \in \mathcal{C}[0, 1], \quad (\text{A.29})$$

where the $\beta_j^{(n)}$ are the Bernstein polynomials from Def. A.7.

Remark A.11. The Bernstein operators B_n are linear operators on $\mathcal{C}[0, 1]$. They are also positive (and hence monotone) on $\mathcal{C}[0, 1]$, since the $\beta_j^{(n)}$ are positive, cf. Remark A.9 b).

Remark A.12. The Bernstein operators $B_n : \mathcal{C}[0, 1] \rightarrow \mathcal{P}_n$ are also bounded (in the sense of Def. 7.3) wrt $\|\cdot\|_\infty$ ($C = 1$ is one upper bound). By Theorem 7.5, they are thus also continuous on $\mathcal{C}[0, 1]$.

Proof. We show the boundedness of the Bernstein operators $B_n : \mathcal{C}[0, 1] \rightarrow \mathcal{P}_n$ directly. Let $f \in \mathcal{C}[0, 1]$ and $n \in \mathbb{N}$ be arbitrary, then

$$\|B_n f\|_\infty = \left\| \sum_{j=0}^n f\left(\frac{j}{n}\right) \beta_j^{(n)}(x) \right\|_\infty = \sup_{x \in [0, 1]} \left\{ \left| \sum_{j=0}^n f\left(\frac{j}{n}\right) \beta_j^{(n)}(x) \right| \right\} \quad (\text{A.30})$$

$$\leq \sup_{x \in [0, 1]} \left\{ \left| \sup_{x \in [0, 1]} \{ |f(x)| \} \sum_{j=0}^n \beta_j^{(n)}(x) \right| \right\} = \|f\|_\infty \sup_{x \in [0, 1]} \left\{ \sum_{j=0}^n \beta_j^{(n)}(x) \right\} = \|f\|_\infty, \quad (\text{A.31})$$

since the Bernstein polynomials form a partition of unity, cf. Remark A.9 d).

Hence, we have shown that $\|B_n f\|_\infty \leq \|f\|_\infty$, which proves the boundedness of B_n and that $C = 1$ is one upper bound for B_n . ■

Theorem A.13. The sequence $(B_n)_{n \in \mathbb{N}}$ with $B_n : \mathcal{C}[0, 1] \rightarrow \mathcal{P}_n$ according to Def. A.7 forms a Korovkin sequence on $\mathcal{C}[0, 1]$, cf. Def. A.4.

Proof. The Bernstein operators B_n for $n \in \mathbb{N}$ reproduce linear polynomials: Let $1 \in \mathcal{C}[0, 1]$ be the constant function that assigns each $x \in [0, 1]$ the value 1, then

$$B_n 1 = 1, \quad (\text{A.32})$$

and by the linearity of B_n we have $B_n(c) = c$ for any $c \in \mathbb{R}$.

We also have $B_n p_1 = p_1$, where $p_1 \in \mathcal{C}[0, 1]$ is the identity function:

$$B_n p_1 = \sum_{j=0}^n p_1 \left(\frac{j}{n} \right) \beta_j^{(n)}(x) = \sum_{j=0}^n \frac{j}{n} \beta_j^{(n)}(x) = \sum_{j=1}^n \frac{j}{n} \beta_j^{(n)}(x) \quad (\text{A.33})$$

$$= \sum_{j=1}^n \binom{n-1}{j-1} x^j (1-x)^{n-j} = \sum_{j=0}^{n-1} \binom{n-1}{j} x^{j+1} (1-x)^{n-j-1} \quad (\text{A.34})$$

$$= x \sum_{j=0}^{n-1} \beta_j^{(n-1)}(x) = x = p_1(x), \quad (\text{A.35})$$

since the $\beta_j^{(n-1)}$ form a partition of unity, cf. Remark A.9 d). This implies that $B_n(\alpha p_1 + c) = \alpha p_1 + c$ for any real α, c .

Now consider $p_2 \in \mathcal{P}_2$, where $p_2(x) := x^2$, and consider the following sequence of polynomials of degree 2 for $n \geq 2$:

$$f_n(x) := \frac{x(nx-1)}{n-1} = \frac{n}{n-1}x^2 - \frac{1}{n-1}x \in \mathcal{P}_2. \quad (\text{A.36})$$

For $n \geq 2$, we have

$$\begin{aligned} (B_n f_n)(x) &= \sum_{j=0}^n f_n \left(\frac{j}{n} \right) \beta_j^{(n)}(x) = \sum_{j=0}^n f_n \left(\frac{j}{n} \right) \binom{n}{j} x^j (1-x)^{n-j} \\ &= \sum_{j=0}^n \frac{j(j-1)}{n(n-1)} \binom{n}{j} x^j (1-x)^{n-j} = \sum_{j=2}^n \frac{j(j-1)}{n(n-1)} \binom{n}{j} x^j (1-x)^{n-j} \\ &= \sum_{j=2}^n \frac{(n-2)!}{(n-j)!(j-2)!} x^j (1-x)^{n-j} = x^2 \sum_{j=0}^{n-2} \binom{n-2}{j} x^j (1-x)^{n-j-2} \\ &= x^2 \sum_{j=0}^{n-2} \beta_j^{(n-2)} = x^2 = p_2(x), \end{aligned} \quad (\text{A.37})$$

since the Bernstein polynomials $\{\beta_j^{(n-2)}\}_{0 \leq j \leq n-2}$ form a partition of unity, cf. Remark A.9 d).

Now note that any polynomial $p \in \mathcal{P}_2$ can be written as $p = \alpha p_2 + \beta p_1 + \gamma$ for some $\alpha, \beta, \gamma \in \mathbb{R}$: Using the linearity and boundedness of the Bernstein operators (where $C = 1$ is one upper bound), we have:

$$B_n p = \alpha B_n p_2 + \beta B_n p_1 + B_n \gamma = \alpha B_n p_2 + \beta p_1 + \gamma, \quad (\text{A.38})$$

since $B_n p_1 = p_1$ and $B_n \gamma = \gamma$, as we have previously shown. Further:

$$\begin{aligned}
\|B_n p - p\|_\infty &= \|\alpha B_n p_2 - \alpha p_2\|_\infty = |\alpha| \|B_n p_2 - p_2\|_\infty \stackrel{(A.37)}{=} |\alpha| \|B_n(p_2 - f_n)\|_\infty \\
&\leq |\alpha| \|p_2 - f_n\|_\infty = |\alpha| \sup_{x \in [0,1]} \{|p_2(x) - f_n(x)|\} \\
&= |\alpha| \sup_{x \in [0,1]} \left\{ \left| x^2 - \frac{nx^2}{n-1} - \frac{x}{n-1} \right| \right\} = |\alpha| \sup_{x \in [0,1]} \left\{ \left| -\frac{x^2}{n-1} - \frac{x}{n-1} \right| \right\} \\
&= |\alpha| \sup_{x \in [0,1]} \left\{ \frac{x^2 + x}{n-1} \right\} \stackrel{x=1}{=} \frac{2|\alpha|}{n-1} \xrightarrow{n \rightarrow \infty} 0.
\end{aligned} \tag{A.39}$$

■

Proof. (Weierstrass Approximation Theorem) Consider the sequence $(B_n)_{n \in \mathbb{N}}$ of Bernstein operators, which are a Korovkin sequence on $\mathcal{C}[0,1]$, cf. Theorem A.13. Let $f \in \mathcal{C}[0,1]$ and $\epsilon > 0$, then by Theorem A.5 there exists an $N(\epsilon) \in \mathbb{N}$ s.t. for all $n \geq N(\epsilon)$ we have $d_\infty(B_n f, f) < \epsilon$. Noting that $B_n f \in \mathcal{P}_n \subset \mathcal{P}$ completes the proof. ■

Example A.14. Let $f : \mathcal{C}[0, \pi] \rightarrow \mathcal{C}[0, \pi], x \mapsto \sin x$. To approximate this function arbitrarily well, we will consider $(B_n f)_{n \in \mathbb{N}}$, where $(B_n)_{n \in \mathbb{N}}$ is the sequence of Bernstein operators. Since in Def. A.10 we considered the B_n to be defined on $\mathcal{C}[0,1]$, we need an affine-linear transformation to map $[0, \pi]$ to $[0,1]$. For this, consider $x \mapsto x/\pi$. And since we considered f to be defined on $[0,1]$ in Def. A.10, consider the mapping $x \mapsto \pi x$ to get to $[0, \pi]$. With this mapping, the Bernstein operators B_n are defined on $\mathcal{C}[0, \pi]$ as follows:

$$(B_n f)(x) = \sum_{j=0}^n \sin\left(\pi \frac{j}{n}\right) \beta_j^{(n)}(x/\pi) = \frac{1}{\pi^j} \sum_{j=0}^n \sin\left(\pi \frac{j}{n}\right) \binom{n}{j} x^j (1 - x/\pi)^{n-j}. \tag{A.40}$$

For different $n \in \mathbb{N}$, Fig. 7 shows some approximating polynomials $B_n f \in \mathcal{P}_n$.




Figure 7: Approximation of $\sin(x)$ by linear combination of Bernstein polynomials on $[a, b]$.

B Construction of \mathbb{R}

There are two possibilities to construct the real numbers \mathbb{R} :

1. Dedekind completion: Every non-empty subset has a least upper bound (wrt \leq).
2. Cauchy completion: Every Cauchy sequence converges wrt $|\cdot|$.

We will use the second definition of completion, and follow Ref. [102, 103].

Definition B.1. The (archimedean) absolute value of \mathbb{Q} is the function

$$|\cdot| : \mathbb{Q} \rightarrow \mathbb{Q}_{\geq 0}, |x| := \begin{cases} x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases} \quad (\text{B.1})$$

Definition B.2. A *Cauchy sequence* of rational numbers is a sequence $(x_n)_{n \in \mathbb{N}}$ s.t. for every $\epsilon \in \mathbb{Q}_{>0}$ there exists an $N = N(\epsilon) \in \mathbb{N}$ s.t. for all $m, n \geq N$, $|x_m - x_n| < \epsilon$.

Theorem B.3. Let $(x_n)_{n \in \mathbb{N}}$ and $(y_n)_{n \in \mathbb{N}}$ be two Cauchy sequences in \mathbb{Q} , then their sum $(x_n + y_n)_{n \in \mathbb{N}}$ is also Cauchy.

Proof. First, we note that the sum of two rational numbers is also rational. Since both $(x_n)_{n \in \mathbb{N}}$ and $(y_n)_{n \in \mathbb{N}}$ are Cauchy sequences, for all $\epsilon \in \mathbb{Q}_{>0}$ there is an $N' \in \mathbb{N}$ s.t. for all $m', n' \geq N'$, we have $|x_{m'} - x_{n'}| < \epsilon$ and an $\tilde{N} \in \mathbb{N}$ s.t. for all $\tilde{m}, \tilde{n} \geq \tilde{N}$, we have $|y_{\tilde{m}} - y_{\tilde{n}}| < \epsilon$. Let $N := \max\{N', \tilde{N}\}$, then for all $m, n \geq N$, we have

$$|x_m + y_m - (x_n + y_n)| = |x_m - x_n + y_m - y_n| \leq |x_m - x_n| + |y_m - y_n| < 2\epsilon,$$

which proves that $(x_n + y_n)_{n \in \mathbb{N}}$ is also a Cauchy sequence. ■

Theorem B.4. Let $(x_n)_{n \in \mathbb{N}}$ and $(y_n)_{n \in \mathbb{N}}$ be two Cauchy sequences in \mathbb{Q} , then their product $(x_n \cdot y_n)_{n \in \mathbb{N}}$ is also Cauchy.

Proof. First, we note that the multiplication and division of two rational numbers is rational. By Theorem 5.59, Cauchy sequences are bounded, hence choose $M \in \mathbb{Q}$ s.t. $|x_n| \leq M$ and $|y_n| \leq M$ for all $n \in \mathbb{N}$, and note that for any $\epsilon \in \mathbb{Q}_{>0}$ there exists $N' \in \mathbb{N}$ s.t. for all $m', n' \geq N'$, we have $|x_{m'} - x_{n'}| < \epsilon/M$ and an $\tilde{N} \in \mathbb{N}$ s.t. for all $\tilde{m}, \tilde{n} \geq \tilde{N}$, $|y_{\tilde{m}} - y_{\tilde{n}}| < \epsilon/M$. Let $N := \max\{N', \tilde{N}\}$, then for all $m, n \geq N$, we have

$$\begin{aligned} |x_m y_m - x_n y_n| &= |x_m y_m + x_m y_n - x_m y_n - x_n y_n| = |x_m (y_m - y_n) + y_n (x_m - x_n)| \\ &\leq |x_m (y_m - y_n)| + |y_n (x_m - x_n)| = |x_m| |y_m - y_n| + |y_n| |x_m - x_n| \\ &\leq M \cdot \epsilon/M + M \cdot \epsilon/M = 2\epsilon, \end{aligned}$$

where we used that $|xy| = |x| |y|$. ■

Remark B.5. The set of Cauchy sequences in \mathbb{Q} forms a ring, cf. Def. 1.12, where $0 = (0)_{n \in \mathbb{N}}$ and $1 = (1)_{n \in \mathbb{N}}$ are the neutral elements for the additive and multiplicative operation respectively. Since multiplication is commutative, we have a commutative ring.

Remark B.6. The set of Cauchy sequences in \mathbb{Q} does not form a field, cf. Def. 1.13, though, since many Cauchy sequences do not have a multiplicative inverse, such as $(1, 0, 0, 0, \dots)$. Thus, the set of Cauchy sequences in \mathbb{Q} does not form a field.

Definition B.7. We say that a rational Cauchy sequence $(x_n)_{n \in \mathbb{N}}$ is *equivalent to zero* if $\lim_{n \rightarrow \infty} |x_n| = 0$. We say that two rational Cauchy sequences $(x_n)_{n \in \mathbb{N}}$ and $(y_n)_{n \in \mathbb{N}}$ are rational, denoted by $(x_n)_{n \in \mathbb{N}} \sim (y_n)_{n \in \mathbb{N}}$, if $(x_n - y_n)_{n \in \mathbb{N}}$ is equivalent to 0.

Theorem B.8. The operation \sim is an equivalence relation on the set of all Cauchy sequences in \mathbb{Q} .

Proof. Note that any rational Cauchy sequence is equivalent to itself, since $(x_n - x_n)_{n \in \mathbb{N}} = (0)_{n \in \mathbb{N}}$, and $\lim_{n \rightarrow \infty} |0| = 0$. We also have symmetry, since $|x_n - y_n| = |y_n - x_n|$. Finally, if $(x_n)_{n \in \mathbb{N}} \sim (y_n)_{n \in \mathbb{N}}$ and $(y_n)_{n \in \mathbb{N}} \sim (z_n)_{n \in \mathbb{N}}$, then $(x_n)_{n \in \mathbb{N}} \sim (z_n)_{n \in \mathbb{N}}$, since $|x_n - z_n| = |x_n - y_n + y_n - z_n| \leq |x_n - y_n| + |y_n - z_n|$, thus the operation \sim is also transitive.

Thus, according to Def. 8.2, \sim forms an equivalence relation. ■

Definition B.9 (Real numbers). We define $\mathbb{R} := \mathbb{Q}/\sim$. We extend the absolute value of \mathbb{Q} to \mathbb{R} via

$$|[(x_n)_{n \in \mathbb{N}}]| := [(|x_n|)_{n \in \mathbb{N}}], \quad (\text{B.2})$$

where $(x_n)_{n \in \mathbb{N}}$ is a Cauchy sequence in \mathbb{Q} .

Remark B.10. We embed \mathbb{Q} in \mathbb{R} via the map $x \mapsto [(x)_{n \in \mathbb{N}}]$, i.e. the equivalence class of a constant sequence, whose elements are always x . This map is injective. It is standard to abuse notation and write x for $[(x)_{n \in \mathbb{N}}]$. That way, we can see \mathbb{Q} as a subset of \mathbb{R} .

Before proving that \mathbb{Q} is dense in \mathbb{R} and that \mathbb{R} is a field, we need to look at the algebraic structure of \mathbb{R} , and at what it means to say $[(a_n)_{n \in \mathbb{N}}] < [(b_n)_{n \in \mathbb{N}}]$.

Definition B.11. Let $s, t \in \mathbb{R}$, i.e. there are rational Cauchy sequences $(s_n)_{n \in \mathbb{N}}$ and $(t_n)_{n \in \mathbb{N}}$ s.t. $s = [(s_n)_{n \in \mathbb{N}}]$ and $t = [(t_n)_{n \in \mathbb{N}}]$.

a) Define the map $+: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, $s + t := [(s_n + t_n)_{n \in \mathbb{N}}]$,

b) and define $\cdot: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, $s \cdot t := [(s_n \cdot t_n)_{n \in \mathbb{N}}]$.

Remark B.12. Since we are dealing with equivalence classes, and $(s_n)_{n \in \mathbb{N}}$ is merely a representation of $[(s_n)_{n \in \mathbb{N}}]$, we need to verify that the above definitions are well-defined, i.e. representation-independent.

Theorem B.13. The addition and multiplication of equivalence classes as defined in Def. B.11 are well-defined.

Proof. Let $s, t \in \mathbb{R}$, i.e. there exist rational Cauchy sequences $(s_n)_{n \in \mathbb{N}}$ and $(t_n)_{n \in \mathbb{N}}$ s.t. $s = [(s_n)_{n \in \mathbb{N}}]$ and $t = [(t_n)_{n \in \mathbb{N}}]$. Now let $(s'_n)_{n \in \mathbb{N}}, (t'_n)_{n \in \mathbb{N}}$ be rational Cauchy sequences s.t. $(s_n)_{n \in \mathbb{N}} \sim (s'_n)_{n \in \mathbb{N}}$ and $(t_n)_{n \in \mathbb{N}} \sim (t'_n)_{n \in \mathbb{N}}$, i.e. $[(s_n)_{n \in \mathbb{N}}] = [(s'_n)_{n \in \mathbb{N}}]$ and $[(t_n)_{n \in \mathbb{N}}] = [(t'_n)_{n \in \mathbb{N}}]$.

Since

$$|s'_n + t'_n - (s_n + t_n)| \leq |s'_n - s_n| + |t'_n - t_n| \xrightarrow{n \rightarrow \infty} 0,$$

we have $(s'_n + t'_n)_{n \in \mathbb{N}} \sim (s_n + t_n)_{n \in \mathbb{N}}$, and thus $[(s'_n + t'_n)_{n \in \mathbb{N}}] = [(s_n + t_n)_{n \in \mathbb{N}}]$, which shows the well-definedness of $+$.

For the multiplication \cdot , note that

$$\begin{aligned} |s'_n t'_n - s_n t_n| &= |s'_n t'_n + t'_n s_n - t'_n s_n - s_n t_n| = |t'_n (s'_n - s_n) + s_n (t'_n - t_n)| \\ &\leq |t'_n| |s'_n - s_n| + |s_n| |t'_n - t_n| \\ &\leq M |s'_n - s_n| + M |t'_n - t_n| \xrightarrow{n \rightarrow \infty} 0, \end{aligned}$$

where we proceeded as in the proof of Theorem B.4 and chose an $M \in \mathbb{Q}$ s.t. $|t'_n| \leq M$ and $|s_n| \leq M$ for all $n \in \mathbb{N}$, cf. Theorem 5.59. Thus, we have shown that $(s'_n t'_n)_{n \in \mathbb{N}} \sim (s_n t_n)_{n \in \mathbb{N}}$, and hence $[(s'_n t'_n)_{n \in \mathbb{N}}] = [(s_n t_n)_{n \in \mathbb{N}}]$, showing the well-definedness of \cdot . \blacksquare

Theorem B.14. \mathbb{R} , equipped with the two operations $+$ and \cdot , is a field.

Proof. Here, we will only prove that every $s \in \mathbb{R} \setminus \{0\}$ has a multiplicative inverse (the rest is generally easier to prove than this one).

We shall first understand what the existence of the multiplicative inverse means. Let $(s_n)_{n \in \mathbb{N}}$ be a rational Cauchy sequence s.t. $s = [(s_n)_{n \in \mathbb{N}}]$. Since $s \neq 0$, we know that $(s_n)_{n \in \mathbb{N}} \not\sim (0)_{n \in \mathbb{N}}$, i.e. $(s_n)_{n \in \mathbb{N}}$ does not converge to 0. We now need to find a multiplicative inverse t s.t. $s \cdot t = 1$. While this might seem trivial (since every $s_n \in \mathbb{Q}$ has a multiplicative inverse, i.e. we could choose $t_n = s_n^{-1}$), there is a subtle difficulty. The fact that s is non-zero does *not* mean that all sequence elements of $(s_n)_{n \in \mathbb{N}}$ are unequal to zero. For example, let $s = 1$, then $(s_n)_{n \in \mathbb{N}} = (0, 0, 0, 1, 1, 1, 1, 1, 1, 1, \dots)$ is in the equivalence class of $[(1)_{n \in \mathbb{N}}]$. However, the point is that *eventually*, the sequence elements must be non-zero (since s is non-zero), and thus there is an $N \in \mathbb{N}$ s.t. for $n \geq N$, we have $s_n \neq 0$. Thus, define a sequence $(t_n)_{n \in \mathbb{N}}$ that for $n < N$ is zero and for $n \geq N$ let $t_n := s_n^{-1}$. Now the multiplicative inverse of s is $t = [(t_n)_{n \in \mathbb{N}}]$. \blacksquare

Definition B.15 (Order of \mathbb{R}). Let $s \in \mathbb{R}$, i.e. there is a rational Cauchy sequence $(s_n)_{n \in \mathbb{N}}$ s.t. $[(s_n)_{n \in \mathbb{N}}] = s$. We say that s is *positive*, denoted by $s > 0$, if $s \neq 0$ and if there is an $N \in \mathbb{N}$ s.t. $s_n > 0$ for all $n \geq N$.

Let $t \in \mathbb{R}$, i.e. there is a rational Cauchy sequence $(t_n)_{n \in \mathbb{N}}$ s.t. $[(t_n)_{n \in \mathbb{N}}] = t$. Then we write $s > t$ if $s - t$ is positive.

Theorem B.16. The order of the real numbers is well-defined.

Proof. Fix $s \in \mathbb{R}$, i.e. $s = [(s_n)_{n \in \mathbb{N}}]$ for a rational Cauchy sequence $(s_n)_{n \in \mathbb{N}}$. Also, let s be positive, i.e. there is an $N \in \mathbb{N}$ s.t. $s_n > 0$ for all $n \geq N$. We need to show that for any other representative of $[(s_n)_{n \in \mathbb{N}}]$, i.e. $(s'_n)_{n \in \mathbb{N}} \in [(s_n)_{n \in \mathbb{N}}]$ with $(s'_n)_{n \in \mathbb{N}} \sim (s_n)_{n \in \mathbb{N}}$, there is an $N' \in \mathbb{N}$ s.t. $s'_n > 0$ for all $n \geq N'$.

We first show that there exists a rational number $r \in \mathbb{Q}_{>0}$ and an $\tilde{N} \in \mathbb{N}$ s.t. $s_n > r$ for all $n \geq \tilde{N}$ by contradiction. Suppose that for all $r \in \mathbb{Q}_{>0}$ and all $\tilde{N} \in \mathbb{N}$ there exists an $n = n(r, \tilde{N}) \geq \tilde{N}$ s.t. $s_n \leq r$. Then there is a subsequence $(s_{n_k})_{k \in \mathbb{N}}$ of $(s_n)_{n \in \mathbb{N}}$ s.t. $0 < s_{n_k} \leq 1/k$. We will prove this via induction. Set $n_1 := n(1, N)$. Assume we have constructed n_1, n_2, \dots, n_k s.t. $n_1 < n_2 < \dots < n_k$ and $0 < s_{n_j} \leq 1/j$ for $j \in \{1, \dots, k\}$. Then set $n_{k+1} := n(1/(k+1), n_k + 1)$. We have $n_k < n_{k+1}$ and $0 < s_{n_{k+1}} \leq 1/(k+1)$.

Since $(s_n)_{n \in \mathbb{N}}$ is Cauchy, there exists an $M_k \in \mathbb{N}$ s.t. $|s_n - s_m| < 1/k$ for all $n, m \geq M_k$. Choose $l \geq k$ s.t. $n_l \geq M_k$. Then for $n \geq M_k$, we have

$$|s_n| = |s_n - s_{n_l} + s_{n_l}| \leq |s_n - s_{n_l}| + |s_{n_l}| < \frac{1}{k} + \frac{1}{l} \leq \frac{2}{k} \xrightarrow{k \rightarrow \infty} 0. \quad (\text{B.3})$$

This means that $(s_n)_{n \in \mathbb{N}}$ converges to 0, and thus $s = 0$, which is a contradiction. Hence, there exists a rational number $r \in \mathbb{Q}_{>0}$ and an $\tilde{N} \in \mathbb{N}$ s.t. $s_n > r$ for all $n \geq \tilde{N}$. Let $(s'_n)_{n \in \mathbb{N}}$ be another representative of $[(s_n)_{n \in \mathbb{N}}]$, i.e. $(s_n)_{n \in \mathbb{N}} \sim (s'_n)_{n \in \mathbb{N}}$, and thus $s'_n - s_n \xrightarrow{n \rightarrow \infty} 0$. This in turn implies that $|s'_n - s_n| < r$ for $n \geq N'$. Let $n \geq \max\{\tilde{N}, N'\}$, then we have

$$s'_n = s_n - (s_n - s'_n) \geq s_n - |s_n - s'_n| > r - r = 0,$$

since $s_n - s'_n \leq |s_n - s'_n|$, which implies $-(s_n - s'_n) \geq -|s_n - s'_n|$. ■

Theorem B.17. Let $s, t, r \in \mathbb{R}$, then if $s > t$, we also have $s + r > t + r$.

Proof. Let $(s_n)_{n \in \mathbb{N}}, (t_n)_{n \in \mathbb{N}}, (r_n)_{n \in \mathbb{N}}$ be rational Cauchy sequences s.t. $s = [(s_n)_{n \in \mathbb{N}}]$, $t = [(t_n)_{n \in \mathbb{N}}]$ and $r = [(r_n)_{n \in \mathbb{N}}]$. Since $s > t$, we know that there is an $N \in \mathbb{N}$ s.t. for all $n \geq N$, we have

$$s_n - t_n > 0 \Leftrightarrow s_n > t_n \Leftrightarrow s_n + r_n > t_n + r_n \Leftrightarrow \underbrace{(s_n + r_n) - (t_n + r_n)}_{=s_n - t_n} > 0$$

Since $s_n - t_n$ does not go to 0 for $n \rightarrow \infty$ (since $s > t$), we have that $(s_n + r_n) - (t_n + r_n)$ does not either, and thus $s + r > t + r$. ■

The density of \mathbb{Q} in \mathbb{R} follows almost immediately from the construction of \mathbb{R} from \mathbb{Q} .

Theorem B.18. \mathbb{Q} is dense in \mathbb{R} , i.e. for any $\epsilon \in \mathbb{Q}_{>0}$ and $s \in \mathbb{R}$ there is a rational number $r \in \mathbb{Q}$ s.t. $|s - r| < \epsilon$.

Proof. Since $s \in \mathbb{R}$, there is a rational Cauchy sequence $(s_n)_{n \in \mathbb{N}}$ s.t. $s = [(s_n)_{n \in \mathbb{N}}]$. Since $(s_n)_{n \in \mathbb{N}}$ is Cauchy, there is an $N \in \mathbb{N}$ s.t. for all $m, n \geq N$, we have $|s_n - s_m| < \epsilon$. Let $r := s_N \in \mathbb{Q}$, and note that we can embed it into \mathbb{R} as $[(a_N)_{n \in \mathbb{N}}]$. Thus, $|s - r| = [(|s_n - s_N|)_{n \in \mathbb{N}}]$. For $n \geq N$, we have $|s_n - s_N| < \epsilon$ and thus $s_n - s_N < \epsilon$ and $s_N - s_n < \epsilon$, since for any rational number a , $a \leq |a|$. Thus, $(s_n - s_N) - \epsilon$ and $(s_N - s_n) - \epsilon$ are negative, i.e. $s - r < \epsilon$ and $r - s < \epsilon$, which can be summarized to $|s - r| < \epsilon$. ■

Remark B.19. The density of \mathbb{Q} in \mathbb{R} implies that we could replace $\epsilon \in \mathbb{Q}_{>0}$ with $\epsilon \in \mathbb{R}_{>0}$ throughout.

Theorem B.20. \mathbb{R} has the Archimedean property, i.e. for all $s, t \in \mathbb{R}$ with $s > 0$, there exists a natural number $m \in \mathbb{N}$ s.t. $m \cdot s > t$.

Proof. Since $s, t \in \mathbb{R}$, there exist rational Cauchy sequences $(s_n)_{n \in \mathbb{N}}$ and $(t_n)_{n \in \mathbb{N}}$ s.t. $s = [(s_n)_{n \in \mathbb{N}}]$ and $t = [(t_n)_{n \in \mathbb{N}}]$.

Recally that by $m \in \mathbb{N} \subset \mathbb{Q}$, we mean the m that is embedded into \mathbb{R} , i.e. $[(m)_{n \in \mathbb{N}}]$. Thus, we need to show that there exists an $m, N \in \mathbb{N}$ s.t. $ms_n - t_n > 0$ for all $n \geq N$ and that $(ms_n - t_n)_{n \in \mathbb{N}}$ does not converge to 0, cf. Definition B.15.

We will prove the first part by contradiction. Assume that for all $m, N \in \mathbb{N}$, there exists an $n \geq N$ s.t. $ms_n - t_n \leq 0$. For this, note that $(t_n)_{n \in \mathbb{N}}$ is bounded, cf. Theorem 5.59, i.e. there is an $M \in \mathbb{Q}$ s.t. $t_n \leq M$ for all $n \in \mathbb{N}$. By the Archimedean property of \mathbb{Q} , for any $\epsilon \in \mathbb{Q}_{>0}$ there is an $m \in \mathbb{N}$ s.t. $m\epsilon/2 > M$, or equivalently $M/m < \epsilon/2$. Thus,

$$ms_n - t_n \leq 0 \Leftrightarrow s_n \leq \frac{t_n}{m} \leq \frac{M}{m} < \frac{\epsilon}{2}. \quad (\text{B.4})$$

Since $(s_n)_{n \in \mathbb{N}}$ is Cauchy, there exists an $N \in \mathbb{N}$ s.t. for all $n', k \geq N$, $|s_k - s_{n'}| < \epsilon/2$, i.e. $s_k - s_{n'} < \epsilon/2$. Now choose an $n' \geq n \geq N$, then for all $k \geq N$, we have

$$s_k < s_{n'} + \frac{\epsilon}{2} \stackrel{\text{(B.4)}}{<} \epsilon.$$

Thus, s cannot be positive, contradicting the assumption that $s > 0$. Hence, we have shown that there exists an $m, N \in \mathbb{N}$ s.t. $ms_n - t_n > 0$ for all $n \geq N$.

If $(ms_n - t_n)_{n \in \mathbb{N}}$ converges to 0, e.g. for $(s_n)_{n \in \mathbb{N}} = (1)_{n \in \mathbb{N}}$ and $(t_n)_{n \in \mathbb{N}} = (m)_{n \in \mathbb{N}}$, then we can choose $m+1$ instead of m , since $(m+1)s_n - t_n = ms_n - t_n + s_n \xrightarrow{n \rightarrow \infty} \lim_{n \rightarrow \infty} s_n > 0$, since $s \neq 0$. ■

We can now prove that \mathbb{R} is complete.

Proposition B.21. Let $(x_n)_{n \in \mathbb{N}}$ be a Cauchy sequence of *real numbers*, then it converges to an $x \in \mathbb{R}$, i.e. \mathbb{R} is complete.

Proof. Fix an x_n of the sequence. By the density of \mathbb{Q} in \mathbb{R} , cf. Theorem B.18, we know there exists a $q_n \in \mathbb{Q}$ s.t. $|x_n - q_n| < 1/n$. We will now show that $(q_n)_{n \in \mathbb{N}}$ is Cauchy in \mathbb{Q} . Let $\epsilon \in \mathbb{Q}_{>0}$, then by the Archimedean property of \mathbb{Q} , there exists an $N \in \mathbb{N}$ s.t. $N\epsilon/3 > 1$, or equivalently $1/N < \epsilon/3$. Since $(x_n)_{n \in \mathbb{N}}$ is Cauchy in \mathbb{R} , there exists an $M \in \mathbb{R}$ s.t. for all $n, m \geq M$, we have $|x_n - x_m| < \epsilon/3$. Now, if $n, m \geq \max\{N, M\}$, we have

$$\begin{aligned} |q_n - q_m| &= |q_n - x_n + x_n - x_m + x_m - q_m| \leq |q_n - x_n| + |x_n - x_m| + |x_m - q_m| \\ &< \frac{1}{n} + \frac{\epsilon}{3} + \frac{1}{m} \leq \frac{2}{N} + \frac{\epsilon}{3} < \epsilon, \end{aligned}$$

which proves that $(q_n)_{n \in \mathbb{N}}$ is a Cauchy sequence, and thus it represents a real number, which we shall denote by x . We now want to show that $(x_n)_{n \in \mathbb{N}}$ converges to x . First of all, by definition of x , $q_n - x \xrightarrow{n \rightarrow \infty} 0$. We also know that $|x_n - q_n| < 1/n$, cf. above. Thus,

$$|x_n - x| = |x_n - q_n + q_n - x| \leq |x_n - q_n| + |q_n - x| < \frac{1}{n} + |q_n - x| \xrightarrow{n \rightarrow \infty} 0.$$

■

Remark B.22. For the proof of the uniqueness of the completion of \mathbb{Q} , we refer to step 5 of the proof of Proposition 5.92.

Theorem B.23. Let $a, b \in \mathbb{R}$ s.t. $a < b$, then there is a rational number $q \in \mathbb{Q}$ s.t. $q \in (a, b)$.

Proof [104, 105]. Since \mathbb{R} has the Archimedean property, cf. Theorem B.20, there exists an $N \in \mathbb{N}$ s.t.

$$N \cdot (b - a) > 1 \Leftrightarrow Nb - Na > 1 \Leftrightarrow Nb - 1 > Na. \quad (\text{B.5})$$

If $Nb \in \mathbb{Z}$, set $m := Nb - 1$, otherwise set $m := \lfloor Nb \rfloor \in \mathbb{Z}$, and we have

$$Nb > m \geq Nb - 1 \stackrel{\text{(B.5)}}{>} Na \Rightarrow b > \frac{m}{N} > a,$$

where $m/N \in \mathbb{Q}$. ■

Corollary B.24. Let $a, b \in \mathbb{R}$ s.t. $a < b$, then there are infinitely many rational numbers that are contained in (a, b) .

Proof. Let $q_1 \in \mathbb{Q}$ be s.t. $a < q_1 < b$, then we can construct a $q_2 \in \mathbb{Q}$ s.t. $a < q_1 < q_2 < b$, now repeat this procedure infinitely many times. ■

References

- [1] Michael M. Bronstein et al. *Geometric Deep Learning: Grids, Groups, Graphs, Geodesics, and Gauges*. 2021. arXiv: 2104.13478 [cs.LG] (cit. on pp. 2, 3).
- [2] Darij Grinberg. *An introduction to the algebra of rings and fields. Text for Math 332 Winter 2023 and Winter 2025 at Drexel University*. accessed on 06/02/25. 2025. URL: <https://www.cip.ifi.lmu.de/~grinberg/t/23wa/23wa.pdf> (cit. on p. 4).
- [3] *Chapter 1. Sigma-Algebras*. URL: <https://www.math.lsu.edu/~sengupta/7312s02/sigmaalg.pdf> (cit. on p. 5).
- [4] John K. Hunter. *CHAPTER 3. Measurable functions*. URL: https://www.math.ucdavis.edu/~hunter/measure_theory/measure_notes_ch3.pdf (cit. on p. 5).
- [5] Davide Giraudo (<https://math.stackexchange.com/users/9849/davide-giraudo>). *Finite measures are σ -finite*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com> (version: 2015-11-23). eprint: <https://math.stackexchange.com/q/144687>. URL: <https://math.stackexchange.com/q/144687> (cit. on p. 6).
- [6] Harkrishan Lal Vasudeva Satish Shirali. *Measure and Integration*. 1st ed., p. 598. DOI: 10.1007/978-3-030-18747-7. URL: <https://katalogplus.sub.uni-hamburg.de/vufind/Record/1678681113?rank=1> (cit. on p. 6).
- [7] Joel Feldman. *The Bolzano-Weierstrass Theorem*. online. accessed on 07/17/25. 2014. URL: <https://personal.math.ubc.ca/~feldman/m320/bolzano.pdf> (cit. on p. 8).
- [8] Ben Frankel (<https://math.stackexchange.com/users/112897/ben-frankel>). *Bolzano-Weierstrass theorem (complex case)*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/1250430> (version: 2015-04-24). eprint: <https://math.stackexchange.com/q/1250430>. URL: <https://math.stackexchange.com/q/1250430> (cit. on p. 8).
- [9] Dirk Werner. *Funktionalanalysis*. Springer Spektrum, Berlin, Heidelberg, 2018, p. 586. ISBN: 978-3-662-55406-7. DOI: <https://link.springer.com/book/10.1007%2F978-3-662-55407-4> (cit. on pp. 13, 81).
- [10] Wikipedia contributors. *Equivalence of metrics*. 2020. URL: https://en.wikipedia.org/wiki/Equivalence_of_metrics (cit. on p. 14).
- [11] Armin Iske. “Funktionalanalysis. Universität Hamburg, Sommersemester 2019”. unpublished lecture notes. 2019 (cit. on pp. 15, 18).
- [12] Fraïssé. *Follow up to a question, why does proof $\rho(x, y) = \frac{d(x, y)}{1+d(x, y)}$ work*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/1860725> (version: 2021-10-23). eprint: <https://math.stackexchange.com/q/1860725>. URL: <https://math.stackexchange.com/q/1860725> (cit. on p. 15).
- [13] Tigran Hakobyan (<https://math.stackexchange.com/users/18140/tigran-hakobyan>). *Showing $\rho(x, y) = \frac{d(x, y)}{1+d(x, y)}$ is a metric*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/686797> (version: 2014-02-23). eprint: <https://math.stackexchange.com/q/686797>. URL: <https://math.stackexchange.com/q/686797> (cit. on p. 15).

- [14] user160823 (<https://math.stackexchange.com/users/160823/user160823>). *Prove intersection of open balls is another open ball*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/886810> (version: 2022-01-18). eprint: <https://math.stackexchange.com/q/886810>. URL: <https://math.stackexchange.com/q/886810> (cit. on p. 16).
- [15] *Notes 3. UNIFORM CONVERGENCE*. online. accessed on 07/18/2025. URL: https://www.math.cuhk.edu.hk/course_builder/1718/math2060a/Notes%5C%203.%5C%20Uniform%5C%20convergence%5C%202018.pdf (cit. on pp. 19, 20).
- [16] *Uniform continuity and convergence*. online. accessed on 07/18/2025. URL: <https://www.dpmms.cam.ac.uk/~cmw50/resources/M2PM1/M2PM1Ch1.pdf> (cit. on p. 19).
- [17] CodeKingPlusPlus (<https://math.stackexchange.com/users/38879/codekingplusplus>). *Prove: If a sequence converges, then every subsequence converges to the same limit*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/213285> (version: 2012-10-13). eprint: <https://math.stackexchange.com/q/213285>. URL: <https://math.stackexchange.com/q/213285> (cit. on p. 20).
- [18] Jungleshrimp (<https://math.stackexchange.com/users/484248/jungleshrimp>). *A closed bounded set contains its supremum*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/2752236> (version: 2018-04-24). eprint: <https://math.stackexchange.com/q/2752236>. URL: <https://math.stackexchange.com/q/2752236> (cit. on p. 20).
- [19] user22805. *If A is a subset of B , then the closure of A is contained in the closure of B* . Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/121238> (version: 2012-03-17). eprint: <https://math.stackexchange.com/q/121238>. URL: <https://math.stackexchange.com/q/121238> (cit. on p. 21).
- [20] Tej Bahadur Singh. *Introduction to Topology*. Springer, Singapore, 2019, p. 452. ISBN: 978-3-662-55406-7. DOI: <https://doi.org/10.1007/978-981-13-6954-4> (cit. on pp. 22, 23, 37, 49–53, 56, 58, 60, 62, 64, 66, 68, 69, 72, 73, 75, 76, 78, 87).
- [21] Martin Sleziak (<https://math.stackexchange.com/users/8297/martin-sleziak>). *Show that $\text{Int}(A) = X \setminus \overline{X \setminus A}$* . Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/872424> (version: 2014-07-20). eprint: <https://math.stackexchange.com/q/872424>. URL: <https://math.stackexchange.com/q/872424> (cit. on p. 22).
- [22] user2345215 (<https://math.stackexchange.com/users/131872/user2345215>). *Prove that the closure of complement, is the complement of the interior*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/970841> (version: 2014-10-12). eprint: <https://math.stackexchange.com/q/970841>. URL: <https://math.stackexchange.com/q/970841> (cit. on p. 22).
- [23] Dome (<https://math.stackexchange.com/users/70644/dome>). *Cauchy sequence is convergent iff it has a convergent subsequence*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/354965> (version: 2014-03-24). eprint: <https://math.stackexchange.com/q/354965>. URL: <https://math.stackexchange.com/q/354965> (cit. on p. 24).

- [24] Mark Bennet (<https://math.stackexchange.com/users/2906/mark-bennet>). *Show that a bounded, monotone increasing sequence is a Cauchy sequence*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/2169936> (version: 2017-03-03). eprint: <https://math.stackexchange.com/q/2169936>. URL: <https://math.stackexchange.com/q/2169936> (cit. on p. 24).
- [25] Wikipedia. accessed on 16.05.2025. URL: https://en.wikipedia.org/wiki/AM%E2%80%93GM_inequality#Proof_by_induction_#1 (cit. on p. 25).
- [26] Srivatsan (<https://math.stackexchange.com/users/13425/srivatsan>). *Proving $(1 + \frac{1}{n+1})^{n+1} \geq (1 + \frac{1}{n})^n$* . Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/64864> (version: 2011-12-10). eprint: <https://math.stackexchange.com/q/64864>. URL: <https://math.stackexchange.com/q/64864> (cit. on p. 25).
- [27] Thomas Andrews (<https://math.stackexchange.com/users/7933/thomas-andrews>). *Proof that $(1 + 1/n)^n$ is Cauchy*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/5065619> (version: 2025-05-15). eprint: <https://math.stackexchange.com/q/5065619>. URL: <https://math.stackexchange.com/q/5065619> (cit. on p. 26).
- [28] Kavi Rama Murthy (<https://math.stackexchange.com/users/142385/kavi-rama-murthy>). *Is there any metric on \mathbb{R} with which it is incomplete*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/3789234> (version: 2020-08-13). eprint: <https://math.stackexchange.com/q/3789234>. URL: <https://math.stackexchange.com/q/3789234> (cit. on p. 27).
- [29] Armin Iske. *Approximation Theory and Algorithms for Data Analysis*. 1st ed. Springer Cham, 2018, p. 358. URL: <https://link.springer.com/book/10.1007/978-3-030-05228-7> (cit. on pp. 32, 98).
- [30] Žan Grad (<https://math.stackexchange.com/users/497464/%c5%bdan-grad>). *Show that a subset A is dense in $X \Leftrightarrow$ for every nonempty open set U we have $A \cap U \neq \emptyset$* . Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/2497036> (version: 2017-10-30). eprint: <https://math.stackexchange.com/q/2497036>. URL: <https://math.stackexchange.com/q/2497036> (cit. on p. 32).
- [31] Lääne-Viru (<https://math.stackexchange.com/users/144375/1%c3%a4%c3%a4ne-viru>). *Proving that a set A is dense in M iff A^c has empty interior*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/937581> (version: 2017-04-13). eprint: <https://math.stackexchange.com/q/937581>. URL: <https://math.stackexchange.com/q/937581> (cit. on p. 32).
- [32] kobe (<https://math.stackexchange.com/users/190421/kobe>). *Intersection of two open dense sets is dense*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/1143218> (version: 2024-09-12). eprint: <https://math.stackexchange.com/q/1143218>. URL: <https://math.stackexchange.com/q/1143218> (cit. on p. 32).
- [33] Henno Brandsma (<https://math.stackexchange.com/users/4280/henno-brandsma>). *Why need open in the Baire Category Theorem*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/1506081> (version: 2015-10-31). eprint: <https://math.stackexchange.com/q/1506081>. URL: <https://math.stackexchange.com/q/1506081> (cit. on p. 32).

- [34] *Math 396. Completions.* online. accessed on 05/29/25. URL: <https://virtualmath1.stanford.edu/~conrad/diffgeomPage/handouts/completion.pdf> (cit. on p. 33).
- [35] Miles Wheeler. *4.4 Relative Compactness.* online. accessed on 06/11/2025. URL: <https://www.mileshwheeler.com/ma30252/sec-rel.html> (cit. on p. 41).
- [36] Joel H. Shapiro. *The Arzela-Ascoli Theorem.* online. accessed on 07/22/25. 1999. URL: <https://users.math.msu.edu/users/shapiro/pubvit/Downloads/ArzNotes/ArzNotes.pdf> (cit. on p. 43).
- [37] Rudy the Reindeer (<https://math.stackexchange.com/users/5798/rudy-the-reindeer>). *Your favourite application of the Baire Category Theorem.* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/165696> (version: 2017-04-13). eprint: <https://math.stackexchange.com/q/165696>. URL: <https://math.stackexchange.com/q/165696> (cit. on p. 44).
- [38] Miel Sharf (<https://math.stackexchange.com/users/193503/miel-sharf>). *The intersection of infinitely many dense open sets may not be dense.* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/1070418> (version: 2014-12-16). eprint: <https://math.stackexchange.com/q/1070418>. URL: <https://math.stackexchange.com/q/1070418> (cit. on p. 45).
- [39] Henno Brandsma (<https://math.stackexchange.com/users/4280/henno-brandsma>). *Baire Category Theorem without Completeness.* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/2369429> (version: 2017-07-23). eprint: <https://math.stackexchange.com/q/2369429>. URL: <https://math.stackexchange.com/q/2369429> (cit. on p. 45).
- [40] Ralf Holtkamp. *Mathematik I für Studierende der Geophysik/Ozeanographie, Meteorologie und Physik. Vorlesungsskript.* 2021. URL: <https://www.math.uni-hamburg.de/home/holtkamp/mfp1/mfp1rh.pdf> (cit. on p. 45).
- [41] George Law (<https://math.stackexchange.com/users/141584/george-law>). *A function which is continuous with respect to some metric but not continuous with respect to some other metric.* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/2087322> (version: 2017-01-07). eprint: <https://math.stackexchange.com/q/2087322>. URL: <https://math.stackexchange.com/q/2087322> (cit. on p. 45).
- [42] *Uniform continuity.* online. accessed on 05/23/25. URL: https://en.wikipedia.org/wiki/Uniform_continuity#Definition_for_functions_on_metric_spaces (cit. on p. 45).
- [43] nicomezi (<https://math.stackexchange.com/users/316579/nicomezi>). *What's the difference between uniformly equicontinuous and uniformly continuous?* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/1674834> (version: 2016-02-27). eprint: <https://math.stackexchange.com/q/1674834>. URL: <https://math.stackexchange.com/q/1674834> (cit. on p. 46).
- [44] Martin Argerami (<https://math.stackexchange.com/users/22857/martin-argerami>). *Why uniform continuity does not imply equicontinuity?* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/4841004> (version: 2024-01-08). eprint: <https://math.stackexchange.com/q/4841004>. URL: <https://math.stackexchange.com/q/4841004> (cit. on p. 46).

- [45] Jeff Diller. *Continuous Functions and Open Sets*. 2014. URL: https://www3.nd.edu/~jdiller/teaching/archive/fall14_20850/continuity.pdf (cit. on p. 47).
- [46] Ink (<https://math.stackexchange.com/users/34881/ink>). *The preimage of continuous function on a closed set is closed*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/437837> (version: 2013-07-07). eprint: <https://math.stackexchange.com/q/437837>. URL: <https://math.stackexchange.com/q/437837> (cit. on p. 47).
- [47] layman (<https://math.stackexchange.com/users/131740/layman>). *Continuous mapping on a compact metric space is uniformly continuous*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/1605098> (version: 2016-01-09). eprint: <https://math.stackexchange.com/q/1605098>. URL: <https://math.stackexchange.com/q/1605098> (cit. on p. 48).
- [48] user61527. *Proving $f(x) = x^2$ is not uniformly continuous on the real line*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/503101> (version: 2013-09-24). eprint: <https://math.stackexchange.com/q/503101>. URL: <https://math.stackexchange.com/q/503101> (cit. on p. 48).
- [49] Daniel Wohlrath (<https://math.stackexchange.com/users/842628/daniel-wohlrath>). *Show that a subset V of X is open iff every point $x \in V$ has a neighborhood $V_x \subseteq V$* . Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/3964428> (version: 2020-12-28). eprint: <https://math.stackexchange.com/q/3964428>. URL: <https://math.stackexchange.com/q/3964428> (cit. on p. 51).
- [50] WaterDrop (<https://math.stackexchange.com/users/641655/waterdrop>). *If X is first countable and S is a subspace of X , show that S is first countable*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/4548419> (version: 2022-10-08). eprint: <https://math.stackexchange.com/q/4548419>. URL: <https://math.stackexchange.com/q/4548419> (cit. on p. 52).
- [51] Ittay Weiss (<https://math.stackexchange.com/users/30953/ittay-weiss>). *Empty intersection and empty union*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/370201> (version: 2020-05-24). eprint: <https://math.stackexchange.com/q/370201>. URL: <https://math.stackexchange.com/q/370201> (cit. on p. 53).
- [52] Henno Brandsma (<https://math.stackexchange.com/users/4280/henno-brandsma>). *Uniqueness of Topology and Basis*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/713096> (version: 2014-03-15). eprint: <https://math.stackexchange.com/q/713096>. URL: <https://math.stackexchange.com/q/713096> (cit. on p. 56).
- [53] *Basis, Subbasis, Subspace*. online. last accessed on 06/22/2025. URL: https://www.nsm.buffalo.edu/~badzioch/MTH427/_static/mth427_notes_4.pdf (cit. on p. 57).
- [54] Henno Brandsma (<https://math.stackexchange.com/users/4280/henno-brandsma>). *Why is the lower limit topology not metrizable?* Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/790629> (version: 2014-05-11). eprint: <https://math.stackexchange.com/q/790629>. URL: <https://math.stackexchange.com/q/790629> (cit. on p. 58).

- [55] Lin (<https://math.stackexchange.com/users/249606/lin>). *May two or more bases of different sizes generate a same topology?* Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/1333309> (version: 2015-06-21). eprint: <https://math.stackexchange.com/q/1333309>. URL: <https://math.stackexchange.com/q/1333309> (cit. on p. 59).
- [56] Conrado Costa (<https://math.stackexchange.com/users/226425/conrado-costa>). *Strongly equivalent metrics are equivalent.* Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/1379634> (version: 2015-07-30). eprint: <https://math.stackexchange.com/q/1379634>. URL: <https://math.stackexchange.com/q/1379634> (cit. on p. 60).
- [57] Bruno Andrades (<https://math.stackexchange.com/users/578498/bruno-andrades>). *Topological Equivalence of Metrics Does Not Imply Strong Equivalence.* Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/5065146> (version: 2025-05-14). eprint: <https://math.stackexchange.com/q/5065146>. URL: <https://math.stackexchange.com/q/5065146> (cit. on p. 61).
- [58] *Math 396. Product topology.* online. last accessed on 06/25/25. URL: <https://virtualmath1.stanford.edu/~conrad/diffgeomPage/handouts/prodmetric.pdf> (cit. on pp. 66, 67).
- [59] k.stm (<https://math.stackexchange.com/users/42242/k-stm>). *How to prove the continuity of the metric function?* Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/287313> (version: 2025-06-25). eprint: <https://math.stackexchange.com/q/287313>. URL: <https://math.stackexchange.com/q/287313> (cit. on p. 67).
- [60] Ruairi (<https://math.stackexchange.com/users/581780/ruairi>). *How to prove the continuity of the metric function?* Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/4228334> (version: 2021-09-03). eprint: <https://math.stackexchange.com/q/4228334>. URL: <https://math.stackexchange.com/q/4228334> (cit. on p. 67).
- [61] Set (<https://math.stackexchange.com/users/26920/set>). *Show that the countable product of metric spaces is metrizable.* Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/361778> (version: 2018-01-16). eprint: <https://math.stackexchange.com/q/361778>. URL: <https://math.stackexchange.com/q/361778> (cit. on p. 71).
- [62] Henno Brandsma (<https://math.stackexchange.com/users/4280/henno-brandsma>). *Show that the countable product of metric spaces is metrizable.* Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/362319> (version: 2019-08-01). eprint: <https://math.stackexchange.com/q/362319>. URL: <https://math.stackexchange.com/q/362319> (cit. on p. 71).
- [63] Cm7F7Bb (<https://math.stackexchange.com/users/23249/cm7f7bb>). *How to interpret the sum of two series?* Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/2163556> (version: 2017-02-27). eprint: <https://math.stackexchange.com/q/2163556>. URL: <https://math.stackexchange.com/q/2163556> (cit. on p. 71).
- [64] user12345 (<https://math.stackexchange.com/users/1108514/user12345>). *Help Understanding the Heine-Borel Theorem Proof.* Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/4561719> (version: 2022-10-26). eprint: <https://math.stackexchange.com/q/4561719>. URL: <https://math.stackexchange.com/q/4561719> (cit. on p. 72).

- [65] 311411 (<https://math.stackexchange.com/users/688046/311411>). *Help Understanding the Heine-Borel Theorem Proof*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/4561736> (version: 2022-10-26). eprint: <https://math.stackexchange.com/q/4561736>. URL: <https://math.stackexchange.com/q/4561736> (cit. on p. 72).
- [66] user895986. *Stuck understanding proof of Heine-Borel Theorem*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/4203835> (version: 2025-07-14). eprint: <https://math.stackexchange.com/q/4203835>. URL: <https://math.stackexchange.com/q/4203835> (cit. on p. 72).
- [67] Daniel Smania (<https://math.stackexchange.com/users/1594236/daniel-smania>). *Finite product of compact topological spaces is compact - understanding notation in proof*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/5082194> (version: 2025-07-11). eprint: <https://math.stackexchange.com/q/5082194>. URL: <https://math.stackexchange.com/q/5082194> (cit. on p. 73).
- [68] Ivo Terek (<https://math.stackexchange.com/users/118056/ivo-terek>). *A subset of a compact set is compact?* Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/877547> (version: 2014-07-25). eprint: <https://math.stackexchange.com/q/877547>. URL: <https://math.stackexchange.com/q/877547> (cit. on p. 73).
- [69] Learner (<https://math.stackexchange.com/users/48763/learner>). *Convergence of sequences in topological spaces*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/289740> (version: 2022-06-30). eprint: <https://math.stackexchange.com/q/289740>. URL: <https://math.stackexchange.com/q/289740> (cit. on p. 74).
- [70] Brian M. Scott (<https://math.stackexchange.com/users/12042/brian-m-scott>). *Proving that a compact subset of a Hausdorff space is closed*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/1329787> (version: 2015-06-18). eprint: <https://math.stackexchange.com/q/1329787>. URL: <https://math.stackexchange.com/q/1329787> (cit. on p. 75).
- [71] *Topology: Show restriction of continuous function is continuous, and restriction of a homeomorphism is a homeomorphism*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/1826827> (version: 2016-06-15). eprint: <https://math.stackexchange.com/q/1826827>. URL: <https://math.stackexchange.com/q/1826827> (cit. on p. 76).
- [72] cdhanson (<https://math.stackexchange.com/users/24500/cdhanson>). *Continuous image of compact sets are compact*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/226328> (version: 2024-02-06). eprint: <https://math.stackexchange.com/q/226328>. URL: <https://math.stackexchange.com/q/226328> (cit. on p. 77).
- [73] Brian M. Scott (<https://math.stackexchange.com/users/12042/brian-m-scott>). *X compact metric space, $f : X \rightarrow \mathbb{R}$ continuous attains max/min*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/109552> (version: 2012-02-15). eprint: <https://math.stackexchange.com/q/109552>. URL: <https://math.stackexchange.com/q/109552> (cit. on p. 78).

- [74] Semih Kara (<https://math.stackexchange.com/users/658227/semih-kara>). *Identity function is a homeomorphism iff topologies are equal*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/3164142> (version: 2019-03-27). eprint: <https://math.stackexchange.com/q/3164142>. URL: <https://math.stackexchange.com/q/3164142> (cit. on p. 79).
- [75] Haha (<https://math.stackexchange.com/users/94689/haha>). *is bounded linear operator necessarily continuous?* Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/556667> (version: 2024-03-16). eprint: <https://math.stackexchange.com/q/556667>. URL: <https://math.stackexchange.com/q/556667> (cit. on p. 81).
- [76] *February 18, 2021*. online. accessed on 05/23/25. URL: https://ocw.mit.edu/courses/18-102-introduction-to-functional-analysis-spring-2021/22bf2f774fb161776032491568148707/MIT18_102s21_lec2.pdf (cit. on p. 81).
- [77] Stanford University. *Binary Relations*. 2007. URL: <https://web.stanford.edu/class/archive/cs/cs103/cs103.1142/lectures/07/Small07.pdf> (cit. on p. 83).
- [78] Wikipedia contributors. *Equivalence relation*. 2021. URL: https://en.wikipedia.org/wiki/Equivalence_relation (cit. on p. 83).
- [79] *Equivalence class*. online. accessed on 05/28/25. URL: https://en.wikipedia.org/wiki/Equivalence_class (cit. on p. 83).
- [80] Chee Han Tan. *Zorn's Lemma*. URL: https://www.math.utah.edu/~tan/6710_FA/Hahn%20Banach%20Theorem.pdf (cit. on p. 83).
- [81] John K. Hunter. *The supremum and infimum*. online. accessed on 07/09/25. URL: <https://www.math.ucdavis.edu/~hunter/m125b/ch2.pdf> (cit. on pp. 84, 97).
- [82] Brian M. Scott (<https://math.stackexchange.com/users/12042/brian-m-scott>). *Am I allowed to use distributive law for infinitely many sets?* Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/481036> (version: 2013-09-01). eprint: <https://math.stackexchange.com/q/481036>. URL: <https://math.stackexchange.com/q/481036> (cit. on p. 84).
- [83] user165987 (<https://math.stackexchange.com/users/165987/user165987>). *Proving the properties of big union of unions for indexed sets*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/878108> (version: 2017-03-14). eprint: <https://math.stackexchange.com/q/878108>. URL: <https://math.stackexchange.com/q/878108> (cit. on p. 85).
- [84] ASB (<https://math.stackexchange.com/users/111607/asb>). *Image of the union and intersection of sets*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/1132766> (version: 2015-02-04). eprint: <https://math.stackexchange.com/q/1132766>. URL: <https://math.stackexchange.com/q/1132766> (cit. on p. 86).
- [85] Manos (<https://math.stackexchange.com/users/11921/manos>). *Union of preimages and preimage of union*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/141357> (version: 2016-11-24). eprint: <https://math.stackexchange.com/q/141357>. URL: <https://math.stackexchange.com/q/141357> (cit. on p. 87).

- [86] user228113. *Function is continuous if and only if preimages of elements of a sub-base are open*. Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/2190480> (version: 2017-03-17). eprint: <https://math.stackexchange.com/q/2190480>. URL: <https://math.stackexchange.com/q/2190480> (cit. on p. 87).
- [87] David Lindemann. *Differential geometry. Lecture 1: Smooth manifolds*. 2020. URL: https://www.math.uni-hamburg.de/home/lindemann/material/DG2020L1_slides.pdf (cit. on p. 90).
- [88] Ralf Holtkamp. *Mathematik III für Studierende der Physik Vorlesungsskript*. 2021. URL: https://www.math.uni-hamburg.de/home/holtkamp/mfp3/mfp3rh_druckskript.pdf (cit. on p. 91).
- [89] Nolan Wallach. *MATH 23b, SPRING 2005. THEORETICAL LINEAR ALGEBRA AND MULTIVARIABLE CALCULUS*. 2005. URL: <http://www.math.ucsd.edu/~nwallach/inverse%5B1%5D.pdf> (cit. on p. 91).
- [90] *5.1 Roots and Radicals*. URL: https://saylordotorg.github.io/text_intermediate-algebra/s08-01-roots-and-radicals.html (cit. on p. 91).
- [91] Wikipedia contributors. *Jacobian matrix and determinant*. 2021. URL: https://en.wikipedia.org/wiki/Jacobian_matrix_and_determinant (cit. on p. 92).
- [92] Khan Academy. *Upper triangular determinant*. URL: <https://www.khanacademy.org/math/linear-algebra/matrix-transformations/determinant-depth/v/linear-algebra-upper-triangular-determinant> (cit. on p. 92).
- [93] Wikipedia contributors. *Determinant*. 2021. URL: <https://en.wikipedia.org/wiki/Determinant> (cit. on p. 92).
- [94] Zilin J. (<https://math.stackexchange.com/users/94898/zilin-j>). *Determinant of a block lower triangular matrix*. Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/1221066> (version: 2015-08-04). eprint: <https://math.stackexchange.com/q/1221066>. URL: <https://math.stackexchange.com/q/1221066> (cit. on p. 93).
- [95] *ECE 5630 Lecture Notes*. https://people.ece.cornell.edu/zivg/ECE_5630_Lectures10.pdf. Accessed on 11/26/23 (cit. on p. 95).
- [96] *ECE 5630 Lecture Notes 6*. https://people.ece.cornell.edu/zivg/ECE_5630_Lectures6.pdf. Accessed on 11/26/23 (cit. on p. 95).
- [97] Ennar (<https://math.stackexchange.com/users/122131/ennar>). *Prove that $\max\{a + b, c + d\} \leq \max\{a, c\} + \max\{b, d\}$* . Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/1975962> (version: 2016-10-19). eprint: <https://math.stackexchange.com/q/1975962>. URL: <https://math.stackexchange.com/q/1975962> (cit. on p. 96).
- [98] flyve (<https://math.stackexchange.com/users/43639/flyve>). *Prove $\sup(f+g) \leq \sup f + \sup g$* . Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/207335> (version: 2019-12-26). eprint: <https://math.stackexchange.com/q/207335>. URL: <https://math.stackexchange.com/q/207335> (cit. on p. 96).

- [99] Hagen von Eitzen (<https://math.stackexchange.com/users/39174/hagen-von-eitzen>). *Prove $\sup(f + g) \leq \sup f + \sup g$* . Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/207339> (version: 2012-10-04). eprint: <https://math.stackexchange.com/q/207339>. URL: <https://math.stackexchange.com/q/207339> (cit. on p. 96).
- [100] Tianyi Hu. “Bernstein Polynomials and their Application in the Finite Element Method”. Bachelor’s Thesis (cit. on p. 100).
- [101] *The Fubini Principle in Discrete Math*. online. accessed on 05/22/25. URL: <https://web.math.ucsb.edu/~cmart07/fubini.pdf> (cit. on p. 100).
- [102] Andrew Sutherland. *18.095 Lecture Series in Mathematics*. online. accessed on 05/29/25. 2015. URL: <https://math.mit.edu/classes/18.095/2015IAP/lecture1/padic.pdf> (cit. on p. 105).
- [103] Todd Kemp. *Cauchy’s Construction of \mathbb{R}* . accessed on 06/02/2025. 2025. URL: <https://mathweb.ucsd.edu/~tkemp/140A/Construction.of.R.pdf> (cit. on p. 105).
- [104] joeA (<https://math.stackexchange.com/users/79432/joea>). *Is there a rational number between any two irrationals?* Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/421600> (version: 2014-01-29). eprint: <https://math.stackexchange.com/q/421600>. URL: <https://math.stackexchange.com/q/421600> (cit. on p. 109).
- [105] Max Herrmann (<https://math.stackexchange.com/users/172142/max-herrmann>). *Terence Tao, Analysis I, Ex. 5.4.5: There is a rational between any two reals*. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/3434503> (version: 2019-11-13). eprint: <https://math.stackexchange.com/q/3434503>. URL: <https://math.stackexchange.com/q/3434503> (cit. on p. 109).