CS2302 - Data Structures

Spring 2018

Lab # 8

Algorithm Design Techniques

Deadline: Thursday, May 3, 11:59 p.m.

1. The partition problem consists of determining if there is a way to partition a set of integers S into two subsets S_1 and S_2 such that $\sum S_1 = \sum S_2$. Recall that S_1 and S_2 are a partition of S if and only if $S_1 \cup S_2 = S$ and $S_1 \cap S_2 = \{\}$. Write a method that solves the partition problem using backtracking.

2. Write a randomized algorithm to check probabilistically whether two trigonometric expressions are identities. Your expression can be written as java methods. For example, to check whether $tan(\theta) = sin(\theta)/cos(\theta)$ you can define methods:

```
public static double f(double theta){
   return Math.sin(theta)/Math.cos(theta);
}

public static double g(double theta){
   return Math.tan(theta);
}
```

   And then implement a method that tests the equality using a large number of random values in the $-\pi$ to π range. Try your method on as many of the identities in the following image as you can. Try it also on expressions that are not identities to make sure those are correctly detected.

Formulas and Identities

Tangent and Cotangent Identities

$$\tan\theta = \frac{\sin\theta}{\cos\theta} \qquad \cot\theta = \frac{\cos\theta}{\sin\theta}$$

Reciprocal Identities

$$\csc\theta = \frac{1}{\sin\theta} \qquad \sin\theta = \frac{1}{\csc\theta}$$

$$\sec\theta = \frac{1}{\cos\theta} \qquad \cos\theta = \frac{1}{\sec\theta}$$

$$\cot\theta = \frac{1}{\tan\theta} \qquad \tan\theta = \frac{1}{\cot\theta}$$

Pythagorean Identities

$$\sin^2\theta + \cos^2\theta = 1$$

$$\tan^2\theta + 1 = \sec^2\theta$$

$$1 + \cot^2\theta = \csc^2\theta$$

Even/Odd Formulas

$$\sin(-\theta) = -\sin\theta \qquad \csc(-\theta) = -\csc\theta$$

$$\cos(-\theta) = \cos\theta \qquad \sec(-\theta) = \sec\theta$$

$$\tan(-\theta) = -\tan\theta \qquad \cot(-\theta) = -\cot\theta$$

Periodic Formulas

If n is an integer.

$$\sin(\theta + 2\pi n) = \sin\theta \qquad \csc(\theta + 2\pi n) = \csc\theta$$

$$\cos(\theta + 2\pi n) = \cos\theta \qquad \sec(\theta + 2\pi n) = \sec\theta$$

$$\tan(\theta + \pi n) = \tan\theta \qquad \cot(\theta + \pi n) = \cot\theta$$

Double Angle Formulas

$$\sin(2\theta) = 2\sin\theta\cos\theta$$

$$\cos(2\theta) = \cos^2\theta - \sin^2\theta$$

$$= 2\cos^2\theta - 1$$

$$= 1 - 2\sin^2\theta$$

$$\tan(2\theta) = \frac{2\tan\theta}{1 - \tan^2\theta}$$

Degrees to Radians Formulas

If x is an angle in degrees and t is an angle in radians then

$$\frac{\pi}{180} = \frac{t}{x} \quad \Rightarrow \quad t = \frac{\pi x}{180} \quad \text{and} \quad x = \frac{180t}{\pi}$$

Half Angle Formulas (alternate form)

$$\sin\frac{\theta}{2} = \pm\sqrt{\frac{1-\cos\theta}{2}} \qquad \sin^2\theta = \frac{1}{2}(1-\cos(2\theta))$$

$$\cos\frac{\theta}{2} = \pm\sqrt{\frac{1+\cos\theta}{2}} \qquad \cos^2\theta = \frac{1}{2}(1+\cos(2\theta))$$

$$\tan\frac{\theta}{2} = \pm\sqrt{\frac{1-\cos\theta}{1+\cos\theta}} \qquad \tan^2\theta = \frac{1-\cos(2\theta)}{1+\cos(2\theta)}$$

Sum and Difference Formulas

$$\sin(\alpha \pm \beta) = \sin\alpha\cos\beta \pm \cos\alpha\sin\beta$$

$$\cos(\alpha \pm \beta) = \cos\alpha\cos\beta \mp \sin\alpha\sin\beta$$

$$\tan(\alpha \pm \beta) = \frac{\tan\alpha \pm \tan\beta}{1 \mp \tan\alpha\tan\beta}$$

Product to Sum Formulas

$$\sin\alpha\sin\beta = \frac{1}{2}\left[\cos(\alpha-\beta) - \cos(\alpha+\beta)\right]$$

$$\cos\alpha\cos\beta = \frac{1}{2}\left[\cos(\alpha-\beta) + \cos(\alpha+\beta)\right]$$

$$\sin\alpha\cos\beta = \frac{1}{2}\left[\sin(\alpha+\beta) + \sin(\alpha-\beta)\right]$$

$$\cos\alpha\sin\beta = \frac{1}{2}\left[\sin(\alpha+\beta) - \sin(\alpha-\beta)\right]$$

Sum to Product Formulas

$$\sin\alpha + \sin\beta = 2\sin\left(\frac{\alpha+\beta}{2}\right)\cos\left(\frac{\alpha-\beta}{2}\right)$$

$$\sin\alpha - \sin\beta = 2\cos\left(\frac{\alpha+\beta}{2}\right)\sin\left(\frac{\alpha-\beta}{2}\right)$$

$$\cos\alpha + \cos\beta = 2\cos\left(\frac{\alpha+\beta}{2}\right)\cos\left(\frac{\alpha-\beta}{2}\right)$$

$$\cos\alpha - \cos\beta = -2\sin\left(\frac{\alpha+\beta}{2}\right)\sin\left(\frac{\alpha-\beta}{2}\right)$$

Cofunction Formulas

$$\sin\left(\frac{\pi}{2}-\theta\right) = \cos\theta \qquad \cos\left(\frac{\pi}{2}-\theta\right) = \sin\theta$$

$$\csc\left(\frac{\pi}{2}-\theta\right) = \sec\theta \qquad \sec\left(\frac{\pi}{2}-\theta\right) = \csc\theta$$

$$\tan\left(\frac{\pi}{2}-\theta\right) = \cot\theta \qquad \cot\left(\frac{\pi}{2}-\theta\right) = \tan\theta$$

As usual, write a report describing your work.