Given an array of integers, calculate the ratios of its elements that are positive, negative, and zero. Print the decimal value of each fraction on a new line with 6 places after the decimal.

Note: This challenge introduces precision problems. The test cases are scaled to six decimal places, though answers with absolute error of up to 10^{-4} are acceptable.

Example

$$arr = [1, 1, 0, -1, -1]$$

There are n=5 elements, two positive, two negative and one zero. Their ratios are $\frac{2}{5}=0.400000$, $\frac{2}{5}=0.400000$ and $\frac{1}{5}=0.200000$. Results are printed as:

- 0.400000
- 0.400000
- 0.200000

Function Description

Complete the plusMinus function in the editor below.

plusMinus has the following parameter(s):

• int arr[n]: an array of integers

Print

Print the ratios of positive, negative and zero values in the array. Each value should be printed on a separate line with 6 digits after the decimal. The function should not return a value.

Input Format

The first line contains an integer, n, the size of the array.

The second line contains n space-separated integers that describe arr[n].

Constraints

$$0 < n \le 100 \ -100 \le arr[i] \le 100$$

Output Format

Print the following 3 lines, each to 6 decimals:

- 1. proportion of positive values
- 2. proportion of negative values
- 3. proportion of zeros

Sample Input

Sample Output

0.500000

0.333333

0.166667

Explanation

There are 3 positive numbers, 2 negative numbers, and 1 zero in the array.

The proportions of occurrence are positive: $\frac{3}{6}=0.500000$, negative: $\frac{2}{6}=0.333333$ and zeros: $\frac{1}{6}=0.166667$.