

# Lower Bounds for the Polynomial Calculus via the “Pigeon Dance”

Imogen Hergeth

January 2023

# Overview

We will present the result from A.A. Razborov, “Lower Bounds for the Polynomial Calculus”, in: Computational Complexity 7.4 (Dec. 2, 1998).

1 Introduction

2 The Pigeonhole Principle

3 The Pigeon Dance

4 Conclusion

# Background

- Lower bounds for proofs in various systems

# Background

- Lower bounds for proofs in various systems
- In particular for the pigeonhole principle

# Background

- Lower bounds for proofs in various systems
- In particular for the pigeonhole principle
- Polynomial calculus is a strong proof system

# Background

- Lower bounds for proofs in various systems
- In particular for the pigeonhole principle
- Polynomial calculus is a strong proof system
- Provide a lower bound for it with the pigeonhole principle

# Definition

- Similar to sequent calculus, but lines are polynomials

# Definition

- Similar to sequent calculus, but lines are polynomials
- We use multilinear polynomials  $S_n(\mathbb{K})$   
 $(xy + xz + v \equiv x^2y + x^3z^5 + v)$



# Definition

- Similar to sequent calculus, but lines are polynomials
- We use multilinear polynomials  $S_n(\mathbb{K})$   
 $(xy + xz + v \equiv x^2y + x^3z^5 + v)$
- Addition

$$\frac{f \quad g}{af + bg}$$

# Definition

- Similar to sequent calculus, but lines are polynomials
- We use multilinear polynomials  $S_n(\mathbb{K})$   
 $(xy + xz + v \equiv x^2y + x^3z^5 + v)$
- Addition

$$\frac{f \quad g}{af + bg}$$

- Multiplication

$$\frac{f}{f \cdot x}$$

# Refutations

- $g$  is provable from  $f_1, \dots, f_n$  if and only if it is in the ideal generated by them

# Refutations

- $g$  is provable from  $f_1, \dots, f_n$  if and only if it is in the ideal generated by them
- A proof of 1 exists if and only if  $f_1, \dots, f_n$  have no common zeroes

# Refutations

- $g$  is provable from  $f_1, \dots, f_n$  if and only if it is in the ideal generated by them
- A proof of 1 exists if and only if  $f_1, \dots, f_n$  have no common zeroes
- We construct polynomials such that their zeroes correspond to satisfying assignments

# Refutations

- $g$  is provable from  $f_1, \dots, f_n$  if and only if it is in the ideal generated by them
- A proof of 1 exists if and only if  $f_1, \dots, f_n$  have no common zeroes
- We construct polynomials such that their zeroes correspond to satisfying assignments
- Proving 1 from them is a *refutation*

# Example proof

- Try to prove  $xy + z$  from  $x + 1$  and  $z$

# Example proof

- Try to prove  $xy + z$  from  $x + 1$  and  $z$

$$x + 1$$



# Example proof

- Try to prove  $xy + z$  from  $x + 1$  and  $z$

$$(\cdot y) \frac{x + 1}{xy + y}$$

# Example proof

- Try to prove  $xy + z$  from  $x + 1$  and  $z$

$$(\cdot y) \frac{x+1}{xy+y} \quad z \quad (+) \\ \hline xy+y+z$$

# Example proof

- Try to prove  $xy + z$  from  $x + 1$  and  $z$

$$(\cdot y) \frac{x+1}{xy+y} \quad z \quad (+) \\ \hline xy+y+z$$

- We now want to subtract  $y$

# Example proof

- Try to prove  $xy + z$  from  $x + 1$  and  $z$

$$(\cdot y) \frac{x+1}{xy+y} \quad z \quad (+) \\ \hline xy + y + z$$

- We now want to subtract  $y$
- There is no way to prove  $y$  from  $x + 1$  and  $z$

# Example proof

- Try to prove  $xy + z$  from  $x + 1$  and  $z$

$$(\cdot y) \frac{x+1}{xy+y} \quad z \quad (+) \\ \hline xy + y + z$$

- We now want to subtract  $y$
- There is no way to prove  $y$  from  $x + 1$  and  $z$
- Closest to  $xy + z$  we can prove is  $xy + y + z$

# Algebraic view of proofs

- $V$  are polynomials we can prove
- $\Delta$  are leading terms of ones we cannot prove
- $S_n(\mathbb{K}) \cong \mathbb{K}\Delta \oplus V$
- Similarly  $V_I, \Delta_I$  using subset of variables  $I$

$$xy + y + z$$

$$y$$

$$xy + z = -y + xy + y + z$$

# The pigeonhole principle

- If there are  $m$  pigeons,  $n$  pigeon holes, and  $m > n$  then at least two pigeons have to share a hole

# The pigeonhole principle

- If there are  $m$  pigeons,  $n$  pigeon holes, and  $m > n$  then at least two pigeons have to share a hole
- Formally: if  $m > n$  there is no injection  $[m] \hookrightarrow [n]$



# The pigeonhole principle

- If there are  $m$  pigeons,  $n$  pigeon holes, and  $m > n$  then at least two pigeons have to share a hole
- Formally: if  $m > n$  there is no injection  $[m] \hookrightarrow [n]$
- Variables:  $x_{ij}, i \in [m], n \in [n]$

# The pigeonhole principle

- If there are  $m$  pigeons,  $n$  pigeon holes, and  $m > n$  then at least two pigeons have to share a hole
- Formally: if  $m > n$  there is no injection  $[m] \hookrightarrow [n]$
- Variables:  $x_{ij}, i \in [m], n \in [n]$
- Assignment of  $x_{3,5}$  corresponds to pigeon 3 being in hole 5

# The pigeonhole principle

- If there are  $m$  pigeons,  $n$  pigeon holes, and  $m > n$  then at least two pigeons have to share a hole
- Formally: if  $m > n$  there is no injection  $[m] \hookrightarrow [n]$
- Variables:  $x_{ij}, i \in [m], n \in [n]$
- Assignment of  $x_{3,5}$  corresponds to pigeon 3 being in hole 5

## Definition ( $\neg\mathcal{PHP}_n^m$ )

$$Q_i := 1 - \sum_{j \in [n]} x_{ij} \quad \text{for each } i \in [m]$$

$$Q_{i_1, i_2, j} := x_{i_1 j} x_{i_2 j} \quad \text{for each } i_1 \neq i_2 \in [m], j \in [n]$$

# Main result

## Theorem

*Every polynomial calculus refutation of  $\neg\mathcal{PHP}_n^m$  must have degree at least  $n/2 + 1$ .*

# Main result

## Theorem

*Every polynomial calculus refutation of  $\neg\mathcal{PHP}_n^m$  must have degree at least  $n/2 + 1$ .*

- $\neg\mathcal{PHP}_n^m$  are constraints on pigeon assignments

# Main result

## Theorem

*Every polynomial calculus refutation of  $\neg\mathcal{PHP}_n^m$  must have degree at least  $n/2 + 1$ .*

- $\neg\mathcal{PHP}_n^m$  are constraints on pigeon assignments
- Proofs combine them into more complex ones

# Main result

## Theorem

*Every polynomial calculus refutation of  $\neg\mathcal{PHP}_n^m$  must have degree at least  $n/2 + 1$ .*

- $\neg\mathcal{PHP}_n^m$  are constraints on pigeon assignments
- Proofs combine them into more complex ones
- Can only derive local constraints with small degrees

# Main result

## Theorem

*Every polynomial calculus refutation of  $\neg\mathcal{PHP}_n^m$  must have degree at least  $n/2 + 1$ .*

- $\neg\mathcal{PHP}_n^m$  are constraints on pigeon assignments
- Proofs combine them into more complex ones
- Can only derive local constraints with small degrees
- Pigeons can fly away to escape local contradictions



# $\neg \mathcal{PHP}_n^m$ constraints

- Pigeons cannot share holes

$\neg \mathcal{PHP}_n^m$  constraints

- Pigeons cannot share holes
- Possible for pigeons  $I \subseteq [m]$  with  $|I| \leq n$

$\neg \mathcal{PHP}_n^m$  constraints

- Pigeons cannot share holes
- Possible for pigeons  $I \subseteq [m]$  with  $|I| \leq n$
- Locally valid assignments are injections  $I \hookrightarrow [n]$

$\neg \mathcal{PHP}_n^m$  constraints

- Pigeons cannot share holes
- Possible for pigeons  $I \subseteq [m]$  with  $|I| \leq n$
- Locally valid assignments are injections  $I \hookrightarrow [n]$
- Corresponding variable assignments are  $M_I$

# Locally valid assignments

- Characterize  $V_I$  as the set of all polynomials with  $a(f) = 0$  for all  $a \in M_I$

# Locally valid assignments

- Characterize  $V_I$  as the set of all polynomials with  $a(f) = 0$  for all  $a \in M_I$
- We're done! Clearly not all polynomials are identically zero on  $M_I$

# Locally valid assignments

- Characterize  $V_I$  as the set of all polynomials with  $a(f) = 0$  for all  $a \in M_I$
- We're done! Clearly not all polynomials are identically zero on  $M_I$
- This definition completely ignores the degrees of the proofs!

# Locally valid assignments

- Characterize  $V_I$  as the set of all polynomials with  $a(f) = 0$  for all  $a \in M_I$
- We're done! Clearly not all polynomials are identically zero on  $M_I$
- This definition completely ignores the degrees of the proofs!
- Only works if whether  $t \in \Delta_I$  is independent of  $I \supseteq \text{dom}(t)$



# Overview

- Goal:

# Overview

- Goal:
  - Use pigeon dance to characterize  $\Delta_I$

# Overview

- Goal:
  - Use pigeon dance to characterize  $\Delta_I$
  - Prove this  $\Delta_I$  is exactly the leading terms of underivable polynomials

# Overview

- Goal:
  - Use pigeon dance to characterize  $\Delta_I$
  - Prove this  $\Delta_I$  is exactly the leading terms of underivable polynomials
- Start with pigeons sitting in their holes

# Overview

- Goal:
  - Use pigeon dance to characterize  $\Delta_I$
  - Prove this  $\Delta_I$  is exactly the leading terms of underivable polynomials
- Start with pigeons sitting in their holes
- The first pigeon flies to an unoccupied hole to its right




# Overview

- Goal:
  - Use pigeon dance to characterize  $\Delta_I$
  - Prove this  $\Delta_I$  is exactly the leading terms of underivable polynomials
- Start with pigeons sitting in their holes
- The first pigeon flies to an unoccupied hole to its right
- Repeat until all pigeons have moved once

# Overview


- Goal:
  - Use pigeon dance to characterize  $\Delta_I$
  - Prove this  $\Delta_I$  is exactly the leading terms of underivable polynomials
- Start with pigeons sitting in their holes
- The first pigeon flies to an unoccupied hole to its right
- Repeat until all pigeons have moved once
- If a pigeon cannot find an empty hole, the dance is aborted

# Example









# Example



# Example




# Example




# Example



# Example



# Example



# Formalization

- Consider partial injections  $I \hookrightarrow [m]$        $(1 \mapsto 4, 2 \mapsto 1, 3 \mapsto 2)$

# Formalization

- Consider partial injections  $I \hookrightarrow [m]$   $(1 \mapsto 4, 2 \mapsto 1, 3 \mapsto 2)$
- Encode pigeon positions as terms  $(x_{1,4} x_{2,1} x_{3,2})$



# Formalization

- Consider partial injections  $I \hookrightarrow [m]$        $(1 \mapsto 4, 2 \mapsto 1, 3 \mapsto 2)$
- Encode pigeon positions as terms       $(x_{1,4} x_{2,1} x_{3,2})$
- $\Delta_I$  is the set of terms that let pigeons complete the dance

# Formalization

- Consider partial injections  $I \hookrightarrow [m]$        $(1 \mapsto 4, 2 \mapsto 1, 3 \mapsto 2)$
- Encode pigeon positions as terms       $(x_{1,4} x_{2,1} x_{3,2})$
- $\Delta_I$  is the set of terms that let pigeons complete the dance
- Whether  $t \in \Delta_I$  is independent of  $I$  since pigeons not in the dance do not affect it

# The Kill operator

- We need to further understand the dance to prove it correct

# The Kill operator

- We need to further understand the dance to prove it correct
- Idea: a way to block specific pigeon holes

# The Kill operator

- We need to further understand the dance to prove it correct
- Idea: a way to block specific pigeon holes
- Kill the first pigeon and moves its hole to the left

# The Kill operator

- We need to further understand the dance to prove it correct
- Idea: a way to block specific pigeon holes
- Kill the first pigeon and moves its hole to the left
- $\text{Kill}(x_{i_1 j_1} \cdots x_{i_d, j_d}) = x_{i_2 j'_2} \cdots x_{i_d j'_d}$  with

$$j'_k := \begin{cases} j_k + 1, & \text{if } j_k < j_1 \\ j_k, & \text{if } j_k > j_1. \end{cases}$$

# Simulating the pigeon dance with Kill

# Simulating the pigeon dance with Kill

## Theorem

$x_{i_1 j_1} \cdots x_{i_d j_d} \in \Delta_I$  if and only if there is a  $j' > j_1$  such that  $\text{Kill}(x_{i_1 j'} \cdots x_{i_d j_d}) \in \Delta_I$ .



# Simulating the pigeon dance with Kill

## Theorem

$x_{i_1 j_1} \cdots x_{i_d j_d} \in \Delta_I$  if and only if there is a  $j' > j_1$  such that  $\text{Kill}(x_{i_1 j'} \cdots x_{i_d j_d}) \in \Delta_I$ .

## Proof sketch

$\text{Kill}(x_{i_1 j'} \cdots x_{i_d j_d})$  effectively moves the first pigeon to an empty hole to its right and then kills it. This is the same as each step in the dance, where the first pigeon flies to some free hole to its right and then occupies it. □

Closure of  $\Delta_I$ 

## Theorem

$\Delta_I$  is closed under Kill.

# Closure of $\Delta_I$

## Theorem

$\Delta_I$  is closed under Kill.

## Proof sketch

If  $t \in \Delta_I$  then the pigeons can complete their dance. During this the first pigeon will start at  $j$  and fly to  $j'$ . Killing the pigeon frees up  $j'$  so any other pigeon that wanted to use  $j$  can use it instead. □

# The lower bound

## Theorem

*If  $|I| \leq (n + 1)/2$  and pigeons can complete the dance, then we can introduce a new pigeon such that  $\text{Kill}(x_{ij}t) \in \Delta_I$ .*

# The lower bound

## Theorem

*If  $|I| \leq (n + 1)/2$  and pigeons can complete the dance, then we can introduce a new pigeon such that  $\text{Kill}(x_{ij}t) \in \Delta_I$ .*

## Proof sketch

At most

$$|\text{dom}(t)| \leq |I \setminus \{i\}| \leq \frac{n-1}{2}$$

pigeons involved in the dance, each occupying two holes. Place pigeon at unused hole and kill it there. The remaining pigeons can complete the dance since the moved hole was not used. □

## Putting things together

- Goal: show pigeon dance correctly characterizes underivable polynomials

## Putting things together

- Goal: show pigeon dance correctly characterizes underivable polynomials
- Formally: each  $f \in \mathbb{K}\Delta_I$  has an  $a \in M_I$  with  $a(f) \neq 0$

## Putting things together

- Goal: show pigeon dance correctly characterizes underivable polynomials
- Formally: each  $f \in \mathbb{K}\Delta_I$  has an  $a \in M_I$  with  $a(f) \neq 0$
- Induction over  $|I|$



## Putting things together

- Goal: show pigeon dance correctly characterizes underivable polynomials
- Formally: each  $f \in \mathbb{K}\Delta_I$  has an  $a \in M_I$  with  $a(f) \neq 0$
- Induction over  $|I|$
- Remove variables  $x_{ij}$  for minimal  $i \in I$  from  $f$

## Putting things together

- Goal: show pigeon dance correctly characterizes underivable polynomials
- Formally: each  $f \in \mathbb{K}\Delta_I$  has an  $a \in M_I$  with  $a(f) \neq 0$
- Induction over  $|I|$
- Remove variables  $x_{ij}$  for minimal  $i \in I$  from  $f$
- Inductive assumption gives us  $a' \in M_{I \setminus \{i\}}$  with  $a'(f') \neq 0$

closure of  $\Delta_I$

## Putting things together

- Goal: show pigeon dance correctly characterizes underivable polynomials
- Formally: each  $f \in \mathbb{K}\Delta_I$  has an  $a \in M_I$  with  $a(f) \neq 0$
- Induction over  $|I|$
- Remove variables  $x_{ij}$  for minimal  $i \in I$  from  $f$
- Inductive assumption gives us  $a' \in M_{I \setminus \{i\}}$  with  $a'(f') \neq 0$
- Pick a  $j$  such that  $\text{Kill}(x_{ij}t) \in \Delta_I$

closure of  $\Delta_I$

lower bound

## Putting things together

- Goal: show pigeon dance correctly characterizes underivable polynomials
- Formally: each  $f \in \mathbb{K}\Delta_I$  has an  $a \in M_I$  with  $a(f) \neq 0$
- Induction over  $|I|$
- Remove variables  $x_{ij}$  for minimal  $i \in I$  from  $f$
- Inductive assumption gives us  $a' \in M_{I \setminus \{i\}}$  with  $a'(f') \neq 0$
- Pick a  $j$  such that  $\text{Kill}(x_{ij}t) \in \Delta_I$
- Extend assignment to  $I$  with  $a(f) \neq 0$

closure of  $\Delta_I$

lower bound

# Summary

- Proven a lower bound  $d \geq n/2 + 1$  for polynomial calculus proofs of  $\neg \mathcal{PHP}_n^m$

# Summary

- Proven a lower bound  $d \geq n/2 + 1$  for polynomial calculus proofs of  $\neg \mathcal{PHP}_n^m$
- Characterize derivable polynomials  $V_I$  through locally consistent assignments  $M_I$

# Summary

- Proven a lower bound  $d \geq n/2 + 1$  for polynomial calculus proofs of  $\neg \mathcal{PHP}_n^m$
- Characterize derivable polynomials  $V_I$  through locally consistent assignments  $M_I$
- This only works if membership in  $\Delta_I$  is independent of  $I$

# Summary

- Proven a lower bound  $d \geq n/2 + 1$  for polynomial calculus proofs of  $\neg \mathcal{PHP}_n^m$
- Characterize derivable polynomials  $V_I$  through locally consistent assignments  $M_I$
- This only works if membership in  $\Delta_I$  is independent of  $I$
- Characterize  $\Delta_I$  through pigeon dance



# Summary

- Proven a lower bound  $d \geq n/2 + 1$  for polynomial calculus proofs of  $\neg \mathcal{PHP}_n^m$
- Characterize derivable polynomials  $V_I$  through locally consistent assignments  $M_I$
- This only works if membership in  $\Delta_I$  is independent of  $I$
- Characterize  $\Delta_I$  through pigeon dance
- This works since  $\Delta_I$  is linearly independent over  $M_I$