

Generating Function And Its Applications

518030910422 Ruihang Lai
Probability Theory, 2020 Spring
ACM Class, Shanghai Jiao Tong University

What is Generating Function?

What is Generating Function?

Assume you have a sequence $a = (a_n)_{n=0}^{\infty}$.

What is Generating Function?

Assume you have a sequence $a = (a_n)_{n=0}^{\infty}$.

The (ordinary) generating function $G_a(s)$ of sequence a is

$$G_a(s) = \sum_{n=0}^{\infty} a_n s^n.$$

What is Generating Function?

Assume you have a sequence $a = (a_n)_{n=0}^{\infty}$.

The (ordinary) generating function $G_a(s)$ of sequence a is

$$G_a(s) = \sum_{n=0}^{\infty} a_n s^n.$$

Sequence a : contains information

Generating function $G_a(s)$: encodes, reflects, stores the information

What is Generating Function?

Assume you have a sequence $a = (a_n)_{n=0}^{\infty}$.

The (ordinary) generating function $G_a(s)$ of sequence a is

$$G_a(s) = \sum_{n=0}^{\infty} a_n s^n.$$

Sequence a : contains information

Generating function $G_a(s)$: encodes, reflects, stores the information

To reconstruct a , just note that $a_n = \frac{G_a^{(n)}(0)}{n!}$, where $G_a^{(n)}$ denotes the n -th derivative of G_a .

What is Generating Function?

Example.

The generating function of $[1, 2, 3, 4, \dots]$ is

$$\sum_{n=0}^{\infty} (n+1)x^n$$

What is Generating Function?

Example.

The generating function of $[1, 2, 3, 4, \dots]$ is

$$\sum_{n=0}^{\infty} (n+1)x^n = \frac{1}{(1-x)^2}$$


knowledge from calculus course



What is Generating Function?


Example.

The generating function of $[1, 2, 3, 4, \dots]$ is

$$\sum_{n=0}^{\infty} (n+1)x^n = \frac{1}{(1-x)^2}$$


knowledge from calculus course

Theorem. (Uniqueness) If $G_a(s) = G_b(s)$ for all $|s| < R'$ where $0 < R' \leq R$, then $a_n = b_n$ for all n .



R is called *radius of convergence*.
However for simplicity we don't
discuss it in slides.

Probability Generating Function

Definition.

The probability generating function of a *discrete* random variable X is defined to be the generating function of its probability mass function:

$$G_X(s) = \mathbb{E}(s^X) = \sum_{i=0}^{\infty} \Pr(X = i) s^i$$

Probability Generating Function

Definition.

The probability generating function of a *discrete* random variable X is defined to be the generating function of its probability mass function:

$$G_X(s) = \mathbb{E}(s^X) = \sum_{i=0}^{\infty} \Pr(X = i) s^i$$



X should be discrete!

Most of the time we assume that X takes non-negative integers. But actually it can take negative integers, and then the form of $G_X(s)$ needs to be changed a bit.

Probability Generating Function

Definition.

The probability generating function of a *discrete* random variable X is defined to be the generating function of its probability mass function:

$$G_X(s) = \mathbb{E}(s^X) = \sum_{i=0}^{\infty} \Pr(X = i) s^i$$

Examples.

- Constant r.v.: $\Pr(X = c) = 1 \longrightarrow G_X(s) = \mathbb{E}(s^X) = s^c$
- Bernoulli r.v.: $\Pr(X = 1) = p, \Pr(X = 0) = 1 - p$
 $\longrightarrow G_X(s) = \mathbb{E}(s^X) = (1 - p)s^0 + ps^1 = (1 - p) + ps$
- Geometric r.v.: $\Pr(X = k) = p(1 - p)^{k-1}$ for $0 < p \leq 1$ and $k \geq 1$
 $\longrightarrow G_X(s) = \mathbb{E}(s^X) = \sum_{k=1}^{\infty} p(1 - p)^{k-1} s^k = \frac{ps}{1 - s(1 - p)}$

Probability Generating Function

Probability generating functions have many important and interesting properties.

Properties.

- (1) Expectations: $\mathbb{E}[X(X-1)\dots(X-k+1)] = G_X^{(k)}(1)$
More specifically, $\mathbb{E}(X) = G'_X(1)$.

Probability Generating Function

Probability generating functions have many important and interesting properties.

Properties.

(1) Expectations: $\mathbb{E}[X(X-1)\dots(X-k+1)] = G_X^{(k)}(1)$

More specifically, $\mathbb{E}(X) = G'_X(1)$.

(2) Variances:
$$\begin{aligned}\text{Var}(X) &= \mathbb{E}(X^2) - \mathbb{E}(X)^2 = \mathbb{E}(X(X-1)) + \mathbb{E}(X) - \mathbb{E}(X)^2 \\ &= G''_X(1) + G'_X(1) - G'_X(1)^2\end{aligned}$$

Probability Generating Function

Probability generating functions have many important and interesting properties.

Properties.

(1) Expectations: $\mathbb{E}[X(X-1)\dots(X-k+1)] = G_X^{(k)}(1)$

More specifically, $\mathbb{E}(X) = G'_X(1)$.

(2) Variances: $\text{Var}(X) = \mathbb{E}(X^2) - \mathbb{E}(X)^2 = \mathbb{E}(X(X-1)) + \mathbb{E}(X) - \mathbb{E}(X)^2$
 $= G''_X(1) + G'_X(1) - G'_X(1)^2$

(3) Generating function of the sum of two independent r.v.:

$$G_{X+Y}(s) = G_X(s)G_Y(s) \text{ if } X \text{ and } Y \text{ are independent.}$$

More over, if $S = X_1 + X_2 + \dots + X_n$ with (X_i) being independent, then

$$G_S = G_{X_1}G_{X_2}\dots G_{X_n}.$$

Probability Generating Function

Properties.

(4) Convolution of sequences:

Given two sequences (a_n) and (b_n) , the convolution sequence (c_n) is defined as

$$c_n = \sum_{i=0}^n a_i b_{n-i},$$

and we denote the convolution by $c = a * b$.

Probability Generating Function

Properties.

(4) Convolution of sequences:

Given two sequences (a_n) and (b_n) , the convolution sequence (c_n) is defined as

$$c_n = \sum_{i=0}^n a_i b_{n-i},$$

and we denote the convolution by $c = a * b$.

Take a look at the generating function of (c_n) :

$$G_c(s) = \sum_{n=0}^{\infty} c_n s^n$$

Probability Generating Function

Properties.

(4) Convolution of sequences:

Given two sequences (a_n) and (b_n) , the convolution sequence (c_n) is defined as

$$c_n = \sum_{i=0}^n a_i b_{n-i},$$

and we denote the convolution by $c = a * b$.

Take a look at the generating function of (c_n) :

$$G_c(s) = \sum_{n=0}^{\infty} c_n s^n = \sum_{n=0}^{\infty} s^n \sum_{i=0}^n a_i b_{n-i}$$

Probability Generating Function

Properties.

(4) Convolution of sequences:

Given two sequences (a_n) and (b_n) , the convolution sequence (c_n) is defined as

$$c_n = \sum_{i=0}^n a_i b_{n-i},$$

and we denote the convolution by $c = a * b$.

Take a look at the generating function of (c_n) :

$$\begin{aligned} G_c(s) &= \sum_{n=0}^{\infty} c_n s^n = \sum_{n=0}^{\infty} s^n \sum_{i=0}^n a_i b_{n-i} \\ &= \sum_{i=0}^{\infty} s^i a_i \sum_{n=i}^{\infty} s^{n-i} b_{n-i} \end{aligned}$$

Probability Generating Function

Properties.

(4) Convolution of sequences:

Given two sequences (a_n) and (b_n) , the convolution sequence (c_n) is defined as

$$c_n = \sum_{i=0}^n a_i b_{n-i},$$

and we denote the convolution by $c = a * b$.

Take a look at the generating function of (c_n) :

$$\begin{aligned} G_c(s) &= \sum_{n=0}^{\infty} c_n s^n = \sum_{n=0}^{\infty} s^n \sum_{i=0}^n a_i b_{n-i} \\ &= \sum_{i=0}^{\infty} s^i a_i \sum_{n=i}^{\infty} s^{n-i} b_{n-i} \\ &= \left(\sum_{i=0}^{\infty} s^i a_i \right) \left(\sum_{j=0}^{\infty} s^j b_j \right) = G_a(s) G_b(s) \end{aligned}$$

Probability Generating Function

Properties.

(4) Convolution of sequences:

Given two sequences (a_n) and (b_n) , the convolution sequence (c_n) is defined as

$$c_n = \sum_{i=0}^n a_i b_{n-i},$$

and we denote the convolution by $c = a * b$.

Take a look at the generating function of (c_n) :

$$\begin{aligned} G_c(s) &= \sum_{n=0}^{\infty} c_n s^n = \sum_{n=0}^{\infty} s^n \sum_{i=0}^n a_i b_{n-i} \\ &= \sum_{i=0}^{\infty} s^i a_i \sum_{n=i}^{\infty} s^{n-i} b_{n-i} \\ &= \left(\sum_{i=0}^{\infty} s^i a_i \right) \left(\sum_{j=0}^{\infty} s^j b_j \right) = G_a(s) G_b(s) \end{aligned}$$

$G_c(s) = G_a(s)G_b(s)!$

Probability Generating Function

Similar to joint probability distribution function, we can take a look at the *joint probability generating function* of two r.v. X_1 and X_2 :

Probability Generating Function

Similar to joint probability distribution function, we can take a look at the *joint probability generating function* of two r.v. X_1 and X_2 :

Definition. The joint probability generating function of X_1 and X_2 is defined as

$$G_{X_1, X_2}(s_1, s_2) = \mathbb{E}(s_1^{X_1} s_2^{X_2})$$

Probability Generating Function

Similar to joint probability distribution function, we can take a look at the *joint probability generating function* of two r.v. X_1 and X_2 :

Definition. The joint probability generating function of X_1 and X_2 is defined as

$$G_{X_1, X_2}(s_1, s_2) = \mathbb{E}(s_1^{X_1} s_2^{X_2})$$

The joint probability generating function can help us characterize “independence of random variables”

Probability Generating Function

Theorem. X_1 and X_2 are independent *if and only if*

$$G_{X_1, X_2}(s_1, s_2) = G_{X_1}(s_1)G_{X_2}(s_2) \text{ for all } s_1 \text{ and } s_2.$$

Probability Generating Function

Theorem. X_1 and X_2 are independent *if and only if*

$$G_{X_1, X_2}(s_1, s_2) = G_{X_1}(s_1)G_{X_2}(s_2) \text{ for all } s_1 \text{ and } s_2.$$

Proof.

X_1 and X_2 are independent.



$$\Pr[X_1 = n, X_2 = m] = \Pr[X_1 = n]\Pr[X_2 = m] \text{ for all } n, m.$$

Probability Generating Function

Theorem. X_1 and X_2 are independent *if and only if*

$$G_{X_1, X_2}(s_1, s_2) = G_{X_1}(s_1)G_{X_2}(s_2) \text{ for all } s_1 \text{ and } s_2.$$

Proof.

X_1 and X_2 are independent.



$$\Pr[X_1 = n, X_2 = m] = \Pr[X_1 = n]\Pr[X_2 = m] \text{ for all } n, m.$$



$$\begin{aligned} \sum_n \sum_m \Pr[X_1 = n, X_2 = m] s_1^n s_2^m &= \sum_n \sum_m \Pr[X_1 = n] \Pr[X_2 = m] s_1^n s_2^m \\ &= \left(\sum_n \Pr[X_1 = n] s_1^n \right) \left(\sum_m \Pr[X_2 = m] s_2^m \right) \quad (\text{for all } s_1, s_2) \end{aligned}$$

Probability Generating Function

Theorem. X_1 and X_2 are independent *if and only if*

$$G_{X_1, X_2}(s_1, s_2) = G_{X_1}(s_1)G_{X_2}(s_2) \text{ for all } s_1 \text{ and } s_2.$$

Proof.

X_1 and X_2 are independent.



$$\Pr[X_1 = n, X_2 = m] = \Pr[X_1 = n]\Pr[X_2 = m] \text{ for all } n, m.$$



$$\begin{aligned} \sum_n \sum_m \Pr[X_1 = n, X_2 = m] s_1^n s_2^m &= \sum_n \sum_m \Pr[X_1 = n] \Pr[X_2 = m] s_1^n s_2^m \\ &= \left(\sum_n \Pr[X_1 = n] s_1^n \right) \left(\sum_m \Pr[X_2 = m] s_2^m \right) \quad (\text{for all } s_1, s_2) \end{aligned}$$

$$G_{X_1, X_2}(s_1, s_2)$$

Probability Generating Function

Theorem. X_1 and X_2 are independent *if and only if*

$$G_{X_1, X_2}(s_1, s_2) = G_{X_1}(s_1)G_{X_2}(s_2) \text{ for all } s_1 \text{ and } s_2.$$

Proof.


X_1 and X_2 are independent.




$$\Pr[X_1 = n, X_2 = m] = \Pr[X_1 = n]\Pr[X_2 = m] \text{ for all } n, m.$$



$$\begin{aligned} \sum_n \sum_m \Pr[X_1 = n, X_2 = m] s_1^n s_2^m &= \sum_n \sum_m \Pr[X_1 = n] \Pr[X_2 = m] s_1^n s_2^m \\ &= \left(\sum_n \Pr[X_1 = n] s_1^n \right) \left(\sum_m \Pr[X_2 = m] s_2^m \right) \quad (\text{for all } s_1, s_2) \end{aligned}$$


$$G_{X_1, X_2}(s_1, s_2)$$


$$G_{X_1}(s_1)G_{X_2}(s_2)$$

Probability Generating Function

Theorem. X_1 and X_2 are independent *if and only if*

$$G_{X_1, X_2}(s_1, s_2) = G_{X_1}(s_1)G_{X_2}(s_2) \text{ for all } s_1 \text{ and } s_2.$$

Proof.

X_1 and X_2 are independent.



$$\Pr[X_1 = n, X_2 = m] = \Pr[X_1 = n]\Pr[X_2 = m] \text{ for all } n, m.$$



$$\begin{aligned} \sum_n \sum_m \Pr[X_1 = n, X_2 = m] s_1^n s_2^m &= \sum_n \sum_m \Pr[X_1 = n]\Pr[X_2 = m] s_1^n s_2^m \\ &= \left(\sum_n \Pr[X_1 = n] s_1^n \right) \left(\sum_m \Pr[X_2 = m] s_2^m \right) \quad (\text{for all } s_1, s_2) \end{aligned}$$

$$G_{X_1, X_2}(s_1, s_2) = G_{X_1}(s_1)G_{X_2}(s_2)$$



Example: Matching Envelope Problem

Here is an example of applying generating functions:

Example. An absent-minded secretary prepares n letters and n matching envelopes to send to n different people, but then but then randomly stuffs the letters into the envelopes. What is the probability p_n that all letters are inserted into the wrong envelopes.

Example: Matching Envelope Problem

Here is an example of applying generating functions:

Example. An absent-minded secretary prepares n letters and n matching envelopes to send to n different people, but then randomly stuffs the letters into the envelopes. What is the probability p_n that all letters are inserted into the wrong envelopes.

Solution.

Let F denote the event: “the first letter is *not* placed into the correct envelope”.

Obviously $\Pr(F) = 1 - \frac{1}{n}$.

Example: Matching Envelope Problem

Here is an example of applying generating functions:

Example. An absent-minded secretary prepares n letters and n matching envelopes to send to n different people, but then randomly stuffs the letters into the envelopes. What is the probability p_n that all letters are inserted into the wrong envelopes.

Solution.

Let F denote the event: “the first letter is *not* placed into the correct envelope”.

Obviously $\Pr(F) = 1 - \frac{1}{n}$.

Let E denote that “all letters are mismatched”, and then we have

$$p_n = \Pr(E) = \Pr[E \mid F]\Pr(F).$$

Example: Matching Envelope Problem

Solution. $p_n = \Pr(E) = \Pr[E \mid F]\Pr(F)$

Consider $\alpha_n := \Pr[E \mid F]$. What can we get from α_n ?

Example: Matching Envelope Problem

Solution. $p_n = \Pr(E) = \Pr[E \mid F]\Pr(F)$

Consider $\alpha_n := \Pr[E \mid F]$. What can we get from α_n ?


α_n = under the condition that “the first letter is mismatched”, the probability that “all the rest letters are mismatched”.

Example: Matching Envelope Problem

Solution. $p_n = \Pr(E) = \Pr[E \mid F]\Pr(F)$

Consider $\alpha_n := \Pr[E \mid F]$. What can we get from α_n ?

α_n = under the condition that “the first letter is mismatched”, the probability that “all the rest letters are mismatched”.




Example: Matching Envelope Problem

Solution. $p_n = \Pr(E) = \Pr[E \mid F]\Pr(F)$

Consider $\alpha_n := \Pr[E \mid F]$. What can we get from α_n ?

α_n = under the condition that “the first letter is mismatched”, the probability that “all the rest letters are mismatched”.



Example: Matching Envelope Problem

Solution. $p_n = \Pr(E) = \Pr[E \mid F]\Pr(F)$

Consider $\alpha_n := \Pr[E \mid F]$. What can we get from α_n ?

α_n = under the condition that “the first letter is mismatched”, the probability that “all the rest letters are mismatched”.

(1) ● ■ [1]

(2) ● ■ [2]

● ■

● ■

● ■

Letters

Envelopes

Letter (1) is matched to env [2].

Now consider letter (2):

- If (2) is matched to [1], there are 3 letters & envelopes left.


The contribution is $\frac{1}{n-1}p_{n-2}$.

Example: Matching Envelope Problem

Solution. $p_n = \Pr(E) = \Pr[E \mid F]\Pr(F)$

Consider $\alpha_n := \Pr[E \mid F]$. What can we get from α_n ?

α_n = under the condition that “the first letter is mismatched”, the probability that “all the rest letters are mismatched”.



Letter (1) is matched to env [2].

Now consider letter (2):

- If (2) is matched to [1], there are 3 letters & envelopes left.

The contribution is $\frac{1}{n-1}p_{n-2}$.

- If (2) is matched to some other envelope (e.g., [5]), then the question becomes calculating α_{n-1} , since we shall consider (5) now.


The contribution is $\frac{n-2}{n-1}\alpha_{n-1}$.

Example: Matching Envelope Problem

Solution. $p_n = \Pr(E) = \Pr[E \mid F]\Pr(F)$

Consider $\alpha_n := \Pr[E \mid F]$. What can we get from α_n ?

α_n = under the condition that “the first letter is mismatched”, the probability that “all the rest letters are mismatched”.



Letter (1) is matched to env [2].

Now consider letter (2):

- If (2) is matched to [1], there are 3 letters & envelopes left.

The contribution is $\frac{1}{n-1}p_{n-2}$.

- If (2) is matched to some other envelope (e.g., [5]), then the question becomes calculating α_{n-1} , since we shall consider (5) now.

The contribution is $\frac{n-2}{n-1}\alpha_{n-1}$.

Thus, $\alpha_n = \frac{1}{n-1}p_{n-2} + \frac{n-2}{n-1}\alpha_{n-1}$.

Example: Matching Envelope Problem

Solution.

$$p_n = \Pr(E) = \Pr[E \mid F]\Pr(F)$$

Example: Matching Envelope Problem

Solution.

$$\begin{aligned} p_n &= \Pr(E) = \Pr[E \mid F] \Pr(F) \\ &= \left(1 - \frac{1}{n}\right) \alpha_n \\ &= \left(1 - \frac{1}{n}\right) \left(\frac{1}{n-1} p_{n-2} + \left(1 - \frac{1}{n-1}\right) \alpha_{n-1} \right) \end{aligned}$$

Example: Matching Envelope Problem


Solution.

$$p_n = \Pr(E) = \Pr[E \mid F] \Pr(F)$$

$$= \left(1 - \frac{1}{n}\right) \alpha_n$$

$$= \left(1 - \frac{1}{n}\right) \left(\frac{1}{n-1} p_{n-2} + \left(1 - \frac{1}{n-1}\right) \alpha_{n-1} \right)$$

$p_n = \left(1 - \frac{1}{n}\right) \alpha_n$, so $p_{n-1} = \left(1 - \frac{1}{n-1}\right) \alpha_{n-1}$.



Example: Matching Envelope Problem

Solution.

$$\begin{aligned} p_n &= \Pr(E) = \Pr[E \mid F] \Pr(F) \\ &= \left(1 - \frac{1}{n}\right) \alpha_n \\ &= \left(1 - \frac{1}{n}\right) \left(\frac{1}{n-1} p_{n-2} + \left(1 - \frac{1}{n-1}\right) \alpha_{n-1} \right) \\ &= \left(1 - \frac{1}{n}\right) \left(\frac{1}{n-1} p_{n-2} + p_{n-1} \right) \\ &= \frac{1}{n} p_{n-2} + \left(1 - \frac{1}{n}\right) p_{n-1}, \text{ with } p_1 = 0 \text{ and } p_2 = \frac{1}{2}. \end{aligned}$$

Example: Matching Envelope Problem

Solution.

$$\begin{aligned} p_n &= \Pr(E) = \Pr[E \mid F]\Pr(F) \\ &= \left(1 - \frac{1}{n}\right) \alpha_n \\ &= \left(1 - \frac{1}{n}\right) \left(\frac{1}{n-1}p_{n-2} + \left(1 - \frac{1}{n-1}\right) \alpha_{n-1}\right) \\ &= \left(1 - \frac{1}{n}\right) \left(\frac{1}{n-1}p_{n-2} + p_{n-1}\right) \\ &= \frac{1}{n}p_{n-2} + \left(1 - \frac{1}{n}\right) p_{n-1}, \text{ with } p_1 = 0 \text{ and } p_2 = \frac{1}{2}. \end{aligned}$$

Let $G_p(s) = \sum_{n=1}^{\infty} p_n s^n$. By multiplying the equation above by ns^{n-1} and then taking sum, we get

$$\sum_{n=3}^{\infty} ns^{n-1}p_n = s \sum_{n=3}^{\infty} s^{n-2}p_{n-2} + s \sum_{n=3}^{\infty} (n-1)s^{n-2}p_{n-1}.$$

Using generating function, we have

$$(1-s)G'(s) = sG(s) + s.$$

Example: Matching Envelope Problem

Solution.

$$(1 - s)G'(s) = sG(s) + s, \text{ and } G(0) = 0.$$

Example: Matching Envelope Problem

Solution.

$$(1 - s)G'(s) = sG(s) + s, \text{ and } G(0) = 0.$$



solve this equation

$$G(s) = \frac{1}{1-s}e^{-s} - 1$$

Example: Matching Envelope Problem

Solution.

$$(1 - s)G'(s) = sG(s) + s, \text{ and } G(0) = 0.$$

↓ solve this equation

$$G(s) = \frac{1}{1-s}e^{-s} - 1$$

↓ expand as a power series

$$p_n = 1 + \frac{(-1)}{1!} + \dots + \frac{(-1)^n}{n!}$$



Random Walks

We can use generating functions to study random walks.

Random Walks

We can use generating functions to study random walks.

Let X_1, X_2, \dots be *i.i.d.*. Each variable takes value 1 with probability p and -1 with probability $1 - p$.

$S_n := \sum_{i=1}^n X_i$. And $S := \{S_i \mid i \geq 0\}$ is the simplest random walk which starts at the origin.

Random Walks

We can use generating functions to study random walks.

Let X_1, X_2, \dots be *i.i.d.*. Each variable takes value 1 with probability p and -1 with probability $1 - p$.

$S_n := \sum_{i=1}^n X_i$. And $S := \{S_i \mid i \geq 0\}$ is the simplest random walk which starts at the origin.

What we concern is “when will the man return back to the origin”.

Random Walks

We can use generating functions to study random walks.

Let X_1, X_2, \dots be *i.i.d.*. Each variable takes value 1 with probability p and -1 with probability $1 - p$.

$S_n := \sum_{i=1}^n X_i$. And $S := \{S_i \mid i \geq 0\}$ is the simplest random walk which starts at the origin.

What we concern is “when will the man return back to the origin”.

Since the process between two consecutive are pairwise independent, we can only focus on the time that the man returns the origin *for the first time*.

Random Walks

Let $p_0(n) := \Pr[S_n = 0]$ be the probability that he returns the origin at timestamp n .

Let $f_0(n) := \Pr[S_1 \neq 0, \dots, S_{n-1} \neq 0, S_n = 0]$ be the probability that he returns the origin for the first time at timestamp n .

And write down the generating function of p_0 and f_0 :

$$P_0(s) = \sum_{n=0}^{\infty} p_0(n) s^n$$

$$F_0(s) = \sum_{n=1}^{\infty} f_0(n) s^n$$

Random Walks

Let $p_0(n) := \Pr[S_n = 0]$ be the probability that he returns the origin at timestamp n .

Let $f_0(n) := \Pr[S_1 \neq 0, \dots, S_{n-1} \neq 0, S_n = 0]$ be the probability that he returns the origin for the first time at timestamp n .

And write down the generating function of p_0 and f_0 :

$$P_0(s) = \sum_{n=0}^{\infty} p_0(n) s^n$$

$$F_0(s) = \sum_{n=1}^{\infty} f_0(n) s^n$$

Let T_0 be the r.v. denoting the time of the first return.

Note that F_0 is the probability generating function of T_0 , and that is, $F_0(s) = \mathbb{E}(s^{T_0})$. (Since $f_0(n) = \Pr[T_0 = n]$.)

Random Walks

Let $p_0(n) := \Pr[S_n = 0]$ be the probability that he returns the origin at timestamp n .
Let $f_0(n) := \Pr[S_1 \neq 0, \dots, S_{n-1} \neq 0, S_n = 0]$ be the probability that he returns the origin for the first time at timestamp n .

And write down the generating function of p_0 and f_0 :

$$P_0(s) = \sum_{n=0}^{\infty} p_0(n) s^n$$
$$F_0(s) = \sum_{n=1}^{\infty} f_0(n) s^n$$

Let T_0 be the r.v. denoting the time of the first return.
Note that F_0 is the probability generating function of T_0 ,
and that is, $F_0(s) = \mathbb{E}(s^{T_0})$. (Since $f_0(n) = \Pr[T_0 = n]$.)

Then we have:

$$(1) \ P_0(s) = (1 - 4p(1 - p)s^2)^{-\frac{1}{2}},$$
$$(2) \ P_0(s) = 1 + P_0(s)F_0(s).$$

From (1), (2), it directly follows that $F_0(s) = 1 - (1 - 4p(1 - p)s^2)^{\frac{1}{2}}$.

Random Walks

Proof of (1).

Assume at some timestamp n , $S_n = 0$.

It means that the man walked $\frac{n}{2}$ steps left, and $\frac{n}{2}$ right. (Obviously n must be even.)

Random Walks

Proof of (1).

Assume at some timestamp n , $S_n = 0$.

It means that the man walked $\frac{n}{2}$ steps left, and $\frac{n}{2}$ right. (Obviously n must be even.)

$$p_0(n) = \binom{n}{\frac{n}{2}} p^{\frac{n}{2}} (1-p)^{\frac{n}{2}}$$

Random Walks

Proof of (1).

Assume at some timestamp n , $S_n = 0$.

It means that the man walked $\frac{n}{2}$ steps left, and $\frac{n}{2}$ right. (Obviously n must be even.)

$$p_0(n) = \binom{n}{\frac{n}{2}} p^{\frac{n}{2}} (1-p)^{\frac{n}{2}} \quad \text{if } n \text{ is even.}$$

$$p_0(n) = 0 \quad \text{if } n \text{ is odd.}$$

Random Walks

Proof of (1).

Assume at some timestamp n , $S_n = 0$.

It means that the man walked $\frac{n}{2}$ steps left, and $\frac{n}{2}$ right. (Obviously n must be even.)

$$p_0(n) = \binom{n}{\frac{n}{2}} p^{\frac{n}{2}} (1-p)^{\frac{n}{2}} \quad \text{if } n \text{ is even.}$$

$$p_0(n) = 0 \quad \text{if } n \text{ is odd.}$$

$$P_0(s) = \sum_{n=0}^{\infty} \binom{2n}{n} p^n (1-p)^n s^{2n}$$

Random Walks

Proof of (1).

Assume at some timestamp n , $S_n = 0$.

It means that the man walked $\frac{n}{2}$ steps left, and $\frac{n}{2}$ right. (Obviously n must be even.)

$$p_0(n) = \binom{n}{\frac{n}{2}} p^{\frac{n}{2}} (1-p)^{\frac{n}{2}} \quad \text{if } n \text{ is even.}$$

$$p_0(n) = 0 \quad \text{if } n \text{ is odd.}$$

$$P_0(s) = \sum_{n=0}^{\infty} \binom{2n}{n} p^n (1-p)^n s^{2n} = (1 - 4p(1-p)s^2)^{-\frac{1}{2}}$$



Random Walks

Proof of (2).

$E_n :=$ the event that $S_n = 0$

The events $([T_0 = k])$ are pairwise disjoint.

Random Walks

Proof of (2).

$E_n :=$ the event that $S_n = 0$

The events $([T_0 = k])$ are pairwise disjoint.



$$p_0(n) = \Pr(E_n) = \sum_{k=1}^n \Pr[T_0 = k] \Pr[E_n \mid T_0 = k] \quad (\text{assuming } n \geq k)$$

Random Walks

Proof of (2).

$E_n :=$ the event that $S_n = 0$

The events $([T_0 = k])$ are pairwise disjoint.



$$\begin{aligned} p_0(n) = \Pr(E_n) &= \sum_{k=1}^n \Pr[T_0 = k] \Pr[E_n \mid T_0 = k] && \text{(assuming } n \geq k) \\ &= \sum_{k=1}^n f_0(k) p_0(n - k) \end{aligned}$$

Random Walks

Proof of (2).

$E_n :=$ the event that $S_n = 0$

The events $([T_0 = k])$ are pairwise disjoint.



$$p_0(n) = \Pr(E_n) = \sum_{k=1}^n \Pr[T_0 = k] \Pr[E_n \mid T_0 = k] \quad (\text{assuming } n \geq k)$$

multiply by s^n
and then sum up

$$= \sum_{k=1}^n f_0(k) p_0(n - k)$$



$$\sum_{n=1}^{\infty} p_0(n) s^n = \sum_{n=1}^{\infty} \left(\sum_{k=1}^n f_0(k) p_0(n - k) \right) s^n$$

Random Walks

Proof of (2).

$E_n :=$ the event that $S_n = 0$

The events $([T_0 = k])$ are pairwise disjoint.



$$p_0(n) = \Pr(E_n) = \sum_{k=1}^n \Pr[T_0 = k] \Pr[E_n \mid T_0 = k] \quad (\text{assuming } n \geq k)$$

multiply by s^n
and then sum up

$$= \sum_{k=1}^n f_0(k) p_0(n - k)$$



$$\sum_{n=1}^{\infty} p_0(n) s^n = \sum_{n=1}^{\infty} \left(\sum_{k=1}^n f_0(k) p_0(n - k) \right) s^n$$

\parallel

$$P_0(s) - 1$$

\parallel by convolution

$$F_0(s) P_0(s)$$



Random Walks

We can get two corollary from (3), which are consistent with our intuition:

- The man will return the origin at least once with probability

$$\sum_{n=1}^{\infty} f_0(n) = F_0(1) = 1 - \sqrt{1 - 4p(1 - p)} = 1 - |2p - 1|.$$

Specially, he almost surely returns the origin *if and only if* $p = \frac{1}{2}$.

- If he returns the origin almost surely (i.e., $p = \frac{1}{2}$), the expected time of the first return is

$$\sum_{n=1}^{\infty} n f_0(n) = F'_0(1) = \infty.$$

Random Walks

There is a relation between f_0 and p_0 :

Proposition. For a symmetric random walk (“symmetric” means that $p = 1 - p = 1/2$), we have $2kf_0(2k) = p_0(2k - 2)$.

Random Walks

There is a relation between f_0 and p_0 :

Proposition. For a symmetric random walk (“symmetric” means that $p = 1 - p = 1/2$), we have $2kf_0(2k) = p_0(2k - 2)$.

Proof.

Consider the generating function of LHS:

$$\sum_{k=1}^{\infty} 2kf_0(2k)s^{2k} = s \sum_{k=1}^{\infty} f_0(2k)(2ks^{2k-1}) = sF'_0(s) = \frac{s^2}{\sqrt{1-s^2}}$$

Note that a man must have walked even steps if he returns the origin.
So $f_0(k) = 0$ if k is odd.

Random Walks

There is a relation between f_0 and p_0 :

Proposition. For a symmetric random walk (“symmetric” means that $p = 1 - p = 1/2$), we have $2kf_0(2k) = p_0(2k - 2)$.

Proof.

Consider the generating function of LHS:

$$\sum_{k=1}^{\infty} 2kf_0(2k)s^{2k} = s \sum_{k=1}^{\infty} f_0(2k)(2ks^{2k-1}) = sF'_0(s) = \frac{s^2}{\sqrt{1-s^2}}$$

Meanwhile, from RHS we have:

$$\sum_{k=1}^{\infty} p_0(2k - 2)s^{2k} = s^2 \sum_{k=1}^{\infty} p_0(2k - 2)s^{2k-2} = s^2 P_0(s) = \frac{s^2}{\sqrt{1-s^2}}$$

Random Walks

There is a relation between f_0 and p_0 :

Proposition. For a symmetric random walk (“symmetric” means that $p = 1 - p = 1/2$), we have $2kf_0(2k) = p_0(2k - 2)$.

Proof.

Consider the generating function of LHS:

$$\sum_{k=1}^{\infty} 2kf_0(2k)s^{2k} = s \sum_{k=1}^{\infty} f_0(2k)(2ks^{2k-1}) = sF'_0(s) = \frac{s^2}{\sqrt{1-s^2}}$$

Meanwhile, from RHS we have:

$$\sum_{k=1}^{\infty} p_0(2k - 2)s^{2k} = s^2 \sum_{k=1}^{\infty} p_0(2k - 2)s^{2k-2} = s^2 P_0(s) = \frac{s^2}{\sqrt{1-s^2}}$$

Therefore $2kf_0(2k) = p_0(2k - 2)$.



Random Walks

And another proposition.

Proposition. For a symmetric random walk, $p_0(2n) = \Pr[S_1 \neq 0, S_2 \neq 0, \dots, S_{2n} \neq 0]$.

Random Walks

And another proposition.

Proposition. For a symmetric random walk, $p_0(2n) = \Pr[S_1 \neq 0, S_2 \neq 0, \dots, S_{2n} \neq 0]$.

Proof.

$$\text{Let } a_n := \Pr[S_1 \neq 0, S_2 \neq 0, \dots, S_{2n} \neq 0] = \sum_{\substack{k=2n+2 \\ k \text{ is even}}} f_0(k).$$

Random Walks

And another proposition.

Proposition. For a symmetric random walk, $p_0(2n) = \Pr[S_1 \neq 0, S_2 \neq 0, \dots, S_{2n} \neq 0]$.

Proof.

$$\text{Let } a_n := \Pr[S_1 \neq 0, S_2 \neq 0, \dots, S_{2n} \neq 0] = \sum_{\substack{k=2n+2 \\ k \text{ is even}}} f_0(k).$$

Consider the generating function of a_n :

$$\begin{aligned} \sum_{n=0}^{\infty} a_n s^{2n} &= \sum_{n=0}^{\infty} s^{2n} \sum_{\substack{k=2n+2 \\ k \text{ is even}}} f_0(k) = \sum_{\substack{k=2 \\ k \text{ is even}}}^{\infty} f_0(k) \sum_{n=0}^{\frac{1}{2}k-1} s^{2n} = \sum_{\substack{k=2 \\ k \text{ is even}}}^{\infty} f_0(k) \frac{1-s^k}{1-s^2} \\ &= (1 - F_0(s)) \frac{1}{1-s^2} = \frac{1}{\sqrt{1-s^2}} \end{aligned}$$

Random Walks

And another proposition.

Proposition. For a symmetric random walk, $p_0(2n) = \Pr[S_1 \neq 0, S_2 \neq 0, \dots, S_{2n} \neq 0]$.

Proof.

$$\text{Let } a_n := \Pr[S_1 \neq 0, S_2 \neq 0, \dots, S_{2n} \neq 0] = \sum_{\substack{k=2n+2 \\ k \text{ is even}}} f_0(k).$$

Consider the generating function of a_n :

$$\begin{aligned} \sum_{n=0}^{\infty} a_n s^{2n} &= \sum_{n=0}^{\infty} s^{2n} \sum_{\substack{k=2n+2 \\ k \text{ is even}}} f_0(k) = \sum_{\substack{k=2 \\ k \text{ is even}}}^{\infty} f_0(k) \sum_{n=0}^{\frac{1}{2}k-1} s^{2n} = \sum_{\substack{k=2 \\ k \text{ is even}}}^{\infty} f_0(k) \frac{1-s^k}{1-s^2} \\ &= (1 - F_0(s)) \frac{1}{1-s^2} = \frac{1}{\sqrt{1-s^2}} \end{aligned}$$

On the other hand, we have

$$\sum_{n=0}^{\infty} p_0(2n) s^{2n} = P_0(s) = \frac{1}{\sqrt{1-s^2}}.$$

Random Walks

And another proposition.

Proposition. For a symmetric random walk, $p_0(2n) = \Pr[S_1 \neq 0, S_2 \neq 0, \dots, S_{2n} \neq 0]$.

Proof.

$$\text{Let } a_n := \Pr[S_1 \neq 0, S_2 \neq 0, \dots, S_{2n} \neq 0] = \sum_{\substack{k=2n+2 \\ k \text{ is even}}} f_0(k).$$

Consider the generating function of a_n :

$$\begin{aligned} \sum_{n=0}^{\infty} a_n s^{2n} &= \sum_{n=0}^{\infty} s^{2n} \sum_{\substack{k=2n+2 \\ k \text{ is even}}} f_0(k) = \sum_{\substack{k=2 \\ k \text{ is even}}}^{\infty} f_0(k) \sum_{n=0}^{\frac{1}{2}k-1} s^{2n} = \sum_{\substack{k=2 \\ k \text{ is even}}}^{\infty} f_0(k) \frac{1-s^k}{1-s^2} \\ &= (1 - F_0(s)) \frac{1}{1-s^2} = \frac{1}{\sqrt{1-s^2}} \end{aligned}$$

On the other hand, we have

$$\sum_{n=0}^{\infty} p_0(2n) s^{2n} = P_0(s) = \frac{1}{\sqrt{1-s^2}}.$$

Therefore $p_0(2n) = a_n$. ■

Random Walks

Now consider the time that the man first arrives at position r .

Random Walks

Now consider the time that the man first arrives at position r .

Define $f_r(n) = \Pr[S_1 \neq r, \dots, S_{n-1} \neq r, S_n = r]$.

And the corresponding generating function $F_r(s) = \sum_{n=1}^{\infty} f_r(n)s^n$.

Random Walks

Now consider the time that the man first arrives at position r .

Define $f_r(n) = \Pr[S_1 \neq r, \dots, S_{n-1} \neq r, S_n = r]$.

And the corresponding generating function $F_r(s) = \sum_{n=1}^{\infty} f_r(n)s^n$.

We have

- (1) $F_r(s) = [F_1(s)]^r$ for $r \geq 1$,
- (2) $F_1(s) = [1 - (1 - 4p(1 - p)s^2)^{\frac{1}{2}}]/(2(1 - p)s)$.

Random Walks

Proof of (1).

$$f_r(n) = \sum_{k=1}^{n-1} f_1(k) f_{r-1}(n-k) \text{ for } r > 1.$$

Then the generating function is

$$F_r(s) = F_1(s) F_{r-1}(s)$$

Random Walks

Proof of (1).

$$f_r(n) = \sum_{k=1}^{n-1} f_1(k) f_{r-1}(n-k) \text{ for } r > 1.$$

Then the generating function is

$$F_r(s) = F_1(s) F_{r-1}(s) = [F_1(s)]^r.$$



Random Walks

Proof of (1).

$$f_r(n) = \sum_{k=1}^{n-1} f_1(k) f_{r-1}(n-k) \text{ for } r > 1.$$

Then the generating function is

$$F_r(s) = F_1(s) F_{r-1}(s) = [F_1(s)]^r.$$



Proof of (2).

Define T_r to be $\min\{n: S_n = r\}$, that is, the first time he arrives at position r .

Random Walks

Proof of (1).

$$f_r(n) = \sum_{k=1}^{n-1} f_1(k) f_{r-1}(n-k) \text{ for } r > 1.$$

Then the generating function is

$$F_r(s) = F_1(s) F_{r-1}(s) = [F_1(s)]^r.$$



Proof of (2).

Define T_r to be $\min\{n: S_n = r\}$, that is, the first time he arrives at position r .

By this definition, T_n may be ∞ , but that's not a big deal.

Random Walks

Proof of (1).

$$f_r(n) = \sum_{k=1}^{n-1} f_1(k) f_{r-1}(n-k) \text{ for } r > 1.$$

Then the generating function is

$$F_r(s) = F_1(s) F_{r-1}(s) = [F_1(s)]^r.$$



Proof of (2).

Define T_r to be $\min\{n: S_n = r\}$, that is, the first time he arrives at position r .

$$\begin{aligned} f_1(n) &= \Pr[T_1 = n] = \Pr[T_1 = n \mid X_1 = 1]p + \Pr[T_1 = n \mid X_1 = -1](1-p) \\ &= 0 \cdot p + \Pr[T_2 = n-1](1-p) \\ &= (1-p)f_2(n-1), \text{ for } n > 1. \end{aligned}$$

Random Walks

Proof of (2).

$$f_1(n) = (1 - p)f_2(n - 1) \text{ for } n > 1$$

Again by multiplying and summing, we have

$$(1 - p)sF_1(s)^2 - F_1(s) + ps = 0.$$

Random Walks

Proof of (2).

$$f_1(n) = (1 - p)f_2(n - 1) \text{ for } n > 1$$

Again by multiplying and summing, we have

$$(1 - p)sF_1(s)^2 - F_1(s) + ps = 0.$$

Since $F_1(0) = 0$, by solving the equation above we get

$$F_1(s) = \frac{1 - \sqrt{1 - 4p(1 - p)s^2}}{2(1 - p)s}.$$



Random Walks

- (1) $F_r(s) = [F_1(s)]^r$ for $r \geq 1$,
- (2) $F_1(s) = [1 - (1 - 4p(1 - p)s^2)^{\frac{1}{2}}] / (2(1 - p)s)$.

Similar to “returning to the origin”, we can also conclude a corollary:

Corollary. The man will arrive at position 1 at least once with probability

$$F_1(1) = \frac{1 - |2p - 1|}{2(1 - p)} = \min\left\{\frac{p}{1 - p}, 1\right\}.$$

Random Walks

Next we consider a more general case: *right-continuous random walks*.

Random Walks


Next we consider a more general case: *right-continuous random walks*.

Definition. Let X_1, X_2, \dots be i.i.d. random variables taking values in integers and $S_n := X_1 + \dots + X_n$. A random walk is called *right-continuous* if $\Pr[X_i > 1] = 0$.

Random Walks

Next we consider a more general case: *right-continuous random walks*.

Definition. Let X_1, X_2, \dots be i.i.d. random variables taking values in integers and $S_n := X_1 + \dots + X_n$. A random walk is called *right-continuous* if $\Pr[X_i > 1] = 0$.



In this discrete setting, “continuous” means that “the difference is no more than 1”, i.e., the step distribution puts no mass on integers ≥ 2 .

Random Walks

Next we consider a more general case: *right-continuous random walks*.

Definition. Let X_1, X_2, \dots be i.i.d. random variables taking values in integers and $S_n := X_1 + \dots + X_n$. A random walk is called *right-continuous* if $\Pr[X_i > 1] = 0$.

For generality, assume $\Pr[X_i = 1] > 0$. We want to prove the following theorem.

Theorem. (Hitting time theorem) Let S be a right-continuous random walk and T_r be the time the man first arrives at position r . Then

$$\Pr[T_r = n] = \frac{r}{n} \Pr[S_n = r] \text{ for } n, r \geq 1.$$

Random Walks

Next we consider a more general case: *right-continuous random walks*.

Definition. Let X_1, X_2, \dots be i.i.d. random variables taking values in integers and $S_n := X_1 + \dots + X_n$. A random walk is called *right-continuous* if $\Pr[X_i > 1] = 0$.

For generality, assume $\Pr[X_i = 1] > 0$. We want to prove the following theorem.

Theorem. (Hitting time theorem) Let S be a right-continuous random walk and T_r be the time the man first arrives at position r . Then

$$\Pr[T_r = n] = \frac{r}{n} \Pr[S_n = r] \text{ for } n, r \geq 1.$$



Quite an interesting statement!

Random Walks

Theorem. (Hitting time theorem) Let S be a right-continuous random walk and T_r be the time the man first arrives at position r . Then

$$\Pr[T_r = n] = \frac{r}{n} \Pr[S_n = r] \text{ for } n, r \geq 1.$$

Proof.

$$\text{Define } G(z) := \mathbb{E}(z^{-X_1}) = \sum_{n=-1}^{\infty} \Pr[X_1 = -n]z^n, \quad F_r(z) := \mathbb{E}(z^{T_r}) = \sum_{n=1}^{\infty} \Pr[T_r = n]z^n. \quad (z \in \mathbb{C})$$

Random Walks

Theorem. (Hitting time theorem) Let S be a right-continuous random walk and T_r be the time the man first arrives at position r . Then

$$\Pr[T_r = n] = \frac{r}{n} \Pr[S_n = r] \text{ for } n, r \geq 1.$$

Proof.

$$\text{Define } G(z) := \mathbb{E}(z^{-X_1}) = \sum_{n=-1}^{\infty} \Pr[X_1 = -n]z^n, F_r(z) := \mathbb{E}(z^{T_r}) = \sum_{n=1}^{\infty} \Pr[T_r = n]z^n. \quad (z \in \mathbb{C})$$

We first focus on $F_1(z)$:

At timestamp 1, the man is at position X_1 .

Since S is right-continuous, we have $1 - X_1 \geq 0$.

$$F_1(z) = \mathbb{E}(z^{T_1})$$

Random Walks

Theorem. (Hitting time theorem) Let S be a right-continuous random walk and T_r be the time the man first arrives at position r . Then

$$\Pr[T_r = n] = \frac{r}{n} \Pr[S_n = r] \text{ for } n, r \geq 1.$$

Proof.

Define $G(z) := \mathbb{E}(z^{-X_1}) = \sum_{n=-1}^{\infty} \Pr[X_1 = -n]z^n$, $F_r(z) := \mathbb{E}(z^{T_r}) = \sum_{n=1}^{\infty} \Pr[T_r = n]z^n$. $(z \in \mathbb{C})$

We first focus on $F_1(z)$:

At timestamp 1, the man is at position X_1 .

Since S is right-continuous, we have $1 - X_1 \geq 0$.

$$F_1(z) = \mathbb{E}(z^{T_1})$$

The distance between the man and position 1 at timestamp 1.

Random Walks

Theorem. (Hitting time theorem) Let S be a right-continuous random walk and T_r be the time the man first arrives at position r . Then

$$\Pr[T_r = n] = \frac{r}{n} \Pr[S_n = r] \text{ for } n, r \geq 1.$$

Proof.

Define $G(z) := \mathbb{E}(z^{-X_1}) = \sum_{n=-1}^{\infty} \Pr[X_1 = -n]z^n$, $F_r(z) := \mathbb{E}(z^{T_r}) = \sum_{n=1}^{\infty} \Pr[T_r = n]z^n$. $(z \in \mathbb{C})$


We first focus on $F_1(z)$:

At timestamp 1, the man is at position X_1 .

Since S is right-continuous, we have $1 - X_1 \geq 0$.

$F_1(z) = \mathbb{E}(z^{T_1}) = \mathbb{E}(z^{1+T_J})$, where $J = 1 - X_1$ for shorthand.

The distance between the man and position 1 at timestamp 1.



Random Walks

Theorem. (Hitting time theorem) Let S be a right-continuous random walk and T_r be the time the man first arrives at position r . Then

$$\Pr[T_r = n] = \frac{r}{n} \Pr[S_n = r] \text{ for } n, r \geq 1.$$

Proof.

Define $G(z) := \mathbb{E}(z^{-X_1}) = \sum_{n=-1}^{\infty} \Pr[X_1 = -n]z^n$, $F_r(z) := \mathbb{E}(z^{T_r}) = \sum_{n=1}^{\infty} \Pr[T_r = n]z^n$. $(z \in \mathbb{C})$

We first focus on $F_1(z)$:

At timestamp 1, the man is at position X_1 .

Since S is right-continuous, we have $1 - X_1 \geq 0$.

The distance between the man and position 1 at timestamp 1.

$$\begin{aligned} F_1(z) &= \mathbb{E}(z^{T_1}) = \mathbb{E}(z^{1+T_J}), \text{ where } J = 1 - X_1 \text{ for shorthand.} \\ &= z\mathbb{E}(z^{T_J}) = z\mathbb{E}(F_J(z)) = z\mathbb{E}(F_{1-X_1}(z)) \end{aligned}$$

Random Walks

Theorem. (Hitting time theorem) Let S be a right-continuous random walk and T_r be the time the man first arrives at position r . Then

$$\Pr[T_r = n] = \frac{r}{n} \Pr[S_n = r] \text{ for } n, r \geq 1.$$

Proof.

Define $G(z) := \mathbb{E}(z^{-X_1}) = \sum_{n=-1}^{\infty} \Pr[X_1 = -n]z^n$, $F_r(z) := \mathbb{E}(z^{T_r}) = \sum_{n=1}^{\infty} \Pr[T_r = n]z^n$. $(z \in \mathbb{C})$

We first focus on $F_1(z)$:

At timestamp 1, the man is at position X_1 .

Since S is right-continuous, we have $1 - X_1 \geq 0$.

The distance between the man and position 1 at timestamp 1.

$F_1(z) = \mathbb{E}(z^{T_1}) = \mathbb{E}(z^{1+T_J})$, where $J = 1 - X_1$ for shorthand.

$$= z\mathbb{E}(z^{T_J}) = z\mathbb{E}(F_J(z)) = z\mathbb{E}(F_{1-X_1}(z))$$

$$= z\mathbb{E}(F_1(z)^{1-X_1})$$

Random Walks

Theorem. (Hitting time theorem) Let S be a right-continuous random walk and T_r be the time the man first arrives at position r . Then

$$\Pr[T_r = n] = \frac{r}{n} \Pr[S_n = r] \text{ for } n, r \geq 1.$$

Proof.

Define $G(z) := \mathbb{E}(z^{-X_1}) = \sum_{n=-1}^{\infty} \Pr[X_1 = -n]z^n$, $F_r(z) := \mathbb{E}(z^{T_r}) = \sum_{n=1}^{\infty} \Pr[T_r = n]z^n$. $(z \in \mathbb{C})$

We first focus on $F_1(z)$:

At timestamp 1, the man is at position X_1 .

Since S is right-continuous, we have $1 - X_1 \geq 0$.

The distance between the man and position 1 at timestamp 1.

$F_1(z) = \mathbb{E}(z^{T_1}) = \mathbb{E}(z^{1+T_J})$, where $J = 1 - X_1$ for shorthand.

$$= z\mathbb{E}(z^{T_J}) = z\mathbb{E}(F_J(z)) = z\mathbb{E}(F_{1-X_1}(z))$$

$$= z\mathbb{E}(F_1(z)^{1-X_1}) = zF_1(z)\mathbb{E}(F_1(z)^{-X_1})$$

Random Walks

Theorem. (Hitting time theorem) Let S be a right-continuous random walk and T_r be the time the man first arrives at position r . Then

$$\Pr[T_r = n] = \frac{r}{n} \Pr[S_n = r] \text{ for } n, r \geq 1.$$

Proof.

Define $G(z) := \mathbb{E}(z^{-X_1}) = \sum_{n=-1}^{\infty} \Pr[X_1 = -n]z^n$, $F_r(z) := \mathbb{E}(z^{T_r}) = \sum_{n=1}^{\infty} \Pr[T_r = n]z^n$. $(z \in \mathbb{C})$

We first focus on $F_1(z)$:

At timestamp 1, the man is at position X_1 .

Since S is right-continuous, we have $1 - X_1 \geq 0$.

The distance between the man and position 1 at timestamp 1.

$F_1(z) = \mathbb{E}(z^{T_1}) = \mathbb{E}(z^{1+T_J})$, where $J = 1 - X_1$ for shorthand.

$$= z\mathbb{E}(z^{T_J}) = z\mathbb{E}(F_J(z)) = z\mathbb{E}(F_{1-X_1}(z))$$

$$= z\mathbb{E}(F_1(z)^{1-X_1}) = zF_1(z)\mathbb{E}(F_1(z)^{-X_1})$$

$$= zF_1(z)G(F_1(z))$$

Random Walks

Theorem. (Hitting time theorem) Let S be a right-continuous random walk and T_r be the time the man first arrives at position r . Then

$$\Pr[T_r = n] = \frac{r}{n} \Pr[S_n = r] \text{ for } n, r \geq 1.$$

Proof.

Define $G(z) := \mathbb{E}(z^{-X_1}) = \sum_{n=-1}^{\infty} \Pr[X_1 = -n]z^n$, $F_r(z) := \mathbb{E}(z^{T_r}) = \sum_{n=1}^{\infty} \Pr[T_r = n]z^n$. $(z \in \mathbb{C})$

We first focus on $F_1(z)$:

At timestamp 1, the man is at position X_1 .

Since S is right-continuous, we have $1 - X_1 \geq 0$.

The distance between the man and position 1 at timestamp 1.

$F_1(z) = \mathbb{E}(z^{T_1}) = \mathbb{E}(z^{1+T_J})$, where $J = 1 - X_1$ for shorthand.

$$= z\mathbb{E}(z^{T_J}) = z\mathbb{E}(F_J(z)) = z\mathbb{E}(F_{1-X_1}(z))$$

$$= z\mathbb{E}(F_1(z)^{1-X_1}) = zF_1(z)\mathbb{E}(F_1(z)^{-X_1})$$

$$= zF_1(z)G(F_1(z)) \longrightarrow z = \frac{1}{G(w)}, (w = w(z) := F_1(z))$$

Random Walks

Proof.

$$z = \frac{1}{G(w)}, \quad w = w(z) = F_1(z)$$

Random Walks

Proof.

$$z = \frac{1}{G(w)}, \quad w = w(z) = F_1(z)$$

To proceed with our proof, we need to introduce Lagrange's inversion theorem:

Theorem. (Lagrange inversion theorem) Let $z = \frac{w}{f(w)}$ with $\frac{w}{f(w)}$ being an analytical function of w on a neighborhood of the origin. Suppose that g is infinite differentiable, then

$$g(w(z)) = g(0) + \sum_{n=1}^{\infty} \frac{1}{n!} z^n \left[\frac{d^{n-1}}{du^{n-1}} [g'(u) f(u)^n] \right]_{u=0}.$$

Random Walks

Proof.

$$z = \frac{1}{G(w)}, \quad w = w(z) = F_1(z)$$

To proceed with our proof, we need to introduce Lagrange's inversion theorem:

Theorem. (Lagrange inversion theorem) Let $z = \frac{w}{f(w)}$ with $\frac{w}{f(w)}$ being an analytical function of w on a neighborhood of the origin. Suppose that g is infinite differentiable, then

$$g(w(z)) = g(0) + \sum_{n=1}^{\infty} \frac{1}{n!} z^n \left[\frac{d^{n-1}}{du^{n-1}} [g'(u) f(u)^n] \right]_{u=0}.$$

Now let $f(w) := wG(w)$ so that $z = \frac{1}{G(w)} = \frac{w}{f(w)}$.

Random Walks

Proof.


$$z = \frac{1}{G(w)}, \quad w = w(z) = F_1(z)$$

To proceed with our proof, we need to introduce Lagrange's inversion theorem:

Theorem. (Lagrange inversion theorem) Let $z = \frac{w}{f(w)}$ with $\frac{w}{f(w)}$ being an analytical function of w on a neighborhood of the origin. Suppose that g is infinite differentiable, then

$$g(w(z)) = g(0) + \sum_{n=1}^{\infty} \frac{1}{n!} z^n \left[\frac{d^{n-1}}{du^{n-1}} [g'(u) f(u)^n] \right]_{u=0}.$$

Now let $f(w) := wG(w)$ so that $z = \frac{1}{G(w)} = \frac{w}{f(w)}$.



Not too hard to verify that it is analytic on the neighborhood of the origin.

Random Walks

Proof.

$$z = \frac{1}{G(w)} = \frac{w}{f(w)}, \quad w = w(z) = F_1(z)$$

Let $g(w) := w^r = F_1(z)^r = F_r(z)$, and the inversion theorem tells us that

$$g(w(z)) = F_r(z) = g(0) + \sum_{n=1}^{\infty} \frac{1}{n!} z^n D_n,$$

where $D_n = \left[\frac{d^{n-1}}{du^{n-1}} \left[r u^{r-1} u^n G(u)^n \right] \right]_{u=0}$.

Random Walks

Proof.

$$z = \frac{1}{G(w)} = \frac{w}{f(w)}, \quad w = w(z) = F_1(z)$$

Let $g(w) := w^r = F_1(z)^r = F_r(z)$, and the inversion theorem tells us that

$$g(w(z)) = F_r(z) = g(0) + \sum_{n=1}^{\infty} \frac{1}{n!} z^n D_n,$$

where $D_n = \left[\frac{d^{n-1}}{du^{n-1}} [ru^{r-1}u^n G(u)^n] \right]_{u=0}$.



So $\Pr[T_r = n]$ is the coefficient of z^n : $\Pr[T_r = n] = \frac{1}{n!} D_n$.

Random Walks

Proof.

$$z = \frac{1}{G(w)} = \frac{w}{f(w)}, \quad w = w(z) = F_1(z)$$

Let $g(w) := w^r = F_1(z)^r = F_r(z)$, and the inversion theorem tells us that

$$g(w(z)) = F_r(z) = g(0) + \sum_{n=1}^{\infty} \frac{1}{n!} z^n D_n,$$

$$\text{where } D_n = \left[\frac{d^{n-1}}{du^{n-1}} \left[r u^{r-1} u^n G(u)^n \right] \right]_{u=0}.$$

So $\Pr[T_r = n]$ is the coefficient of z^n : $\Pr[T_r = n] = \frac{1}{n!} D_n$.

$$\begin{aligned} \text{Moreover, } D_n &= \left[\frac{d^{n-1}}{du^{n-1}} \left[r u^{r-1} u^n G(u)^n \right] \right]_{u=0} = \left[\frac{d^{n-1}}{du^{n-1}} \left[r u^{n+r-1} \mathbb{E}(u^{-S_n}) \right] \right]_{u=0} \\ &= \left[\frac{d^{n-1}}{du^{n-1}} \left[r \sum_{i=-n}^{\infty} u^{n+r-1+i} \Pr[S_i = -i] \right] \right]_{u=0} \end{aligned}$$

Random Walks

Proof.

$$z = \frac{1}{G(w)} = \frac{w}{f(w)}, \quad w = w(z) = F_1(z)$$

Let $g(w) := w^r = F_1(z)^r = F_r(z)$, and the inversion theorem tells us that

$$g(w(z)) = F_r(z) = g(0) + \sum_{n=1}^{\infty} \frac{1}{n!} z^n D_n,$$

$$\text{where } D_n = \left[\frac{d^{n-1}}{du^{n-1}} \left[r u^{r-1} u^n G(u)^n \right] \right]_{u=0}.$$

So $\Pr[T_r = n]$ is the coefficient of z^n : $\Pr[T_r = n] = \frac{1}{n!} D_n$.

$$\begin{aligned} \text{Moreover, } D_n &= \left[\frac{d^{n-1}}{du^{n-1}} \left[r u^{r-1} u^n G(u)^n \right] \right]_{u=0} = \left[\frac{d^{n-1}}{du^{n-1}} \left[r u^{n+r-1} \mathbb{E}(u^{-S_n}) \right] \right]_{u=0} \\ &= \left[\frac{d^{n-1}}{du^{n-1}} \left[r \sum_{i=-n}^{\infty} u^{n+r-1+i} \Pr[S_i = -i] \right] \right]_{u=0} = r(n-1)! \Pr[S_n = r]. \end{aligned}$$

Random Walks

Proof.

$$z = \frac{1}{G(w)} = \frac{w}{f(w)}, \quad w = w(z) = F_1(z)$$

Let $g(w) := w^r = F_1(z)^r = F_r(z)$, and the inversion theorem tells us that

$$g(w(z)) = F_r(z) = g(0) + \sum_{n=1}^{\infty} \frac{1}{n!} z^n D_n,$$

where $D_n = \left[\frac{d^{n-1}}{du^{n-1}} \left[r u^{r-1} u^n G(u)^n \right] \right]_{u=0}.$

$$\Pr[T_r = n] = \frac{r}{n} \Pr[S_n = r]$$

So $\Pr[T_r = n]$ is the coefficient of z^n : $\Pr[T_r = n] = \frac{1}{n!} D_n.$

$$\begin{aligned} \text{Moreover, } D_n &= \left[\frac{d^{n-1}}{du^{n-1}} \left[r u^{r-1} u^n G(u)^n \right] \right]_{u=0} = \left[\frac{d^{n-1}}{du^{n-1}} \left[r u^{n+r-1} \mathbb{E}(u^{-S_n}) \right] \right]_{u=0} \\ &= \left[\frac{d^{n-1}}{du^{n-1}} \left[r \sum_{i=-n}^{\infty} u^{n+r-1+i} \Pr[S_i = -i] \right] \right]_{u=0} = r(n-1)! \Pr[S_n = r]. \end{aligned}$$



Random Walks

Here is an exercise which will use the hitting time theorem to prove:

Exercise. Let $\{S_n : n \geq 0\}$ be a simple symmetric random walk with $S_0 = 0$. And let $T = \min\{n > 0 : S_n = 0\}$. Show that

$$\mathbb{E}(T \wedge 2m) = 2\mathbb{E}|S_{2m}| = 4m\Pr[S_{2m} = 0]$$

for $m \geq 0$.

Random Walks

Here is an exercise which will use the hitting time theorem to prove:

Exercise. Let $\{S_n : n \geq 0\}$ be a simple symmetric random walk with $S_0 = 0$. And let $T = \min\{n > 0 : S_n = 0\}$. Show that

$$\mathbb{E}(T \wedge 2m) = 2\mathbb{E}|S_{2m}| = 4m\Pr[S_{2m} = 0]$$

for $m \geq 0$.

(a) (b) (c)

Proof.

(a), (b) and (c) can be regarded as sequences related to m .

Random Walks

Here is an exercise which will use the hitting time theorem to prove:

Exercise. Let $\{S_n : n \geq 0\}$ be a simple symmetric random walk with $S_0 = 0$. And let $T = \min\{n > 0 : S_n = 0\}$. Show that

$$\mathbb{E}(T \wedge 2m) = 2\mathbb{E}|S_{2m}| = 4m\Pr[S_{2m} = 0]$$

for $m \geq 0$.

(a) (b) (c)

Proof.

(a), (b) and (c) can be regarded as sequences related to m .

So we can consider their generating functions respectively.

Random Walks

Proof.

(a): $\mathbb{E}(T \wedge 2m)$

Random Walks

Proof.

$$(a): \quad \mathbb{E}(T \wedge 2m) = \sum_{k=1}^m f_0(2k) \cdot (2k) + (2m) \cdot \Pr[T > 2m]$$

Random Walks

Proof.

$$(a): \quad \mathbb{E}(T \wedge 2m) = \underbrace{\sum_{k=1}^m f_0(2k) \cdot (2k)}_{T \leq 2m} + \underbrace{(2m) \cdot \Pr[T > 2m]}_{T > 2m}$$

Random Walks

Proof.

$$\begin{aligned} \text{(a): } \mathbb{E}(T \wedge 2m) &= \sum_{k=1}^m f_0(2k) \cdot (2k) + (2m) \cdot \Pr[T > 2m] \\ &\quad \parallel \quad \text{by previous} \quad \parallel \\ &\quad \text{propositions} \\ &= \sum_{k=1}^m p_0(2k-2) + 2mp_0(2m) \end{aligned}$$

Random Walks

Proof.

$$\begin{aligned}
 \text{(a): } \mathbb{E}(T \wedge 2m) &= \sum_{k=1}^m f_0(2k) \cdot (2k) + (2m) \cdot \Pr[T > 2m] \\
 &\quad \parallel \quad \text{by previous} \quad \parallel \\
 &\quad \text{propositions} \\
 &= \sum_{k=1}^m p_0(2k - 2) + 2mp_0(2m)
 \end{aligned}$$

Therefore

$$\begin{aligned}
 \sum_{m=0}^{\infty} \mathbb{E}(T \wedge 2m) s^{2m} &= \sum_{m=0}^{\infty} \sum_{k=1}^m p_0(2k - 2) s^{2m} + \sum_{m=0}^{\infty} 2mp_0(2m) s^{2m} \\
 &= \sum_{k=1}^{\infty} p_0(2k - 2) \sum_{m=k}^{\infty} s^{2m} + s \sum_{m=0}^{\infty} 2mp_0(2m) s^{2m-1} \\
 &= \sum_{k=1}^{\infty} p_0(2k - 2) \frac{s^{2k}}{1-s^2} + sP'_0(s) \\
 &= \frac{s^2}{1-s^2} P_0(s) + sP'_0(s) = \frac{2s^2}{(1-s^2)^{\frac{3}{2}}}.
 \end{aligned}$$

Random Walks


Proof.

(b):
$$\sum_{m=0}^{\infty} 2\mathbb{E}|S_{2m}|s^{2m}$$

Random Walks

Proof.

(b):
$$\sum_{m=0}^{\infty} 2\mathbb{E}|S_{2m}|s^{2m} = \sum_{m=0}^{\infty} 2 \cdot 2 \left(\sum_{k=1}^m 2k \Pr[S_{2m} = 2k] \right) s^{2m}$$




by symmetry of random walk

Random Walks

Proof.

$$(b): \sum_{m=0}^{\infty} 2\mathbb{E}|S_{2m}| s^{2m} = \sum_{m=0}^{\infty} 2 \cdot 2 \left(\sum_{k=1}^m 2k \Pr[S_{2m} = 2k] \right) s^{2m}$$



$$= 4 \sum_{m=0}^{\infty} \left(\sum_{k=1}^m 2mf_{2k}(2m) \right) s^{2m}$$

by hitting time theorem

Random Walks

Proof.

$$\begin{aligned} \text{(b): } \sum_{m=0}^{\infty} 2\mathbb{E}|S_{2m}|s^{2m} &= \sum_{m=0}^{\infty} 2 \cdot 2 \left(\sum_{k=1}^m 2k \Pr[S_{2m} = 2k] \right) s^{2m} \\ &= 4 \sum_{m=0}^{\infty} \left(\sum_{k=1}^m 2m f_{2k}(2m) \right) s^{2m} \\ &= 4s \left[\sum_{m=0}^{\infty} \left(\sum_{k=1}^m f_{2k}(2m) \right) s^{2m} \right]' \end{aligned}$$

Random Walks


Proof.

$$(b): \sum_{m=0}^{\infty} 2\mathbb{E}|S_{2m}|s^{2m} = \sum_{m=0}^{\infty} 2 \cdot 2 \left(\sum_{k=1}^m 2k \Pr[S_{2m} = 2k] \right) s^{2m}$$

$$= 4 \sum_{m=0}^{\infty} \left(\sum_{k=1}^m 2m f_{2k}(2m) \right) s^{2m}$$

when $m < k$, $f_{2k}(2m) = 0$

$$= 4s \left[\sum_{m=0}^{\infty} \left(\sum_{k=1}^m f_{2k}(2m) \right) s^{2m} \right]'$$

$$= 4s \left[\sum_{k=1}^{\infty} \sum_{m=0}^{\infty} f_{2k}(2m) s^{2m} \right]'$$


Random Walks

Proof.

$$\begin{aligned} \text{(b): } \sum_{m=0}^{\infty} 2\mathbb{E}|S_{2m}|s^{2m} &= \sum_{m=0}^{\infty} 2 \cdot 2 \left(\sum_{k=1}^m 2k \Pr[S_{2m} = 2k] \right) s^{2m} \\ &= 4 \sum_{m=0}^{\infty} \left(\sum_{k=1}^m 2m f_{2k}(2m) \right) s^{2m} \\ &= 4s \left[\sum_{m=0}^{\infty} \left(\sum_{k=1}^m f_{2k}(2m) \right) s^{2m} \right]' \\ &= 4s \left[\sum_{k=1}^{\infty} \sum_{m=0}^{\infty} f_{2k}(2m) s^{2m} \right]' \\ &= 4s \left[\sum_{k=1}^{\infty} F_{2k}(s) \right]' = 4s \left[\sum_{k=1}^{\infty} F_1(s)^{2k} \right]' = 4s \left[\frac{F_1(s)^2}{1 - F_1(s)^2} \right]' \end{aligned}$$

Random Walks

Proof.

$$\begin{aligned} \text{(b): } \sum_{m=0}^{\infty} 2\mathbb{E}|S_{2m}|s^{2m} &= \sum_{m=0}^{\infty} 2 \cdot 2 \left(\sum_{k=1}^m 2k \Pr[S_{2m} = 2k] \right) s^{2m} \\ &= 4 \sum_{m=0}^{\infty} \left(\sum_{k=1}^m 2m f_{2k}(2m) \right) s^{2m} \\ &= 4s \left[\sum_{m=0}^{\infty} \left(\sum_{k=1}^m f_{2k}(2m) \right) s^{2m} \right]' \\ &= 4s \left[\sum_{k=1}^{\infty} \sum_{m=0}^{\infty} f_{2k}(2m) s^{2m} \right]' \\ &= 4s \left[\sum_{k=1}^{\infty} F_{2k}(s) \right]' = 4s \left[\sum_{k=1}^{\infty} F_1(s)^{2k} \right]' = 4s \left[\frac{F_1(s)^2}{1 - F_1(s)^2} \right]' \\ &= \frac{2s^2}{(1-s^2)^{\frac{3}{2}}} \end{aligned}$$

Random Walks

Proof.

(c):
$$\sum_{m=0}^{\infty} 4mp_0(2m)s^{2m}$$

Random Walks

Proof.

$$(c): \sum_{m=0}^{\infty} 4mp_0(2m)s^{2m} = 2s \sum_{m=0}^{\infty} p_0(2m)(2m)s^{2m-1}$$

Random Walks

Proof.

$$\begin{aligned} \text{(c): } \sum_{m=0}^{\infty} 4mp_0(2m)s^{2m} &= 2s \sum_{m=0}^{\infty} p_0(2m)(2m)s^{2m-1} \\ &= 2sP'_0(s) \\ &= \frac{2s^2}{(1-s^2)^{\frac{3}{2}}} \end{aligned}$$

Random Walks

Proof.


$$\begin{aligned} \text{(c): } \sum_{m=0}^{\infty} 4mp_0(2m)s^{2m} &= 2s \sum_{m=0}^{\infty} p_0(2m)(2m)s^{2m-1} \\ &= 2sP'_0(s) \\ &= \frac{2s^2}{(1-s^2)^{\frac{3}{2}}} \end{aligned}$$

We see that $\sum_{m=0}^{\infty} \mathbb{E}(T \wedge 2m)s^{2m} = \sum_{m=0}^{\infty} 2\mathbb{E}|S_{2m}|s^{2m} = \sum_{m=0}^{\infty} 4mp_0(2m)s^{2m}$, and it follows by uniqueness that $\mathbb{E}(T \wedge 2m) = 2\mathbb{E}|S_{2m}| = 4m\Pr[S_{2m} = 0]$.



Branching Processes

Generating function is a powerful tool for studying branching processes.



Branching Processes

Branching processes mainly focus on “reproduction”.

Branching Processes

Branching processes mainly focus on “reproduction”.

Let Z_n be the number of members of the n -th generation. Assume that $Z_0 = 1$.


Each member of the n -th generation will reproduce some children (may be none), which are its family and the members of the $(n + 1)$ -th generation.

Branching Processes

Branching processes mainly focus on “reproduction”.

Let Z_n be the number of members of the n -th generation. Assume that $Z_0 = 1$.

Each member of the n -th generation will reproduce some children (may be none), which are its family and the members of the $(n + 1)$ -th generation.




The size of a family is a random variable.

Branching Processes

Branching processes mainly focus on “reproduction”.

Let Z_n be the number of members of the n -th generation. Assume that $Z_0 = 1$.

Each member of the n -th generation will reproduce some children (may be none), which are its family and the members of the $(n + 1)$ -th generation.



The size of a family is a random variable.

We have the assumption:

the family size of each individual is identically independent distributed.

Let G be the generating function of “family size of a individual”.

And let $G_n(s) = \mathbb{E}(s^{Z_n})$ be the generating function of Z_n .

Branching Processes

As proved in textbook *Probability with Martingales*, section 0.3, using simple conditional expectation we can prove an important result for branching process:

Proposition. $G_{n+1}(s) = G_n(G(s))$, and thus $G_n(s) = G(G(\dots(G(s))\dots))$ is the n -fold iterate of G .

Branching Processes

As proved in textbook *Probability with Martingales*, section 0.3, using simple conditional expectation we can prove an important result for branching process:

Proposition. $G_{n+1}(s) = G_n(G(s))$, and thus $G_n(s) = G(G(\dots(G(s))\dots))$ is the n -fold iterate of G .

Using this we can prove:

Let $\mu := \mathbb{E}(Z_1)$, $\sigma^2 = \text{Var}(Z_1)$, then we have

$$\mathbb{E}(Z_n) = \mu^n, \quad \text{Var}(Z_n) = \begin{cases} n\sigma^2, & \text{if } \mu = 1, \\ \frac{\sigma^2(\mu^n - 1)\mu^{n-1}}{\mu - 1}, & \text{if } \mu \neq 1. \end{cases}$$

Branching Processes

Proof.

$$G_{n+1}(s) = G_n(G(s))$$

Branching Processes

Proof.

$$G_{n+1}(s) = G_n(G(s))$$

take differentiation,
let $s = 1$

$$\mathbb{E}(Z_n) = \mu \mathbb{E}(Z_{n-1})$$

Branching Processes

Proof.

$$\begin{array}{lcl} G_{n+1}(s) = G_n(G(s)) & \xrightarrow{\substack{\text{take two differentiations,} \\ \text{let } s = 1}} & G_n''(1) = G''(1)G_{n-1}'(1)^2 + G'(1)G_{n-1}''(1) \\ \downarrow \substack{\text{take differentiation,} \\ \text{let } s = 1} & & \downarrow \\ \mathbb{E}(Z_n) = \mu \mathbb{E}(Z_{n-1}) & & \text{Var}(Z_n) = G_n''(1) + G_n'(1) - G_n'(1)^2 \\ & & \text{(we mentioned it before)} \end{array}$$



Branching Processes

For branching process, an important topic is about the “ultimate extinction probability”, which is already discussed in textbook. So I don’t want to discuss more here.

Generating Functions

Not only in probability, generating functions have various applications in many other fields, more in combinatorics.

Generating Functions

Not only in probability, generating functions have various applications in many other fields, more in combinatorics.

We only discussed ordinary generating function there. However, there are other types of generating functions, such as *exponential generating functions*: $E_a(s) = \sum_{i=0}^{\infty} \frac{a_i s^i}{i!}$,

Poisson generating functions: $P_a(s) = \sum_{i=0}^{\infty} a_i e^{-s} \frac{s^i}{i!} = e^{-x} E_a(x)$, etc. They are all powerful math tools and all have widely usage.

Why Generating Functions?

Why Generating Functions?

At the beginning, I have no
idea....

Why Generating Functions?

At the beginning, I have no
idea....




Branching process? Seems good...

Why Generating Functions?

At the beginning, I have no
idea....

Branching process? Seems good...

The material says generating functions are a
good tool. Our textbook also mentioned
generating functions when dealing with
branching process.



Why Generating Functions?

At the beginning, I have no idea....

Branching process? Seems good...

The material says generating functions are a good tool. Our textbook also mentioned generating functions when dealing with branching process.

I knowed a little bit about generating functions before, in high school.

Why Generating Functions?

At the beginning, I have no idea....

Branching process? Seems good...

The material says generating functions are a good tool. Our textbook also mentioned generating functions when dealing with branching process.

I knowed a little bit about generating functions before, in high school.

Okay, generating functions. That's it!

My Contributions

- (1) Make this “dynamic slides” so that the contents are more understandable. And maybe it can be used in a lecture.
- (2) Fully understand what I make.
- (3) Solve two not-too-hard exercises in the book I referenced to.

References

- [1] Geoffrey R. Grimmett, David R. Stirzaker, *Probability and Random Processes*, 3rd Edition, Oxford University Press, 2001
- [2] Wikipedia Generating Functions
- [3] Wikipedia Probability Generating Functions
- [4] Hitting time theorem from University of Nebraska–Lincoln
- [5] Krishna B. Athreya, Peter E. Ney, *Branching Processes*, Springer-Verlag Berlin Heidelberg, 1972
- [6] David Williams, *Probability with Martingales*, Cambridge University Press, 1991

Thank you!