

Solution to Exercise 1 of March 13

金弘义 518030910333

March 17, 2020

Exercise 1. 1) Show that the Cantor Set C is nowhere dense in $[0, 1]$.

2) Find a meager set T in \mathbb{R} such that $T + T = \mathbb{R}$.

3) Show that every subset of the real line \mathbb{R} can be partitioned into two sets, one being of first category and the other being negligible.

Proof. 1) Recall the the explicit closed formulas for the Cantor set are:

$$C = [0, 1] \setminus \bigcup_{n=0}^{\infty} \bigcup_{k=0}^{3^n-1} \left(\frac{3k+1}{3^{n+1}}, \frac{3k+2}{3^{n+1}} \right)$$

It's obvious that $(\frac{3k+1}{3^{n+1}}, \frac{3k+2}{3^{n+1}})$ is an open set. Since the union of open sets is an open set, the complement of C is also open, which leads to the fact that C is a closed set. So $\bar{C} = C$.

Note that Cantor set can also be characterized as the set of all number in $[0, 1]$ whose base-3 expansion doesn't contain any 1s. Assume there exists an interior point x of C , then there exists a ball B , with radius $r > 0$, which is contained in C . Since $C \subseteq [0, 1]$, B is an interval, whose length is $2r$. Arbitrarily choose $x_1 \in B$ and $x_2 \in B$ such that $x_1 < x_2$ and $x_2 - x_1 \geq r$. Let $n = \lceil \log_{\frac{1}{3}} r \rceil + 2$. Consider $x_3 = x_1 + (\frac{1}{3})^n$, $x_4 = x_1 + 2 \cdot (\frac{1}{3})^n$. Since $(\frac{1}{3})^n < (\frac{1}{3})^{\log_{\frac{1}{3}} r + 1} = \frac{1}{3}r$, $x_3, x_4 \in B$. It's obvious that one of them contains 1 in its base-3 expansion, which means one of them doesn't belong to Cantor set. This leads to a contradiction, or equivalently, that there is no interior point of C .

Now we can safely conclude that C is nowhere dense.

2) Let

$$T(n) = \{a + n | a \in C\}$$

$$T = \bigcup_{n \in \mathbb{Z}} T(n)$$

First We will prove T is a meager set. Since $T(n)$ is constructed by translation of all points in C by n , $T(n)$ is also nowhere dense. This leads to the fact that T is a meager set, because \mathbb{Z} is countable.

Then we prove $T + T = \mathbb{R}$. From $C + C = [0, 2]$, we can similarly get $T(n) + T(n) = [2n, 2n+2]$. Note that:

$$\bigcup_{n \in \mathbb{Z}} (T(n) + T(n)) \subseteq T + T \subseteq \mathbb{R}$$

However,

$$\bigcup_{n \in \mathbb{Z}} (T(n) + T(n)) = \bigcup_{n \in \mathbb{Z}} [2n, 2n + 2] = \mathbb{R}$$

Now we conclude that $T + T = \mathbb{R}$.

3) Pick an enumeration of \mathbb{Q} , named q_n . Let:

$$\begin{aligned} I_{i,j} &= (q_i - \frac{1}{2^{i+j+2}}, q_i + \frac{1}{2^{i+j+2}}) \\ A_j &= \bigcup_{i \in \mathbb{N}} I_{i,j} \\ B &= \bigcap_{j \in \mathbb{N}} A_j \end{aligned}$$

For any $\epsilon > 0$, exist $j_0 > 0$ such that $|\frac{1}{2^{j_0}}| < \epsilon$. So we have :

$$\begin{aligned} |A_j| &\leq \sum_{i=0}^{\infty} \frac{1}{2^{i+j+1}} = \frac{1}{2^j} \\ |B| &\leq |A_{j_0}| \leq \frac{1}{2^{j_0}} < \epsilon \end{aligned}$$

This leads to that B is negligible. Then we prove B^C is a meager set.

$$B^C = \bigcup_{j \in \mathbb{N}} A_j^C$$

Since $\mathbb{Q} \subseteq A_j$ and \mathbb{Q} is dense, we can know that A_j is dense. A_j is also open because A_j is the union of open intervals. These 2 properties of A_j can lead to that A_j^C is nowhere dense. So B^C is a meager set, which finishes the proof. \square