

MA 322: Scientific Computing

Lecture - 1



Course Structure

Course Structure

- Unit-I

Course Structure

● Unit-I

- Numerical methods for solving scalar nonlinear equations.

Course Structure

● Unit-I

- Numerical methods for solving scalar nonlinear equations.

● Unit-II

Course Structure

● Unit-I

- Numerical methods for solving scalar nonlinear equations.

● Unit-II

- Interpolation: Polynomial approximation for continuous functions.

Course Structure

● Unit-I

- Numerical methods for solving scalar nonlinear equations.

● Unit-II

- Interpolation: Polynomial approximation for continuous functions.

● Unit-III

Course Structure

● Unit-I

- Numerical methods for solving scalar nonlinear equations.

● Unit-II

- Interpolation: Polynomial approximation for continuous functions.

● Unit-III

- Numerical Integration.

Course Structure

● Unit-I

- Numerical methods for solving scalar nonlinear equations.

● Unit-II

- Interpolation: Polynomial approximation for continuous functions.

● Unit-III

- Numerical Integration.

● Unit-IV

Course Structure

● Unit-I

- Numerical methods for solving scalar nonlinear equations.

● Unit-II

- Interpolation: Polynomial approximation for continuous functions.

● Unit-III

- Numerical Integration.

● Unit-IV

- Initial value problems for ordinary differential equations and its numerical approximations.

Course Structure

● Unit-I

- Numerical methods for solving scalar nonlinear equations.

● Unit-II

- Interpolation: Polynomial approximation for continuous functions.

● Unit-III

- Numerical Integration.

● Unit-IV

- Initial value problems for ordinary differential equations and its numerical approximations.

● Unit-V

Course Structure

● Unit-I

- Numerical methods for solving scalar nonlinear equations.

● Unit-II

- Interpolation: Polynomial approximation for continuous functions.

● Unit-III

- Numerical Integration.

● Unit-IV

- Initial value problems for ordinary differential equations and its numerical approximations.

● Unit-V

- Finite difference schemes for partial differential equations.

Course Structure

● Unit-I

- Numerical methods for solving scalar nonlinear equations.

● Unit-II

- Interpolation: Polynomial approximation for continuous functions.

● Unit-III

- Numerical Integration.

● Unit-IV

- Initial value problems for ordinary differential equations and its numerical approximations.

● Unit-V

- Finite difference schemes for partial differential equations.

Motivation

Motivation

- Why do we need Numerical Methods?

Motivation

- Why do we need Numerical Methods?
- Can we solve following problems exactly?

Motivation

- Why do we need Numerical Methods?
- Can we solve following problems exactly?
 - Problem-1: Evaluate $\sqrt{2}$ without using calculator?

Motivation

- Why do we need Numerical Methods?
- Can we solve following problems exactly?
 - Problem-1: Evaluate $\sqrt{2}$ without using calculator?
 - Problem-2: Without using calculator, can we find $\sin 3^\circ$ from the known data

Motivation

- Why do we need Numerical Methods?
- Can we solve following problems exactly?
 - Problem-1: Evaluate $\sqrt{2}$ without using calculator?
 - Problem-2: Without using calculator, can we find $\sin 3^\circ$ from the known data

θ	0°	30°	45°	60°	90°
$\sin \theta$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1

- Problem-3: Find the integration

$$I = \int_0^1 e^{x^2} dx.$$

Motivation

- Why do we need Numerical Methods?
- Can we solve following problems exactly?
 - Problem-1: Evaluate $\sqrt{2}$ without using calculator?
 - Problem-2: Without using calculator, can we find $\sin 3^\circ$ from the known data

θ	0°	30°	45°	60°	90°
$\sin \theta$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1

- Problem-3: Find the integration

$$I = \int_0^1 e^{x^2} dx.$$

- Problem-4: Find y such that

$$y' = x^2 + y^2, \quad y(0) = 1.$$

Motivation

- Why do we need Numerical Methods?
- Can we solve following problems exactly?
 - Problem-1: Evaluate $\sqrt{2}$ without using calculator?
 - Problem-2: Without using calculator, can we find $\sin 3^\circ$ from the known data

θ	0°	30°	45°	60°	90°
$\sin \theta$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1

- Problem-3: Find the integration

$$I = \int_0^1 e^{x^2} dx.$$

- Problem-4: Find y such that

$$y' = x^2 + y^2, \quad y(0) = 1.$$

More precisely, can we solve non-linear IVP exactly?

Motivation Contd....

Motivation Contd....

- Consider the first problem.

Motivation Contd....

- Consider the first problem.
 - Let $x = \sqrt{2}$,

Motivation Contd....

- Consider the first problem.
 - Let $x = \sqrt{2}$, so that $x^2 - 2 = 0$.

Motivation Contd....

- Consider the first problem.
 - Let $x = \sqrt{2}$, so that $x^2 - 2 = 0$. Then, by setting $f(x) = x^2 - 2$,

Motivation Contd....

- Consider the first problem.
 - Let $x = \sqrt{2}$, so that $x^2 - 2 = 0$. Then, by setting $f(x) = x^2 - 2$, we may conclude following:


Motivation Contd....

- Consider the first problem.
 - Let $x = \sqrt{2}$, so that $x^2 - 2 = 0$. Then, by setting $f(x) = x^2 - 2$, we may conclude following:
 - Thus, finding $\sqrt{2}$ is equivalent to find $c \in \mathbb{R}$ such that $f(c) = 0$.

Motivation Contd....

- Consider the first problem.


- Let $x = \sqrt{2}$, so that $x^2 - 2 = 0$. Then, by setting $f(x) = x^2 - 2$, we may conclude following:
- Thus, finding $\sqrt{2}$ is equivalent to find $c \in \mathbb{R}$ such that $f(c) = 0$.
- In general, c is a zero of a function $y = f(x)$, or root of the equation $f(x) = 0$ provided $f(c) = 0$.



Motivation Contd....

- Consider the first problem.

- Let $x = \sqrt{2}$, so that $x^2 - 2 = 0$. Then, by setting $f(x) = x^2 - 2$, we may conclude following:
- Thus, finding $\sqrt{2}$ is equivalent to find $c \in \mathbb{R}$ such that $f(c) = 0$.
- In general, c is a zero of a function $y = f(x)$, or root of the equation $f(x) = 0$ provided $f(c) = 0$.




- This leads to following question

Motivation Contd....

- Consider the first problem.

- Let $x = \sqrt{2}$, so that $x^2 - 2 = 0$. Then, by setting $f(x) = x^2 - 2$, we may conclude following:
- Thus, finding $\sqrt{2}$ is equivalent to find $c \in \mathbb{R}$ such that $f(c) = 0$.
- In general, c is a zero of a function $y = f(x)$, or root of the equation $f(x) = 0$ provided $f(c) = 0$.




- This leads to following question

"Under what assumptions a function has a zero?"

Motivation Contd....

- Consider the first problem.

- Let $x = \sqrt{2}$, so that $x^2 - 2 = 0$. Then, by setting $f(x) = x^2 - 2$, we may conclude following:
- Thus, finding $\sqrt{2}$ is equivalent to find $c \in \mathbb{R}$ such that $f(c) = 0$.
- In general, c is a zero of a function $y = f(x)$, or root of the equation $f(x) = 0$ provided $f(c) = 0$.



- This leads to following question


“Under what assumptions a function has a zero?”

This is answered by the properties of ‘Continuous Functions’ in the interval $[a, b] \subset \mathbb{R}$.

Motivation Contd....

- Consider the first problem.

- Let $x = \sqrt{2}$, so that $x^2 - 2 = 0$. Then, by setting $f(x) = x^2 - 2$, we may conclude following:
- Thus, finding $\sqrt{2}$ is equivalent to find $c \in \mathbb{R}$ such that $f(c) = 0$.
- In general, c is a zero of a function $y = f(x)$, or root of the equation $f(x) = 0$ provided $f(c) = 0$.



- This leads to following question

“Under what assumptions a function has a zero?”


This is answered by the properties of ‘Continuous Functions’ in the interval $[a, b] \subset \mathbb{R}$.

- The numerical methods for finding the roots are called *iterative methods*,

Motivation Contd....

- Consider the first problem.

- Let $x = \sqrt{2}$, so that $x^2 - 2 = 0$. Then, by setting $f(x) = x^2 - 2$, we may conclude following:
- Thus, finding $\sqrt{2}$ is equivalent to find $c \in \mathbb{R}$ such that $f(c) = 0$.
- In general, c is a zero of a function $y = f(x)$, or root of the equation $f(x) = 0$ provided $f(c) = 0$.



- This leads to following question

“Under what assumptions a function has a zero?”

This is answered by the properties of ‘Continuous Functions’ in the interval $[a, b] \subset \mathbb{R}$.

- The numerical methods for finding the roots are called *iterative methods*, and they are the main subject of next few lectures.

Properties of Continuous Functions Contd..

Properties of Continuous Functions Contd..

Location of Roots Theorem:

Properties of Continuous Functions Contd..

Location of Roots Theorem: Let $I = [a, b]$ and let $f : I \rightarrow \mathbb{R}$ be a continuous on I .

Properties of Continuous Functions Contd..

Location of Roots Theorem: Let $I = [a, b]$ and let $f : I \rightarrow \mathbb{R}$ be a continuous on I . If $f(a) < 0 < f(b)$ or if $f(a) > 0 > f(b)$, then there exists a number $c \in (a, b)$ such that $f(c) = 0$.

Properties of Continuous Functions Contd..

Location of Roots Theorem: Let $I = [a, b]$ and let $f : I \rightarrow \mathbb{R}$ be a continuous on I . If $f(a) < 0 < f(b)$ or if $f(a) > 0 > f(b)$, then there exists a number $c \in (a, b)$ such that $f(c) = 0$.

● **Remark:**

Properties of Continuous Functions Contd..

Location of Roots Theorem: Let $I = [a, b]$ and let $f : I \rightarrow \mathbb{R}$ be a continuous on I . If $f(a) < 0 < f(b)$ or if $f(a) > 0 > f(b)$, then there exists a number $c \in (a, b)$ such that $f(c) = 0$.

● **Remark:** $f(c) = 0$ means c is a root of the function $f(x)$.

Properties of Continuous Functions Contd..

Location of Roots Theorem: Let $I = [a, b]$ and let $f : I \rightarrow \mathbb{R}$ be a continuous on I . If $f(a) < 0 < f(b)$ or if $f(a) > 0 > f(b)$, then there exists a number $c \in (a, b)$ such that $f(c) = 0$.

● **Remark:** $f(c) = 0$ means c is a root of the function $f(x)$.

Consider the function $f(x) = x^2 - 2$.

Properties of Continuous Functions Contd..

Location of Roots Theorem: Let $I = [a, b]$ and let $f : I \rightarrow \mathbb{R}$ be a continuous on I . If $f(a) < 0 < f(b)$ or if $f(a) > 0 > f(b)$, then there exists a number $c \in (a, b)$ such that $f(c) = 0$.

● **Remark:** $f(c) = 0$ means c is a root of the function $f(x)$.

Consider the function $f(x) = x^2 - 2$.

Clearly $f(1) = -1 < 0$

Properties of Continuous Functions Contd..

Location of Roots Theorem: Let $I = [a, b]$ and let $f : I \rightarrow \mathbb{R}$ be a continuous on I . If $f(a) < 0 < f(b)$ or if $f(a) > 0 > f(b)$, then there exists a number $c \in (a, b)$ such that $f(c) = 0$.

● **Remark:** $f(c) = 0$ means c is a root of the function $f(x)$.

Consider the function $f(x) = x^2 - 2$.

Clearly $f(1) = -1 < 0$ and $f(2) = 2 > 0$.

Properties of Continuous Functions Contd..

Location of Roots Theorem: Let $I = [a, b]$ and let $f : I \rightarrow \mathbb{R}$ be a continuous on I . If $f(a) < 0 < f(b)$ or if $f(a) > 0 > f(b)$, then there exists a number $c \in (a, b)$ such that $f(c) = 0$.

● **Remark:** $f(c) = 0$ means c is a root of the function $f(x)$.

Consider the function $f(x) = x^2 - 2$.

Clearly $f(1) = -1 < 0$ and $f(2) = 2 > 0$.

Therefore, $f(x) = x^2 - 2$ has a root in the interval $(1, 2)$.

Properties of Continuous Functions Contd..

Location of Roots Theorem: Let $I = [a, b]$ and let $f : I \rightarrow \mathbb{R}$ be a continuous on I . If $f(a) < 0 < f(b)$ or if $f(a) > 0 > f(b)$, then there exists a number $c \in (a, b)$ such that $f(c) = 0$.

● **Remark:** $f(c) = 0$ means c is a root of the function $f(x)$.

Consider the function $f(x) = x^2 - 2$.

Clearly $f(1) = -1 < 0$ and $f(2) = 2 > 0$.

Therefore, $f(x) = x^2 - 2$ has a root in the interval $(1, 2)$. That is there exists a $c \in (1, 2)$ such that $f(c) = 0$ or $c^2 - 2 = 0$.

Properties of Continuous Functions Contd..

Location of Roots Theorem: Let $I = [a, b]$ and let $f : I \rightarrow \mathbb{R}$ be a continuous on I . If $f(a) < 0 < f(b)$ or if $f(a) > 0 > f(b)$, then there exists a number $c \in (a, b)$ such that $f(c) = 0$.

● **Remark:** $f(c) = 0$ means c is a root of the function $f(x)$.

Consider the function $f(x) = x^2 - 2$.

Clearly $f(1) = -1 < 0$ and $f(2) = 2 > 0$.


Therefore, $f(x) = x^2 - 2$ has a root in the interval $(1, 2)$. That is there exists a $c \in (1, 2)$ such that $f(c) = 0$ or $c^2 - 2 = 0$. This means there exists a $c \in (1, 2)$ such that $c = \sqrt{2}$.

Properties of Continuous Functions Contd..

Location of Roots Theorem: Let $I = [a, b]$ and let $f : I \rightarrow \mathbb{R}$ be a continuous on I . If $f(a) < 0 < f(b)$ or if $f(a) > 0 > f(b)$, then there exists a number $c \in (a, b)$ such that $f(c) = 0$.

● **Remark:** $f(c) = 0$ means c is a root of the function $f(x)$.

Consider the function $f(x) = x^2 - 2$. Clearly $f(1) = -1 < 0$ and $f(2) = 2 > 0$. Therefore, $f(x) = x^2 - 2$ has a root in the interval $(1, 2)$. That is there exists a $c \in (1, 2)$ such that $f(c) = 0$ or $c^2 - 2 = 0$. This means there exists a $c \in (1, 2)$ such that $c = \sqrt{2}$.




Properties of Continuous Functions Contd..

Location of Roots Theorem: Let $I = [a, b]$ and let $f : I \rightarrow \mathbb{R}$ be a continuous on I . If $f(a) < 0 < f(b)$ or if $f(a) > 0 > f(b)$, then there exists a number $c \in (a, b)$ such that $f(c) = 0$.

● **Remark:** $f(c) = 0$ means c is a root of the function $f(x)$.

Consider the function $f(x) = x^2 - 2$. Clearly $f(1) = -1 < 0$ and $f(2) = 2 > 0$. Therefore, $f(x) = x^2 - 2$ has a root in the interval $(1, 2)$. That is there exists a $c \in (1, 2)$ such that $f(c) = 0$ or $c^2 - 2 = 0$. This means there exists a $c \in (1, 2)$ such that $c = \sqrt{2}$.




● **Next question arises** 'How do you locate c in $(1, 2)$?'

Properties of Continuous Functions Contd..

Location of Roots Theorem: Let $I = [a, b]$ and let $f : I \rightarrow \mathbb{R}$ be a continuous on I . If $f(a) < 0 < f(b)$ or if $f(a) > 0 > f(b)$, then there exists a number $c \in (a, b)$ such that $f(c) = 0$.

● **Remark:** $f(c) = 0$ means c is a root of the function $f(x)$.

Consider the function $f(x) = x^2 - 2$. Clearly $f(1) = -1 < 0$ and $f(2) = 2 > 0$. Therefore, $f(x) = x^2 - 2$ has a root in the interval $(1, 2)$. That is there exists a $c \in (1, 2)$ such that $f(c) = 0$ or $c^2 - 2 = 0$. This means there exists a $c \in (1, 2)$ such that $c = \sqrt{2}$.




● Next question arises 'How do you locate c in $(1, 2)$?' To locate the root, say c , of a non-linear equation $f(x) = 0$, we follow a very simple iterative method, known as the bisection method,

Properties of Continuous Functions Contd..

Location of Roots Theorem: Let $I = [a, b]$ and let $f : I \rightarrow \mathbb{R}$ be a continuous on I . If $f(a) < 0 < f(b)$ or if $f(a) > 0 > f(b)$, then there exists a number $c \in (a, b)$ such that $f(c) = 0$.

● **Remark:** $f(c) = 0$ means c is a root of the function $f(x)$.

Consider the function $f(x) = x^2 - 2$. Clearly $f(1) = -1 < 0$ and $f(2) = 2 > 0$. Therefore, $f(x) = x^2 - 2$ has a root in the interval $(1, 2)$. That is there exists a $c \in (1, 2)$ such that $f(c) = 0$ or $c^2 - 2 = 0$. This means there exists a $c \in (1, 2)$ such that $c = \sqrt{2}$.



● Next question arises 'How do you locate c in $(1, 2)$?' To locate the root, say c , of a non-linear equation $f(x) = 0$, we follow a very simple iterative method, known as the bisection method, which is based on the following fact: If f is continuous at c and $c_n \rightarrow c$, then

$$f(c_n) \rightarrow f(c) = 0.$$

The Bisection Method

The Bisection Method

Algorithm:

The Bisection Method

Algorithm:

- Fix a and b such that $f(a) \cdot f(b) < 0$.

The Bisection Method

Algorithm:

- Fix a and b such that $f(a) \cdot f(b) < 0$. Set $a_1 = a$ and $b_1 = b$.

The Bisection Method

Algorithm:

- Fix a and b such that $f(a) \cdot f(b) < 0$. Set $a_1 = a$ and $b_1 = b$.
Define $c_1 = \frac{1}{2}(a_1 + b_1)$.

The Bisection Method

Algorithm:

- Fix a and b such that $f(a) \cdot f(b) < 0$. Set $a_1 = a$ and $b_1 = b$.
Define $c_1 = \frac{1}{2}(a_1 + b_1)$. Check $f(a_1) \cdot f(c_1)$ and $f(c_1) \cdot f(b_1)$.

The Bisection Method

Algorithm:

- Fix a and b such that $f(a) \cdot f(b) < 0$. Set $a_1 = a$ and $b_1 = b$.
Define $c_1 = \frac{1}{2}(a_1 + b_1)$. Check $f(a_1) \cdot f(c_1)$ and $f(c_1) \cdot f(b_1)$. Define

$$(a_2, b_2) = \begin{cases} (a_1, c_1), & \text{if } f(a_1) \cdot f(c_1) < 0, \\ (c_1, b_1), & \text{if } f(c_1) \cdot f(b_1) < 0. \end{cases}$$

The Bisection Method

Algorithm:

- Fix a and b such that $f(a) \cdot f(b) < 0$. Set $a_1 = a$ and $b_1 = b$.
Define $c_1 = \frac{1}{2}(a_1 + b_1)$. Check $f(a_1) \cdot f(c_1)$ and $f(c_1) \cdot f(b_1)$. Define

$$(a_2, b_2) = \begin{cases} (a_1, c_1), & \text{if } f(a_1) \cdot f(c_1) < 0, \\ (c_1, b_1), & \text{if } f(c_1) \cdot f(b_1) < 0. \end{cases}$$

Observe that $f(a_2) \cdot f(b_2) < 0$, so that root $c \in [a_2, b_2]$.

The Bisection Method

Algorithm:

- Fix a and b such that $f(a) \cdot f(b) < 0$. Set $a_1 = a$ and $b_1 = b$.
Define $c_1 = \frac{1}{2}(a_1 + b_1)$. Check $f(a_1) \cdot f(c_1)$ and $f(c_1) \cdot f(b_1)$. Define

$$(a_2, b_2) = \begin{cases} (a_1, c_1), & \text{if } f(a_1) \cdot f(c_1) < 0, \\ (c_1, b_1), & \text{if } f(c_1) \cdot f(b_1) < 0. \end{cases}$$

Observe that $f(a_2) \cdot f(b_2) < 0$, so that root $c \in [a_2, b_2]$. Define $c_2 = \frac{1}{2}(a_2 + b_2)$.

The Bisection Method

Algorithm:

- Fix a and b such that $f(a) \cdot f(b) < 0$. Set $a_1 = a$ and $b_1 = b$.
Define $c_1 = \frac{1}{2}(a_1 + b_1)$. Check $f(a_1) \cdot f(c_1)$ and $f(c_1) \cdot f(b_1)$. Define

$$(a_2, b_2) = \begin{cases} (a_1, c_1), & \text{if } f(a_1) \cdot f(c_1) < 0, \\ (c_1, b_1), & \text{if } f(c_1) \cdot f(b_1) < 0. \end{cases}$$

Observe that $f(a_2) \cdot f(b_2) < 0$, so that root $c \in [a_2, b_2]$. Define $c_2 = \frac{1}{2}(a_2 + b_2)$. If $f(c_2) = 0$, then stop.

The Bisection Method

Algorithm:

- Fix a and b such that $f(a) \cdot f(b) < 0$. Set $a_1 = a$ and $b_1 = b$.
Define $c_1 = \frac{1}{2}(a_1 + b_1)$. Check $f(a_1) \cdot f(c_1)$ and $f(c_1) \cdot f(b_1)$. Define

$$(a_2, b_2) = \begin{cases} (a_1, c_1), & \text{if } f(a_1) \cdot f(c_1) < 0, \\ (c_1, b_1), & \text{if } f(c_1) \cdot f(b_1) < 0. \end{cases}$$

Observe that $f(a_2) \cdot f(b_2) < 0$, so that root $c \in [a_2, b_2]$. Define $c_2 = \frac{1}{2}(a_2 + b_2)$. If $f(c_2) = 0$, then stop.

- In general, for $n \geq 1$, we define (a_{n+1}, b_{n+1}) by

$$(a_{n+1}, b_{n+1}) = \begin{cases} (a_n, c_n), & \text{if } f(a_n) \cdot f(c_n) < 0, \\ (c_n, b_n), & \text{if } f(c_n) \cdot f(b_n) < 0. \end{cases}$$

The Bisection Method

Algorithm:

- Fix a and b such that $f(a) \cdot f(b) < 0$. Set $a_1 = a$ and $b_1 = b$. Define $c_1 = \frac{1}{2}(a_1 + b_1)$. Check $f(a_1) \cdot f(c_1)$ and $f(c_1) \cdot f(b_1)$. Define

$$(a_2, b_2) = \begin{cases} (a_1, c_1), & \text{if } f(a_1) \cdot f(c_1) < 0, \\ (c_1, b_1), & \text{if } f(c_1) \cdot f(b_1) < 0. \end{cases}$$

Observe that $f(a_2) \cdot f(b_2) < 0$, so that root $c \in [a_2, b_2]$. Define $c_2 = \frac{1}{2}(a_2 + b_2)$. If $f(c_2) = 0$, then stop.

- In general, for $n \geq 1$, we define (a_{n+1}, b_{n+1}) by

$$(a_{n+1}, b_{n+1}) = \begin{cases} (a_n, c_n), & \text{if } f(a_n) \cdot f(c_n) < 0, \\ (c_n, b_n), & \text{if } f(c_n) \cdot f(b_n) < 0. \end{cases}$$

Thus, we can construct sequences $\langle a_{n+1} \rangle$ and $\langle b_{n+1} \rangle$ such that $f(a_{n+1}) \cdot f(b_{n+1}) < 0$, so that root $c \in [a_{n+1}, b_{n+1}]$.

The Bisection Method

Algorithm:

- Fix a and b such that $f(a) \cdot f(b) < 0$. Set $a_1 = a$ and $b_1 = b$. Define $c_1 = \frac{1}{2}(a_1 + b_1)$. Check $f(a_1) \cdot f(c_1)$ and $f(c_1) \cdot f(b_1)$. Define

$$(a_2, b_2) = \begin{cases} (a_1, c_1), & \text{if } f(a_1) \cdot f(c_1) < 0, \\ (c_1, b_1), & \text{if } f(c_1) \cdot f(b_1) < 0. \end{cases}$$

Observe that $f(a_2) \cdot f(b_2) < 0$, so that root $c \in [a_2, b_2]$. Define $c_2 = \frac{1}{2}(a_2 + b_2)$. If $f(c_2) = 0$, then stop.

- In general, for $n \geq 1$, we define (a_{n+1}, b_{n+1}) by

$$(a_{n+1}, b_{n+1}) = \begin{cases} (a_n, c_n), & \text{if } f(a_n) \cdot f(c_n) < 0, \\ (c_n, b_n), & \text{if } f(c_n) \cdot f(b_n) < 0. \end{cases}$$

Thus, we can construct sequences $\langle a_{n+1} \rangle$ and $\langle b_{n+1} \rangle$ such that $f(a_{n+1}) \cdot f(b_{n+1}) < 0$, so that root $c \in [a_{n+1}, b_{n+1}]$.

The Bisection Method

Algorithm:

- Fix a and b such that $f(a) \cdot f(b) < 0$. Set $a_1 = a$ and $b_1 = b$. Define $c_1 = \frac{1}{2}(a_1 + b_1)$. Check $f(a_1) \cdot f(c_1)$ and $f(c_1) \cdot f(b_1)$. Define

$$(a_2, b_2) = \begin{cases} (a_1, c_1), & \text{if } f(a_1) \cdot f(c_1) < 0, \\ (c_1, b_1), & \text{if } f(c_1) \cdot f(b_1) < 0. \end{cases}$$

Observe that $f(a_2) \cdot f(b_2) < 0$, so that root $c \in [a_2, b_2]$. Define $c_2 = \frac{1}{2}(a_2 + b_2)$. If $f(c_2) = 0$, then stop.

- In general, for $n \geq 1$, we define (a_{n+1}, b_{n+1}) by

$$(a_{n+1}, b_{n+1}) = \begin{cases} (a_n, c_n), & \text{if } f(a_n) \cdot f(c_n) < 0, \\ (c_n, b_n), & \text{if } f(c_n) \cdot f(b_n) < 0. \end{cases}$$

Thus, we can construct sequences $\langle a_{n+1} \rangle$ and $\langle b_{n+1} \rangle$ such that $f(a_{n+1}) \cdot f(b_{n+1}) < 0$, so that root $c \in [a_{n+1}, b_{n+1}]$. Define

$$c_{n+1} = \frac{1}{2}(a_{n+1} + b_{n+1}).$$

The Bisection Method

Algorithm:

- Fix a and b such that $f(a) \cdot f(b) < 0$. Set $a_1 = a$ and $b_1 = b$. Define $c_1 = \frac{1}{2}(a_1 + b_1)$. Check $f(a_1) \cdot f(c_1)$ and $f(c_1) \cdot f(b_1)$. Define

$$(a_2, b_2) = \begin{cases} (a_1, c_1), & \text{if } f(a_1) \cdot f(c_1) < 0, \\ (c_1, b_1), & \text{if } f(c_1) \cdot f(b_1) < 0. \end{cases}$$

Observe that $f(a_2) \cdot f(b_2) < 0$, so that root $c \in [a_2, b_2]$. Define $c_2 = \frac{1}{2}(a_2 + b_2)$. If $f(c_2) = 0$, then stop.

- In general, for $n \geq 1$, we define (a_{n+1}, b_{n+1}) by

$$(a_{n+1}, b_{n+1}) = \begin{cases} (a_n, c_n), & \text{if } f(a_n) \cdot f(c_n) < 0, \\ (c_n, b_n), & \text{if } f(c_n) \cdot f(b_n) < 0. \end{cases}$$


Thus, we can construct sequences $\langle a_{n+1} \rangle$ and $\langle b_{n+1} \rangle$ such that $f(a_{n+1}) \cdot f(b_{n+1}) < 0$, so that root $c \in [a_{n+1}, b_{n+1}]$. Define

$$c_{n+1} = \frac{1}{2}(a_{n+1} + b_{n+1}).$$


If $f(c_{n+1}) = 0$, then stop, else repeat this procedure.

Bisection Method Contd.....

Bisection Method Contd.....




Bisection Method Contd.....



Observations:


Bisection Method Contd.....



Observations: First observe that

$$b_2 - a_2 = \frac{1}{2}(b_1 - a_1),$$

Bisection Method Contd.....




Observations: First observe that

$$b_2 - a_2 = \frac{1}{2}(b_1 - a_1),$$

$$b_3 - a_3 = \frac{1}{2}(b_2 - a_2) =$$

Bisection Method Contd.....




Observations: First observe that

$$b_2 - a_2 = \frac{1}{2}(b_1 - a_1),$$

$$b_3 - a_3 = \frac{1}{2}(b_2 - c_2) = \frac{1}{2}\left(b_2 - \frac{1}{2}(a_2 + b_2)\right) =$$

Bisection Method Contd.....




Observations: First observe that

$$b_2 - a_2 = \frac{1}{2}(b_1 - a_1),$$

$$\begin{aligned} b_3 - a_3 &= \frac{1}{2}(b_2 - c_2) = \frac{1}{2}\left(b_2 - \frac{1}{2}(a_2 + b_2)\right) = \left(\frac{1}{2}\right)^2(b_2 - a_2) \\ &= \left(\frac{1}{2}\right)^3(b_1 - a_1) = \left(\frac{1}{2}\right)^3(b - a), \end{aligned}$$

Bisection Method Contd.....



Observations: First observe that

$$b_2 - a_2 = \frac{1}{2}(b_1 - a_1),$$

$$\begin{aligned} b_3 - a_3 &= \frac{1}{2}(b_2 - c_2) = \frac{1}{2}\left(b_2 - \frac{1}{2}(a_2 + b_2)\right) = \left(\frac{1}{2}\right)^2(b_2 - a_2) \\ &= \left(\frac{1}{2}\right)^3(b_1 - a_1) = \left(\frac{1}{2}\right)^3(b - a), \end{aligned}$$

which implies

$$|c_3 - c| \leq |b_3 - a_3| \leq \left(\frac{1}{2}\right)^3(b - a).$$

Bisection Method Contd.....

Bisection Method Contd.....

- In general, we have

$$|c - c_n| \leq \left(\frac{1}{2}\right)^n (b - a).$$

Bisection Method Contd.....

- In general, we have

$$|c - c_n| \leq \left(\frac{1}{2}\right)^n (b - a).$$

Hence,

$$c_n \rightarrow c \text{ as } n \rightarrow \infty$$

Bisection Method Contd.....

- In general, we have

$$|c - c_n| \leq \left(\frac{1}{2}\right)^n (b - a).$$

Hence,

$$c_n \rightarrow c \text{ as } n \rightarrow \infty \quad f(c_n) \rightarrow f(c) = 0.$$

Bisection Method Contd.....

- In general, we have

$$|c - c_n| \leq \left(\frac{1}{2}\right)^n (b - a).$$

Hence,

$$c_n \rightarrow c \text{ as } n \rightarrow \infty \quad f(c_n) \rightarrow f(c) = 0.$$

- Therefore, the sequence $\langle c_n \rangle$ gives approximation to the exact root c .

Bisection Method Contd.....

- In general, we have

$$|c - c_n| \leq \left(\frac{1}{2}\right)^n (b - a).$$

Hence,

$$c_n \rightarrow c \text{ as } n \rightarrow \infty \quad f(c_n) \rightarrow f(c) = 0.$$

- Therefore, the sequence $\langle c_n \rangle$ gives approximation to the exact root c .

Example: Find an approximation for the largest root $\alpha \approx 1.1347241$ of the equation

$$x^6 - x - 1 = 0.$$

Solution:

Bisection Method Contd.....

- In general, we have

$$|c - c_n| \leq \left(\frac{1}{2}\right)^n (b - a).$$

Hence,

$$c_n \rightarrow c \text{ as } n \rightarrow \infty \quad f(c_n) \rightarrow f(c) = 0.$$

- Therefore, the sequence $\langle c_n \rangle$ gives approximation to the exact root c .

Example: Find an approximation for the largest root $\alpha \approx 1.1347241$ of the equation

$$x^6 - x - 1 = 0.$$

Solution: It is straightforward to show that $1 < c < 2$.

Bisection Method Contd.....

- In general, we have

$$|c - c_n| \leq \left(\frac{1}{2}\right)^n (b - a).$$

Hence,

$$c_n \rightarrow c \text{ as } n \rightarrow \infty \quad f(c_n) \rightarrow f(c) = 0.$$

- Therefore, the sequence $\langle c_n \rangle$ gives approximation to the exact root c .

Example: Find an approximation for the largest root $\alpha \approx 1.1347241$ of the equation

$$x^6 - x - 1 = 0.$$

Solution: It is straightforward to show that $1 < c < 2$. Therefore, there exists at least one root $c \in [1, 2]$.

Bisection Method Contd.....

- In general, we have

$$|c - c_n| \leq \left(\frac{1}{2}\right)^n (b - a).$$

Hence,

$$c_n \rightarrow c \text{ as } n \rightarrow \infty \quad f(c_n) \rightarrow f(c) = 0.$$

- Therefore, the sequence $\langle c_n \rangle$ gives approximation to the exact root c .

Example: Find an approximation for the largest root $\alpha \approx 1.1347241$ of the equation

$$x^6 - x - 1 = 0.$$

Solution: It is straightforward to show that $1 < c < 2$. Therefore, there exists at least one root $c \in [1, 2]$. The approximation of the actual root c is given by

$$c_{n+1} = \frac{1}{2}(a_{n+1} + b_{n+1}), \quad a_1 = 1 \text{ \& } b_1 = 2,$$

where

$$(a_{n+1}, b_{n+1}) = \begin{cases} (a_n, c_n), & \text{if } f(a_n) \cdot f(c_n) < 0, \\ (c_n, b_n), & \text{if } f(c_n) \cdot f(b_n) < 0. \end{cases}$$

Bisection Method Contd.....

Bisection Method Contd.....

Example of bisection method

n	c_n	n	c_n
	$2.0 = b$	8	1.13672
	$1.0 = a$	9	1.13477
1	1.5	10	1.13379
2	1.25	11	1.13428
3	1.125	12	1.13452
4	1.1875	13	1.13464
5	1.15625	14	1.13470
6	1.14063	15	1.13474
7	1.13281		

(Source: An Introduction to Numerical Analysis by K. E. Atkinson)

Bisection Method Contd.....

Example of bisection method

n	c_n	n	c_n
	$2.0 = b$	8	1.13672
	$1.0 = a$	9	1.13477
1	1.5	10	1.13379
2	1.25	11	1.13428
3	1.125	12	1.13452
4	1.1875	13	1.13464
5	1.15625	14	1.13470
6	1.14063	15	1.13474
7	1.13281		

(Source: An Introduction to Numerical Analysis by K. E. Atkinson)

● Next question arises “What is the guarantee that the calculated sequence $\langle c_n \rangle$ converges to actual root c with desired accuracy?”

Bisection Method Contd.....

Example of bisection method

n	c_n	n	c_n
	$2.0 = b$	8	1.13672
	$1.0 = a$	9	1.13477
1	1.5	10	1.13379
2	1.25	11	1.13428
3	1.125	12	1.13452
4	1.1875	13	1.13464
5	1.15625	14	1.13470
6	1.14063	15	1.13474
7	1.13281		

(Source: An Introduction to Numerical Analysis by K. E. Atkinson)

- Next question arises “What is the guarantee that the calculated sequence $\langle c_n \rangle$ converges to actual root c with desired accuracy?”
- We call $c_n \rightarrow c$ if for given $\epsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that

$$|c_n - c| < \epsilon \quad \forall n \geq n_0. \quad (1)$$

Bisection Method Contd.....

Example of bisection method

n	c_n	n	c_n
	$2.0 = b$	8	1.13672
	$1.0 = a$	9	1.13477
1	1.5	10	1.13379
2	1.25	11	1.13428
3	1.125	12	1.13452
4	1.1875	13	1.13464
5	1.15625	14	1.13470
6	1.14063	15	1.13474
7	1.13281		

(Source: An Introduction to Numerical Analysis by K. E. Atkinson)

- Next question arises “What is the guarantee that the calculated sequence $\langle c_n \rangle$ converges to actual root c with desired accuracy?”
 - We call $c_n \rightarrow c$ if for given $\epsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that

$$|c_n - c| < \epsilon \quad \forall n \geq n_0. \quad (1)$$

In scientific computing, we call above condition as ‘Stopping Criterion’

Bisection Method Contd.....

Example of bisection method

n	c_n	n	c_n
	$2.0 = b$	8	1.13672
	$1.0 = a$	9	1.13477
1	1.5	10	1.13379
2	1.25	11	1.13428
3	1.125	12	1.13452
4	1.1875	13	1.13464
5	1.15625	14	1.13470
6	1.14063	15	1.13474
7	1.13281		

(Source: An Introduction to Numerical Analysis by K. E. Atkinson)

- Next question arises “What is the guarantee that the calculated sequence $\langle c_n \rangle$ converges to actual root c with desired accuracy?”
 - We call $c_n \rightarrow c$ if for given $\epsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that

$$|c_n - c| < \epsilon \quad \forall n \geq n_0. \quad (1)$$

In scientific computing, we call above condition as ‘Stopping Criterion’ and $\epsilon > 0$ is known as a ‘tolerance’.

Stopping Criterion for Bisection Method

Stopping Criterion for Bisection Method

- For the Bisection method, let us discuss the 'Stopping Criterion'.

Stopping Criterion for Bisection Method

- For the Bisection method, let us discuss the 'Stopping Criterion'.
- First, recall following error relation

$$|c_n - c| \leq \left(\frac{1}{2}\right)^n (b - a). \quad (2)$$

Stopping Criterion for Bisection Method

- For the Bisection method, let us discuss the 'Stopping Criterion'.
- First, recall following error relation

$$|c_n - c| \leq \left(\frac{1}{2}\right)^n (b - a). \quad (2)$$

Now, for given 'tolerance' $\epsilon > 0$, the error

$$|c_n - c| < \epsilon \text{ if we set } \left(\frac{1}{2}\right)^n (b - a) < \epsilon. \quad (3)$$

Stopping Criterion for Bisection Method

● For the Bisection method, let us discuss the 'Stopping Criterion'.

- First, recall following error relation

$$|c_n - c| \leq \left(\frac{1}{2}\right)^n (b - a). \quad (2)$$

Now, for given 'tolerance' $\epsilon > 0$, the error

$$|c_n - c| < \epsilon \text{ if we set } \left(\frac{1}{2}\right)^n (b - a) < \epsilon. \quad (3)$$

- Therefore, for $|c_n - c| < \epsilon$, it suffices to take

$$n \geq \frac{\log(b - a) - \log \epsilon}{\log 2}. \quad (4)$$

Stopping Criterion for Bisection Method

● For the Bisection method, let us discuss the 'Stopping Criterion'.

- First, recall following error relation

$$|c_n - c| \leq \left(\frac{1}{2}\right)^n (b - a). \quad (2)$$

Now, for given 'tolerance' $\epsilon > 0$, the error

$$|c_n - c| < \epsilon \text{ if we set } \left(\frac{1}{2}\right)^n (b - a) < \epsilon. \quad (3)$$

- Therefore, for $|c_n - c| < \epsilon$, it suffices to take

$$n \geq \frac{\log(b - a) - \log \epsilon}{\log 2}. \quad (4)$$

● What is the advantage of such boundes?

Stopping Criterion for Bisection Method

● For the Bisection method, let us discuss the 'Stopping Criterion'.

- First, recall following error relation

$$|c_n - c| \leq \left(\frac{1}{2}\right)^n (b - a). \quad (2)$$

Now, for given 'tolerance' $\epsilon > 0$, the error

$$|c_n - c| < \epsilon \text{ if we set } \left(\frac{1}{2}\right)^n (b - a) < \epsilon. \quad (3)$$

- Therefore, for $|c_n - c| < \epsilon$, it suffices to take

$$n \geq \frac{\log(b - a) - \log \epsilon}{\log 2}. \quad (4)$$

● What is the advantage of such boundes?

- Before calculation, for a given tolerance ϵ , we know how many steps we need to take to achieve a desired accuracy.

Algorithm for Bisection Method

Algorithm for Bisection Method

- Algorithm:

Algorithm for Bisection Method

- Algorithm:

Step-1: Input a , b & ϵ .

Algorithm for Bisection Method

- Algorithm:

Step-1: Input a , b & ϵ .

Step-2: Set n such that

$$n \geq \frac{\log(b - a) - \log \epsilon}{\log 2}.$$

Algorithm for Bisection Method

- Algorithm:

Step-1: Input a , b & ϵ .

Step-2: Set n such that

$$n \geq \frac{\log(b - a) - \log \epsilon}{\log 2}.$$

Step-3: Define

$$c = \frac{a + b}{2}.$$

Algorithm for Bisection Method

- Algorithm:

Step-1: Input a , b & ϵ .

Step-2: Set n such that

$$n \geq \frac{\log(b - a) - \log \epsilon}{\log 2}.$$

Step-3: Define

$$c = \frac{a + b}{2}.$$

Step-4: If $\text{sign}(f(b)) \cdot \text{sign}(f(c)) \leq 0$, then define $a = c$, otherwise set $b = c$.

Algorithm for Bisection Method

- Algorithm:

Step-1: Input a , b & ϵ .

Step-2: Set n such that

$$n \geq \frac{\log(b - a) - \log \epsilon}{\log 2}.$$

Step-3: Define

$$c = \frac{a + b}{2}.$$

Step-4: If $\text{sign}(f(b)) \cdot \text{sign}(f(c)) \leq 0$, then define $a = c$, otherwise set $b = c$.

Step-5: Return to Step-3 and repeat the iteration for n steps.

Homework: Consider the function $f(x) = x^3 - 2$ with initial interval $[0, 2]$.

Algorithm for Bisection Method

- Algorithm:

Step-1: Input a , b & ϵ .

Step-2: Set n such that

$$n \geq \frac{\log(b - a) - \log \epsilon}{\log 2}.$$

Step-3: Define

$$c = \frac{a + b}{2}.$$

Step-4: If $\text{sign}(f(b)) \cdot \text{sign}(f(c)) \leq 0$, then define $a = c$, otherwise set $b = c$.

Step-5: Return to Step-3 and repeat the iteration for n steps.

Homework: Consider the function $f(x) = x^3 - 2$ with initial interval $[0, 2]$. Determine the number of steps necessary for the root of $f(x) = 0$ to be accurate to within 10^{-6} .

Order of Convergence

Order of Convergence

Definition: A sequence of iterates $\langle x_n \rangle$ is said to converge with order $p \geq 1$ to the actual root α if

$$|\alpha - x_{n+1}| \leq C|\alpha - x_n|^p \quad n \geq 0 \quad (5)$$

for some $C > 0$.

Order of Convergence

Definition: A sequence of iterates $\langle x_n \rangle$ is said to converge with order $p \geq 1$ to the actual root α if

$$|\alpha - x_{n+1}| \leq C|\alpha - x_n|^p \quad n \geq 0 \quad (5)$$

for some $C > 0$. If $p = 1$, the sequence is said to converge linearly to α .

Order of Convergence

Definition: A sequence of iterates $\langle x_n \rangle$ is said to converge with order $p \geq 1$ to the actual root α if

$$|\alpha - x_{n+1}| \leq C|\alpha - x_n|^p \quad n \geq 0 \quad (5)$$

for some $C > 0$. If $p = 1$, the sequence is said to converge linearly to α .

Remark:

- For the linear convergence, the constant C must satisfy $C < 1$.
(Homework)

Order of Convergence

Definition: A sequence of iterates $\langle x_n \rangle$ is said to converge with order $p \geq 1$ to the actual root α if

$$|\alpha - x_{n+1}| \leq C |\alpha - x_n|^p \quad n \geq 0 \quad (5)$$

for some $C > 0$. If $p = 1$, the sequence is said to converge linearly to α .

Remark:

- For the linear convergence, the constant C must satisfy $C < 1$.
(Homework)

Equivalent Definition: Let $\langle x_n \rangle$ be a sequence of values converging to α , such that

$$\lim_{n \rightarrow \infty} \frac{(\alpha - x_{n+1})}{(\alpha - x_n)^p} = C \quad (6)$$

for some non-zero (finite) constant C , and some p .

Order of Convergence

Definition: A sequence of iterates $\langle x_n \rangle$ is said to converge with order $p \geq 1$ to the actual root α if

$$|\alpha - x_{n+1}| \leq C|\alpha - x_n|^p \quad n \geq 0 \quad (5)$$

for some $C > 0$. If $p = 1$, the sequence is said to converge linearly to α .

Remark:

- For the linear convergence, the constant C must satisfy $C < 1$.
(Homework)

Equivalent Definition: Let $\langle x_n \rangle$ be a sequence of values converging to α , such that

$$\lim_{n \rightarrow \infty} \frac{(\alpha - x_{n+1})}{(\alpha - x_n)^p} = C \quad (6)$$

for some non-zero (finite) constant C , and some p . Then p is called the order of convergence for the sequence.

Order of Convergence

Definition: A sequence of iterates $\langle x_n \rangle$ is said to converge with order $p \geq 1$ to the actual root α if

$$|\alpha - x_{n+1}| \leq C|\alpha - x_n|^p \quad n \geq 0 \quad (5)$$

for some $C > 0$. If $p = 1$, the sequence is said to converge linearly to α .

Remark:

- For the linear convergence, the constant C must satisfy $C < 1$.
(Homework)

Equivalent Definition: Let $\langle x_n \rangle$ be a sequence of values converging to α , such that

$$\lim_{n \rightarrow \infty} \frac{(\alpha - x_{n+1})}{(\alpha - x_n)^p} = C \quad (6)$$

for some non-zero (finite) constant C , and some p . Then p is called the order of convergence for the sequence. Bisection method has linear convergence.

Order of Convergence Contd....

Order of Convergence Contd....

Remark:

- Generally speaking, having p larger means that convergence is more rapid.

Order of Convergence Contd....

Remark:

- Generally speaking, having p larger means that convergence is more rapid. For the p th order convergence methods, we obtain

$$\lim_{n \rightarrow \infty} \frac{(\alpha - x_{n+1})}{(\alpha - x_n)^p} = C, \quad (7)$$

Order of Convergence Contd....

Remark:

- Generally speaking, having p larger means that convergence is more rapid. For the p th order convergence methods, we obtain

$$\lim_{n \rightarrow \infty} \frac{(\alpha - x_{n+1})}{(\alpha - x_n)^p} = C, \quad (7)$$

which implies

$$(\alpha - x_{n+1}) \approx C(\alpha - x_n)^p. \quad (8)$$

Order of Convergence Contd....

Remark:

- Generally speaking, having p larger means that convergence is more rapid. For the p th order convergence methods, we obtain

$$\lim_{n \rightarrow \infty} \frac{(\alpha - x_{n+1})}{(\alpha - x_n)^p} = C, \quad (7)$$

which implies

$$(\alpha - x_{n+1}) \approx C(\alpha - x_n)^p. \quad (8)$$

Taking common logarithms of both sides, we have

$$\log_{10}(\alpha - x_{n+1}) \approx \log_{10} C + p \log_{10}(\alpha - x_n). \quad (9)$$

Again, the quantity

$$\log_{10}(\alpha - x_n)$$

can be interpreted as the number of correct decimal digits in the approximation (as long as the error is less than 1),

Order of Convergence Contd....

Remark:

- Generally speaking, having p larger means that convergence is more rapid. For the p th order convergence methods, we obtain

$$\lim_{n \rightarrow \infty} \frac{(\alpha - x_{n+1})}{(\alpha - x_n)^p} = C, \quad (7)$$

which implies

$$(\alpha - x_{n+1}) \approx C(\alpha - x_n)^p. \quad (8)$$

Taking common logarithms of both sides, we have

$$\log_{10}(\alpha - x_{n+1}) \approx \log_{10} C + p \log_{10}(\alpha - x_n). \quad (9)$$

Again, the quantity

$$\log_{10}(\alpha - x_n)$$

can be interpreted as the number of correct decimal digits in the approximation (as long as the error is less than 1), and relation (9) shows that (ignoring the constant term) this p times each iteration.

Summary

Summary

Regarding the Bisection method, we observe following:

Summary

Regarding the Bisection method, we observe following:

- Disadvantages:

Summary

Regarding the Bisection method, we observe following:

- Disadvantages:
 - The major problem with bisection method is that it converges very slowly when compared with the methods defined in next few lectures.

Summary

Regarding the Bisection method, we observe following:

- Disadvantages:
 - The major problem with bisection method is that it converges very slowly when compared with the methods defined in next few lectures.
- Advantages:

Summary

Regarding the Bisection method, we observe following:

- Disadvantages:
 - The major problem with bisection method is that it converges very slowly when compared with the methods defined in next few lectures.
- Advantages:
 - The method is guaranteed to converge (provided f is continuous on $[a, b]$).

Summary

Regarding the Bisection method, we observe following:

- Disadvantages:
 - The major problem with bisection method is that it converges very slowly when compared with the methods defined in next few lectures.
- Advantages:
 - The method is guaranteed to converge (provided f is continuous on $[a, b]$).
 - At every step, we get upper and lower bounds on the root.