

MA 322: Scientific Computing

Lecture - 8



Interpolation

Interpolation

Recall that interpolation is a mathematical technique used to

Interpolation

Recall that interpolation is a mathematical technique used to find (or estimate) value of a function at intermediate points with the help of a given set of data points.

Interpolation

Recall that interpolation is a mathematical technique used to find (or estimate) value of a function at intermediate points with the help of a given set of data points.

- Technically, from a given set of $(n + 1)$ data points

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

Interpolation

Recall that interpolation is a mathematical technique used to find (or estimate) value of a function at intermediate points with the help of a given set of data points.

- Technically, from a given set of $(n + 1)$ data points

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

we find a polynomial P_n of degree n

Interpolation

Recall that interpolation is a mathematical technique used to find (or estimate) value of a function at intermediate points with the help of a given set of data points.

- Technically, from a given set of $(n + 1)$ data points

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

we find a polynomial P_n of degree n such that

$$P_n(x_i) = f(x_i), \quad 0 \leq i \leq n.$$

Interpolation

Recall that interpolation is a mathematical technique used to find (or estimate) value of a function at intermediate points with the help of a given set of data points.

- Technically, from a given set of $(n + 1)$ data points

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

we find a polynomial P_n of degree n such that

$$P_n(x_i) = f(x_i), \quad 0 \leq i \leq n.$$

- Our objective is to discuss few algorithm which are capable of constructing polynomial function $P_n(x)$ with minimum computational cost.

Interpolation

Recall that interpolation is a mathematical technique used to find (or estimate) value of a function at intermediate points with the help of a given set of data points.

- Technically, from a given set of $(n + 1)$ data points

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

we find a polynomial P_n of degree n such that

$$P_n(x_i) = f(x_i), \quad 0 \leq i \leq n.$$

- Our objective is to discuss few algorithm which are capable of constructing polynomial function $P_n(x)$ with minimum computational cost. Such few algorithms are Lagrange's Interpolation and Newton's Interpolation etc.

Interpolation

Recall that interpolation is a mathematical technique used to find (or estimate) value of a function at intermediate points with the help of a given set of data points.

- Technically, from a given set of $(n + 1)$ data points

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

we find a polynomial P_n of degree n such that

$$P_n(x_i) = f(x_i), \quad 0 \leq i \leq n.$$

● Our objective is to discuss few algorithm which are capable of constructing polynomial function $P_n(x)$ with minimum computational cost. Such few algorithms are Lagrange's Interpolation and Newton's Interpolation etc.

- Recall the advantages and disadvantages of Lagrange's and Newton's backward/forward interpolation.

Interpolation

Recall that interpolation is a mathematical technique used to find (or estimate) value of a function at intermediate points with the help of a given set of data points.

- Technically, from a given set of $(n + 1)$ data points

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

we find a polynomial P_n of degree n such that

$$P_n(x_i) = f(x_i), \quad 0 \leq i \leq n.$$

● Our objective is to discuss few algorithm which are capable of constructing polynomial function $P_n(x)$ with minimum computational cost. Such few algorithms are Lagrange's Interpolation and Newton's Interpolation etc.

- Recall the advantages and disadvantages of Lagrange's and Newton's backward/forward interpolation. Now, our objective is to develop an interpolation method which can accommodate new data and does not require equidistant arguments.

Newton Divided Differences

Newton Divided Differences

Motivation:

Newton Divided Differences

Motivation:

- Suppose P_{n-1} is a polynomial of degree $n - 1$ such that

$$P_{n-1}(x_i) = f(x_i), \quad 0 \leq i \leq n - 1.$$

Newton Divided Differences

Motivation:

- Suppose P_{n-1} is a polynomial of degree $n - 1$ such that

$$P_{n-1}(x_i) = f(x_i), \quad 0 \leq i \leq n - 1.$$

- Now, with the information of polynomial P_{n-1} , we wish to construct a polynomial P_n of degree n

Newton Divided Differences

Motivation:

- Suppose P_{n-1} is a polynomial of degree $n - 1$ such that

$$P_{n-1}(x_i) = f(x_i), \quad 0 \leq i \leq n - 1.$$

- Now, with the information of polynomial P_{n-1} , we wish to construct a polynomial P_n of degree n such that

$$P_n(x_i) = f(x_i), \quad 0 \leq i \leq n.$$

Newton Divided Differences

Motivation:

- Suppose P_{n-1} is a polynomial of degree $n - 1$ such that

$$P_{n-1}(x_i) = f(x_i), \quad 0 \leq i \leq n - 1.$$

- Now, with the information of polynomial P_{n-1} , we wish to construct a polynomial P_n of degree n such that

$$P_n(x_i) = f(x_i), \quad 0 \leq i \leq n.$$

Idea:

Newton Divided Differences

Motivation:

- Suppose P_{n-1} is a polynomial of degree $n - 1$ such that

$$P_{n-1}(x_i) = f(x_i), \quad 0 \leq i \leq n - 1.$$

- Now, with the information of polynomial P_{n-1} , we wish to construct a polynomial P_n of degree n such that

$$P_n(x_i) = f(x_i), \quad 0 \leq i \leq n.$$

Idea: This is done by introducing a correction term $C(x)$ such that

$$P_n(x) = P_{n-1}(x) + C(x). \quad (1)$$

Newton Divided Differences

Motivation:

- Suppose P_{n-1} is a polynomial of degree $n - 1$ such that

$$P_{n-1}(x_i) = f(x_i), \quad 0 \leq i \leq n - 1.$$

- Now, with the information of polynomial P_{n-1} , we wish to construct a polynomial P_n of degree n such that

$$P_n(x_i) = f(x_i), \quad 0 \leq i \leq n.$$

Idea: This is done by introducing a correction term $C(x)$ such that

$$P_n(x) = P_{n-1}(x) + C(x). \quad (1)$$

Observe that $C(x)$ is a polynomial of degree n such that

$$C(x_i) = P_n(x_i) - P_{n-1}(x_i) = f(x_i) - f(x_i) = 0, \quad 0 \leq i \leq n - 1. \quad (2)$$

Newton Divided Differences

Motivation:

- Suppose P_{n-1} is a polynomial of degree $n - 1$ such that

$$P_{n-1}(x_i) = f(x_i), \quad 0 \leq i \leq n - 1.$$

- Now, with the information of polynomial P_{n-1} , we wish to construct a polynomial P_n of degree n such that

$$P_n(x_i) = f(x_i), \quad 0 \leq i \leq n.$$

Idea: This is done by introducing a correction term $C(x)$ such that

$$P_n(x) = P_{n-1}(x) + C(x). \quad (1)$$

Observe that $C(x)$ is a polynomial of degree n such that

$$C(x_i) = P_n(x_i) - P_{n-1}(x_i) = f(x_i) - f(x_i) = 0, \quad 0 \leq i \leq n - 1. \quad (2)$$

Thus, $C(x) = a_n(x - x_0)(x - x_1)(x - x_2) \dots (x - x_{n-1})$.

Newton Divided Differences

Motivation:

- Suppose P_{n-1} is a polynomial of degree $n - 1$ such that

$$P_{n-1}(x_i) = f(x_i), \quad 0 \leq i \leq n - 1.$$

- Now, with the information of polynomial P_{n-1} , we wish to construct a polynomial P_n of degree n such that

$$P_n(x_i) = f(x_i), \quad 0 \leq i \leq n.$$

Idea: This is done by introducing a correction term $C(x)$ such that

$$P_n(x) = P_{n-1}(x) + C(x). \quad (1)$$

Observe that $C(x)$ is a polynomial of degree n such that

$$C(x_i) = P_n(x_i) - P_{n-1}(x_i) = f(x_i) - f(x_i) = 0, \quad 0 \leq i \leq n - 1. \quad (2)$$

Thus, $C(x) = a_n(x - x_0)(x - x_1)(x - x_2) \dots (x - x_{n-1})$. Then unknown quantity a_n is calculated from (1) using the condition $P_n(x_n) = f(x_n)$, so that

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \dots (x_n - x_{n-1})}. \quad (3)$$

Divided Differences

Divided Differences

First order Newton's divided difference:

$$f[x_0, x_1] = \frac{f(x_1) - f(x_0)}{x_1 - x_0}, \quad f[x_1, x_2] = \frac{f(x_2) - f(x_1)}{x_2 - x_1}, \dots$$

Divided Differences

First order Newton's divided difference:

$$f[x_0, x_1] = \frac{f(x_1) - f(x_0)}{x_1 - x_0}, \quad f[x_1, x_2] = \frac{f(x_2) - f(x_1)}{x_2 - x_1}, \dots$$

Second order Newton's divided difference:

$$f[x_0, x_1, x_2] = \frac{f[x_1, x_2] - f[x_0, x_1]}{x_2 - x_0}, \quad f[x_1, x_2, x_3] = \frac{f[x_2, x_3] - f[x_1, x_2]}{x_3 - x_1}, \dots$$

Divided Differences

First order Newton's divided difference:

$$f[x_0, x_1] = \frac{f(x_1) - f(x_0)}{x_1 - x_0}, \quad f[x_1, x_2] = \frac{f(x_2) - f(x_1)}{x_2 - x_1}, \dots$$

Second order Newton's divided difference:

$$f[x_0, x_1, x_2] = \frac{f[x_1, x_2] - f[x_0, x_1]}{x_2 - x_0}, \quad f[x_1, x_2, x_3] = \frac{f[x_2, x_3] - f[x_1, x_2]}{x_3 - x_1}, \dots$$

\vdots
 \vdots
 \vdots

n -th order Newton's divided difference:

$$f[x_0, x_1, \dots, x_n] = \frac{f[x_1, x_2, \dots, x_n] - f[x_0, x_1, \dots, x_{n-1}]}{x_n - x_0}, \dots$$

Divided Differences Contd....

Divided Differences Contd....

Let us recall the coefficients a_n , which is given by

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \dots (x_n - x_{n-1})}. \quad (4)$$

Divided Differences Contd....

Let us recall the coefficients a_n , which is given by

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \cdots (x_n - x_{n-1})}. \quad (4)$$

In fact, the coefficient a_n can be characterized as the n -th order Newton divided difference $f[x_0, x_1, \dots, x_n]$.

Divided Differences Contd....

Let us recall the coefficients a_n , which is given by

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \dots (x_n - x_{n-1})}. \quad (4)$$

In fact, the coefficient a_n can be characterized as the n -th order Newton divided difference $f[x_0, x_1, \dots, x_n]$. For a_n , we note that

$$a_1 = \frac{f(x_1) - P_0(x_1)}{x_1 - x_0} =$$

Divided Differences Contd....

Let us recall the coefficients a_n , which is given by

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \dots (x_n - x_{n-1})}. \quad (4)$$

In fact, the coefficient a_n can be characterized as the n -th order Newton divided difference $f[x_0, x_1, \dots, x_n]$. For a_n , we note that

$$a_1 = \frac{f(x_1) - P_0(x_1)}{x_1 - x_0} = \frac{f(x_1) - f(x_0)}{x_1 - x_0} =: f[x_0, x_1]$$

Divided Differences Contd....

Let us recall the coefficients a_n , which is given by

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \dots (x_n - x_{n-1})}. \quad (4)$$

In fact, the coefficient a_n can be characterized as the n -th order Newton divided difference $f[x_0, x_1, \dots, x_n]$. For a_n , we note that

$$a_1 = \frac{f(x_1) - P_0(x_1)}{x_1 - x_0} = \frac{f(x_1) - f(x_0)}{x_1 - x_0} =: f[x_0, x_1]$$

$$a_2 = \frac{f(x_2) - P_1(x_2)}{(x_2 - x_0)(x_2 - x_1)} = \frac{f(x_2) - f(x_0) - \frac{f(x_1) - f(x_0)}{x_1 - x_0}(x_2 - x_0)}{(x_2 - x_0)(x_2 - x_1)}$$

Divided Differences Contd....

Let us recall the coefficients a_n , which is given by

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \dots (x_n - x_{n-1})}. \quad (4)$$

In fact, the coefficient a_n can be characterized as the n -th order Newton divided difference $f[x_0, x_1, \dots, x_n]$. For a_n , we note that

$$\begin{aligned} a_1 &= \frac{f(x_1) - P_0(x_1)}{x_1 - x_0} = \frac{f(x_1) - f(x_0)}{x_1 - x_0} =: f[x_0, x_1] \\ a_2 &= \frac{f(x_2) - P_1(x_2)}{(x_2 - x_0)(x_2 - x_1)} = \frac{f(x_2) - f(x_0) - \frac{f(x_1) - f(x_0)}{x_1 - x_0}(x_2 - x_0)}{(x_2 - x_0)(x_2 - x_1)} \\ &= \left(\frac{f(x_2) - f(x_0)}{x_2 - x_0} - f[x_0, x_1] \right) / (x_2 - x_1) \end{aligned}$$

Divided Differences Contd....

Let us recall the coefficients a_n , which is given by

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \dots (x_n - x_{n-1})}. \quad (4)$$

In fact, the coefficient a_n can be characterized as the n -th order Newton divided difference $f[x_0, x_1, \dots, x_n]$. For a_n , we note that

$$\begin{aligned} a_1 &= \frac{f(x_1) - P_0(x_1)}{x_1 - x_0} = \frac{f(x_1) - f(x_0)}{x_1 - x_0} =: f[x_0, x_1] \\ a_2 &= \frac{f(x_2) - P_1(x_2)}{(x_2 - x_0)(x_2 - x_1)} = \frac{f(x_2) - f(x_0) - \frac{f(x_1) - f(x_0)}{x_1 - x_0}(x_2 - x_0)}{(x_2 - x_0)(x_2 - x_1)} \\ &= \left(\frac{f(x_2) - f(x_0)}{x_2 - x_0} - f[x_0, x_1] \right) / (x_2 - x_1) \\ &= \left(\frac{f(x_2) - f(x_1)}{x_2 - x_0} + \frac{f(x_1) - f(x_0)}{x_2 - x_0} - f[x_0, x_1] \right) / (x_2 - x_1) \end{aligned}$$

Divided Differences Contd....

Let us recall the coefficients a_n , which is given by

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \dots (x_n - x_{n-1})}. \quad (4)$$

In fact, the coefficient a_n can be characterized as the n -th order Newton divided difference $f[x_0, x_1, \dots, x_n]$. For a_n , we note that

$$a_1 = \frac{f(x_1) - P_0(x_1)}{x_1 - x_0} = \frac{f(x_1) - f(x_0)}{x_1 - x_0} =: f[x_0, x_1]$$

$$\begin{aligned} a_2 &= \frac{f(x_2) - P_1(x_2)}{(x_2 - x_0)(x_2 - x_1)} = \frac{f(x_2) - f(x_0) - \frac{f(x_1) - f(x_0)}{x_1 - x_0}(x_2 - x_0)}{(x_2 - x_0)(x_2 - x_1)} \\ &= \left(\frac{f(x_2) - f(x_0)}{x_2 - x_0} - f[x_0, x_1] \right) / (x_2 - x_1) \\ &= \left(\frac{f(x_2) - f(x_1)}{x_2 - x_0} + \frac{f(x_1) - f(x_0)}{x_2 - x_0} - f[x_0, x_1] \right) / (x_2 - x_1) \\ &= \frac{f[x_1, x_2]}{x_2 - x_0} + \frac{f[x_0, x_1]}{x_2 - x_1} \left[\frac{x_1 - x_0}{x_2 - x_0} - 1 \right] = \frac{f[x_1, x_2] - f[x_0, x_1]}{x_2 - x_0} =: f[x_0, x_1, x_2] \end{aligned}$$

Newton's Divided Difference Formula

Newton's Divided Difference Formula

Therefore, for the coefficients a_n , we write

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \dots (x_n - x_{n-1})} = f[x_0, x_1, x_2, \dots, x_n]. \quad (5)$$

Newton's Divided Difference Formula

Therefore, for the coefficients a_n , we write

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \dots (x_n - x_{n-1})} = f[x_0, x_1, x_2, \dots, x_n]. \quad (5)$$

Hence, the correction term $C(x)$ can be written as

$$C(x) = (x - x_0)(x - x_1)(x - x_2) \dots (x - x_{n-1})f[x_0, x_1, x_2, \dots, x_n], \quad (6)$$

Newton's Divided Difference Formula

Therefore, for the coefficients a_n , we write

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \dots (x_n - x_{n-1})} = f[x_0, x_1, x_2, \dots, x_n]. \quad (5)$$

Hence, the correction term $C(x)$ can be written as

$$C(x) = (x - x_0)(x - x_1)(x - x_2) \dots (x - x_{n-1})f[x_0, x_1, x_2, \dots, x_n], \quad (6)$$

so that

$$P_n(x) = P_{n-1}(x) + C(x)$$

Newton's Divided Difference Formula

Therefore, for the coefficients a_n , we write

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \dots (x_n - x_{n-1})} = f[x_0, x_1, x_2, \dots, x_n]. \quad (5)$$

Hence, the correction term $C(x)$ can be written as

$$C(x) = (x - x_0)(x - x_1)(x - x_2) \dots (x - x_{n-1})f[x_0, x_1, x_2, \dots, x_n], \quad (6)$$

so that

$$\begin{aligned} P_n(x) &= P_{n-1}(x) + C(x) \\ &= P_{n-1} + (x - x_0)(x - x_1)(x - x_2) \dots (x - x_{n-1})f[x_0, x_1, x_2, \dots, x_n]. \end{aligned}$$

Newton's Divided Difference Formula

Therefore, for the coefficients a_n , we write

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \dots (x_n - x_{n-1})} = f[x_0, x_1, x_2, \dots, x_n]. \quad (5)$$

Hence, the correction term $C(x)$ can be written as

$$C(x) = (x - x_0)(x - x_1)(x - x_2) \dots (x - x_{n-1})f[x_0, x_1, x_2, \dots, x_n], \quad (6)$$

so that

$$\begin{aligned} P_n(x) &= P_{n-1}(x) + C(x) \\ &= P_{n-1} + (x - x_0)(x - x_1)(x - x_2) \dots (x - x_{n-1})f[x_0, x_1, x_2, \dots, x_n]. \end{aligned}$$

Recursively, we now construct Newton's divided difference formula for the interpolating polynomial

$$P_0(x) = f(x_0),$$

Newton's Divided Difference Formula

Therefore, for the coefficients a_n , we write

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \dots (x_n - x_{n-1})} = f[x_0, x_1, x_2, \dots, x_n]. \quad (5)$$

Hence, the correction term $C(x)$ can be written as

$$C(x) = (x - x_0)(x - x_1)(x - x_2) \dots (x - x_{n-1})f[x_0, x_1, x_2, \dots, x_n], \quad (6)$$

so that

$$\begin{aligned} P_n(x) &= P_{n-1}(x) + C(x) \\ &= P_{n-1} + (x - x_0)(x - x_1)(x - x_2) \dots (x - x_{n-1})f[x_0, x_1, x_2, \dots, x_n]. \end{aligned}$$

Recursively, we now construct Newton's divided difference formula for the interpolating polynomial

$$P_0(x) = f(x_0), \quad P_1(x) = f(x_0) + (x - x_0)f[x_0, x_1],$$

Newton's Divided Difference Formula

Therefore, for the coefficients a_n , we write

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \dots (x_n - x_{n-1})} = f[x_0, x_1, x_2, \dots, x_n]. \quad (5)$$

Hence, the correction term $C(x)$ can be written as

$$C(x) = (x - x_0)(x - x_1)(x - x_2) \dots (x - x_{n-1})f[x_0, x_1, x_2, \dots, x_n], \quad (6)$$

so that

$$\begin{aligned} P_n(x) &= P_{n-1}(x) + C(x) \\ &= P_{n-1} + (x - x_0)(x - x_1)(x - x_2) \dots (x - x_{n-1})f[x_0, x_1, x_2, \dots, x_n]. \end{aligned}$$

Recursively, we now construct Newton's divided difference formula for the interpolating polynomial

$$\begin{aligned} P_0(x) &= f(x_0), \quad P_1(x) = f(x_0) + (x - x_0)f[x_0, x_1], \\ P_2(x) &= f(x_0) + (x - x_0)f[x_0, x_1] + (x - x_0)(x - x_1)f[x_0, x_1, x_2] \end{aligned}$$

Newton's Divided Difference Formula

Therefore, for the coefficients a_n , we write

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \dots (x_n - x_{n-1})} = f[x_0, x_1, x_2, \dots, x_n]. \quad (5)$$

Hence, the correction term $C(x)$ can be written as

$$C(x) = (x - x_0)(x - x_1)(x - x_2) \dots (x - x_{n-1})f[x_0, x_1, x_2, \dots, x_n], \quad (6)$$

so that

$$\begin{aligned} P_n(x) &= P_{n-1}(x) + C(x) \\ &= P_{n-1} + (x - x_0)(x - x_1)(x - x_2) \dots (x - x_{n-1})f[x_0, x_1, x_2, \dots, x_n]. \end{aligned}$$

Recursively, we now construct Newton's divided difference formula for the interpolating polynomial

$$\begin{aligned} P_0(x) &= f(x_0), \quad P_1(x) = f(x_0) + (x - x_0)f[x_0, x_1], \\ P_2(x) &= f(x_0) + (x - x_0)f[x_0, x_1] + (x - x_0)(x - x_1)f[x_0, x_1, x_2] \\ &\dots \end{aligned}$$

Newton's Divided Difference Formula

Therefore, for the coefficients a_n , we write

$$a_n = \frac{f(x_n) - P_{n-1}(x_n)}{(x_n - x_0) \dots (x_n - x_{n-1})} = f[x_0, x_1, x_2, \dots, x_n]. \quad (5)$$

Hence, the correction term $C(x)$ can be written as

$$C(x) = (x - x_0)(x - x_1)(x - x_2) \dots (x - x_{n-1})f[x_0, x_1, x_2, \dots, x_n], \quad (6)$$

so that

$$\begin{aligned} P_n(x) &= P_{n-1}(x) + C(x) \\ &= P_{n-1} + (x - x_0)(x - x_1)(x - x_2) \dots (x - x_{n-1})f[x_0, x_1, x_2, \dots, x_n]. \end{aligned}$$

Recursively, we now construct Newton's divided difference formula for the interpolating polynomial

[illegible]

Divided Difference Table

Divided Difference Table

Format for constructing divided differences of $f(x)$

x_i	$f(x_i)$	$f[x_i, x_{i+1}]$	$f[x_i, x_{i+1}, x_{i+2}] \cdots$
x_0	f_0		
		$f[x_0, x_1]$	
x_1	f_1		$f[x_0, x_1, x_2] \cdots$
		$f[x_1, x_2]$	
x_2	f_2		$f[x_1, x_2, x_3] \cdots$
		$f[x_2, x_3]$	
x_3	f_3		$f[x_2, x_3, x_4] \cdots$
		$f[x_3, x_4]$	
x_4	f_4		$f[x_3, x_4, x_5] \cdots$
		$f[x_4, x_5]$	
x_5	f_5		
\vdots	\vdots	\vdots	\vdots

Divided differences table

Consider the nodes $\mathbf{x} := [x_0, \dots, x_n]$ and the values $\mathbf{f} := [f_0, \dots, f_n]$.

Divided differences table

Consider the nodes $\mathbf{x} := [x_0, \dots, x_n]$ and the values $\mathbf{f} := [f_0, \dots, f_n]$.

Then, we generate following table of divided differences:

\mathbf{x}	\mathbf{f}			
x_0	f_0			
x_1	f_1	$f[x_0, x_1]$		
x_2	f_2	$f[x_1, x_2]$	$f[x_0, x_1, x_2]$	
x_3	f_3	$f[x_2, x_3]$	$f[x_1, x_2, x_3]$	$f[x_0, x_1, x_2, x_3]$
\cdot	\cdot	\dots	\dots	\dots

Divided differences table

Consider the nodes $\mathbf{x} := [x_0, \dots, x_n]$ and the values $\mathbf{f} := [f_0, \dots, f_n]$. Then, we generate following table of divided differences:

\mathbf{x}	\mathbf{f}			
x_0	f_0			
x_1	f_1	$f[x_0, x_1]$		
x_2	f_2	$f[x_1, x_2]$	$f[x_0, x_1, x_2]$	
x_3	f_3	$f[x_2, x_3]$	$f[x_1, x_2, x_3]$	$f[x_0, x_1, x_2, x_3]$
\cdot	\cdot	\dots	\dots	\dots

The $n + 1$ diagonal entries in the table of divided differences are the coefficients of the Newton interpolating polynomial $p(x)$.

Divided differences table

Consider the nodes $\mathbf{x} := [x_0, \dots, x_n]$ and the values $\mathbf{f} := [f_0, \dots, f_n]$.

Then, we generate following table of divided differences:

\mathbf{x}	\mathbf{f}				
x_0	f_0				
x_1	f_1	$f[x_0, x_1]$			
x_2	f_2	$f[x_1, x_2]$	$f[x_0, x_1, x_2]$		
x_3	f_3	$f[x_2, x_3]$	$f[x_1, x_2, x_3]$	$f[x_0, x_1, x_2, x_3]$	
\cdot	\cdot	\dots	\dots	\dots	

The $n + 1$ diagonal entries in the table of divided differences are the coefficients of the Newton interpolating polynomial $p(x)$.

\mathbf{x}	\mathbf{f}				
0	3				
1	4	$(4 - 3)/(1 - 0) = 1$			
2	7	$(7 - 4)/(2 - 1) = 3$	$(3 - 1)/(2 - 0) = 1$		
4	19	$(19 - 7)/(4 - 2) = 6$	$(6 - 3)/(4 - 1) = 1$	$(1 - 1)/(4 - 0) = 0$	

Divided differences table

Consider the nodes $\mathbf{x} := [x_0, \dots, x_n]$ and the values $\mathbf{f} := [f_0, \dots, f_n]$. Then, we generate following table of divided differences:

\mathbf{x}	\mathbf{f}				
x_0	f_0				
x_1	f_1	$f[x_0, x_1]$			
x_2	f_2	$f[x_1, x_2]$	$f[x_0, x_1, x_2]$		
x_3	f_3	$f[x_2, x_3]$	$f[x_1, x_2, x_3]$	$f[x_0, x_1, x_2, x_3]$	
\vdots	\vdots	\dots	\dots	\dots	

The $n + 1$ diagonal entries in the table of divided differences are the coefficients of the Newton interpolating polynomial $p(x)$.

\mathbf{x}	\mathbf{f}				
0	3				
1	4	$(4 - 3)/(1 - 0) = 1$			
2	7	$(7 - 4)/(2 - 1) = 3$	$(3 - 1)/(2 - 0) = 1$		
4	19	$(19 - 7)/(4 - 2) = 6$	$(6 - 3)/(4 - 1) = 1$	$(1 - 1)/(4 - 0) = 0$	

$$p(x) = 3 + 1 \cdot (x - 0) + 1 \cdot (x - 0)(x - 1) + 0 \cdot (x - 0)(x - 1)(x - 2) = 3 + x^2.$$

Divided differences at repeated nodes

Theorem: If $f \in C^n[a, b]$ and $[x_0, \dots, x_n]$ are distinct nodes then

$$f[x_0, \dots, x_n] = \frac{f^{(n)}(\theta)}{n!}$$

for some θ between the smallest and the largest nodes $[x_0, \dots, x_n]$.

Divided differences at repeated nodes

Theorem: If $f \in C^n[a, b]$ and $[x_0, \dots, x_n]$ are distinct nodes then

$$f[x_0, \dots, x_n] = \frac{f^{(n)}(\theta)}{n!}$$

for some θ between the smallest and the largest nodes $[x_0, \dots, x_n]$.

If x_j tends to x_0 for $j = 1 : n$, then θ , being trapped between the nodes, must tend to x_0 . Hence taking limit as $x_j \rightarrow x_0$ for $j = 1 : n$, we have

$$f[x_0, \dots, x_0] = \frac{f^{(n)}(x_0)}{n!}.$$

Divided differences at repeated nodes

Theorem: If $f \in C^n[a, b]$ and $[x_0, \dots, x_n]$ are distinct nodes then

$$f[x_0, \dots, x_n] = \frac{f^{(n)}(\theta)}{n!}$$

for some θ between the smallest and the largest nodes $[x_0, \dots, x_n]$.

If x_j tends to x_0 for $j = 1 : n$, then θ , being trapped between the nodes, must tend to x_0 . Hence taking limit as $x_j \rightarrow x_0$ for $j = 1 : n$, we have

$$f[x_0, \dots, x_0] = \frac{f^{(n)}(x_0)}{n!}.$$

This defines the n -th order divided difference of f at $n + 1$ times repeated node x_0 .