

# MA 322: Scientific Computing

## Lecture - 11



# Method of undetermined coefficients

The weights in the Simpson's rule

$$S(f) = \left[ w_0 f(a) + w_1 f\left(\frac{a+b}{2}\right) + w_2 f(b) \right] \approx \int_a^b f(x) dx$$

can be determined by utilizing the fact that  $S(f)$  is exact for  $f(x) = 1$ ,  $f(x) = x$ , and  $f(x) = x^2$ .

# Method of undetermined coefficients

The weights in the Simpson's rule

$$S(f) = \left[ w_0 f(a) + w_1 f\left(\frac{a+b}{2}\right) + w_2 f(b) \right] \approx \int_a^b f(x) dx$$

can be determined by utilizing the fact that  $S(f)$  is exact for  $f(x) = 1$ ,  $f(x) = x$ , and  $f(x) = x^2$ . Hence

$$w_0 + w_1 + w_2 = \int_a^b dx = b - a$$

# Method of undetermined coefficients

The weights in the Simpson's rule

$$S(f) = \left[ w_0 f(a) + w_1 f\left(\frac{a+b}{2}\right) + w_2 f(b) \right] \approx \int_a^b f(x) dx$$

can be determined by utilizing the fact that  $S(f)$  is exact for  $f(x) = 1$ ,  $f(x) = x$ , and  $f(x) = x^2$ . Hence

$$\begin{aligned} w_0 + w_1 + w_2 &= \int_a^b dx = b - a \\ w_0 a + w_1 \left(\frac{a+b}{2}\right) + w_2 b &= \int_a^b x dx = \frac{(b^2 - a^2)}{2} \end{aligned}$$

# Method of undetermined coefficients

The weights in the Simpson's rule

$$S(f) = \left[ w_0 f(a) + w_1 f\left(\frac{a+b}{2}\right) + w_2 f(b) \right] \approx \int_a^b f(x) dx$$

can be determined by utilizing the fact that  $S(f)$  is exact for  $f(x) = 1$ ,  $f(x) = x$ , and  $f(x) = x^2$ . Hence

$$\begin{aligned} w_0 + w_1 + w_2 &= \int_a^b dx = b - a \\ w_0 a + w_1 \left(\frac{a+b}{2}\right) + w_2 b &= \int_a^b x dx = \frac{(b^2 - a^2)}{2} \\ w_0 a^2 + w_1 \left(\frac{a+b}{2}\right)^2 + w_2 b^2 &= \int_a^b x^2 dx = \frac{(b^3 - a^3)}{3} \end{aligned}$$

# Method of undetermined coefficients

Solving the Vandermonde system

$$\begin{bmatrix} 1 & 1 & 1 \\ a & \frac{a+b}{2} & b \\ a^2 & \frac{(a+b)^2}{4} & b^2 \end{bmatrix} \begin{bmatrix} w_0 \\ w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} b-a \\ \frac{b^2-a^2}{2} \\ \frac{b^3-a^3}{3} \end{bmatrix}$$

we obtain  $w_0 = (b-a)/6 = w_2$  and  $w_1 = 2(b-a)/3$ .

# Method of undetermined coefficients

Solving the Vandermonde system

$$\begin{bmatrix} 1 & 1 & 1 \\ a & \frac{a+b}{2} & b \\ a^2 & \frac{(a+b)^2}{4} & b^2 \end{bmatrix} \begin{bmatrix} w_0 \\ w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} b-a \\ \frac{b^2-a^2}{2} \\ \frac{b^3-a^3}{3} \end{bmatrix}$$

we obtain  $w_0 = (b-a)/6 = w_2$  and  $w_1 = 2(b-a)/3$ .

Now consider an  $(n+1)$ -point quadrature rule

$$\int_b^a f(x) dx \approx \sum_{j=0}^n w_j f(x_j).$$

# Method of undetermined coefficients

Solving the Vandermonde system

$$\begin{bmatrix} 1 & 1 & 1 \\ a & \frac{a+b}{2} & b \\ a^2 & \frac{(a+b)^2}{4} & b^2 \end{bmatrix} \begin{bmatrix} w_0 \\ w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} b-a \\ \frac{b^2-a^2}{2} \\ \frac{b^3-a^3}{3} \end{bmatrix}$$

we obtain  $w_0 = (b-a)/6 = w_2$  and  $w_1 = 2(b-a)/3$ .

Now consider an  $(n+1)$ -point quadrature rule

$$\int_b^a f(x) dx \approx \sum_{j=0}^n w_j f(x_j).$$

If the quadrature rule is exact for polynomials of degree  $\leq n$ , then the weights can be determined by solving the Vandermonde system



# Method of undetermined coefficients

Solving the Vandermonde system

$$\begin{bmatrix} 1 & 1 & 1 \\ a & \frac{a+b}{2} & b \\ a^2 & \frac{(a+b)^2}{4} & b^2 \end{bmatrix} \begin{bmatrix} w_0 \\ w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} b-a \\ \frac{b^2-a^2}{2} \\ \frac{b^3-a^3}{3} \end{bmatrix}$$

we obtain  $w_0 = (b-a)/6 = w_2$  and  $w_1 = 2(b-a)/3$ .

Now consider an  $(n+1)$ -point quadrature rule

$$\int_b^a f(x) dx \approx \sum_{j=0}^n w_j f(x_j).$$

If the quadrature rule is exact for polynomials of degree  $\leq n$ , then the weights can be determined by solving the Vandermonde system

$$\begin{bmatrix} 1 & 1 & \cdots & 1 \\ x_0 & x_1 & \cdots & x_n \\ \vdots & \vdots & \cdots & \vdots \\ x_0^n & x_1^n & \cdots & x_n^n \end{bmatrix} \begin{bmatrix} w_0 \\ w_1 \\ \vdots \\ w_n \end{bmatrix} = \begin{bmatrix} b-a \\ \frac{b^2-a^2}{2} \\ \vdots \\ \frac{b^{n+1}-a^{n+1}}{n+1} \end{bmatrix}$$

## Example

Once again consider the Simpson rule

$$\int_0^1 f(x)dx \approx w_0f(0) + w_1f(1/2) + w_2f(1).$$

Since Simpson's rule is exact for polynomials of degree  $\leq 2$ ,

## Example

Once again consider the Simpson rule

$$\int_0^1 f(x)dx \approx w_0 f(0) + w_1 f(1/2) + w_2 f(1).$$

Since Simpson's rule is exact for polynomials of degree  $\leq 2$ , the Vandermonde system is given by

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & \frac{1}{2} & 1 \\ 0 & \frac{1}{4} & 1 \end{bmatrix} \begin{bmatrix} w_0 \\ w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} 1 \\ \frac{1}{2} \\ \frac{1}{3} \end{bmatrix}.$$

## Example

Once again consider the Simpson rule

$$\int_0^1 f(x)dx \approx w_0 f(0) + w_1 f(1/2) + w_2 f(1).$$

Since Simpson's rule is exact for polynomials of degree  $\leq 2$ , the Vandermonde system is given by

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & \frac{1}{2} & 1 \\ 0 & \frac{1}{4} & 1 \end{bmatrix} \begin{bmatrix} w_0 \\ w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} 1 \\ \frac{1}{2} \\ \frac{1}{3} \end{bmatrix}.$$

Solving the system, we have  $w_0 = 1/6 = w_2$  and  $w_1 = 2/3$ . Hence the Simpson's rule is given by

## Example

Once again consider the Simpson rule

$$\int_0^1 f(x)dx \approx w_0 f(0) + w_1 f(1/2) + w_2 f(1).$$

Since Simpson's rule is exact for polynomials of degree  $\leq 2$ , the Vandermonde system is given by

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & \frac{1}{2} & 1 \\ 0 & \frac{1}{4} & 1 \end{bmatrix} \begin{bmatrix} w_0 \\ w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} 1 \\ \frac{1}{2} \\ \frac{1}{3} \end{bmatrix}.$$

Solving the system, we have  $w_0 = 1/6 = w_2$  and  $w_1 = 2/3$ . Hence the Simpson's rule is given by

$$\int_0^1 f(x)dx \approx \frac{1}{6}[f(0) + 4f(1/2) + f(1)].$$

# Error estimation of Simpson's rule

**Theorem:** Suppose that  $f \in C^4[a, b]$ . Then

$$\int_a^b f(x)dx = \frac{(b-a)}{6} \left[ f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] + E_2(f),$$

where

# Error estimation of Simpson's rule

**Theorem:** Suppose that  $f \in C^4[a, b]$ . Then

$$\int_a^b f(x) dx = \frac{(b-a)}{6} \left[ f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] + E_2(f),$$

where  $E_2(f) = -\frac{1}{90} \left( \frac{b-a}{2} \right)^5 f^{(4)}(\theta)$  for some  $\theta \in [a, b]$ .

# Error estimation of Simpson's rule

**Theorem:** Suppose that  $f \in C^4[a, b]$ . Then

$$\int_a^b f(x) dx = \frac{(b-a)}{6} \left[ f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] + E_2(f),$$

where  $E_2(f) = -\frac{1}{90} \left( \frac{b-a}{2} \right)^5 f^{(4)}(\theta)$  for some  $\theta \in [a, b]$ .

**Proof:** Set  $h := (b-a)/2$ . Expanding  $f(a+h)$  and  $f(a+2h)$  by Taylor series, we have



# Error estimation of Simpson's rule

**Theorem:** Suppose that  $f \in C^4[a, b]$ . Then

$$\int_a^b f(x) dx = \frac{(b-a)}{6} \left[ f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] + E_2(f),$$

where  $E_2(f) = -\frac{1}{90} \left( \frac{b-a}{2} \right)^5 f^{(4)}(\theta)$  for some  $\theta \in [a, b]$ .

**Proof:** Set  $h := (b-a)/2$ . Expanding  $f(a+h)$  and  $f(a+2h)$  by Taylor series, we have

$$S(f) = \frac{h}{3} [f(a) + 4f(a+h) + f(a+2h)]$$

# Error estimation of Simpson's rule

**Theorem:** Suppose that  $f \in C^4[a, b]$ . Then

$$\int_a^b f(x) dx = \frac{(b-a)}{6} \left[ f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] + E_2(f),$$

where  $E_2(f) = -\frac{1}{90} \left( \frac{b-a}{2} \right)^5 f^{(4)}(\theta)$  for some  $\theta \in [a, b]$ .

**Proof:** Set  $h := (b-a)/2$ . Expanding  $f(a+h)$  and  $f(a+2h)$  by Taylor series, we have

$$\begin{aligned} S(f) &= \frac{h}{3} [f(a) + 4f(a+h) + f(a+2h)] \\ &= 2hf(a) + 2h^2 f'(a) + \frac{4}{3} h^3 f''(a) + \frac{2}{3} h^4 f'''(a) + \frac{100}{3 \cdot 5!} h^5 f^{(4)}(a) + \dots \end{aligned}$$

# Error estimation of Simpson's rule

**Theorem:** Suppose that  $f \in C^4[a, b]$ . Then

$$\int_a^b f(x)dx = \frac{(b-a)}{6} \left[ f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] + E_2(f),$$

where  $E_2(f) = -\frac{1}{90} \left( \frac{b-a}{2} \right)^5 f^{(4)}(\theta)$  for some  $\theta \in [a, b]$ .

**Proof:** Set  $h := (b-a)/2$ . Expanding  $f(a+h)$  and  $f(a+2h)$  by Taylor series, we have

$$\begin{aligned} S(f) &= \frac{h}{3} [f(a) + 4f(a+h) + f(a+2h)] \\ &= 2hf(a) + 2h^2f'(a) + \frac{4}{3}h^3f''(a) + \frac{2}{3}h^4f'''(a) + \frac{100}{3 \cdot 5!}h^5f^{(4)}(a) + \dots \end{aligned}$$

Let  $F(a+2h) := \int_a^{a+2h} f(x)dx$ . Then

# Error estimation of Simpson's rule

**Theorem:** Suppose that  $f \in C^4[a, b]$ . Then

$$\int_a^b f(x) dx = \frac{(b-a)}{6} \left[ f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] + E_2(f),$$

where  $E_2(f) = -\frac{1}{90} \left( \frac{b-a}{2} \right)^5 f^{(4)}(\theta)$  for some  $\theta \in [a, b]$ .

**Proof:** Set  $h := (b-a)/2$ . Expanding  $f(a+h)$  and  $f(a+2h)$  by Taylor series, we have

$$\begin{aligned} S(f) &= \frac{h}{3} [f(a) + 4f(a+h) + f(a+2h)] \\ &= 2hf(a) + 2h^2f'(a) + \frac{4}{3}h^3f''(a) + \frac{2}{3}h^4f'''(a) + \frac{100}{3 \cdot 5!}h^5f^{(4)}(a) + \dots \end{aligned}$$

Let  $F(a+2h) := \int_a^{a+2h} f(x) dx$ . Then

$$F(a+2h) = 2hf(a) + 2h^2f'(a) + \frac{4}{3}h^3f''(a) + \frac{2}{3}h^4f'''(a) + \frac{32}{5!}h^5f^{(4)}(a) + \dots$$

# Error estimation of Simpson's rule

**Theorem:** Suppose that  $f \in C^4[a, b]$ . Then

$$\int_a^b f(x) dx = \frac{(b-a)}{6} \left[ f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] + E_2(f),$$

where  $E_2(f) = -\frac{1}{90} \left( \frac{b-a}{2} \right)^5 f^{(4)}(\theta)$  for some  $\theta \in [a, b]$ .

**Proof:** Set  $h := (b-a)/2$ . Expanding  $f(a+h)$  and  $f(a+2h)$  by Taylor series, we have

$$\begin{aligned} S(f) &= \frac{h}{3} [f(a) + 4f(a+h) + f(a+2h)] \\ &= 2hf(a) + 2h^2f'(a) + \frac{4}{3}h^3f''(a) + \frac{2}{3}h^4f'''(a) + \frac{100}{3 \cdot 5!}h^5f^{(4)}(a) + \dots \end{aligned}$$

Let  $F(a+2h) := \int_a^{a+2h} f(x) dx$ . Then

$$F(a+2h) = 2hf(a) + 2h^2f'(a) + \frac{4}{3}h^3f''(a) + \frac{2}{3}h^4f'''(a) + \frac{32}{5!}h^5f^{(4)}(a) + \dots$$

Combining the two expansions, the desired result follows.

# Degree of exactness

The error  $E_2(f) = -\frac{1}{90} \left( \frac{b-a}{2} \right)^5 f^{(4)}(\theta)$  shows that Simpson's rule  $S(f)$  is exact for polynomials of degree  $\leq 3$ , which is **one degree more than a 3-points Newton-Cotes quadrature formula can guarantee.**

# Degree of exactness

The error  $E_2(f) = -\frac{1}{90} \left( \frac{b-a}{2} \right)^5 f^{(4)}(\theta)$  shows that Simpson's rule  $S(f)$  is exact for polynomials of degree  $\leq 3$ , which is **one degree more than a 3-points Newton-Cotes quadrature formula can guarantee.**

Also, the midpoint rule  $M(f)$ , which corresponds to Newton-Cotes quadrature for  $n = 0$ , is exact for polynomial of degree  $\leq 1$ . Again this is **one degree more than a 1-point Newton-Cotes formula can guarantee.**

# Degree of exactness

The error  $E_2(f) = -\frac{1}{90} \left( \frac{b-a}{2} \right)^5 f^{(4)}(\theta)$  shows that Simpson's rule  $S(f)$  is exact for polynomials of degree  $\leq 3$ , which is **one degree more than a 3-points Newton-Cotes quadrature formula can guarantee.**

Also, the midpoint rule  $M(f)$ , which corresponds to Newton-Cotes quadrature for  $n = 0$ , is exact for polynomial of degree  $\leq 1$ . Again this is **one degree more than a 1-point Newton-Cotes formula can guarantee.**

For example,  $\int_0^1 f(x) dx \approx f(1/2)$  is exact for  $f(x) = 1$  and  $f(x) = x$ .



## Degree of exactness

The error  $E_2(f) = -\frac{1}{90} \left( \frac{b-a}{2} \right)^5 f^{(4)}(\theta)$  shows that Simpson's rule  $S(f)$  is exact for polynomials of degree  $\leq 3$ , which is **one degree more than a 3-points Newton-Cotes quadrature formula can guarantee.**

Also, the midpoint rule  $M(f)$ , which corresponds to Newton-Cotes quadrature for  $n = 0$ , is exact for polynomial of degree  $\leq 1$ . Again this is **one degree more than a 1-point Newton-Cotes formula can guarantee.**

For example,  $\int_0^1 f(x)dx \approx f(1/2)$  is exact for  $f(x) = 1$  and  $f(x) = x$ .

In general, if  $n$  is even then an  **$(n+1)$ -point Newton-Cotes rule gains an extra degree of exactness** beyond that of the polynomial interpolant on which it is based.

## Degree of exactness

The error  $E_2(f) = -\frac{1}{90} \left( \frac{b-a}{2} \right)^5 f^{(4)}(\theta)$  shows that Simpson's rule  $S(f)$  is exact for polynomials of degree  $\leq 3$ , which is **one degree more than a 3-points Newton-Cotes quadrature formula can guarantee.**

Also, the midpoint rule  $M(f)$ , which corresponds to Newton-Cotes quadrature for  $n = 0$ , is exact for polynomial of degree  $\leq 1$ . Again this is **one degree more than a 1-point Newton-Cotes formula can guarantee.**

For example,  $\int_0^1 f(x)dx \approx f(1/2)$  is exact for  $f(x) = 1$  and  $f(x) = x$ .

In general, if  $n$  is even then an  **$(n+1)$ -point Newton-Cotes rule gains an extra degree of exactness** beyond that of the polynomial interpolant on which it is based.

This phenomenon is due to cancellation of positive and negative errors.

## Degree of exactness

The error  $E_2(f) = -\frac{1}{90} \left( \frac{b-a}{2} \right)^5 f^{(4)}(\theta)$  shows that Simpson's rule  $S(f)$  is exact for polynomials of degree  $\leq 3$ , which is **one degree more than a 3-points Newton-Cotes quadrature formula can guarantee.**

Also, the midpoint rule  $M(f)$ , which corresponds to Newton-Cotes quadrature for  $n = 0$ , is exact for polynomial of degree  $\leq 1$ . Again this is **one degree more than a 1-point Newton-Cotes formula can guarantee.**

For example,  $\int_0^1 f(x) dx \approx f(1/2)$  is exact for  $f(x) = 1$  and  $f(x) = x$ .

In general, if  $n$  is even then an  **$(n+1)$ -point Newton-Cotes rule gains an extra degree of exactness** beyond that of the polynomial interpolant on which it is based.

This phenomenon is due to cancellation of positive and negative errors. Such cancellation does not occur, however, when  $n$  is odd.

## Degree of exactness

The error  $E_2(f) = -\frac{1}{90} \left( \frac{b-a}{2} \right)^5 f^{(4)}(\theta)$  shows that Simpson's rule  $S(f)$  is exact for polynomials of degree  $\leq 3$ , which is **one degree more than a 3-points Newton-Cotes quadrature formula can guarantee.**

Also, the midpoint rule  $M(f)$ , which corresponds to Newton-Cotes quadrature for  $n = 0$ , is exact for polynomial of degree  $\leq 1$ . Again this is **one degree more than a 1-point Newton-Cotes formula can guarantee.**

For example,  $\int_0^1 f(x) dx \approx f(1/2)$  is exact for  $f(x) = 1$  and  $f(x) = x$ .

In general, if  $n$  is even then an  **$(n+1)$ -point Newton-Cotes rule gains an extra degree of exactness** beyond that of the polynomial interpolant on which it is based.

This phenomenon is due to cancellation of positive and negative errors. Such cancellation does not occur, however, when  $n$  is odd.

Thus, **the degree of exactness of an  $(n+1)$ -point Newton-Cotes rule is  $n$  if  $n$  is odd, but the degree of exactness is  $n+1$  if  $n$  is even.**

## Example

Consider the integral

$$I := \int_0^1 e^{-x^2} dx.$$

The three Newton-Cotes (midpoint, trapezoid, and Simpson's rule) formula yield

$$M(f) = (1 - 0)e^{-0.25} \approx 0.778801$$

## Example

Consider the integral

$$I := \int_0^1 e^{-x^2} dx.$$

The three Newton-Cotes (midpoint, trapezoid, and Simpson's rule) formula yield

$$M(f) = (1 - 0)e^{-0.25} \approx 0.778801$$

$$T(f) = \frac{1}{2}[e^0 + e^{-1}] \approx 0.683940$$

## Example

Consider the integral

$$I := \int_0^1 e^{-x^2} dx.$$

The three Newton-Cotes (midpoint, trapezoid, and Simpson's rule) formula yield

$$M(f) = (1 - 0)e^{-0.25} \approx 0.778801$$

$$T(f) = \frac{1}{2}[e^0 + e^{-1}] \approx 0.683940$$

$$S(f) = \frac{1}{6}[e^0 + 4e^{-0.25} + e^{-1}] \approx 0.747180$$

## Example

Consider the integral

$$I := \int_0^1 e^{-x^2} dx.$$

The three Newton-Cotes (midpoint, trapezoid, and Simpson's rule) formula yield

$$M(f) = (1 - 0)e^{-0.25} \approx 0.778801$$

$$T(f) = \frac{1}{2}[e^0 + e^{-1}] \approx 0.683940$$

$$S(f) = \frac{1}{6}[e^0 + 4e^{-0.25} + e^{-1}] \approx 0.747180$$

The correctly rounded value is  $I = 0.746824$ .



## Example

Consider the integral

$$I := \int_0^1 e^{-x^2} dx.$$

The three Newton-Cotes (midpoint, trapezoid, and Simpson's rule) formula yield

$$M(f) = (1 - 0)e^{-0.25} \approx 0.778801$$

$$T(f) = \frac{1}{2}[e^0 + e^{-1}] \approx 0.683940$$

$$S(f) = \frac{1}{6}[e^0 + 4e^{-0.25} + e^{-1}] \approx 0.747180$$

The correctly rounded value is  $I = 0.746824$ . It is somewhat surprising to see that the error from the trapezoid rule (0.062884) is about twice that from the midpoint rule (0.031977).

## Example

Consider the integral

$$I := \int_0^1 e^{-x^2} dx.$$

The three Newton-Cotes (midpoint, trapezoid, and Simpson's rule) formula yield

$$M(f) = (1 - 0)e^{-0.25} \approx 0.778801$$

$$T(f) = \frac{1}{2}[e^0 + e^{-1}] \approx 0.683940$$

$$S(f) = \frac{1}{6}[e^0 + 4e^{-0.25} + e^{-1}] \approx 0.747180$$

The correctly rounded value is  $I = 0.746824$ . It is somewhat surprising to see that the error from the trapezoid rule (0.062884) is about twice that from the midpoint rule (0.031977).

The Simpson's rule, with an error of only 0.000356, seems remarkably accurate considering the size of the interval over which it is applied.

# Degree of exactness of midpoint rule

Consider the midpoint rule

$$\int_a^b f(x) dx \approx (b-a)f\left(\frac{a+b}{2}\right) = M(f).$$

# Degree of exactness of midpoint rule

Consider the midpoint rule

$$\int_a^b f(x) dx \approx (b-a)f\left(\frac{a+b}{2}\right) = M(f).$$

The error in the midpoint rule  $M(f)$  can be estimated by means of Taylor series expansion about the midpoint  $c := (a+b)/2$  of the interval  $[a, b]$  :

# Degree of exactness of midpoint rule

Consider the midpoint rule

$$\int_a^b f(x) dx \approx (b-a)f\left(\frac{a+b}{2}\right) = M(f).$$

The error in the midpoint rule  $M(f)$  can be estimated by means of Taylor series expansion about the midpoint  $c := (a+b)/2$  of the interval  $[a, b]$  :

$$\begin{aligned} f(x) &= f(c) + f'(c)(x-c) + \frac{1}{2}f''(c)(x-c)^2 + \frac{1}{6}f'''(c)(x-c)^3 \\ &\quad + \frac{1}{24}f^{(4)}(c)(x-c)^4 + \dots \end{aligned}$$

# Degree of exactness of midpoint rule

Consider the midpoint rule

$$\int_a^b f(x) dx \approx (b-a)f\left(\frac{a+b}{2}\right) = M(f).$$

The error in the midpoint rule  $M(f)$  can be estimated by means of Taylor series expansion about the midpoint  $c := (a+b)/2$  of the interval  $[a, b]$  :

$$\begin{aligned} f(x) &= f(c) + f'(c)(x-c) + \frac{1}{2}f''(c)(x-c)^2 + \frac{1}{6}f'''(c)(x-c)^3 \\ &\quad + \frac{1}{24}f^{(4)}(c)(x-c)^4 + \dots \end{aligned}$$

Upon integration from  $a$  to  $b$ , the odd-order terms drops out, yielding

# Degree of exactness of midpoint rule

Consider the midpoint rule

$$\int_a^b f(x)dx \approx (b-a)f\left(\frac{a+b}{2}\right) = M(f).$$

The error in the midpoint rule  $M(f)$  can be estimated by means of Taylor series expansion about the midpoint  $c := (a+b)/2$  of the interval  $[a, b]$  :

$$\begin{aligned} f(x) &= f(c) + f'(c)(x-c) + \frac{1}{2}f''(c)(x-c)^2 + \frac{1}{6}f'''(c)(x-c)^3 \\ &\quad + \frac{1}{24}f^{(4)}(c)(x-c)^4 + \dots \end{aligned}$$

Upon integration from  $a$  to  $b$ , the odd-order terms drops out, yielding

$$\int_a^b f(x)dx = (b-a)f(c) + \frac{1}{6}f''(c)(b-a)^3 + \frac{1}{1920}f^{(4)}(c)(b-a)^5 + \dots$$

# Degree of exactness of midpoint rule

Consider the midpoint rule

$$\int_a^b f(x)dx \approx (b-a)f\left(\frac{a+b}{2}\right) = M(f).$$

The error in the midpoint rule  $M(f)$  can be estimated by means of Taylor series expansion about the midpoint  $c := (a+b)/2$  of the interval  $[a, b]$  :

$$\begin{aligned} f(x) &= f(c) + f'(c)(x-c) + \frac{1}{2}f''(c)(x-c)^2 + \frac{1}{6}f'''(c)(x-c)^3 \\ &\quad + \frac{1}{24}f^{(4)}(c)(x-c)^4 + \dots \end{aligned}$$

Upon integration from  $a$  to  $b$ , the odd-order terms drops out, yielding

$$\int_a^b f(x)dx = (b-a)f(c) + \frac{1}{6}f''(c)(b-a)^3 + \frac{1}{1920}f^{(4)}(c)(b-a)^5 + \dots$$

This shows that the degree of exactness of  $M(f)$  is 1.



# Composite Simpson's rule

Suppose that  $n$  is even. Set  $h := (b - a)/n$  and consider the nodes  $x_j = a + jh$  for  $j = 0 : n$ . Then

$$\int_a^b f(x)dx = \int_{x_0}^{x_2} f(x)dx + \int_{x_2}^{x_4} f(x)dx + \cdots + \int_{x_{n-2}}^{x_n} f(x)dx$$

# Composite Simpson's rule

Suppose that  $n$  is even. Set  $h := (b - a)/n$  and consider the nodes  $x_j = a + jh$  for  $j = 0 : n$ . Then

$$\begin{aligned}\int_a^b f(x)dx &= \int_{x_0}^{x_2} f(x)dx + \int_{x_2}^{x_4} f(x)dx + \cdots + \int_{x_{n-2}}^{x_n} f(x)dx \\ &= \sum_{j=1}^{n/2} \int_{x_{2j-2}}^{x_{2j}} f(x)dx\end{aligned}$$

# Composite Simpson's rule

Suppose that  $n$  is even. Set  $h := (b - a)/n$  and consider the nodes  $x_j = a + jh$  for  $j = 0 : n$ . Then

$$\begin{aligned}\int_a^b f(x)dx &= \int_{x_0}^{x_2} f(x)dx + \int_{x_2}^{x_4} f(x)dx + \cdots + \int_{x_{n-2}}^{x_n} f(x)dx \\ &= \sum_{j=1}^{n/2} \int_{x_{2j-2}}^{x_{2j}} f(x)dx \approx \sum_{j=1}^{n/2} \frac{h}{3} [f(x_{2j-2}) + 4f(x_{2j-1}) + f(x_{2j})]\end{aligned}$$

# Composite Simpson's rule

Suppose that  $n$  is even. Set  $h := (b - a)/n$  and consider the nodes  $x_j = a + jh$  for  $j = 0 : n$ . Then

$$\begin{aligned}\int_a^b f(x)dx &= \int_{x_0}^{x_2} f(x)dx + \int_{x_2}^{x_4} f(x)dx + \cdots + \int_{x_{n-2}}^{x_n} f(x)dx \\&= \sum_{j=1}^{n/2} \int_{x_{2j-2}}^{x_{2j}} f(x)dx \approx \sum_{j=1}^{n/2} \frac{h}{3} [f(x_{2j-2}) + 4f(x_{2j-1}) + f(x_{2j})] \\&\approx \frac{h}{3} [f(x_0) + 2 \sum_{j=2}^{n/2} f(x_{2j-2}) + 4 \sum_{j=1}^{n/2} f(x_{2j-1}) + f(x_n)].\end{aligned}$$

# Composite Simpson's rule

Suppose that  $n$  is even. Set  $h := (b - a)/n$  and consider the nodes  $x_j = a + jh$  for  $j = 0 : n$ . Then

$$\begin{aligned}\int_a^b f(x)dx &= \int_{x_0}^{x_2} f(x)dx + \int_{x_2}^{x_4} f(x)dx + \cdots + \int_{x_{n-2}}^{x_n} f(x)dx \\&= \sum_{j=1}^{n/2} \int_{x_{2j-2}}^{x_{2j}} f(x)dx \approx \sum_{j=1}^{n/2} \frac{h}{3} [f(x_{2j-2}) + 4f(x_{2j-1}) + f(x_{2j})] \\&\approx \frac{h}{3} [f(x_0) + 2 \sum_{j=2}^{n/2} f(x_{2j-2}) + 4 \sum_{j=1}^{n/2} f(x_{2j-1}) + f(x_n)].\end{aligned}$$

Thus the composite Simpson's rule is given by

$$S_n(f) := \frac{h}{3} [f(x_0) + 2 \sum_{j=2}^{n/2} f(x_{2j-2}) + 4 \sum_{j=1}^{n/2} f(x_{2j-1}) + f(x_n)]$$

# Composite Simpson's rule

Suppose that  $n$  is even. Set  $h := (b - a)/n$  and consider the nodes  $x_j = a + jh$  for  $j = 0 : n$ . Then

$$\begin{aligned}\int_a^b f(x)dx &= \int_{x_0}^{x_2} f(x)dx + \int_{x_2}^{x_4} f(x)dx + \cdots + \int_{x_{n-2}}^{x_n} f(x)dx \\&= \sum_{j=1}^{n/2} \int_{x_{2j-2}}^{x_{2j}} f(x)dx \approx \sum_{j=1}^{n/2} \frac{h}{3} [f(x_{2j-2}) + 4f(x_{2j-1}) + f(x_{2j})] \\&\approx \frac{h}{3} [f(x_0) + 2 \sum_{j=2}^{n/2} f(x_{2j-2}) + 4 \sum_{j=1}^{n/2} f(x_{2j-1}) + f(x_n)].\end{aligned}$$

Thus the composite Simpson's rule is given by

$$S_n(f) := \frac{h}{3} [f(x_0) + 2 \sum_{j=2}^{n/2} f(x_{2j-2}) + 4 \sum_{j=1}^{n/2} f(x_{2j-1}) + f(x_n)]$$

with error  $E_n^S(f) = -\frac{1}{90}h^5 \sum_{j=1}^{n/2} f^{(4)}(\theta_j) = -\frac{(b-a)}{180}h^4 f^{(4)}(\theta)$  for some  $\theta \in [a, b]$ .