

# MA 322: Scientific Computing

## Lecture - 2




# The Newton's Method

# The Newton's Method

Idea:

# The Newton's Method

Idea:




**Figure:** Newton's method.

- Assume that an initial estimate  $x_0$  is known for the desired root  $\alpha$  of equation  $f(x) = 0$ .

# The Newton's Method

Idea:




**Figure:** Newton's method.

- Assume that an initial estimate  $x_0$  is known for the desired root  $\alpha$  of equation  $f(x) = 0$ .
- Newton's method will produce a sequence of iterates  $\{x_n : n \geq 1\}$ ,

# The Newton's Method

Idea:




**Figure:** Newton's method.

- Assume that an initial estimate  $x_0$  is known for the desired root  $\alpha$  of equation  $f(x) = 0$ .
- Newton's method will produce a sequence of iterates  $\{x_n : n \geq 1\}$ , which we hope will converge to  $\alpha$ .

# The Newton's Method

Idea:




**Figure:** Newton's method.

Construction of Newton's Method:

- Assume that an initial estimate  $x_0$  is known for the desired root  $\alpha$  of equation  $f(x) = 0$ .
- Newton's method will produce a sequence of iterates  $\{x_n : n \geq 1\}$ , which we hope will converge to  $\alpha$ .

# The Newton's Method

Idea:



**Figure:** Newton's method.

Construction of Newton's Method:


- Slope of the tangent to the curve  $y = f(x)$  at  $(x_0, f(x_0))$  is given by  $f'(x_0)$ ,

- Assume that an initial estimate  $x_0$  is known for the desired root  $\alpha$  of equation  $f(x) = 0$ .
- Newton's method will produce a sequence of iterates  $\{x_n : n \geq 1\}$ , which we hope will converge to  $\alpha$ .



# The Newton's Method

Idea:



**Figure:** Newton's method.

- Assume that an initial estimate  $x_0$  is known for the desired root  $\alpha$  of equation  $f(x) = 0$ .
- Newton's method will produce a sequence of iterates  $\{x_n : n \geq 1\}$ , which we hope will converge to  $\alpha$ .


Construction of Newton's Method:

- Slope of the tangent to the curve  $y = f(x)$  at  $(x_0, f(x_0))$  is given by  $f'(x_0)$ , thus, the tangent line passing through  $(x_0, f(x_0))$  is given by

$$y - f(x_0) = f'(x_0)(x - x_0). \quad (1)$$

# The Newton's Method

Idea:



**Figure:** Newton's method.

- Assume that an initial estimate  $x_0$  is known for the desired root  $\alpha$  of equation  $f(x) = 0$ .
- Newton's method will produce a sequence of iterates  $\{x_n : n \geq 1\}$ , which we hope will converge to  $\alpha$ .

Construction of Newton's Method:

- Slope of the tangent to the curve  $y = f(x)$  at  $(x_0, f(x_0))$  is given by  $f'(x_0)$ , thus, the tangent line passing through  $(x_0, f(x_0))$  is given by

$$y - f(x_0) = f'(x_0)(x - x_0). \quad (1)$$

Since this tangent line expected to cut x-axis at  $(x_1, 0)$ , we obtain

$$-f(x_0) = f'(x_0)(x_1 - x_0) \quad \text{Or} \quad x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}. \quad (2)$$

# Newton's Method Contd.....

# Newton's Method Contd.....

Sequence of Iterates for Newton's Method:

# Newton's Method Contd.....

Sequence of Iterates for Newton's Method:

- In general, we have

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}. \quad (3)$$

# Newton's Method Contd.....

Sequence of Iterates for Newton's Method:

- In general, we have

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}. \quad (3)$$

Example: Find an approximation for the largest root  $\alpha \approx 1.1347241$  of the equation

$$x^6 - x - 1 = 0.$$

**Newton's method**

$n$	$x_n$
0	2.0
1	1.680628273
2	1.430738989
3	1.254970957
4	1.161538433
5	1.136353274
6	1.134730528
7	1.134724139

**Bisection method**

$n$	$c_n$	$n$	$c_n$
$2 = b, 1 = a$		8	1.13672
1	1.5	9	1.13477
2	1.25	10	1.13379
3	1.125	11	1.13428
4	1.1875	12	1.13452
5	1.15625	13	1.13464
6	1.14063	14	1.13470
7	1.13281	15	1.13474

# Newton's Method Contd.....

## Newton's Method Contd.....

Example: Find an approximation for the root  $\alpha \approx 1.414213562373095$  of the equation

$$x^2 - 2 = 0.$$

A Matlab Code:

```
n=5;  x(1)=2;  
for i=1:n  
x(i+1)=x(i)-(x(i)^2-2)/(2*x(i));  
end
```



## Newton's Method Contd.....

Example: Find an approximation for the root  $\alpha \approx 1.414213562373095$  of the equation

$$x^2 - 2 = 0.$$

A Matlab Code:

```
n=5;  x(1)=2;
```

```
for i=1:n
```

```
  x(i+1)=x(i)-(x(i)^2-2)/(2*x(i));
```

```
end
```

$$x_1 = 2, \quad x_2 = 1.5000000000$$

$$x_3 = 1.416666666666667,$$

$$x_4 = 1.4142156862$$

$$x_5 = 1.4142135623,$$

$$x_6 = 1.4142135623$$

## Newton's Method Contd.....

Example: Find an approximation for the root  $\alpha \approx 1.414213562373095$  of the equation

$$x^2 - 2 = 0.$$

A Matlab Code:

```
n=5; x(1)=2;
```

```
for i=1:n
```

```
x(i+1)=x(i)-(x(i)^2-2)/(2*x(i));
```

```
end
```

$$x_1 = 2, \quad x_2 = 1.5000000000$$

$$x_3 = 1.416666666666667,$$

$$x_4 = 1.4142156862$$

$$x_5 = 1.4142135623,$$

$$x_6 = 1.4142135623$$

Convergence of Newton's Method:

## Newton's Method Contd.....

Example: Find an approximation for the root  $\alpha \approx 1.414213562373095$  of the equation

$$x^2 - 2 = 0.$$

A Matlab Code:

```
n=5; x(1)=2;
```

```
for i=1:n
```

```
x(i+1)=x(i)-(x(i)^2-2)/(2*x(i));
```

```
end
```

$$x_1 = 2, \quad x_2 = 1.5000000000$$

$$x_3 = 1.416666666666667,$$

$$x_4 = 1.4142156862$$

$$x_5 = 1.4142135623,$$

$$x_6 = 1.4142135623$$

Convergence of Newton's Method:

- Recall following Taylor series expansion

$$f(x) = f(x_n) + (x - x_n)f'(x_n) + \frac{(x - x_n)^2}{2}f''(\xi_n), \quad (4)$$

## Newton's Method Contd.....

Example: Find an approximation for the root  $\alpha \approx 1.414213562373095$  of the equation

$$x^2 - 2 = 0.$$

A Matlab Code:

```
n=5; x(1)=2;
```

```
for i=1:n
```

```
x(i+1)=x(i)-(x(i)^2-2)/(2*x(i));
```

```
end
```

$$x_1 = 2, \quad x_2 = 1.5000000000$$

$$x_3 = 1.416666666666667,$$

$$x_4 = 1.4142156862$$

$$x_5 = 1.4142135623,$$

$$x_6 = 1.4142135623$$

Convergence of Newton's Method:

- Recall following Taylor series expansion

$$f(x) = f(x_n) + (x - x_n)f'(x_n) + \frac{(x - x_n)^2}{2}f''(\xi_n), \quad (4)$$

with  $\xi_n$  between  $x$  &  $x_n$ .

## Newton's Method Contd.....

Example: Find an approximation for the root  $\alpha \approx 1.414213562373095$  of the equation

$$x^2 - 2 = 0.$$

A Matlab Code:

```
n=5; x(1)=2;
```

```
for i=1:n
```

```
x(i+1)=x(i)-(x(i)^2-2)/(2*x(i));
```

```
end
```

$$x_1 = 2, \quad x_2 = 1.5000000000$$

$$x_3 = 1.416666666666667,$$

$$x_4 = 1.4142156862$$

$$x_5 = 1.4142135623,$$

$$x_6 = 1.4142135623$$

Convergence of Newton's Method:

- Recall following Taylor series expansion

$$f(x) = f(x_n) + (x - x_n)f'(x_n) + \frac{(x - x_n)^2}{2}f''(\xi_n), \quad (4)$$

with  $\xi_n$  between  $x$  &  $x_n$ . Letting  $x = \alpha$  and using  $f(\alpha) = 0$ , we have

$$\alpha = x_n - \frac{f(x_n)}{f'(x_n)} - \frac{(\alpha - x_n)^2}{2} \frac{f''(\xi_n)}{f'(x_n)}, \quad (5)$$

# Newton's Method Contd.....

## Newton's Method Contd.....

In the process of convergence of Newton's method, we arrive at

$$\alpha - x_{n+1} = -\frac{(\alpha - x_n)^2}{2} \frac{f''(\xi_n)}{f'(x_n)}. \quad (6)$$

## Newton's Method Contd.....

In the process of convergence of Newton's method, we arrive at

$$\alpha - x_{n+1} = -\frac{(\alpha - x_n)^2}{2} \frac{f''(\xi_n)}{f'(x_n)}. \quad (6)$$

Hence,

$$\frac{(\alpha - x_{n+1})}{(\alpha - x_n)^2} = -\frac{f''(\xi_n)}{2f'(x_n)}. \quad (7)$$



## Newton's Method Contd.....

In the process of convergence of Newton's method, we arrive at

$$\alpha - x_{n+1} = -\frac{(\alpha - x_n)^2}{2} \frac{f''(\xi_n)}{f'(x_n)}. \quad (6)$$

Hence,

$$\frac{(\alpha - x_{n+1})}{(\alpha - x_n)^2} = -\frac{f''(\xi_n)}{2f'(x_n)}. \quad (7)$$

Convergence of Newton's Method:

## Newton's Method Contd.....

In the process of convergence of Newton's method, we arrive at

$$\alpha - x_{n+1} = -\frac{(\alpha - x_n)^2}{2} \frac{f''(\xi_n)}{f'(\alpha)}. \quad (6)$$

Hence,

$$\frac{(\alpha - x_{n+1})}{(\alpha - x_n)^2} = -\frac{f''(\xi_n)}{2f'(\alpha)}. \quad (7)$$

Convergence of Newton's Method: Assume  $f$ ,  $f'$  and  $f''$  are continuous for all  $x$  in some neighborhood of  $\alpha$ , and assume that  $f(\alpha) = 0$ ,  $f'(\alpha) \neq 0$ . Then if  $x_0$  is chosen sufficiently close to  $\alpha$ , the sequence  $\langle x_n \rangle$  given by

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

will converge to  $\alpha$ . Moreover,

$$\lim_{n \rightarrow \infty} \frac{(\alpha - x_{n+1})}{(\alpha - x_n)^2} = -\frac{f''(\alpha)}{2f'(\alpha)}$$

providing that the iterates have an order of convergence  $p = 2$ .

# Newton's Method Contd.....

# Newton's Method Contd.....

The proof is constructed in the following steps:

## Newton's Method Contd.....

The proof is constructed in the following steps:

- Recall following error relation

$$\alpha - x_{n+1} = -\frac{(\alpha - x_n)^2}{2} \frac{f''(\xi_n)}{f'(x_n)}. \quad (8)$$

## Newton's Method Contd.....

The proof is constructed in the following steps:

- Recall following error relation

$$\alpha - x_{n+1} = -\frac{(\alpha - x_n)^2}{2} \frac{f''(\xi_n)}{f'(x_n)}. \quad (8)$$

- Since  $f' \neq 0$  and  $f$  is  $C^2$  function, so there exists an interval  $I = [\alpha - \epsilon, \alpha + \epsilon]$  such that  $f'(x) \neq 0$  for all  $x \in I$ .

## Newton's Method Contd.....

The proof is constructed in the following steps:

- Recall following error relation

$$\alpha - x_{n+1} = -\frac{(\alpha - x_n)^2}{2} \frac{f''(\xi_n)}{f'(x_n)}. \quad (8)$$

- Since  $f' \neq 0$  and  $f$  is  $C^2$  function, so there exists an interval  $I = [\alpha - \epsilon, \alpha + \epsilon]$  such that  $f'(x) \neq 0$  for all  $x \in I$ . Now, set

$$M = \frac{\max_{x \in I} |f''(x)|}{2 \min_{x \in I} |f'(x)|}.$$

## Newton's Method Contd.....

The proof is constructed in the following steps:

- Recall following error relation

$$\alpha - x_{n+1} = -\frac{(\alpha - x_n)^2}{2} \frac{f''(\xi_n)}{f'(x_n)}. \quad (8)$$

- Since  $f' \neq 0$  and  $f$  is  $C^2$  function, so there exists an interval  $I = [\alpha - \epsilon, \alpha + \epsilon]$  such that  $f'(x) \neq 0$  for all  $x \in I$ . Now, set

$$M = \frac{\max_{x \in I} |f''(x)|}{2 \min_{x \in I} |f'(x)|}.$$

For  $|\alpha - x_0| \leq \epsilon$ , the equation (8) yields

$$|\alpha - x_1| \leq M |\alpha - x_0|^2$$



## Newton's Method Contd.....

The proof is constructed in the following steps:

- Recall following error relation

$$\alpha - x_{n+1} = -\frac{(\alpha - x_n)^2}{2} \frac{f''(\xi_n)}{f'(x_n)}. \quad (8)$$

- Since  $f' \neq 0$  and  $f$  is  $C^2$  function, so there exists an interval  $I = [\alpha - \epsilon, \alpha + \epsilon]$  such that  $f'(x) \neq 0$  for all  $x \in I$ . Now, set

$$M = \frac{\max_{x \in I} |f''(x)|}{2 \min_{x \in I} |f'(x)|}.$$

For  $|\alpha - x_0| \leq \epsilon$ , the equation (8) yields

$$|\alpha - x_1| \leq M|\alpha - x_0|^2 \quad \text{Or} \quad M|\alpha - x_1| \leq (M|\alpha - x_0|)^2. \quad (9)$$

## Newton's Method Contd.....

The proof is constructed in the following steps:

- Recall following error relation

$$\alpha - x_{n+1} = -\frac{(\alpha - x_n)^2}{2} \frac{f''(\xi_n)}{f'(x_n)}. \quad (8)$$

- Since  $f' \neq 0$  and  $f$  is  $C^2$  function, so there exists an interval  $I = [\alpha - \epsilon, \alpha + \epsilon]$  such that  $f'(x) \neq 0$  for all  $x \in I$ . Now, set

$$M = \frac{\max_{x \in I} |f''(x)|}{2 \min_{x \in I} |f'(x)|}.$$

For  $|\alpha - x_0| \leq \epsilon$ , the equation (8) yields

$$|\alpha - x_1| \leq M|\alpha - x_0|^2 \quad \text{Or} \quad M|\alpha - x_1| \leq (M|\alpha - x_0|)^2. \quad (9)$$

- To ensure  $x_1 \in I$ , we set  $M|\alpha - x_0| < 1$ ,

## Newton's Method Contd.....

The proof is constructed in the following steps:

- Recall following error relation

$$\alpha - x_{n+1} = -\frac{(\alpha - x_n)^2}{2} \frac{f''(\xi_n)}{f'(x_n)}. \quad (8)$$

- Since  $f' \neq 0$  and  $f$  is  $C^2$  function, so there exists an interval  $I = [\alpha - \epsilon, \alpha + \epsilon]$  such that  $f'(x) \neq 0$  for all  $x \in I$ . Now, set

$$M = \frac{\max_{x \in I} |f''(x)|}{2 \min_{x \in I} |f'(x)|}.$$

For  $|\alpha - x_0| \leq \epsilon$ , the equation (8) yields

$$|\alpha - x_1| \leq M|\alpha - x_0|^2 \quad \text{Or} \quad M|\alpha - x_1| \leq (M|\alpha - x_0|)^2. \quad (9)$$

- To ensure  $x_1 \in I$ , we set  $M|\alpha - x_0| < 1$ , for which  $M|\alpha - x_1| \leq M|\alpha - x_0|$ .

## Newton's Method Contd.....

The proof is constructed in the following steps:

- Recall following error relation

$$\alpha - x_{n+1} = -\frac{(\alpha - x_n)^2}{2} \frac{f''(\xi_n)}{f'(x_n)}. \quad (8)$$

- Since  $f' \neq 0$  and  $f$  is  $C^2$  function, so there exists an interval  $I = [\alpha - \epsilon, \alpha + \epsilon]$  such that  $f'(x) \neq 0$  for all  $x \in I$ . Now, set

$$M = \frac{\max_{x \in I} |f''(x)|}{2 \min_{x \in I} |f'(x)|}.$$

For  $|\alpha - x_0| \leq \epsilon$ , the equation (8) yields

$$|\alpha - x_1| \leq M|\alpha - x_0|^2 \quad \text{Or} \quad M|\alpha - x_1| \leq (M|\alpha - x_0|)^2. \quad (9)$$

- To ensure  $x_1 \in I$ , we set  $M|\alpha - x_0| < 1$ , for which  $M|\alpha - x_1| \leq M|\alpha - x_0|$ . Thus,  $|\alpha - x_1| \leq \epsilon$ .

## Newton's Method Contd.....

The proof is constructed in the following steps:

- Recall following error relation

$$\alpha - x_{n+1} = -\frac{(\alpha - x_n)^2}{2} \frac{f''(\xi_n)}{f'(x_n)}. \quad (8)$$

- Since  $f' \neq 0$  and  $f$  is  $C^2$  function, so there exists an interval  $I = [\alpha - \epsilon, \alpha + \epsilon]$  such that  $f'(x) \neq 0$  for all  $x \in I$ . Now, set

$$M = \frac{\max_{x \in I} |f''(x)|}{2 \min_{x \in I} |f'(x)|}.$$

For  $|\alpha - x_0| \leq \epsilon$ , the equation (8) yields

$$|\alpha - x_1| \leq M|\alpha - x_0|^2 \quad \text{Or} \quad M|\alpha - x_1| \leq (M|\alpha - x_0|)^2. \quad (9)$$

- To ensure  $x_1 \in I$ , we set  $M|\alpha - x_0| < 1$ , for which  $M|\alpha - x_1| \leq M|\alpha - x_0|$ . Thus,  $|\alpha - x_1| \leq \epsilon$ .
- We can apply the same argument to  $x_1, x_2, \dots$ , inductively, so that  $|\alpha - x_n| \leq \epsilon$  and  $M|\alpha - x_n| < 1$  for all  $n \geq 1$ .

# Newton's Method Contd.....

## Newton's Method Contd.....

- Finally, we obtain

$$|\alpha - x_{n+1}| \leq M(\alpha - x_n)^2 \quad \text{Or} \quad M|\alpha - x_{n+1}| \leq (M|\alpha - x_n|)^2. \quad (10)$$

## Newton's Method Contd.....

- Finally, we obtain

$$|\alpha - x_{n+1}| \leq M(\alpha - x_n)^2 \quad \text{Or} \quad M|\alpha - x_{n+1}| \leq (M|\alpha - x_n|)^2. \quad (10)$$

- Inductively, we obtain

$$M|\alpha - x_n| \leq (M|\alpha - x_{n-1}|)^2$$



## Newton's Method Contd.....

- Finally, we obtain

$$|\alpha - x_{n+1}| \leq M(\alpha - x_n)^2 \quad \text{Or} \quad M|\alpha - x_{n+1}| \leq (M|\alpha - x_n|)^2. \quad (10)$$

- Inductively, we obtain

$$\begin{aligned} M|\alpha - x_n| &\leq (M|\alpha - x_{n-1}|)^2 \\ &\leq \left( (M|\alpha - x_{n-2}|)^2 \right)^2 = (M|\alpha - x_{n-2}|)^{2^2} \end{aligned}$$

## Newton's Method Contd.....

- Finally, we obtain

$$|\alpha - x_{n+1}| \leq M(\alpha - x_n)^2 \quad \text{Or} \quad M|\alpha - x_{n+1}| \leq (M|\alpha - x_n|)^2. \quad (10)$$

- Inductively, we obtain

$$\begin{aligned} M|\alpha - x_n| &\leq (M|\alpha - x_{n-1}|)^2 \\ &\leq \left( (M|\alpha - x_{n-2}|)^2 \right)^2 = (M|\alpha - x_{n-2}|)^{2^2} \\ &\leq \dots\dots\dots \end{aligned}$$

## Newton's Method Contd.....

- Finally, we obtain

$$|\alpha - x_{n+1}| \leq M(\alpha - x_n)^2 \quad \text{Or} \quad M|\alpha - x_{n+1}| \leq (M|\alpha - x_n|)^2. \quad (10)$$

- Inductively, we obtain

$$\begin{aligned} M|\alpha - x_n| &\leq (M|\alpha - x_{n-1}|)^2 \\ &\leq \left( (M|\alpha - x_{n-2}|)^2 \right)^2 = (M|\alpha - x_{n-2}|)^{2^2} \\ &\leq \dots\dots\dots \\ &\leq (M|\alpha - x_0|)^{2^n}. \end{aligned}$$

## Newton's Method Contd.....

- Finally, we obtain

$$|\alpha - x_{n+1}| \leq M(\alpha - x_n)^2 \quad \text{Or} \quad M|\alpha - x_{n+1}| \leq (M|\alpha - x_n|)^2. \quad (10)$$

- Inductively, we obtain

$$\begin{aligned} M|\alpha - x_n| &\leq (M|\alpha - x_{n-1}|)^2 \\ &\leq \left( (M|\alpha - x_{n-2}|)^2 \right)^2 = (M|\alpha - x_{n-2}|)^{2^2} \\ &\leq \dots\dots\dots \\ &\leq (M|\alpha - x_0|)^{2^n}. \end{aligned}$$

Now, apply the fact that  $M|\alpha - x_0| < 1$  to obtain  $x_n \rightarrow \alpha$ .

## Newton's Method Contd.....

- Finally, we obtain

$$|\alpha - x_{n+1}| \leq M(\alpha - x_n)^2 \quad \text{Or} \quad M|\alpha - x_{n+1}| \leq (M|\alpha - x_n|)^2. \quad (10)$$

- Inductively, we obtain

$$\begin{aligned} M|\alpha - x_n| &\leq (M|\alpha - x_{n-1}|)^2 \\ &\leq \left( (M|\alpha - x_{n-2}|)^2 \right)^2 = (M|\alpha - x_{n-2}|)^{2^2} \\ &\leq \dots\dots\dots \\ &\leq (M|\alpha - x_0|)^{2^n}. \end{aligned}$$

Now, apply the fact that  $M|\alpha - x_0| < 1$  to obtain  $x_n \rightarrow \alpha$ .

- Again error relation

$$\frac{(\alpha - x_{n+1})}{(\alpha - x_n)^2} = -\frac{f''(\xi_n)}{2f'(x_n)} \quad (11)$$

yields

$$\lim_{n \rightarrow \infty} \frac{(\alpha - x_{n+1})}{(\alpha - x_n)^2} = -\lim_{n \rightarrow \infty} \frac{f''(\xi_n)}{2f'(x_n)} = -\frac{f''(\alpha)}{2f'(\alpha)} \quad \square \quad (12)$$

# Newton's Method Contd.....

# Newton's Method Contd.....

Remarks:

# Newton's Method Contd.....

Remarks:

- We have proved the convergence of Newton's method under the assumptions that

$$|\alpha - x_0| \leq \epsilon$$



# Newton's Method Contd.....

Remarks:

- We have proved the convergence of Newton's method under the assumptions that

$$|\alpha - x_0| \leq \epsilon \quad \text{and} \quad |\alpha - x_0| < \frac{1}{M}.$$

# Newton's Method Contd.....

Remarks:

- We have proved the convergence of Newton's method under the assumptions that

$$|\alpha - x_0| \leq \epsilon \quad \text{and} \quad |\alpha - x_0| < \frac{1}{M}.$$

- In real application we do not have the exact root  $\alpha$ .

# Newton's Method Contd.....

Remarks:

- We have proved the convergence of Newton's method under the assumptions that

$$|\alpha - x_0| \leq \epsilon \quad \text{and} \quad |\alpha - x_0| < \frac{1}{M}.$$

- In real application we do not have the exact root  $\alpha$ . So, it is not very practical to use above conditions to predict *a priori* what value of  $x_0$  to use.

# Newton's Method Contd.....

Remarks:

- We have proved the convergence of Newton's method under the assumptions that

$$|\alpha - x_0| \leq \epsilon \quad \text{and} \quad |\alpha - x_0| < \frac{1}{M}.$$

- In real application we do not have the exact root  $\alpha$ . So, it is not very practical to use above conditions to predict *a priori* what value of  $x_0$  to use.
- The point of the theorem is not to help us choose  $x_0$ , rather it explains that there exists an  $x_0$  for which the iteration converges quadratically.

# Newton's Method Contd.....

Remarks:

- We have proved the convergence of Newton's method under the assumptions that

$$|\alpha - x_0| \leq \epsilon \quad \text{and} \quad |\alpha - x_0| < \frac{1}{M}.$$

- In real application we do not have the exact root  $\alpha$ . So, it is not very practical to use above conditions to predict *a priori* what value of  $x_0$  to use.
- The point of the theorem is not to help us choose  $x_0$ , rather it explains that there exists an  $x_0$  for which the iteration converges quadratically.
- These are very restrictive assumptions and we would like to remove them, if possible.

# Newton's Method Contd.....

Remarks:

- We have proved the convergence of Newton's method under the assumptions that

$$|\alpha - x_0| \leq \epsilon \quad \text{and} \quad |\alpha - x_0| < \frac{1}{M}.$$

- In real application we do not have the exact root  $\alpha$ . So, it is not very practical to use above conditions to predict *a priori* what value of  $x_0$  to use.
- The point of the theorem is not to help us choose  $x_0$ , rather it explains that there exists an  $x_0$  for which the iteration converges quadratically.
- These are very restrictive assumptions and we would like to remove them, if possible. Suppose  $f \in C^2[a, b]$  and  $f(a) < 0$ ,  $f(b) > 0$ , and there is a unique root  $\alpha$  in  $[a, b]$ .

# Newton's Method Contd.....

Remarks:

- We have proved the convergence of Newton's method under the assumptions that

$$|\alpha - x_0| \leq \epsilon \quad \text{and} \quad |\alpha - x_0| < \frac{1}{M}.$$

- In real application we do not have the exact root  $\alpha$ . So, it is not very practical to use above conditions to predict *a priori* what value of  $x_0$  to use.
- The point of the theorem is not to help us choose  $x_0$ , rather it explains that there exists an  $x_0$  for which the iteration converges quadratically.
- These are very restrictive assumptions and we would like to remove them, if possible. Suppose  $f \in C^2[a, b]$  and  $f(a) < 0$ ,  $f(b) > 0$ , and there is a unique root  $\alpha$  in  $[a, b]$ . Further, we assume

$$f'(x) > 0 \quad \text{and} \quad f''(x) > 0 \quad x \in [a, b],$$

# Newton's Method Contd.....

Remarks:

- We have proved the convergence of Newton's method under the assumptions that

$$|\alpha - x_0| \leq \epsilon \quad \text{and} \quad |\alpha - x_0| < \frac{1}{M}.$$

- In real application we do not have the exact root  $\alpha$ . So, it is not very practical to use above conditions to predict *a priori* what value of  $x_0$  to use.
- The point of the theorem is not to help us choose  $x_0$ , rather it explains that there exists an  $x_0$  for which the iteration converges quadratically.
- These are very restrictive assumptions and we would like to remove them, if possible. Suppose  $f \in C^2[a, b]$  and  $f(a) < 0$ ,  $f(b) > 0$ , and there is a unique root  $\alpha$  in  $[a, b]$ . Further, we assume

$$f'(x) > 0 \quad \text{and} \quad f''(x) > 0 \quad x \in [a, b],$$

then the iterates  $x_n$  of Newton's method with  $x_0 = b$  converges to  $\alpha$ .



# Newton's Method: Stopping Criterion

# Newton's Method: Stopping Criterion

For a given tolerance  $\epsilon > 0$ , can we find  $n_0 \in \mathbb{N}$  such that

$$|\alpha - x_n| \leq \epsilon? \quad (13)$$

# Newton's Method: Stopping Criterion

For a given tolerance  $\epsilon > 0$ , can we find  $n_0 \in \mathbb{N}$  such that

$$|\alpha - x_n| \leq \epsilon? \quad (13)$$

To answer this question, let us see the *a priori* error bounds

$$M|\alpha - x_n| \leq (M|\alpha - x_0|)^{2^n} \quad \text{Or} \quad |\alpha - x_n| \leq \frac{1}{M}(M|\alpha - x_0|)^{2^n}.$$

# Newton's Method: Stopping Criterion

For a given tolerance  $\epsilon > 0$ , can we find  $n_0 \in \mathbb{N}$  such that

$$|\alpha - x_n| \leq \epsilon? \quad (13)$$

To answer this question, let us see the *a priori* error bounds

$$M|\alpha - x_n| \leq (M|\alpha - x_0|)^{2^n} \quad \text{Or} \quad |\alpha - x_n| \leq \frac{1}{M}(M|\alpha - x_0|)^{2^n}.$$

Hence,  $|\alpha - x_n| \leq \epsilon$  provided

$$\frac{1}{M}(M|\alpha - x_0|)^{2^n} \leq \epsilon.$$

# Newton's Method: Stopping Criterion

For a given tolerance  $\epsilon > 0$ , can we find  $n_0 \in \mathbb{N}$  such that

$$|\alpha - x_n| \leq \epsilon? \quad (13)$$

To answer this question, let us see the *a priori* error bounds

$$M|\alpha - x_n| \leq (M|\alpha - x_0|)^{2^n} \quad \text{Or} \quad |\alpha - x_n| \leq \frac{1}{M}(M|\alpha - x_0|)^{2^n}.$$

Hence,  $|\alpha - x_n| \leq \epsilon$  provided

$$\frac{1}{M}(M|\alpha - x_0|)^{2^n} \leq \epsilon.$$

Thus, finding  $n$  such that relation (13) holds depend on  $x_0$ ,  $\alpha$  and  $M$ .

## Newton's Method: Stopping Criterion

For a given tolerance  $\epsilon > 0$ , can we find  $n_0 \in \mathbb{N}$  such that

$$|\alpha - x_n| \leq \epsilon? \quad (13)$$

To answer this question, let us see the *a priori* error bounds

$$M|\alpha - x_n| \leq (M|\alpha - x_0|)^{2^n} \quad \text{Or} \quad |\alpha - x_n| \leq \frac{1}{M}(M|\alpha - x_0|)^{2^n}.$$

Hence,  $|\alpha - x_n| \leq \epsilon$  provided

$$\frac{1}{M}(M|\alpha - x_0|)^{2^n} \leq \epsilon.$$

Thus, finding  $n$  such that relation (13) holds depend on  $x_0$ ,  $\alpha$  and  $M$ .

But,  $\alpha$  might not be available before computation.

## Newton's Method: Stopping Criterion

For a given tolerance  $\epsilon > 0$ , can we find  $n_0 \in \mathbb{N}$  such that

$$|\alpha - x_n| \leq \epsilon? \quad (13)$$

To answer this question, let us see the *a priori* error bounds

$$M|\alpha - x_n| \leq (M|\alpha - x_0|)^{2^n} \quad \text{Or} \quad |\alpha - x_n| \leq \frac{1}{M}(M|\alpha - x_0|)^{2^n}.$$

Hence,  $|\alpha - x_n| \leq \epsilon$  provided

$$\frac{1}{M}(M|\alpha - x_0|)^{2^n} \leq \epsilon.$$

Thus, finding  $n$  such that relation (13) holds depend on  $x_0$ ,  $\alpha$  and  $M$ .  
But,  $\alpha$  might not be available before computation. What is next?

Remarks:

# Newton's Method: Stopping Criterion

For a given tolerance  $\epsilon > 0$ , can we find  $n_0 \in \mathbb{N}$  such that

$$|\alpha - x_n| \leq \epsilon? \quad (13)$$

To answer this question, let us see the *a priori* error bounds

$$M|\alpha - x_n| \leq (M|\alpha - x_0|)^{2^n} \quad \text{Or} \quad |\alpha - x_n| \leq \frac{1}{M}(M|\alpha - x_0|)^{2^n}.$$

Hence,  $|\alpha - x_n| \leq \epsilon$  provided

$$\frac{1}{M}(M|\alpha - x_0|)^{2^n} \leq \epsilon.$$

Thus, finding  $n$  such that relation (13) holds depend on  $x_0$ ,  $\alpha$  and  $M$ .

But,  $\alpha$  might not be available before computation. What is next?

Remarks:

- Suppose Newton's sequence  $\langle x_n \rangle$  converges to actual root  $\alpha$ .



# Newton's Method: Stopping Criterion

For a given tolerance  $\epsilon > 0$ , can we find  $n_0 \in \mathbb{N}$  such that

$$|\alpha - x_n| \leq \epsilon? \quad (13)$$

To answer this question, let us see the *a priori* error bounds

$$M|\alpha - x_n| \leq (M|\alpha - x_0|)^{2^n} \quad \text{Or} \quad |\alpha - x_n| \leq \frac{1}{M}(M|\alpha - x_0|)^{2^n}.$$

Hence,  $|\alpha - x_n| \leq \epsilon$  provided

$$\frac{1}{M}(M|\alpha - x_0|)^{2^n} \leq \epsilon.$$

Thus, finding  $n$  such that relation (13) holds depend on  $x_0$ ,  $\alpha$  and  $M$ . But,  $\alpha$  might not be available before computation. What is next?

Remarks:

- Suppose Newton's sequence  $\langle x_n \rangle$  converges to actual root  $\alpha$ . Then we may observe following

$$|x_n - x_{n-1}| \leq |x_n - \alpha| + |\alpha - x_{n-1}| \rightarrow 0 \quad \text{as} \quad n \rightarrow \infty.$$

# Newton's Method: Stopping Criterion

For a given tolerance  $\epsilon > 0$ , can we find  $n_0 \in \mathbb{N}$  such that

$$|\alpha - x_n| \leq \epsilon? \quad (13)$$

To answer this question, let us see the *a priori* error bounds

$$M|\alpha - x_n| \leq (M|\alpha - x_0|)^{2^n} \quad \text{Or} \quad |\alpha - x_n| \leq \frac{1}{M}(M|\alpha - x_0|)^{2^n}.$$

Hence,  $|\alpha - x_n| \leq \epsilon$  provided

$$\frac{1}{M}(M|\alpha - x_0|)^{2^n} \leq \epsilon.$$

Thus, finding  $n$  such that relation (13) holds depend on  $x_0$ ,  $\alpha$  and  $M$ . But,  $\alpha$  might not be available before computation. What is next?

Remarks:

- Suppose Newton's sequence  $\langle x_n \rangle$  converges to actual root  $\alpha$ . Then we may observe following

$$|x_n - x_{n-1}| \leq |x_n - \alpha| + |\alpha - x_{n-1}| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

- Therefore, for a given tolerance  $\epsilon$ , an alternative stopping criterion could be

$$|x_n - x_{n-1}| \leq \epsilon.$$

# Newton's Method: Algorithm

# Newton's Method: Algorithm

**Algorithm** *Newton* ( $f, df, x_0, \epsilon, \text{root}, \text{itmax}, \text{ier}$ )

1. Remark:  $df$  is the derivative function  $f'(x)$ ,  $\text{itmax}$  is the maximum number of iterates to be computed, and  $\text{ier}$  is an error flag to the user.
2.  $\text{itnum} := 1$
3.  $\text{denom} := df(x_0)$ .
4. If  $\text{denom} = 0$ , then  $\text{ier} := 2$  and exit.
5.  $x_1 := x_0 - f(x_0)/\text{denom}$
6. If  $|x_1 - x_0| \leq \epsilon$ , then set  $\text{ier} := 0$ ,  $\text{root} := x_1$ , and exit.
7. If  $\text{itnum} = \text{itmax}$ , set  $\text{ier} := 1$  and exit.
8. Otherwise,  $\text{itnum} := \text{itnum} + 1$ ,  $x_0 := x_1$ , and go to step 3.

(Source: An Introduction to Numerical Analysis by Atkinson)

# Newton's Method: Algorithm Contd....

# Newton's Method: Algorithm Contd....

Remark:

# Newton's Method: Algorithm Contd....

Remark:

- In the above algorithm, we have used the fact that Newton's sequence  $\langle x_n \rangle$  converges to the actual root  $\alpha$  and then  $x_n - x_{n-1} \rightarrow 0$ .

# Newton's Method: Algorithm Contd....

Remark:

- In the above algorithm, we have used the fact that Newton's sequence  $\langle x_n \rangle$  converges to the actual root  $\alpha$  and then  $x_n - x_{n-1} \rightarrow 0$ .
- Let  $\langle x_n \rangle$  be the sequence defined by  $x_n = \sum_{k=1}^n \frac{1}{k}$ .



# Newton's Method: Algorithm Contd....

Remark:

- In the above algorithm, we have used the fact that Newton's sequence  $\langle x_n \rangle$  converges to the actual root  $\alpha$  and then  $x_n - x_{n-1} \rightarrow 0$ .
- Let  $\langle x_n \rangle$  be the sequence defined by  $x_n = \sum_{k=1}^n \frac{1}{k}$ . Then verify that  $x_n - x_{n-1} \rightarrow 0$ , but,  $\langle x_n \rangle$  is divergent.

# Newton's Method: Algorithm Contd....

Remark:

- In the above algorithm, we have used the fact that Newton's sequence  $\langle x_n \rangle$  converges to the actual root  $\alpha$  and then  $x_n - x_{n-1} \rightarrow 0$ .
- Let  $\langle x_n \rangle$  be the sequence defined by  $x_n = \sum_{k=1}^n \frac{1}{k}$ . Then verify that  $x_n - x_{n-1} \rightarrow 0$ , but,  $\langle x_n \rangle$  is divergent. More precisely,  $x_n - x_{n-1} \rightarrow 0$  does not imply  $\langle x_n \rangle$  is a convergent sequence.

# Newton's Method: Algorithm Contd....

Remark:

- In the above algorithm, we have used the fact that Newton's sequence  $\langle x_n \rangle$  converges to the actual root  $\alpha$  and then  $x_n - x_{n-1} \rightarrow 0$ .
- Let  $\langle x_n \rangle$  be the sequence defined by  $x_n = \sum_{k=1}^n \frac{1}{k}$ . Then verify that  $x_n - x_{n-1} \rightarrow 0$ , but,  $\langle x_n \rangle$  is divergent. More precisely,  $x_n - x_{n-1} \rightarrow 0$  does not imply  $\langle x_n \rangle$  is a convergent sequence. Can we have both way?

# Newton's Method: Algorithm Contd....

Remark:

- In the above algorithm, we have used the fact that Newton's sequence  $\langle x_n \rangle$  converges to the actual root  $\alpha$  and then  $x_n - x_{n-1} \rightarrow 0$ .
- Let  $\langle x_n \rangle$  be the sequence defined by  $x_n = \sum_{k=1}^n \frac{1}{k}$ . Then verify that  $x_n - x_{n-1} \rightarrow 0$ , but,  $\langle x_n \rangle$  is divergent. More precisely,  $x_n - x_{n-1} \rightarrow 0$  does not imply  $\langle x_n \rangle$  is a convergent sequence. Can we have both way?

Motivation:

# Newton's Method: Algorithm Contd....

Remark:

- In the above algorithm, we have used the fact that Newton's sequence  $\langle x_n \rangle$  converges to the actual root  $\alpha$  and then  $x_n - x_{n-1} \rightarrow 0$ .
- Let  $\langle x_n \rangle$  be the sequence defined by  $x_n = \sum_{k=1}^n \frac{1}{k}$ . Then verify that  $x_n - x_{n-1} \rightarrow 0$ , but,  $\langle x_n \rangle$  is divergent. More precisely,  $x_n - x_{n-1} \rightarrow 0$  does not imply  $\langle x_n \rangle$  is a convergent sequence. Can we have both way?

Motivation:

- Using the mean value theorem,

$$f(x_n) = f(x_n) - f(\alpha) =$$

# Newton's Method: Algorithm Contd....

Remark:

- In the above algorithm, we have used the fact that Newton's sequence  $\langle x_n \rangle$  converges to the actual root  $\alpha$  and then  $x_n - x_{n-1} \rightarrow 0$ .
- Let  $\langle x_n \rangle$  be the sequence defined by  $x_n = \sum_{k=1}^n \frac{1}{k}$ . Then verify that  $x_n - x_{n-1} \rightarrow 0$ , but,  $\langle x_n \rangle$  is divergent. More precisely,  $x_n - x_{n-1} \rightarrow 0$  does not imply  $\langle x_n \rangle$  is a convergent sequence. Can we have both way?

Motivation:

- Using the mean value theorem,

$$f(x_n) - f(\alpha) = f'(\xi_n)(x_n - \alpha)$$

# Newton's Method: Algorithm Contd....

Remark:

- In the above algorithm, we have used the fact that Newton's sequence  $\langle x_n \rangle$  converges to the actual root  $\alpha$  and then  $x_n - x_{n-1} \rightarrow 0$ .
- Let  $\langle x_n \rangle$  be the sequence defined by  $x_n = \sum_{k=1}^n \frac{1}{k}$ . Then verify that  $x_n - x_{n-1} \rightarrow 0$ , but,  $\langle x_n \rangle$  is divergent. More precisely,  $x_n - x_{n-1} \rightarrow 0$  does not imply  $\langle x_n \rangle$  is a convergent sequence. Can we have both way?

Motivation:

- Using the mean value theorem,

$$f(x_n) = f(x_n) - f(\alpha) = f'(\xi_n)(x_n - \alpha) \quad \text{Or} \quad \alpha - x_n = \frac{-f(x_n)}{f'(\xi_n)}, \quad (14)$$

where  $\xi_n$  between  $x_n$  and  $\alpha$ .

# Newton's Method: Algorithm Contd....

Remark:

- In the above algorithm, we have used the fact that Newton's sequence  $\langle x_n \rangle$  converges to the actual root  $\alpha$  and then  $x_n - x_{n-1} \rightarrow 0$ .
- Let  $\langle x_n \rangle$  be the sequence defined by  $x_n = \sum_{k=1}^n \frac{1}{k}$ . Then verify that  $x_n - x_{n-1} \rightarrow 0$ , but,  $\langle x_n \rangle$  is divergent. More precisely,  $x_n - x_{n-1} \rightarrow 0$  does not imply  $\langle x_n \rangle$  is a convergent sequence. Can we have both way?

Motivation:

- Using the mean value theorem,

$$f(x_n) = f(x_n) - f(\alpha) = f'(\xi_n)(x_n - \alpha) \quad \text{Or} \quad \alpha - x_n = \frac{-f(x_n)}{f'(\xi_n)}, \quad (14)$$

where  $\xi_n$  between  $x_n$  and  $\alpha$ . Then we can expect to approximate  $f'(\xi_n)$  by  $f'(x_n)$ , and

$$\alpha - x_n \approx \frac{-f(x_n)}{f'(x_n)} =$$



# Newton's Method: Algorithm Contd....

Remark:

- In the above algorithm, we have used the fact that Newton's sequence  $\langle x_n \rangle$  converges to the actual root  $\alpha$  and then  $x_n - x_{n-1} \rightarrow 0$ .
- Let  $\langle x_n \rangle$  be the sequence defined by  $x_n = \sum_{k=1}^n \frac{1}{k}$ . Then verify that  $x_n - x_{n-1} \rightarrow 0$ , but,  $\langle x_n \rangle$  is divergent. More precisely,  $x_n - x_{n-1} \rightarrow 0$  does not imply  $\langle x_n \rangle$  is a convergent sequence. Can we have both way?

Motivation:

- Using the mean value theorem,

$$f(x_n) = f(x_n) - f(\alpha) = f'(\xi_n)(x_n - \alpha) \quad \text{Or} \quad \alpha - x_n = \frac{-f(x_n)}{f'(\xi_n)}, \quad (14)$$

where  $\xi_n$  between  $x_n$  and  $\alpha$ . Then we can expect to approximate  $f'(\xi_n)$  by  $f'(x_n)$ , and

$$\alpha - x_n \approx \frac{-f(x_n)}{f'(x_n)} = x_{n+1} - x_n.$$

# Newton's Method: Algorithm Contd....

Remark:

- In the above algorithm, we have used the fact that Newton's sequence  $\langle x_n \rangle$  converges to the actual root  $\alpha$  and then  $x_n - x_{n-1} \rightarrow 0$ .
- Let  $\langle x_n \rangle$  be the sequence defined by  $x_n = \sum_{k=1}^n \frac{1}{k}$ . Then verify that  $x_n - x_{n-1} \rightarrow 0$ , but,  $\langle x_n \rangle$  is divergent. More precisely,  $x_n - x_{n-1} \rightarrow 0$  does not imply  $\langle x_n \rangle$  is a convergent sequence. Can we have both way?

Motivation:

- Using the mean value theorem,

$$f(x_n) = f(x_n) - f(\alpha) = f'(\xi_n)(x_n - \alpha) \quad \text{Or} \quad \alpha - x_n = \frac{-f(x_n)}{f'(\xi_n)}, \quad (14)$$

where  $\xi_n$  between  $x_n$  and  $\alpha$ . Then we can expect to approximate  $f'(\xi_n)$  by  $f'(x_n)$ , and

$$\alpha - x_n \approx \frac{-f(x_n)}{f'(x_n)} = x_{n+1} - x_n.$$

- Thus, for Newton's method, the standard error is  $\alpha - x_n \approx x_{n+1} - x_n$ .

# Newton's Method: Algorithm Contd....

Remark:

- In the above algorithm, we have used the fact that Newton's sequence  $\langle x_n \rangle$  converges to the actual root  $\alpha$  and then  $x_n - x_{n-1} \rightarrow 0$ .
- Let  $\langle x_n \rangle$  be the sequence defined by  $x_n = \sum_{k=1}^n \frac{1}{k}$ . Then verify that  $x_n - x_{n-1} \rightarrow 0$ , but,  $\langle x_n \rangle$  is divergent. More precisely,  $x_n - x_{n-1} \rightarrow 0$  does not imply  $\langle x_n \rangle$  is a convergent sequence. Can we have both way?

Motivation:

- Using the mean value theorem,

$$f(x_n) = f(x_n) - f(\alpha) = f'(\xi_n)(x_n - \alpha) \quad \text{Or} \quad \alpha - x_n = \frac{-f(x_n)}{f'(\xi_n)}, \quad (14)$$

where  $\xi_n$  between  $x_n$  and  $\alpha$ . Then we can expect to approximate  $f'(\xi_n)$  by  $f'(x_n)$ , and

$$\alpha - x_n \approx \frac{-f(x_n)}{f'(x_n)} = x_{n+1} - x_n.$$

- Thus, for Newton's method, the standard error is  $\alpha - x_n \approx x_{n+1} - x_n$ .

# The Secant Method

# The Secant Method

Regarding the Newton's method, we observe following:

# The Secant Method

Regarding the Newton's method, we observe following:

- When Newton's method converges, it generally does so quite rapidly, an advantage over the bisection method.

# The Secant Method

Regarding the Newton's method, we observe following:

- When Newton's method converges, it generally does so quite rapidly, an advantage over the bisection method. But again, it need not converge.

# The Secant Method

Regarding the Newton's method, we observe following:

- When Newton's method converges, it generally does so quite rapidly, an advantage over the bisection method. But again, it need not converge.
- Another source of difficulty in some cases is the necessity of knowing  $f'(x)$  explicitly.



# The Secant Method

Regarding the Newton's method, we observe following:

- When Newton's method converges, it generally does so quite rapidly, an advantage over the bisection method. But again, it need not converge.
- Another source of difficulty in some cases is the necessity of knowing  $f'(x)$  explicitly. With some root finding problems, this is not possible.

# The Secant Method

Regarding the Newton's method, we observe following:

- When Newton's method converges, it generally does so quite rapidly, an advantage over the bisection method. But again, it need not converge.
- Another source of difficulty in some cases is the necessity of knowing  $f'(x)$  explicitly. With some root finding problems, this is not possible.


Remark: Then we introduce an another method 'Secant Method' remedies this situation, at the cost of a somewhat slower speed of convergence.

# The Secant Method

Regarding the Newton's method, we observe following:

- When Newton's method converges, it generally does so quite rapidly, an advantage over the bisection method. But again, it need not converge.
- Another source of difficulty in some cases is the necessity of knowing  $f'(x)$  explicitly. With some root finding problems, this is not possible.

Remark: Then we introduce an another method 'Secant Method' remedies this situation, at the cost of a somewhat slower speed of convergence.



The secant method.


Assume  $x_0$  and  $x_1$  are two initial estimates of the root  $\alpha$ .

# The Secant Method

Regarding the Newton's method, we observe following:

- When Newton's method converges, it generally does so quite rapidly, an advantage over the bisection method. But again, it need not converge.
- Another source of difficulty in some cases is the necessity of knowing  $f'(x)$  explicitly. With some root finding problems, this is not possible.

Remark: Then we introduce an another method 'Secant Method' remedies this situation, at the cost of a somewhat slower speed of convergence.



The secant method.


Assume  $x_0$  and  $x_1$  are two initial estimates of the root  $\alpha$ . Consider the secant line passing through  $(x_0, f(x_0))$  and  $(x_1, f(x_1))$ .

# The Secant Method

Regarding the Newton's method, we observe following:

- When Newton's method converges, it generally does so quite rapidly, an advantage over the bisection method. But again, it need not converge.
- Another source of difficulty in some cases is the necessity of knowing  $f'(x)$  explicitly. With some root finding problems, this is not possible.

Remark: Then we introduce an another method 'Secant Method' remedies this situation, at the cost of a somewhat slower speed of convergence.



The secant method.

Assume  $x_0$  and  $x_1$  are two initial estimates of the root  $\alpha$ . Consider the secant line passing through  $(x_0, f(x_0))$  and  $(x_1, f(x_1))$ . Then, we obtain

$$x_2 = x_1 - f(x_1) \cdot \frac{x_1 - x_0}{f(x_1) - f(x_0)}.$$

# The Secant Method Contd....

## The Secant Method Contd....

In general,

$$x_{n+1} = x_n - f(x_n) \cdot \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})}. \quad (15)$$

# The Secant Method Contd....

In general,

$$x_{n+1} = x_n - f(x_n) \cdot \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})}. \quad (15)$$

Comparison of Newton's method and Secant method:



# The Secant Method Contd....

In general,

$$x_{n+1} = x_n - f(x_n) \cdot \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})}. \quad (15)$$

Comparison of Newton's method and Secant method:

- Newton's method and the secant method are closely related.

# The Secant Method Contd....

In general,

$$x_{n+1} = x_n - f(x_n) \cdot \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})}. \quad (15)$$

Comparison of Newton's method and Secant method:

- Newton's method and the secant method are closely related. If the approximation

$$f'(x_n) \approx \frac{f(x_n) - f(x_{n-1})}{x_n - x_{n-1}} \quad (16)$$

is used in the Newton formula, we obtain the secant formula.

# The Secant Method Contd....

In general,

$$x_{n+1} = x_n - f(x_n) \cdot \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})}. \quad (15)$$

Comparison of Newton's method and Secant method:

- Newton's method and the secant method are closely related. If the approximation

$$f'(x_n) \approx \frac{f(x_n) - f(x_{n-1})}{x_n - x_{n-1}} \quad (16)$$

is used in the Newton formula, we obtain the secant formula. But, above approximation is not quadratic.

# The Secant Method Contd....

In general,

$$x_{n+1} = x_n - f(x_n) \cdot \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})}. \quad (15)$$

Comparison of Newton's method and Secant method:

- Newton's method and the secant method are closely related. If the approximation

$$f'(x_n) \approx \frac{f(x_n) - f(x_{n-1})}{x_n - x_{n-1}} \quad (16)$$

is used in the Newton formula, we obtain the secant formula. But, above approximation is not quadratic. Hence, we can not expect quadratic convergence for secant method.

# The Secant Method Contd....

In general,

$$x_{n+1} = x_n - f(x_n) \cdot \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})}. \quad (15)$$

Comparison of Newton's method and Secant method:

- Newton's method and the secant method are closely related. If the approximation

$$f'(x_n) \approx \frac{f(x_n) - f(x_{n-1})}{x_n - x_{n-1}} \quad (16)$$

is used in the Newton formula, we obtain the secant formula. But, above approximation is not quadratic. Hence, we can not expect quadratic convergence for secant method.

Convergence of Secant's Method:

# The Secant Method Contd....

In general,

$$x_{n+1} = x_n - f(x_n) \cdot \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})}. \quad (15)$$

Comparison of Newton's method and Secant method:

- Newton's method and the secant method are closely related. If the approximation

$$f'(x_n) \approx \frac{f(x_n) - f(x_{n-1})}{x_n - x_{n-1}} \quad (16)$$

is used in the Newton formula, we obtain the secant formula. But, above approximation is not quadratic. Hence, we can not expect quadratic convergence for secant method.

Convergence of Secant's Method: Assume  $f$ ,  $f'$  and  $f''$  are continuous for all  $x$  in some neighborhood of  $\alpha$ , and assume that  $f(\alpha) = 0$ ,  $f'(\alpha) \neq 0$ . Then if  $x_0$  and  $x_1$  are chosen sufficiently close to  $\alpha$ , the sequence  $\langle x_n \rangle$  given by

$$x_{n+1} = x_n - f(x_n) \cdot \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})}$$

will converge to  $\alpha$ .

# The Secant Method Contd....

# The Secant Method Contd....

Example: Find an approximation for the largest root  $\alpha \approx 1.1347241$  of the equation

$$x^6 - x - 1 = 0.$$

## Secant method

$n$	$x_n$
0	2.0
1	1.0
2	1.016129032
3	1.190577769
4	1.117655831
5	1.132531550
6	1.134816808
7	1.134723646



# The Secant Method Contd....

Example: Find an approximation for the largest root  $\alpha \approx 1.1347241$  of the equation

$$x^6 - x - 1 = 0.$$

Secant method	
$n$	$x_n$
0	2.0
1	1.0
2	1.016129032
3	1.190577769
4	1.117655831
5	1.132531550
6	1.134816808
7	1.134723646

Newton's method	
$n$	$x_n$
0	2.0
1	1.680628273
2	1.430738989
3	1.254970957
4	1.161538433
5	1.136353274
6	1.134730528
7	1.134724139

Bisection method			
$n$	$c_n$	$n$	$c_n$
$2=b, 1=a$		8	1.13672
1	1.5	9	1.13477
2	1.25	10	1.13379
3	1.125	11	1.13428
4	1.1875	12	1.13452
5	1.15625	13	1.13464
6	1.14063	14	1.13470
7	1.13281	15	1.13474

(Source: An Introduction to Numerical Analysis by K. E. Atkinson)

## The Secant Method Contd....

Example: Find an approximation for the largest root  $\alpha \approx 1.1347241$  of the equation

$$x^6 - x - 1 = 0.$$

Secant method	
$n$	$x_n$
0	2.0
1	1.0
2	1.016129032
3	1.190577769
4	1.117655831
5	1.132531550
6	1.134816808
7	1.134723646

Newton's method	
$n$	$x_n$
0	2.0
1	1.680628273
2	1.430738989
3	1.254970957
4	1.161538433
5	1.136353274
6	1.134730528
7	1.134724139

Bisection method			
$n$	$c_n$	$n$	$c_n$
$2 = b, 1 = a$		8	1.13672
1	1.5	9	1.13477
2	1.25	10	1.13379
3	1.125	11	1.13428
4	1.1875	12	1.13452
5	1.15625	13	1.13464
6	1.14063	14	1.13470
7	1.13281	15	1.13474

(Source: An Introduction to Numerical Analysis by K. E. Atkinson)

Remark:

## The Secant Method Contd....

Example: Find an approximation for the largest root  $\alpha \approx 1.1347241$  of the equation

$$x^6 - x - 1 = 0.$$

Secant method	
$n$	$x_n$
0	2.0
1	1.0
2	1.016129032
3	1.190577769
4	1.117655831
5	1.132531550
6	1.134816808
7	1.134723646

Newton's method	
$n$	$x_n$
0	2.0
1	1.680628273
2	1.430738989
3	1.254970957
4	1.161538433
5	1.136353274
6	1.134730528
7	1.134724139

Bisection method			
$n$	$c_n$	$n$	$c_n$
$2 = b, 1 = a$		8	1.13672
1	1.5	9	1.13477
2	1.25	10	1.13379
3	1.125	11	1.13428
4	1.1875	12	1.13452
5	1.15625	13	1.13464
6	1.14063	14	1.13470
7	1.13281	15	1.13474

(Source: An Introduction to Numerical Analysis by K. E. Atkinson)

Remark:

- It is better than Bisection method but not good as Newton's method.

# Order of Convergence for Secant Method

# Order of Convergence for Secant Method

- At this point we simply note, without proof, that an error formula for the secant method can be derived, which says that

$$\alpha - x_{n+1} = C_n(\alpha - x_n)(\alpha - x_{n-1}) \quad \text{Or} \quad |e_{n+1}| = C_n|e_n e_{n-1}|. \quad (17)$$

# Order of Convergence for Secant Method

- At this point we simply note, without proof, that an error formula for the secant method can be derived, which says that

$$\alpha - x_{n+1} = C_n(\alpha - x_n)(\alpha - x_{n-1}) \text{ Or } |e_{n+1}| = C_n|e_n e_{n-1}|. \quad (17)$$

- For the order of convergence, we seek a number  $p > 0$  such that

$$\lim_{n \rightarrow \infty} \frac{|e_{n+1}|}{|e_n|^p} = C \quad (18)$$

## Order of Convergence for Secant Method

● At this point we simply note, without proof, that an error formula for the secant method can be derived, which says that

$$\alpha - x_{n+1} = C_n(\alpha - x_n)(\alpha - x_{n-1}) \text{ Or } |e_{n+1}| = C_n|e_n e_{n-1}|. \quad (17)$$

● For the order of convergence, we seek a number  $p > 0$  such that

$$\lim_{n \rightarrow \infty} \frac{|e_{n+1}|}{|e_n|^p} = C \quad (18)$$

Then equation (17) yields

$$\frac{|e_{n+1}|}{|e_n|^p} = C_n |e_n|^{1-p} |e_{n-1}| = C_n \left( \frac{|e_n|}{|e_{n-1}|^p} \right)^\alpha, \quad (19)$$

provided  $\alpha = 1 - p$  and  $\alpha p = -1$ .

## Order of Convergence for Secant Method

● At this point we simply note, without proof, that an error formula for the secant method can be derived, which says that

$$\alpha - x_{n+1} = C_n(\alpha - x_n)(\alpha - x_{n-1}) \text{ Or } |e_{n+1}| = C_n|e_n e_{n-1}|. \quad (17)$$

● For the order of convergence, we seek a number  $p > 0$  such that

$$\lim_{n \rightarrow \infty} \frac{|e_{n+1}|}{|e_n|^p} = C \quad (18)$$

Then equation (17) yields

$$\frac{|e_{n+1}|}{|e_n|^p} = C_n |e_n|^{1-p} |e_{n-1}| = C_n \left( \frac{|e_n|}{|e_{n-1}|^p} \right)^\alpha, \quad (19)$$

provided  $\alpha = 1 - p$  and  $\alpha p = -1$ . Solving for  $p$ , we obtain

$$p - p^2 = -1$$



## Order of Convergence for Secant Method

● At this point we simply note, without proof, that an error formula for the secant method can be derived, which says that

$$\alpha - x_{n+1} = C_n(\alpha - x_n)(\alpha - x_{n-1}) \text{ Or } |e_{n+1}| = C_n|e_n e_{n-1}|. \quad (17)$$

● For the order of convergence, we seek a number  $p > 0$  such that

$$\lim_{n \rightarrow \infty} \frac{|e_{n+1}|}{|e_n|^p} = C \quad (18)$$

Then equation (17) yields

$$\frac{|e_{n+1}|}{|e_n|^p} = C_n |e_n|^{1-p} |e_{n-1}| = C_n \left( \frac{|e_n|}{|e_{n-1}|^p} \right)^\alpha, \quad (19)$$

provided  $\alpha = 1 - p$  and  $\alpha p = -1$ . Solving for  $p$ , we obtain

$$p - p^2 = -1 \text{ Or } p^2 - p - 1 = 0. \quad (20)$$

## Order of Convergence for Secant Method

● At this point we simply note, without proof, that an error formula for the secant method can be derived, which says that

$$\alpha - x_{n+1} = C_n(\alpha - x_n)(\alpha - x_{n-1}) \text{ Or } |e_{n+1}| = C_n|e_n e_{n-1}|. \quad (17)$$

● For the order of convergence, we seek a number  $p > 0$  such that

$$\lim_{n \rightarrow \infty} \frac{|e_{n+1}|}{|e_n|^p} = C \quad (18)$$

Then equation (17) yields

$$\frac{|e_{n+1}|}{|e_n|^p} = C_n |e_n|^{1-p} |e_{n-1}| = C_n \left( \frac{|e_n|}{|e_{n-1}|^p} \right)^\alpha, \quad (19)$$

provided  $\alpha = 1 - p$  and  $\alpha p = -1$ . Solving for  $p$ , we obtain

$$p - p^2 = -1 \text{ Or } p^2 - p - 1 = 0. \quad (20)$$

Solving above equation we find that

$$p = \frac{1 \pm \sqrt{1+4}}{2}.$$

## Order of Convergence for Secant Method

● At this point we simply note, without proof, that an error formula for the secant method can be derived, which says that

$$\alpha - x_{n+1} = C_n(\alpha - x_n)(\alpha - x_{n-1}) \text{ Or } |e_{n+1}| = C_n|e_n e_{n-1}|. \quad (17)$$

● For the order of convergence, we seek a number  $p > 0$  such that

$$\lim_{n \rightarrow \infty} \frac{|e_{n+1}|}{|e_n|^p} = C \quad (18)$$

Then equation (17) yields

$$\frac{|e_{n+1}|}{|e_n|^p} = C_n |e_n|^{1-p} |e_{n-1}| = C_n \left( \frac{|e_n|}{|e_{n-1}|^p} \right)^\alpha, \quad (19)$$

provided  $\alpha = 1 - p$  and  $\alpha p = -1$ . Solving for  $p$ , we obtain

$$p - p^2 = -1 \text{ Or } p^2 - p - 1 = 0. \quad (20)$$

Solving above equation we find that

$$p = \frac{1 \pm \sqrt{1+4}}{2}.$$

Thus, secant method converges with order  $p = \frac{1+\sqrt{5}}{2} = 1.618\dots$