

Cholesky Decomposition for Positive Definite Matrices

Positive Definite Matrices

Positive Definite Matrices: An $n \times n$ real matrix A is said to be (symmetric) positive definite if the following hold:

1. A is symmetric, that is, $A^T = A$;
2. $x^T A x > 0$ for all $x \in \mathbb{R}^n$ with $x \neq 0$.

Positive Definite Matrices

Positive Definite Matrices: An $n \times n$ real matrix A is said to be (symmetric) positive definite if the following hold:

1. A is symmetric, that is, $A^T = A$;
2. $x^T A x > 0$ for all $x \in \mathbb{R}^n$ with $x \neq 0$.

In the literature very often positive definite matrices are defined as those satisfying only 2. Such matrices need not be symmetric.

Positive Definite Matrices

Positive Definite Matrices: An $n \times n$ real matrix A is said to be (symmetric) positive definite if the following hold:

1. A is symmetric, that is, $A^T = A$;
2. $x^T A x > 0$ for all $x \in \mathbb{R}^n$ with $x \neq 0$.

In the literature very often positive definite matrices are defined as those satisfying only 2. Such matrices need not be symmetric.

Find an example!

Positive Definite Matrices

Positive Definite Matrices: An $n \times n$ real matrix A is said to be (symmetric) positive definite if the following hold:

1. A is symmetric, that is, $A^T = A$;
2. $x^T A x > 0$ for all $x \in \mathbb{R}^n$ with $x \neq 0$.

In the literature very often positive definite matrices are defined as those satisfying only 2. Such matrices need not be symmetric.

Find an example!

However, in application real matrices that satisfy 2 are most often also symmetric. In accordance with the textbook, we will refer to symmetric positive definite matrices as (just) positive definite matrices.

Positive Definite Matrices

Positive Definite Matrices: An $n \times n$ real matrix A is said to be (symmetric) positive definite if the following hold:

1. A is symmetric, that is, $A^T = A$;
2. $x^T A x > 0$ for all $x \in \mathbb{R}^n$ with $x \neq 0$.

In the literature very often positive definite matrices are defined as those satisfying only 2. Such matrices need not be symmetric.

Find an example!

However, in application real matrices that satisfy 2 are most often also symmetric. In accordance with the textbook, we will refer to symmetric positive definite matrices as (just) positive definite matrices.

The counterpart of transpose (T) for complex matrices is complex conjugate transpose (*). If an $n \times n$ matrix is non real, that is it is a complex matrix with non zero imaginary part, then 2 with T replaced by * implies that $A^* = A$, that is, A is Hermitian.

Positive Definite Matrices

Positive Definite Matrices: An $n \times n$ real matrix A is said to be (symmetric) positive definite if the following hold:

1. A is symmetric, that is, $A^T = A$;
2. $x^T A x > 0$ for all $x \in \mathbb{R}^n$ with $x \neq 0$.

In the literature very often positive definite matrices are defined as those satisfying only 2. Such matrices need not be symmetric.

Find an example!

However, in application real matrices that satisfy 2 are most often also symmetric. In accordance with the textbook, we will refer to symmetric positive definite matrices as (just) positive definite matrices.

The counterpart of transpose (T) for complex matrices is complex conjugate transpose (*). If an $n \times n$ matrix is non real, that is it is a complex matrix with non zero imaginary part, then 2 with T replaced by * implies that $A^* = A$, that is, A is Hermitian. Prove this!

Positive Definite Matrices

Positive Definite Matrices: An $n \times n$ real matrix A is said to be (symmetric) positive definite if the following hold:

1. A is symmetric, that is, $A^T = A$;
2. $x^T A x > 0$ for all $x \in \mathbb{R}^n$ with $x \neq 0$.

In the literature very often positive definite matrices are defined as those satisfying only 2. Such matrices need not be symmetric.

Find an example!

However, in application real matrices that satisfy 2 are most often also symmetric. In accordance with the textbook, we will refer to symmetric positive definite matrices as (just) positive definite matrices.

The counterpart of transpose (T) for complex matrices is complex conjugate transpose (*). If an $n \times n$ matrix is non real, that is it is a complex matrix with non zero imaginary part, then 2 with T replaced by * implies that $A^* = A$, that is, A is Hermitian. Prove this!

Therefore if A is an $n \times n$ non real matrix, then A is said to be positive definite if $x^* A x > 0$ for all nonzero $x \in \mathbb{C}^n$.

Properties and Applications

Positive definite matrices arise in many applications involving optimization and discretization of partial differential equations. Consider systems $Ax = b$.

Properties and Applications

Positive definite matrices arise in many applications involving optimization and discretization of partial differential equations. Consider systems $Ax = b$.

- ▶ When associated with electrical circuits with x representing loop currents, $x^T Ax$ is the total power drawn by the resistors.
- ▶ When associated with mass spring systems, $(x^T Ax)/2$ is the strain energy of the system.

Properties and Applications

Positive definite matrices arise in many applications involving optimization and discretization of partial differential equations. Consider systems $Ax = b$.

- ▶ When associated with electrical circuits with x representing loop currents, $x^T Ax$ is the total power drawn by the resistors.
- ▶ When associated with mass spring systems, $(x^T Ax)/2$ is the strain energy of the system.

(For examples see Section 1.2 of *D. S. Watkins, Fundamentals of Matrix Computations (second edition)*)

Properties and Applications

Positive definite matrices arise in many applications involving optimization and discretization of partial differential equations. Consider systems $Ax = b$.

- ▶ When associated with electrical circuits with x representing loop currents, $x^T Ax$ is the total power drawn by the resistors.
- ▶ When associated with mass spring systems, $(x^T Ax)/2$ is the strain energy of the system.

(For examples see Section 1.2 of *D. S. Watkins, Fundamentals of Matrix Computations (second edition)*)

Important properties:

Properties and Applications

Positive definite matrices arise in many applications involving optimization and discretization of partial differential equations. Consider systems $Ax = b$.

- ▶ When associated with electrical circuits with x representing loop currents, $x^T Ax$ is the total power drawn by the resistors.
- ▶ When associated with mass spring systems, $(x^T Ax)/2$ is the strain energy of the system.

(For examples see Section 1.2 of *D. S. Watkins, Fundamentals of Matrix Computations (second edition)*)

Important properties:

- ▶ A positive definite matrix is a nonsingular matrix.

Properties and Applications

Positive definite matrices arise in many applications involving optimization and discretization of partial differential equations. Consider systems $Ax = b$.

- ▶ When associated with electrical circuits with x representing loop currents, $x^T Ax$ is the total power drawn by the resistors.
- ▶ When associated with mass spring systems, $(x^T Ax)/2$ is the strain energy of the system.

(For examples see Section 1.2 of *D. S. Watkins, Fundamentals of Matrix Computations (second edition)*)

Important properties:

- ▶ A positive definite matrix is a nonsingular matrix.
- ▶ If A is an $n \times n$ positive definite matrix, then for any nonsingular $n \times n$ matrix X , $X^T AX$ is also positive definite.

Properties and Applications

Positive definite matrices arise in many applications involving optimization and discretization of partial differential equations. Consider systems $Ax = b$.

- ▶ When associated with electrical circuits with x representing loop currents, $x^T Ax$ is the total power drawn by the resistors.
- ▶ When associated with mass spring systems, $(x^T Ax)/2$ is the strain energy of the system.

(For examples see Section 1.2 of *D. S. Watkins, Fundamentals of Matrix Computations (second edition)*)

Important properties:

- ▶ A positive definite matrix is a nonsingular matrix.
- ▶ If A is an $n \times n$ positive definite matrix, then for any nonsingular $n \times n$ matrix X , $X^T AX$ is also positive definite.
- ▶ All principal submatrices of positive definite matrices are positive definite.

Properties and Applications

Positive definite matrices arise in many applications involving optimization and discretization of partial differential equations. Consider systems $Ax = b$.

- ▶ When associated with electrical circuits with x representing loop currents, $x^T Ax$ is the total power drawn by the resistors.
- ▶ When associated with mass spring systems, $(x^T Ax)/2$ is the strain energy of the system.

(For examples see Section 1.2 of *D. S. Watkins, Fundamentals of Matrix Computations (second edition)*)

Important properties:

- ▶ A positive definite matrix is a nonsingular matrix.
- ▶ If A is an $n \times n$ positive definite matrix, then for any nonsingular $n \times n$ matrix X , $X^T AX$ is also positive definite.
- ▶ All principal submatrices of positive definite matrices are positive definite.

To know more about the theory and applications of positive definite matrices, check out the following link:

[MIT OCW on Symmetric Positive Definite Matrices](#)

Cholesky Decomposition for Positive Definite Matrices

Let A be a positive definite matrix. Then there exists a unique upper triangular matrix G with positive diagonal entries such that $A = G^T G$. This is called the **Cholesky Decomposition** of A and G is called the Cholesky factor of A .

Cholesky Decomposition for Positive Definite Matrices

Let A be a positive definite matrix. Then there exists a unique upper triangular matrix G with positive diagonal entries such that $A = G^T G$. This is called the **Cholesky Decomposition** of A and G is called the Cholesky factor of A .



André-Louis Cholesky
(1875-1918)

Cholesky Decomposition Theorem

Theorem An $n \times n$ matrix A is positive definite if and only if there exists an upper triangular matrix G with positive diagonal entries such that

$$A = G^T G.$$

Cholesky Decomposition Theorem

Theorem An $n \times n$ matrix A is positive definite if and only if there exists an upper triangular matrix G with positive diagonal entries such that

$$A = G^T G.$$

Proof: Suppose $A = G^T G$ for some upper triangular matrix G with positive diagonal entries. Then

$$A^T = (G^T G)^T = G^T G = A.$$

For any $x \neq 0$, $x^T A x = x^T G^T G x = (Gx)^T (Gx) = \|Gx\|_2^2 \geq 0$.

Cholesky Decomposition Theorem

Theorem An $n \times n$ matrix A is positive definite if and only if there exists an upper triangular matrix G with positive diagonal entries such that

$$A = G^T G.$$

Proof: Suppose $A = G^T G$ for some upper triangular matrix G with positive diagonal entries. Then

$$A^T = (G^T G)^T = G^T G = A.$$

For any $x \neq 0$, $x^T A x = x^T G^T G x = (Gx)^T (Gx) = \|Gx\|_2^2 \geq 0$.

As G is upper triangular with positive diagonal entries, it is nonsingular.

Therefore as $x \neq 0$, $Gx \neq 0 \Rightarrow \|Gx\|_2 > 0$ and from above, $x^T A x > 0$ for all $x \neq 0$.

Cholesky Decomposition Theorem

Theorem An $n \times n$ matrix A is positive definite if and only if there exists an upper triangular matrix G with positive diagonal entries such that

$$A = G^T G.$$

Proof: Suppose $A = G^T G$ for some upper triangular matrix G with positive diagonal entries. Then

$$A^T = (G^T G)^T = G^T G = A.$$

For any $x \neq 0$, $x^T A x = x^T G^T G x = (Gx)^T (Gx) = \|Gx\|_2^2 \geq 0$.

As G is upper triangular with positive diagonal entries, it is nonsingular.

Therefore as $x \neq 0$, $Gx \neq 0 \Rightarrow \|Gx\|_2 > 0$ and from above, $x^T A x > 0$ for all $x \neq 0$.

Conversely, suppose A is an $n \times n$ positive definite matrix. Then A is nonsingular and all its leading principal submatrices are also positive definite and hence nonsingular. Moreover $x^T A x > 0$ for all $x \in \mathbb{R}^n \setminus \{0\}$.

Cholesky Decomposition Theorem

Theorem An $n \times n$ matrix A is positive definite if and only if there exists an upper triangular matrix G with positive diagonal entries such that

$$A = G^T G.$$

Proof: Suppose $A = G^T G$ for some upper triangular matrix G with positive diagonal entries. Then

$$A^T = (G^T G)^T = G^T G = A.$$

For any $x \neq 0$, $x^T A x = x^T G^T G x = (Gx)^T (Gx) = \|Gx\|_2^2 \geq 0$.

As G is upper triangular with positive diagonal entries, it is nonsingular. Therefore as $x \neq 0$, $Gx \neq 0 \Rightarrow \|Gx\|_2 > 0$ and from above, $x^T A x > 0$ for all $x \neq 0$.

Conversely, suppose A is an $n \times n$ positive definite matrix. Then A is nonsingular and all its leading principal submatrices are also positive definite and hence nonsingular. Moreover $x^T A x > 0$ for all $x \in \mathbb{R}^n \setminus \{0\}$.

Therefore, there is a unique unit lower triangular matrix L and a matrix $D = \text{diag}(d_{11}, \dots, d_{nn})$ $d_{ii} > 0$, such that $A = LDL^T$.

Cholesky Decomposition Theorem

Theorem An $n \times n$ matrix A is positive definite if and only if there exists an upper triangular matrix G with positive diagonal entries such that

$$A = G^T G.$$

Proof: Suppose $A = G^T G$ for some upper triangular matrix G with positive diagonal entries. Then

$$A^T = (G^T G)^T = G^T G = A.$$

For any $x \neq 0$, $x^T A x = x^T G^T G x = (Gx)^T (Gx) = \|Gx\|_2^2 \geq 0$.

As G is upper triangular with positive diagonal entries, it is nonsingular. Therefore as $x \neq 0$, $Gx \neq 0 \Rightarrow \|Gx\|_2 > 0$ and from above, $x^T A x > 0$ for all $x \neq 0$.

Conversely, suppose A is an $n \times n$ positive definite matrix. Then A is nonsingular and all its leading principal submatrices are also positive definite and hence nonsingular. Moreover $x^T A x > 0$ for all $x \in \mathbb{R}^n \setminus \{0\}$.

Therefore, there is a unique unit lower triangular matrix L and a matrix $D = \text{diag}(d_{11}, \dots, d_{nn})$ $d_{ii} > 0$, such that $A = LDL^T$. Let $G = D^{1/2} L^T$ where, $D^{1/2} = \text{diag}(\sqrt{d_{11}}, \dots, \sqrt{d_{nn}})$.

Cholesky Decomposition Theorem

Theorem An $n \times n$ matrix A is positive definite if and only if there exists an upper triangular matrix G with positive diagonal entries such that

$$A = G^T G.$$

Proof: Suppose $A = G^T G$ for some upper triangular matrix G with positive diagonal entries. Then

$$A^T = (G^T G)^T = G^T G = A.$$

For any $x \neq 0$, $x^T A x = x^T G^T G x = (Gx)^T (Gx) = \|Gx\|_2^2 \geq 0$.

As G is upper triangular with positive diagonal entries, it is nonsingular. Therefore as $x \neq 0$, $Gx \neq 0 \Rightarrow \|Gx\|_2 > 0$ and from above, $x^T A x > 0$ for all $x \neq 0$.

Conversely, suppose A is an $n \times n$ positive definite matrix. Then A is nonsingular and all its leading principal submatrices are also positive definite and hence nonsingular. Moreover $x^T A x > 0$ for all $x \in \mathbb{R}^n \setminus \{0\}$.

Therefore, there is a unique unit lower triangular matrix L and a matrix $D = \text{diag}(d_{11}, \dots, d_{nn})$ $d_{ii} > 0$, such that $A = LDL^T$. Let $G = D^{1/2} L^T$ where, $D^{1/2} = \text{diag}(\sqrt{d_{11}}, \dots, \sqrt{d_{nn}})$. Then G is upper triangular with positive diagonal entries such that $A = G^T G$. Hence the proof. \square

Algorithms for Computing Cholesky Decomposition

Inner Product Form

Suppose

$$\underbrace{\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{12} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1n} & a_{2n} & \cdots & a_{nn} \end{bmatrix}}_{=:A} = \begin{bmatrix} g_{11} & & & \\ g_{12} & g_{22} & & \\ \vdots & \vdots & \ddots & \\ g_{1n} & g_{2n} & \cdots & g_{nn} \end{bmatrix} \underbrace{\begin{bmatrix} g_{11} & g_{12} & \cdots & g_{1n} \\ & g_{22} & \cdots & g_{2n} \\ & & \ddots & \vdots \\ & & & g_{nn} \end{bmatrix}}_{=:G}$$

Equating entries on both sides for $j = 1 : n$,

$$\sum_{k=1}^j g_{kj}^2 = a_{jj} \text{ \& } \sum_{i=1}^j g_{ij} g_{ik} = a_{jk};$$
$$\Rightarrow g_{jj} = \underbrace{\left(a_{jj} - \sum_{k=1}^{j-1} g_{kj}^2 \right)^{1/2}}_{\text{costs } 2(j-1) \text{ flops} + \text{one square root}} \text{ \& } g_{jk} = \underbrace{\left(a_{jk} - \sum_{i=1}^{j-1} g_{ij} g_{ik} \right)}_{\text{costs } 2j-1 \text{ flops for each } k} / g_{jj}; \quad k = j+1 : n$$

This is the *inner product formulation* for finding the Cholesky factor G one row at a time.

Flop count: $n^3/3 + O(n^2)$ flops. (Exercise!)

Outer Product Form

Let

$$b = A(1,2:n)^T, \hat{A} = A(2:n,2:n),$$

$$g = G(1,2:n)^T, \hat{G} = G(2:n,2:n).$$

Then

$$\left[\begin{array}{c|c} a_{11} & b^T \\ \hline b & \hat{A} \end{array} \right] = \left[\begin{array}{c|c} g_{11} & \\ \hline g & \hat{G}^T \end{array} \right] \left[\begin{array}{c|c} g_{11} & g^T \\ \hline & \hat{G} \end{array} \right] \Rightarrow \begin{cases} g_{11} = \sqrt{a_{11}} \\ g = b/g_{11} \\ \hat{G}^T \hat{G} = \hat{A} - gg^T \end{cases}$$

Outer Product Form

Let

$$b = A(1,2:n)^T, \hat{A} = A(2:n,2:n), \\ g = G(1,2:n)^T, \hat{G} = G(2:n,2:n).$$

Then

$$\left[\begin{array}{c|c} a_{11} & b^T \\ \hline b & \hat{A} \end{array} \right] = \left[\begin{array}{c|c} g_{11} & \\ \hline g & \hat{G}^T \end{array} \right] \left[\begin{array}{c|c} g_{11} & g^T \\ \hline & \hat{G} \end{array} \right] \Rightarrow \begin{cases} g_{11} = \sqrt{a_{11}} \\ g = b/g_{11} \\ \hat{G}^T \hat{G} = \hat{A} - gg^T \end{cases}$$

which gives the pseudocode

1. Compute $g_{11} = \sqrt{a_{11}}$.
2. Compute $g = b/g_{11}$.
3. Compute the Cholesky factor \hat{G} of $\hat{A} - gg^T$.

for a recursive algorithm to find the Cholesky factor of A .

This is the *outer product formulation* for finding the Cholesky factor as it involves the outer product gg^T .

Outer Product Form

Let

$$b = A(1,2:n)^T, \hat{A} = A(2:n,2:n), \\ g = G(1,2:n)^T, \hat{G} = G(2:n,2:n).$$

Then

$$\left[\begin{array}{c|c} a_{11} & b^T \\ \hline b & \hat{A} \end{array} \right] = \left[\begin{array}{c|c} g_{11} & \\ \hline g & \hat{G}^T \end{array} \right] \left[\begin{array}{c|c} g_{11} & g^T \\ \hline & \hat{G} \end{array} \right] \Rightarrow \begin{cases} g_{11} = \sqrt{a_{11}} \\ g = b/g_{11} \\ \hat{G}^T \hat{G} = \hat{A} - gg^T \end{cases}$$

which gives the pseudocode

1. Compute $g_{11} = \sqrt{a_{11}}$.
2. Compute $g = b/g_{11}$.
3. Compute the Cholesky factor \hat{G} of $\hat{A} - gg^T$.

for a recursive algorithm to find the Cholesky factor of A .

This is the *outer product formulation* for finding the Cholesky factor as it involves the outer product gg^T .

Verify that this also costs $n^3/3 + O(n^2)$ flops!

Bordered Form

For $j = 2:n$ let

$$A_{j-1} = A(1:j-1, 1:j-1); c = A(1:j-1, j)$$

$$G_{j-1} = G(1:j-1; 1:j-1); h = G(1:j-1, j)$$

Bordered Form

For $j = 2:n$ let

$$A_{j-1} = A(1:j-1, 1:j-1); c = A(1:j-1, j)$$

$$G_{j-1} = G(1:j-1; 1:j-1); h = G(1:j-1, j)$$

Then for $j = 2:n$

$$\begin{aligned} A = G^T G &\Rightarrow \left[\begin{array}{c|c} A_{j-1} & c \\ \hline c^T & a_{jj} \end{array} \right] = \left[\begin{array}{c|c} G_{j-1}^T & \\ \hline h^T & g_{jj} \end{array} \right] \left[\begin{array}{c|c} G_{j-1} & h \\ \hline & g_{jj} \end{array} \right] \\ &\Rightarrow A_{j-1} = G_{j-1}^T G_{j-1}; \quad c = G_{j-1}^T h; \quad a_{jj} = h^T h + g_{jj}^2; \end{aligned}$$

Bordered Form

For $j = 2:n$ let

$$A_{j-1} = A(1:j-1, 1:j-1); c = A(1:j-1, j)$$

$$G_{j-1} = G(1:j-1, 1:j-1); h = G(1:j-1, j)$$

Then for $j = 2:n$

$$A = G^T G \Rightarrow \left[\begin{array}{c|c} A_{j-1} & c \\ \hline c^T & a_{jj} \end{array} \right] = \left[\begin{array}{c|c} G_{j-1}^T & \\ \hline h^T & g_{jj} \end{array} \right] \left[\begin{array}{c|c} G_{j-1} & h \\ \hline & g_{jj} \end{array} \right]$$
$$\Rightarrow A_{j-1} = G_{j-1}^T G_{j-1}; c = G_{j-1}^T h; a_{jj} = h^T h + g_{jj}^2;$$

This gives the following pseudocode for computing G :

1. Set $G = \text{zeros}(n, n)$; & $G(1, 1) = \sqrt{A(1, 1)}$;
2. for $j = 2:n$
 Solve $G(1:j-1, 1:j-1)^T h = A(1:j-1, j)$ for h .
 Set $G(1:j-1, j) = h$;
 Set $G(j, j) = \sqrt{A(j, j) - h^T h}$;
end

This is the *bordered* form of finding G which also costs $n^3/3 + O(n^2)$ flops

Solving a Positive Definite System of Equations

Pseudocode for solving an $n \times n$ system $Ax = b$ where A is a positive definite matrix:

1. Find the Cholesky factor G of A . (costs $n^3/3 + O(n^2)$ flops)
2. Solve $G^T y = b$ for y . (costs n^2 flops)
3. Solve $Gx = y$ for x . (costs n^2 flops)

Total cost is $n^3/3 + O(n^2)$ flops.