

MA 322: Scientific Computing

Lecture - 9



Convergence of Lagrange's Interpolating polynomial

Result

- Suppose x_0, x_1, \dots, x_n are distinct real numbers, and let f be a given real valued function with $n + 1$ continuous derivatives on the interval $I_t = \mathcal{H}\{t, x_0, x_1, \dots, x_n\}$, with t some given real number. Then there exists a number $\xi_t \in I_t$ such that

$$f(t) - p_n(t) = \frac{f^{(n+1)}(\xi_t)}{(n+1)!} (t - x_0)(t - x_1) \dots (t - x_n), \quad (1)$$

where $p_n(t) = \sum_{j=0}^n f(x_j) \ell_j(t)$ is the Lagrange interpolating polynomial of f with degree n .

Proof. We define the error function E by

$$E(t) = f(t) - p_n(t), \quad p_n(t) = \sum_{j=0}^n f(x_j) \ell_j(t) \quad (2)$$

Convergence of Lagrange's Interpolating polynomial

Result

- Suppose x_0, x_1, \dots, x_n are distinct real numbers, and let f be a given real valued function with $n + 1$ continuous derivatives on the interval $I_t = \mathcal{H}\{t, x_0, x_1, \dots, x_n\}$, with t some given real number. Then there exists a number $\xi_t \in I_t$ such that

$$f(t) - p_n(t) = \frac{f^{(n+1)}(\xi_t)}{(n+1)!} (t - x_0)(t - x_1) \dots (t - x_n), \quad (1)$$

where $p_n(t) = \sum_{j=0}^n f(x_j) \ell_j(t)$ is the Lagrange interpolating polynomial of f with degree n .

Proof. We define the error function E by

$$E(t) = f(t) - p_n(t), \quad p_n(t) = \sum_{j=0}^n f(x_j) \ell_j(t) \quad (2)$$

and a user defined function G by

$$G(x) = E(x) - \frac{\Psi(x)}{\Psi(t)} E(t) \quad (3)$$

with $\Psi(x) = (x - x_0)(x - x_1) \dots (x - x_n)$.

Convergence Analysis Contd...

Then for the user defined function

$$G(x) = E(x) - \frac{\Psi(x)}{\Psi(t)} E(t), \quad \Psi(x) = (x - x_0)(x - x_1) \dots (x - x_n), \quad (4)$$

observe that

$$G(x_i) = E(x_i) - \frac{\Psi(x_i)}{\Psi(t)} E(t) = 0, \quad i = 0, 1, 2, \dots, n \quad (5)$$

$$G(t) = E(t) - E(t) = 0. \quad (6)$$

Convergence Analysis Contd...

Then for the user defined function

$$G(x) = E(x) - \frac{\Psi(x)}{\Psi(t)} E(t), \quad \Psi(x) = (x - x_0)(x - x_1) \dots (x - x_n), \quad (4)$$

observe that

$$G(x_i) = E(x_i) - \frac{\Psi(x_i)}{\Psi(t)} E(t) = 0, \quad i = 0, 1, 2, \dots, n \quad (5)$$

$$G(t) = E(t) - E(t) = 0. \quad (6)$$

Thus, the function G has $n + 2$ distinct roots in the interval I_t .

Convergence Analysis Contd...

Then for the user defined function

$$G(x) = E(x) - \frac{\Psi(x)}{\Psi(t)} E(t), \quad \Psi(x) = (x - x_0)(x - x_1) \dots (x - x_n), \quad (4)$$

observe that

$$G(x_i) = E(x_i) - \frac{\Psi(x_i)}{\Psi(t)} E(t) = 0, \quad i = 0, 1, 2, \dots, n \quad (5)$$

$$G(t) = E(t) - E(t) = 0. \quad (6)$$

Thus, the function G has $n + 2$ distinct roots in the interval I_t . Then using the MVT, G' has $n + 1$ distinct roots in the interval I_t .

Convergence Analysis Contd...

Then for the user defined function

$$G(x) = E(x) - \frac{\Psi(x)}{\Psi(t)} E(t), \quad \Psi(x) = (x - x_0)(x - x_1) \dots (x - x_n), \quad (4)$$

observe that

$$G(x_i) = E(x_i) - \frac{\Psi(x_i)}{\Psi(t)} E(t) = 0, \quad i = 0, 1, 2, \dots, n \quad (5)$$

$$G(t) = E(t) - E(t) = 0. \quad (6)$$

Thus, the function G has $n + 2$ distinct roots in the interval I_t . Then using the MVT, G' has $n + 1$ distinct roots in the interval I_t . Inductively, $G^{(n+1)}$ has one root, say ξ , in the interval I_t ,

Convergence Analysis Contd...

Then for the user defined function

$$G(x) = E(x) - \frac{\Psi(x)}{\Psi(t)} E(t), \quad \Psi(x) = (x - x_0)(x - x_1) \dots (x - x_n), \quad (4)$$

observe that

$$G(x_i) = E(x_i) - \frac{\Psi(x_i)}{\Psi(t)} E(t) = 0, \quad i = 0, 1, 2, \dots, n \quad (5)$$

$$G(t) = E(t) - E(t) = 0. \quad (6)$$

Thus, the function G has $n + 2$ distinct roots in the interval I_t . Then using the MVT, G' has $n + 1$ distinct roots in the interval I_t . Inductively, $G^{(n+1)}$ has one root, say ξ , in the interval I_t , so that

$$G^{(n+1)}(\xi) = 0.$$

Convergence Analysis Contd...

Then for the user defined function

$$G(x) = E(x) - \frac{\Psi(x)}{\Psi(t)} E(t), \quad \Psi(x) = (x - x_0)(x - x_1) \dots (x - x_n), \quad (4)$$

observe that

$$G(x_i) = E(x_i) - \frac{\Psi(x_i)}{\Psi(t)} E(t) = 0, \quad i = 0, 1, 2, \dots, n \quad (5)$$

$$G(t) = E(t) - E(t) = 0. \quad (6)$$

Thus, the function G has $n + 2$ distinct roots in the interval I_t . Then using the MVT, G' has $n + 1$ distinct roots in the interval I_t . Inductively, $G^{(n+1)}$ has one root, say ξ , in the interval I_t , so that

$$G^{(n+1)}(\xi) = 0.$$

Then, for the error function $E(x) = f(x) - p_n(x)$, we obtain

$$E^{(n+1)}(x) = f^{(n+1)}(x)$$

Convergence Analysis Contd...

Then for the user defined function

$$G(x) = E(x) - \frac{\Psi(x)}{\Psi(t)} E(t), \quad \Psi(x) = (x - x_0)(x - x_1) \dots (x - x_n), \quad (4)$$

observe that

$$G(x_i) = E(x_i) - \frac{\Psi(x_i)}{\Psi(t)} E(t) = 0, \quad i = 0, 1, 2, \dots, n \quad (5)$$

$$G(t) = E(t) - E(t) = 0. \quad (6)$$

Thus, the function G has $n + 2$ distinct roots in the interval I_t . Then using the MVT, G' has $n + 1$ distinct roots in the interval I_t . Inductively, $G^{(n+1)}$ has one root, say ξ , in the interval I_t , so that

$$G^{(n+1)}(\xi) = 0.$$

Then, for the error function $E(x) = f(x) - p_n(x)$, we obtain

$$E^{(n+1)}(x) = f^{(n+1)}(x) \quad \& \quad \Psi^{(n+1)}(x) = (n + 1)! \quad (7)$$

Convergence Analysis Contd...

Then for the user defined function

$$G(x) = E(x) - \frac{\Psi(x)}{\Psi(t)} E(t), \quad \Psi(x) = (x - x_0)(x - x_1) \dots (x - x_n), \quad (4)$$

observe that

$$G(x_i) = E(x_i) - \frac{\Psi(x_i)}{\Psi(t)} E(t) = 0, \quad i = 0, 1, 2, \dots, n \quad (5)$$

$$G(t) = E(t) - E(t) = 0. \quad (6)$$

Thus, the function G has $n + 2$ distinct roots in the interval I_t . Then using the MVT, G' has $n + 1$ distinct roots in the interval I_t . Inductively, $G^{(n+1)}$ has one root, say ξ , in the interval I_t , so that

$$G^{(n+1)}(\xi) = 0.$$

Then, for the error function $E(x) = f(x) - p_n(x)$, we obtain

$$E^{(n+1)}(x) = f^{(n+1)}(x) \quad \& \quad \Psi^{(n+1)}(x) = (n + 1)! \quad (7)$$

Therefore, we have

$$G^{(n+1)}(x) = f^{(n+1)}(x) - \frac{(n + 1)!}{\Psi(t)} E(t) \quad (8)$$

Convergence Analysis Contd...

Now, substitute $x = \xi$ in the following relation

$$G^{(n+1)}(x) = f^{(n+1)}(x) - \frac{(n+1)!}{\Psi(t)} E(t) \quad (9)$$

to obtain

$$E(t) = \frac{\Psi(t)}{(n+1)!} f^{(n+1)}(\xi). \quad (10)$$

Convergence Analysis Contd...

Now, substitute $x = \xi$ in the following relation

$$G^{(n+1)}(x) = f^{(n+1)}(x) - \frac{(n+1)!}{\Psi(t)} E(t) \quad (9)$$

to obtain

$$E(t) = \frac{\Psi(t)}{(n+1)!} f^{(n+1)}(\xi). \quad (10)$$

Hence, for any real number x , we have following error representations

$$f(x) - p_n(x) = \frac{(x - x_0)(x - x_1) \dots (x - x_n)}{(n+1)!} f^{(n+1)}(\xi_x), \quad (11)$$

where $\xi_x \in \mathcal{H}\{x, x_0, x_1, \dots, x_n\}$.

Convergence Analysis Contd...

Now, substitute $x = \xi$ in the following relation

$$G^{(n+1)}(x) = f^{(n+1)}(x) - \frac{(n+1)!}{\Psi(t)} E(t) \quad (9)$$

to obtain

$$E(t) = \frac{\Psi(t)}{(n+1)!} f^{(n+1)}(\xi). \quad (10)$$

Hence, for any real number x , we have following error representations

$$f(x) - p_n(x) = \frac{(x - x_0)(x - x_1) \dots (x - x_n)}{(n+1)!} f^{(n+1)}(\xi_x), \quad (11)$$

where $\xi_x \in \mathcal{H}\{x, x_0, x_1, \dots, x_n\}$.

Remark:

- Now, we wish to calculate the distance between f and p_n , which is done by introducing norm over function spaces.

Norm: Generalization of $|\cdot|$

Norm: Generalization of $|\cdot|$

- Suppose $(X, +, \cdot)$ is a vector space over \mathbb{R} .

Norm: Generalization of $|\cdot|$

- Suppose $(X, +, \cdot)$ is a vector space over \mathbb{R} . Then, a real-valued function, denoted by $\|\cdot\|$, is called a **norm** if it satisfies the following properties:

Norm: Generalization of $|\cdot|$

● Suppose $(X, +, \cdot)$ is a vector space over \mathbb{R} . Then, a real-valued function, denoted by $\|\cdot\|$, is called a **norm** if it satisfies the following properties:

- $\|x\| \geq 0$ for all $x \in X$, and $\|x\| = 0$ if and only if $x = 0$,

Norm: Generalization of $|\cdot|$

● Suppose $(X, +, \cdot)$ is a vector space over \mathbb{R} . Then, a real-valued function, denoted by $\|\cdot\|$, is called a **norm** if it satisfies the following properties:

- $\|x\| \geq 0$ for all $x \in X$, and $\|x\| = 0$ if and only if $x = 0$,
- $\|\alpha x\| = |\alpha| \|x\|$ for all $x \in X$ and $\alpha \in \mathbb{R}$,

Norm: Generalization of $|\cdot|$

● Suppose $(X, +, \cdot)$ is a vector space over \mathbb{R} . Then, a real-valued function, denoted by $\|\cdot\|$, is called a **norm** if it satisfies the following properties:

- $\|x\| \geq 0$ for all $x \in X$, and $\|x\| = 0$ if and only if $x = 0$,
- $\|\alpha x\| = |\alpha| \|x\|$ for all $x \in X$ and $\alpha \in \mathbb{R}$,
- $\|x + y\| \leq \|x\| + \|y\|$ for all $x, y \in X$.

Norm: Generalization of $|\cdot|$

● Suppose $(X, +, \cdot)$ is a vector space over \mathbb{R} . Then, a real-valued function, denoted by $\|\cdot\|$, is called a **norm** if it satisfies the following properties:

- $\|x\| \geq 0$ for all $x \in X$, and $\|x\| = 0$ if and only if $x = 0$,
- $\|\alpha x\| = |\alpha| \|x\|$ for all $x \in X$ and $\alpha \in \mathbb{R}$,
- $\|x + y\| \leq \|x\| + \|y\|$ for all $x, y \in X$.

● A vector space X equipped with a norm is called normed linear space and it is denoted by pair $(X, \|\cdot\|)$.

Norm: Generalization of $|\cdot|$

● Suppose $(X, +, \cdot)$ is a vector space over \mathbb{R} . Then, a real-valued function, denoted by $\|\cdot\|$, is called a **norm** if it satisfies the following properties:

- $\|x\| \geq 0$ for all $x \in X$, and $\|x\| = 0$ if and only if $x = 0$,
- $\|\alpha x\| = |\alpha| \|x\|$ for all $x \in X$ and $\alpha \in \mathbb{R}$,
- $\|x + y\| \leq \|x\| + \|y\|$ for all $x, y \in X$.

● A vector space X equipped with a norm is called normed linear space and it is denoted by pair $(X, \|\cdot\|)$.

● Remark:

Norm: Generalization of $|\cdot|$

● Suppose $(X, +, \cdot)$ is a vector space over \mathbb{R} . Then, a real-valued function, denoted by $\|\cdot\|$, is called a **norm** if it satisfies the following properties:

- $\|x\| \geq 0$ for all $x \in X$, and $\|x\| = 0$ if and only if $x = 0$,
- $\|\alpha x\| = |\alpha| \|x\|$ for all $x \in X$ and $\alpha \in \mathbb{R}$,
- $\|x + y\| \leq \|x\| + \|y\|$ for all $x, y \in X$.

● A vector space X equipped with a norm is called normed linear space and it is denoted by pair $(X, \|\cdot\|)$.

● Remark:

- A norm $\|\cdot\|$ on a vector space X defines the distance between two vectors in X .

Norm: Generalization of $|\cdot|$

● Suppose $(X, +, \cdot)$ is a vector space over \mathbb{R} . Then, a real-valued function, denoted by $\|\cdot\|$, is called a **norm** if it satisfies the following properties:

- $\|x\| \geq 0$ for all $x \in X$, and $\|x\| = 0$ if and only if $x = 0$,
- $\|\alpha x\| = |\alpha| \|x\|$ for all $x \in X$ and $\alpha \in \mathbb{R}$,
- $\|x + y\| \leq \|x\| + \|y\|$ for all $x, y \in X$.

● A vector space X equipped with a norm is called normed linear space and it is denoted by pair $(X, \|\cdot\|)$.

● Remark:

- A norm $\|\cdot\|$ on a vector space X defines the distance between two vectors in X . How?

Norm: Generalization of $|\cdot|$

● Suppose $(X, +, \cdot)$ is a vector space over \mathbb{R} . Then, a real-valued function, denoted by $\|\cdot\|$, is called a **norm** if it satisfies the following properties:

- $\|x\| \geq 0$ for all $x \in X$, and $\|x\| = 0$ if and only if $x = 0$,
- $\|\alpha x\| = |\alpha| \|x\|$ for all $x \in X$ and $\alpha \in \mathbb{R}$,
- $\|x + y\| \leq \|x\| + \|y\|$ for all $x, y \in X$.

● A vector space X equipped with a norm is called normed linear space and it is denoted by pair $(X, \|\cdot\|)$.

● Remark:

- A norm $\|\cdot\|$ on a vector space X defines the distance between two vectors in X . How?
- Because a normed linear space $(X, \|\cdot\|)$ is also a metric space with respect to the metric

Norm: Generalization of $|\cdot|$

● Suppose $(X, +, \cdot)$ is a vector space over \mathbb{R} . Then, a real-valued function, denoted by $\|\cdot\|$, is called a **norm** if it satisfies the following properties:

- $\|x\| \geq 0$ for all $x \in X$, and $\|x\| = 0$ if and only if $x = 0$,
- $\|\alpha x\| = |\alpha| \|x\|$ for all $x \in X$ and $\alpha \in \mathbb{R}$,
- $\|x + y\| \leq \|x\| + \|y\|$ for all $x, y \in X$.

● A vector space X equipped with a norm is called normed linear space and it is denoted by pair $(X, \|\cdot\|)$.

● Remark:

- A norm $\|\cdot\|$ on a vector space X defines the distance between two vectors in X . How?
- Because a normed linear space $(X, \|\cdot\|)$ is also a metric space with respect to the metric

$$d(x, y) = \|x - y\| \quad \forall x, y \in X.$$

Norm: Generalization of $|\cdot|$

● Suppose $(X, +, \cdot)$ is a vector space over \mathbb{R} . Then, a real-valued function, denoted by $\|\cdot\|$, is called a **norm** if it satisfies the following properties:

- $\|x\| \geq 0$ for all $x \in X$, and $\|x\| = 0$ if and only if $x = 0$,
- $\|\alpha x\| = |\alpha| \|x\|$ for all $x \in X$ and $\alpha \in \mathbb{R}$,
- $\|x + y\| \leq \|x\| + \|y\|$ for all $x, y \in X$.

● A vector space X equipped with a norm is called normed linear space and it is denoted by pair $(X, \|\cdot\|)$.

● Remark:


- A norm $\|\cdot\|$ on a vector space X defines the distance between two vectors in X . How?
- Because a normed linear space $(X, \|\cdot\|)$ is also a metric space with respect to the metric

$$d(x, y) = \|x - y\| \quad \forall x, y \in X.$$

- Therefore, in order to define the distance between two functions in $C[a, b]$, we first try to associate vector space $C[a, b]$ a norm.


Concept of Distance Between two Functions

Concept of Distance Between two Functions




- For $f, g \in C[a, b]$,

Concept of Distance Between two Functions




- For $f, g \in C[a, b]$, the distance between f and g at a particular x is given by $|f(x) - g(x)|$.

Concept of Distance Between two Functions




- For $f, g \in C[a, b]$, the distance between f and g at a particular x is given by $|f(x) - g(x)|$. To measure the distance, which takes care all x , we may try to evaluate $\max_{x \in [a, b]} |f(x) - g(x)|$.

Concept of Distance Between two Functions




- For $f, g \in C[a, b]$, the distance between f and g at a particular x is given by $|f(x) - g(x)|$. To measure the distance, which takes care all x , we may try to evaluate $\max_{x \in [a, b]} |f(x) - g(x)|$. Does it represents the distance between f and g ?

Concept of Distance Between two Functions




- For $f, g \in C[a, b]$, the distance between f and g at a particular x is given by $|f(x) - g(x)|$. To measure the distance, which takes care all x , we may try to evaluate $\max_{x \in [a, b]} |f(x) - g(x)|$. Does it represents the distance between f and g ? It will measure the distance between f and g provided this distance is generated from a norm.

Concept of Distance Between two Functions



- For $f, g \in C[a, b]$, the distance between f and g at a particular x is given by $|f(x) - g(x)|$. To measure the distance, which takes care all x , we may try to evaluate $\max_{x \in [a, b]} |f(x) - g(x)|$. Does it represents the distance between f and g ? It will measure the distance between f and g provided this distance is generated from a norm.
- It is an easy exercise to verify that $\|f\|_{\infty} = \max_{x \in [a, b]} |f(x)|$ defines a norm on $C[a, b]$, known as infinity norm.

Concept of Distance Between two Functions



- For $f, g \in C[a, b]$, the distance between f and g at a particular x is given by $|f(x) - g(x)|$. To measure the distance, which takes care all x , we may try to evaluate $\max_{x \in [a, b]} |f(x) - g(x)|$. Does it represents the distance between f and g ? It will measure the distance between f and g provided this distance is generated from a norm.
- It is an easy exercise to verify that $\|f\|_{\infty} = \max_{x \in [a, b]} |f(x)|$ defines a norm on $C[a, b]$, known as infinity norm. The distance associated with this norm is given by

$$d_{\infty}(f, g) = \max_{x \in [a, b]} |(f - g)(x)| = \max_{x \in [a, b]} |f(x) - g(x)|.$$

Notion of Convergence

● In $(C[a, b], d_\infty)$

- A sequence $\langle f_n \rangle$ in $C[a, b]$ is said to be convergent w.r.t d_∞ , if there exists a $f \in C[a, b]$ s. t.

$$d_\infty(f_n, f) \rightarrow 0 \text{ or equivalently } \|f_n - f\|_\infty \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (12)$$

Remark:

Notion of Convergence

• In $(C[a, b], d_\infty)$

- A sequence $\langle f_n \rangle$ in $C[a, b]$ is said to be convergent w.r.t d_∞ , if there exists a $f \in C[a, b]$ s. t.

$$d_\infty(f_n, f) \rightarrow 0 \text{ or equivalently } \|f_n - f\|_\infty \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (12)$$

Remark:

- Convergence with respect to the $\|\cdot\|_\infty$ norm leads to uniform convergence.

Notion of Convergence

• In $(C[a, b], d_\infty)$

- A sequence $\langle f_n \rangle$ in $C[a, b]$ is said to be convergent w.r.t d_∞ , if there exists a $f \in C[a, b]$ s. t.

$$d_\infty(f_n, f) \rightarrow 0 \text{ or equivalently } \|f_n - f\|_\infty \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (12)$$

Remark:

- Convergence with respect to the $\|\cdot\|_\infty$ norm leads to uniform convergence. Again uniform convergence has following consequences
 - Uniform limit of a sequence of continuous functions is also continuous.

Notion of Convergence

• In $(C[a, b], d_\infty)$

- A sequence $\langle f_n \rangle$ in $C[a, b]$ is said to be convergent w.r.t d_∞ , if there exists a $f \in C[a, b]$ s. t.

$$d_\infty(f_n, f) \rightarrow 0 \text{ or equivalently } \|f_n - f\|_\infty \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (12)$$

Remark:

- Convergence with respect to the $\|\cdot\|_\infty$ norm leads to uniform convergence. Again uniform convergence has following consequences
 - Uniform limit of a sequence of continuous functions is also continuous.
 - If $f_n \rightarrow f$ converges in $(C[a, b], d_\infty)$, then

$$\lim_{n \rightarrow \infty} \int_a^b f_n(x) dx = \int_a^b \lim_{n \rightarrow \infty} f_n(x) dx = \int_a^b f(x) dx. \quad (13)$$

About Uniform Convergence of Interpolating Polynomials

Consider again the error formula

$$f(x) - p_n(x) = \frac{(x - x_0)(x - x_1) \dots (x - x_n)}{(n + 1)!} f^{(n+1)}(\xi_x), \quad (14)$$

where $\xi_x \in \mathcal{H}\{x, x_0, x_1, \dots, x_n\}$.

About Uniform Convergence of Interpolating Polynomials

Consider again the error formula

$$f(x) - p_n(x) = \frac{(x - x_0)(x - x_1) \dots (x - x_n)}{(n+1)!} f^{(n+1)}(\xi_x), \quad (14)$$

where $\xi_x \in \mathcal{H}\{x, x_0, x_1, \dots, x_n\}$. Then for any interval I that contains $\mathcal{H}\{x, x_0, x_1, \dots, x_n\}$ and for $f \in C^{(n+1)}(I)$, we obtain

$$\max_{x \in I} |f(x) - p_n(x)| \leq \frac{1}{(n+1)!} \max_{x \in I} |f^{(n+1)}(\xi_x)| \max_{x \in I} |\Psi_n(x)|, \quad (15)$$

where $\Psi_n(x) = (x - x_0)(x - x_1) \dots (x - x_n)$.

Remark:

- Now, $p_n \rightarrow f$ uniformly provided quantities

$$M_{n+1} = \max_{x \in I} |f^{(n+1)}(\xi_x)| \quad \text{and} \quad \max_{x \in I} |\Psi_n(x)|$$

are uniformly bounded in I .

About Uniform Convergence of Interpolating Polynomials

Consider again the error formula

$$f(x) - p_n(x) = \frac{(x - x_0)(x - x_1) \dots (x - x_n)}{(n + 1)!} f^{(n+1)}(\xi_x), \quad (14)$$

where $\xi_x \in \mathcal{H}\{x, x_0, x_1, \dots, x_n\}$. Then for any interval I that contains $\mathcal{H}\{x, x_0, x_1, \dots, x_n\}$ and for $f \in C^{(n+1)}(I)$, we obtain

$$\max_{x \in I} |f(x) - p_n(x)| \leq \frac{1}{(n + 1)!} \max_{x \in I} |f^{(n+1)}(\xi_x)| \max_{x \in I} |\Psi_n(x)|, \quad (15)$$

where $\Psi_n(x) = (x - x_0)(x - x_1) \dots (x - x_n)$.

Remark:

- Now, $p_n \rightarrow f$ uniformly provided quantities

$$M_{n+1} = \max_{x \in I} |f^{(n+1)}(\xi_x)| \quad \text{and} \quad \max_{x \in I} |\Psi_n(x)|$$

are uniformly bounded in I . More precisely, we may expect $\|f - p_n\|_\infty \rightarrow 0$ as $n \rightarrow \infty$ for C^∞ (infinitely many times differentiable functions) functions when

$$\left(M_{n+1} \cdot \max_{x \in I} |\Psi_n(x)| \right) \text{ are uniformly bounded in } I.$$

About U.C. of Interpolating Polynomials Contd....

Unfortunately, this is not so, since the sequence

$$\left(M_{n+1} \cdot \max_{x \in I} |\Psi_n(x)| \right)$$

may tend to ∞ , as $n \rightarrow \infty$, faster than the sequence $\frac{1}{(n+1)!}$ tends to 0.

About U.C. of Interpolating Polynomials Contd....

Unfortunately, this is not so, since the sequence

$$\left(M_{n+1} \cdot \max_{x \in I} |\Psi_n(x)| \right)$$

may tend to ∞ , as $n \rightarrow \infty$, faster than the sequence $\frac{1}{(n+1)!}$ tends to 0. This is known as *Runge phenomenon*.

About U.C. of Interpolating Polynomials Contd....

Unfortunately, this is not so, since the sequence

$$\left(M_{n+1} \cdot \max_{x \in I} |\Psi_n(x)| \right)$$

may tend to ∞ , as $n \rightarrow \infty$, faster than the sequence $\frac{1}{(n+1)!}$ tends to 0.

This is known as *Runge phenomenon*.

Example:

About U.C. of Interpolating Polynomials Contd....

Unfortunately, this is not so, since the sequence

$$\left(M_{n+1} \cdot \max_{x \in I} |\Psi_n(x)| \right)$$

may tend to ∞ , as $n \rightarrow \infty$, faster than the sequence $\frac{1}{(n+1)!}$ tends to 0.

This is known as *Runge phenomenon*.

Example: Consider the function $f(x) = \frac{1}{1+x^2}$, $x \in [-5, 5]$ and let us try to calculate $\|f - p_n\|$ for different values of n .

About U.C. of Interpolating Polynomials Contd....

Unfortunately, this is not so, since the sequence

$$\left(M_{n+1} \cdot \max_{x \in I} |\Psi_n(x)| \right)$$

may tend to ∞ , as $n \rightarrow \infty$, faster than the sequence $\frac{1}{(n+1)!}$ tends to 0.

This is known as *Runge phenomenon*.

Example: Consider the function $f(x) = \frac{1}{1+x^2}$, $x \in [-5, 5]$ and let us try to calculate $\|f - p_n\|$ for different values of n .

Degree n	Max error
2	0.65
4	0.44
6	0.61
8	1.04
10	1.92
12	3.66
14	7.15
16	14.25
18	28.74
20	58.59
22	121.02
24	252.78