#flo #ref #hw

1 | def of a vector space

- Props of addition and scalar multiplication in F^N
 - +: comutative, associative, identiy
 - * every element has an additive inverse
 - *: associative, identity
 - addition and scalar multiplication, connected by distributive props
- let *V* be a set with an addition and scalar multiplication that satisfy the props,

```
**addition, scalar multiplication**
- addition: assigns an element u+v in V to each pair of elements u, v in V
- scalar multiplication: lv with l in f and v in V

**vector space**
is V with addition and SCMUL with:
- commutativitity
- associativity
- additive idenitity
- additive inverse
- multiplicative identity
- distibutive properties
```

- · no multiplicative inverse?
 - is this how you solve the 0 issue?
- · vec, point
 - elements of vec space are called vecs or points
- simplest vec space: {0}
- f^{infin} is the set of all sequences of elements of F
 - additive identity: sequece of all zeros
- · vector space can include a set of functions? not quite...
 - let S be a set, and FS be the set of functions from S to F
 - what?? #review
- let S be the interval [0,1] and F=R
 - R^[\0, \1] is the set of real valued function on the interval [0,1]
 - ??

- $F^N -> F^{1,2,...,n}$
- F^{infin} -> F^{1,2,...}
- · vector spaces need unique additive inverse
 - 0'=0'+0=0+0'=0
 - * nicer than my proof
- · unique additive inverse
 - w=w+0=w+(v+w')=(w+v)=(w+v)+w'=0+w'=w'

V denotes a vector space over F

- 1. no multiplicative inverse required?
- 2. what does the set of functions from S to F mean?

1.1 | exercises

1. prove that -(-v) = v

(a)
$$-(-v) = -1(-1v) = (-1*(-1))v = 1v = v$$

- 2. ab = 0, prove that a or b = 0
 - (a) a=0/v = 0, v=0/a = 0
- 3. empty set is not a vector space, it fails to satisfy only of the reqs. which one?
 - (a) no additive idenity
 - i. "there exists an element 0 in v" no there doesn't.

homework: KBxSolvingSystems