

1 | 1)

$$\vec{L} = \vec{p} \times m\vec{v}$$

The circle has a circumference of $2\pi R$, and it takes $\frac{2\pi}{\omega}$ seconds to travel that distance, so the tangential velocity must be $\vec{v} = 2\pi R \div \frac{2\pi}{\omega} = R\vec{\omega}$. Therefore, $\vec{L} = \vec{R} \times mR\vec{\omega}$

The vector is pointing out of the page, as the object is rotating counterclockwise. $|\vec{L}| = mR|\vec{R}||\vec{\omega}|$
 $= mR^2\omega$

We know that $\sin \theta$ is 1 because the vectors are perpendicular.

2 | 2)

The angular momentum changes because although the momentum of the object does not change, its position changes.

3 | 3)

$$\begin{aligned} \vec{L} &= \vec{p}\vec{v} \\ \frac{d\vec{L}}{dt} &= \frac{d\vec{p}}{dt} \times m\vec{v} + \frac{dm\vec{v}}{dt} \times \vec{p} \\ \text{We are given: } &= \vec{v} \times m\vec{v} + m\vec{a} \times \vec{p} = 0 + \vec{p} \times m\vec{a} \\ &= \vec{p} \times \vec{F} \\ &= \vec{\tau} \end{aligned}$$