

## Q3

Zhen Zhang

4/24/2020

```
set.seed(1)
x1 = runif(100)
x2 = 0.5 * x1 + rnorm(100) / 10
y = 2 + 2 * x1 + 0.3 * x2 + rnorm(100)
```

(a) Write out the form of the linear model. What are the regression coefficients?


$B_0 = B_1 = 2, B_2 = 0.3, s = 1, y = 2 + 2x_1 + 0.3x_2 + e.$

(b) What is the correlation between  $x_1$  and  $x_2$ ? Create a scatterplot displaying the relationship between the variables.

```
cor(x1, x2)
```

```
## [1] 0.8351212
```

```
plot(x1 ~ x2)
```



```
fit = lm(y ~ x1 + x2)
summary(fit)
```

```
##
## Call:
## lm(formula = y ~ x1 + x2)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -2.8311 -0.7273 -0.0537  0.6338  2.3359
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   2.1305     0.2319   9.188 7.61e-15 ***
## x1             1.4396     0.7212   1.996  0.0487 *
## x2             1.0097     1.1337   0.891  0.3754
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 1.056 on 97 degrees of freedom
## Multiple R-squared:  0.2088, Adjusted R-squared:  0.1925
## F-statistic: 12.8 on 2 and 97 DF, p-value: 1.164e-05
```

(d) Now fit a least squares regression to predict  $y$  using only  $x_1$ . Comment on your results. Can you reject the null hypothesis  $H_0: \beta_1 = 0$ ?

```
fit2 = lm(y ~ x1)
summary(fit2)
```

```
##
## Call:
## lm(formula = y ~ x1)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -2.89495 -0.66874 -0.07785  0.59221  2.45560
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   2.1124     0.2307   9.155 8.27e-15 ***
## x1             1.9759     0.3963   4.986 2.66e-06 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 1.055 on 98 degrees of freedom
## Multiple R-squared:  0.2024, Adjusted R-squared:  0.1942
## F-statistic: 24.86 on 1 and 98 DF,  p-value: 2.661e-06
```

(e) Now fit a least squares regression to predict  $y$  using only  $x_2$ . Comment on your results. Can you reject the null hypothesis  $H_0: \beta_1 = 0$ ?

```
fit3 = lm(y ~ x2)
summary(fit3)
```

```
##
## Call:
## lm(formula = y ~ x2)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -2.62687 -0.75156 -0.03598  0.72383  2.44890
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   2.3899     0.1949  12.26 < 2e-16 ***
## x2             2.8996     0.6330   4.58 1.37e-05 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
##  
## Residual standard error: 1.072 on 98 degrees of freedom  
## Multiple R-squared:  0.1763, Adjusted R-squared:  0.1679  
## F-statistic: 20.98 on 1 and 98 DF,  p-value: 1.366e-05
```

**(f) Do the results obtained in (c)–(e) contradict each other? Explain your answer.**

x1 and x2 are multicollinearity.