

Official websites use .gov A .gov website belongs to an official government organization in the United States. Secure .gov websites use HTTPS A lock () or https:// means you've safely connected to the .gov website. Share sensitive information only on official, secure websites.

General Best Practice Guidelines for Immunization Updated August 1, 2023 Printer friendly version [21 pages] Major changes to the best practice guidance in this section include 1) enhancement of the definition of a “precaution” to include any condition that might confuse diagnostic accuracy and 2) recommendation to vaccinate during a hospitalization if a patient is not acutely moderately or severely ill. National standards for pediatric vaccination practices have been established and include descriptions of valid contraindications and precautions to vaccination (2). Persons who administer vaccines should screen patients for contraindications and precautions to the vaccine before each dose of vaccine is administered (Table 4-1). Screening is facilitated by consistent use of screening questionnaires, which are available from certain state vaccination programs and other sources (e.g., the Immunization Action Coalition). Contraindications (conditions in a recipient that increases the risk for a serious adverse reaction) to vaccination are conditions under which vaccines should not be administered. Because the majority of contraindications are temporary, vaccinations often can be administered later when the condition leading to a contraindication no longer exists. A vaccine should not be administered when a contraindication is present; for example, MMR vaccine should not be administered to severely immunocompromised persons (1). However, certain conditions are commonly misperceived as contraindications (i.e., are not valid reasons to defer vaccination). Severely immunocompromised persons generally should not receive live vaccines (3). Because of the theoretical risk to the fetus, women known to be pregnant generally should not receive live, attenuated virus vaccines (4). Persons who experienced encephalopathy within 7 days after administration of a previous dose of pertussis-containing vaccine not attributable to another identifiable cause should not

receive additional doses of a vaccine that contains pertussis (4, 5). Severe Combined Immunodeficiency (SCID) disease and a history of intussusception are both contraindications to the receipt of rotavirus vaccines (6). A precaution is a condition in a recipient that might increase the risk for a serious adverse reaction, might cause diagnostic confusion, or might compromise the ability of the vaccine to produce immunity (e.g., administering measles vaccine to a person with passive immunity to measles from a blood transfusion administered up to 7 months prior) (7). A person might experience a more severe reaction to the vaccine than would have otherwise been expected; however, the risk for this happening is less than the risk expected with a contraindication. In general, vaccinations should be deferred when a precaution is present. However, a vaccination might be indicated in the presence of a precaution if the benefit of protection from the vaccine outweighs the risk for an adverse reaction. The presence of a moderate or severe acute illness with or without a fever is a precaution to administration of all vaccines (Table 4-1). The decision to administer or delay vaccination because of a current or recent acute illness depends on the severity of symptoms and etiology of the condition. The safety and efficacy of vaccinating persons who have mild illnesses have been documented (8-11). Vaccination should be deferred for persons with a moderate or severe acute illness. This precaution avoids causing diagnostic confusion between manifestations of the underlying illness and possible adverse effects of vaccination or superimposing adverse effects of the vaccine on the underlying illness. After they are screened for contraindications, persons with moderate or severe acute illness should be vaccinated as soon as the acute illness has improved. Studies indicate that failure to vaccinate children with minor illnesses can impede vaccination efforts (12-14). Among persons whose compliance with medical care cannot be ensured, use of every opportunity to administer appropriate vaccines is critical. Hospitalization should be used as an opportunity to provide recommended vaccinations. Health-care facilities are held to standards of offering influenza vaccine

for hospitalized patients, so providers are incentivized to vaccinate these patients at some point during hospitalization (15). Likewise, patients admitted for elective procedures will not be acutely ill during all times during their hospitalization. Most studies that have explored the effect of surgery or anesthesia on the immune system were observational, included only infants and children, and were small and indirect, in that they did not look at the immune effect on the response to vaccination specifically (16-35). They do not provide convincing evidence that recent anesthesia or surgery significantly affect response to vaccines. Current, recent, or upcoming anesthesia/surgery/hospitalization is not a contraindication to vaccination, but certain factors might lead a provider to consider current, recent, or upcoming anesthesia/surgery/hospitalization as a precaution (16-35). Efforts should be made to ensure vaccine administration during the hospitalization or at discharge. For patients who are deemed moderately or severely ill throughout the hospitalization, vaccination should occur at the earliest opportunity (i.e., during immediate post-hospitalization follow-up care, including home or office visits) when patients' clinical symptoms have improved. A personal or family history of seizures is a precaution for MMRV vaccination; this is because a recent study found an increased risk for febrile seizures in children 12-23 months who receive MMRV compared with MMR and varicella vaccine (36). Clinicians or other health-care providers might misperceive certain conditions or circumstances as valid contraindications or precautions to vaccination when they actually do not preclude vaccination (2) (Table 4-2). These misperceptions result in missed opportunities to administer recommended vaccines (37). Routine physical examinations and procedures (e.g., measuring temperatures) are not prerequisites for vaccinating persons who appear to be healthy. The provider should ask the parent or guardian if the child is ill. If the child has a moderate or severe illness, the vaccination should be postponed. Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable cause, within 7 days of

administration of previous dose of DTP or DTaP Hypersensitivity to yeast Age <6 weeks
 Preterm birth (MenACWY-CRM)(i) Pregnancy Known severe immunodeficiency (e.g.,
 from hematologic and solid tumors, receipt of chemotherapy, congenital
 immunodeficiency, long-term immunosuppressive therapy(c) or patients with HIV
 infection who are severely immunocompromised) Family history of altered
 immunocompetence(m) History of thrombocytopenia or thrombocytopenic purpura
 Need for tuberculin skin testing or interferon-gamma release assay (IGRA) testing(l)
 Moderate or severe acute illness with or without fever SCID History of intussusception
 Moderate or severe acute illness with or without fever Encephalopathy (e.g., coma,
 decreased level of consciousness, prolonged seizures), not attributable to another
 identifiable cause, within 7 days of administration of previous dose of DTP, DTaP, or
 Tdap Progressive or unstable neurological disorder, uncontrolled seizures, or
 progressive encephalopathy until a treatment regimen has been established and the
 condition has stabilized History of Arthus-type hypersensitivity reactions after a
 previous dose of diphtheria-toxoid—containing or tetanus-toxoid-containing vaccine;
 defer vaccination until at least 10 years have elapsed since the last
 tetanus-toxoid-containing vaccine Moderate or severe acute illness with or without
 fever Known severe immunodeficiency (e.g., from hematologic and solid tumors, receipt
 of chemotherapy, congenital immunodeficiency, long-term immunosuppressive
 therapy(c) or patients with HIV infection who are severely immunocompromised)(j)
 Pregnancy Family history of altered immunocompetence(m) Abbreviations: DT =
 diphtheria and tetanus toxoids; DTaP = diphtheria and tetanus toxoids and acellular
 pertussis; DTP = diphtheria toxoid, tetanus toxoid, and pertussis; GBS = Guillain-Barré
 syndrome; Hib = Haemophilus influenzae type b; HIV = human immunodeficiency virus;
 HPV = human papillomavirus; IIV = inactivated influenza vaccine; IPV = inactivated
 poliovirus; LAIV = live, attenuated influenza vaccine; MenACWY = quadrivalent
 meningococcal conjugate vaccine; MMR = measles, mumps, and rubella; MPSV4 =

quadrivalent meningococcal polysaccharide vaccine; PCV13 = pneumococcal conjugate vaccine; PPSV23= pneumococcal polysaccharide vaccine; SCID = severe combined immunodeficiency; RIV=recombinant influenza vaccine; Td = tetanus and diphtheria toxoids; Tdap = tetanus toxoid, reduced diphtheria toxoid, and acellular pertussis. (a) Events or conditions listed as precautions should be reviewed carefully. Benefits of and risks for administering a specific vaccine to a person under these circumstances should be considered. If the risk from the vaccine is believed to outweigh the benefit, the vaccine should not be administered. If the benefit of vaccination is believed to outweigh the risk, the vaccine should be administered. Whether and when to administer DTaP to children with proven or suspected underlying neurologic disorders should be decided on a case-by-case basis. (b) Only persons with laboratory confirmation of immunity according to strict guidance at Laboratory Testing Requirements for Vaccination with Dengvaxia Dengue Vaccine should receive dengue vaccination. (c) Substantially immunosuppressive steroid dose is considered to be ≥ 2 weeks of daily receipt of 20 mg or 2 mg/kg body weight of prednisone or equivalent. (d) HPV vaccine is not recommended during pregnancy (e) When applying this contraindication to cclIV, the history of severe allergic reaction (e.g., anaphylaxis) must be specific to the event occurring following a dose of cclIV. Likewise, when applying this contraindication to RIV, the history of severe allergic reaction (e.g., anaphylaxis) must be specific to the event occurring following a dose of RIV. A history of severe allergic reaction (e.g., anaphylaxis) to a non-cclIV vaccine or to a component specific to components not contained in cclIV, is a precaution to cclIV. A history of severe allergic reaction (e.g., anaphylaxis) to a non-RIV vaccine or to a component specific to components not contained in RIV is a precaution to RIV. (f) In addition, ACIP recommends LAIV not be used for pregnant women, immunosuppressed persons, and children aged 2-4 years who have asthma or who have had a wheezing episode noted in the medical record within the past 12 months, or for whom parents report that a health care provider

stated that they had wheezing or asthma within the last 12 months. LAIV should not be administered to persons who have taken influenza antiviral medications within the previous 48 hours. Persons who care for severely immunosuppressed persons who require a protective environment should not receive LAIV, or should avoid contact with such persons for 7 days after receipt. (g) See reference: Grohskopf LA, Alyanak E, Ferdinands JM, et al. Prevention and Control of Seasonal Influenza with Vaccines: Recommendations of the Advisory Committee on Immunization Practices, United States, 2021-2022 Influenza Season. MMWR Recomm Rep 2021;70(No. RR-5):1-30. (h) These values are based on the clearance of the particular antiviral. LAIV4 should not be administered to persons who have taken oseltamivir or zanamivir within the previous 48 hours, peramivir within the previous 5 days, or baloxavir within the previous 17 days. This “contraindication” is due to concern with reduced effectiveness of the vaccine. To obtain specific information, please refer to Grohskopf LA, Alyanak, E, Broder KR, et. al. Prevention and Control of Seasonal Influenza with Vaccines: Recommendations of the Advisory Committee on Immunization Practices — United States, 2020–21 Influenza Season. MMWR Recomm Rep 2020;69 (No. RR-8:1-26. Also at <https://www.cdc.gov/mmwr/volumes/69/rr/pdfs/rr6908a1-H.pdf> (i) This precaution applies to infants younger than 9 months old (j) HIV-infected children may receive varicella vaccine if CD4+ T-lymphocyte count is $\geq 15\%$ and should receive MMR vaccine if they are aged ≥ 12 months and do not have evidence of current severe immunosuppression (i.e., individuals aged ≤ 5 years must have CD4+T lymphocyte [CD4] percentages $\geq 15\%$ for ≥ 6 months; and individuals aged > 5 years must have CD4+percentages $\geq 15\%$ and CD4+ ≥ 200 lymphocytes/mm³ for ≥ 6 months) or other current evidence of measles, rubella, and mumps immunity. In cases when only CD4+cell counts or only CD4+percentages are available for those older than age 5 years, the assessment of severe immunosuppression can be based on the CD4+values (count or percentage) that are available. In cases when CD4+percentages are not

available for those aged ≤ 5 years, the assessment of severe immunosuppression can be based on age-specific CD4+counts at the time CD4+counts were measured; i.e., absence of severe immunosuppression is defined as ≥ 6 months above age-specific CD4+count criteria: CD4+count > 750 lymphocytes/mm³ while aged ≤ 12 months and CD4+count ≥ 500 lymphocytes/mm³ while aged 1 through 5 years. Sources: (1, 50). (k) MMR and varicella-containing vaccines can be administered on the same day. If not administered on the same day, these vaccines should be separated by at least 28 days. (l) If active tuberculosis is suspected, MMR should be delayed. Measles vaccination might suppress tuberculin reactivity temporarily. Measles-containing vaccine can be administered on the same day as tuberculin skin or IGRA testing. If testing cannot be performed until after the day of MMR vaccination, the test should be postponed for ≥ 4 weeks after the vaccination. If an urgent need exists to skin test or IGRA, do so with the understanding that reactivity might be reduced by the vaccine. (m) family history of congenital or hereditary immunodeficiency in first-degree relatives (e.g., parents and siblings), unless the immune competence of the potential vaccine recipient has been substantiated clinically or verified by a laboratory (n) For RV1 only, based on latex in product/packaging. Note that anaphylactic allergy to latex is covered in the contraindication, and would also be isolated to RV 1 in the case of latex. For more details, see (55). (o) No adverse events associated with the use of aspirin or aspirin-containing products after varicella vaccination have been reported; however, the vaccine manufacturer recommends that vaccine recipients avoid using aspirin or aspirin-containing products for 6 weeks after receiving varicella vaccines because of the association between aspirin use and Reye syndrome after varicella. Vaccination with subsequent close monitoring should be considered for children who have rheumatoid arthritis or other conditions requiring therapeutic aspirin. The risk for serious complications associated with aspirin is likely to be greater in children in whom natural varicella develops than it is in children who receive the vaccine containing attenuated

VZV. No association has been documented between Reye syndrome and analgesics or antipyretics that do not contain aspirin.” Top of Page Abbreviations: DT = diphtheria and tetanus toxoids; DTP = diphtheria toxoid, tetanus toxoid, and pertussis; DTaP = diphtheria and tetanus toxoids and acellular pertussis; GBS = Guillain-Barré syndrome; HBsAg = hepatitis B surface antigen; Hib = Haemophilus influenzae type b; HIV = human immunodeficiency virus; HPV = human papillomavirus; IIV = inactivated influenza vaccine; IPV = inactivated poliovirus; LAIV = live, attenuated influenza vaccine; MenACWY = quadrivalent meningococcal conjugate vaccine; MMR = measles, mumps, and rubella; MPSV4 = quadrivalent meningococcal polysaccharide vaccine; PCV = pneumococcal conjugate vaccine; PPSV23= pneumococcal polysaccharide vaccine; Td = tetanus and diphtheria toxoids; Tdap = tetanus toxoid, reduced diphtheria toxoid, and acellular pertussis. (a) Antibacterial drugs might interfere with Ty21a oral typhoid vaccine, and certain antiviral drugs might interfere with varicella-containing vaccines and LAIV4. (b) Hepatitis B vaccination should be deferred for infants weighing <2,000 g if the mother is documented to be HBsAg negative. Vaccination should commence at chronological age 1 month or at hospital discharge. For infants born to HBsAg-positive women, hepatitis B immune globulin and hepatitis B vaccine should be administered within 12 hours after birth, regardless of weight. (c) An exception is Guillain-Barré syndrome within 6 weeks of a dose of influenza vaccine or tetanus-toxoid-containing vaccine, which are precautions for influenza vaccines and tetanus-toxoid containing vaccines, respectively. (d) MMR and varicella vaccines can be administered on the same day. If not administered on the same day, these vaccines should be separated by at least 28 days. (e) HIV-infected children should receive immune globulin after exposure to measles. HIV-infected children can receive varicella and measles vaccine if CD4+ T-lymphocyte count is >15%. (55). (f) Measles vaccination might suppress tuberculin reactivity temporarily. Measles-containing vaccine can be administered on the same day as tuberculin skin or IGRA testing. If testing cannot be

performed until after the day of MMR vaccination, the test should be postponed for at least 4 weeks after the vaccination. If an urgent need exists to skin test or IGRA, do so with the understanding that reactivity might be reduced by the vaccine. (g) If a vaccinee experiences a presumed vaccine-related rash 7-25 days after vaccination, the person should avoid direct contact with immunocompromised persons for the duration of the rash. Top of Page Top of Page To receive email updates about this page, enter your email address:

