

Persons using assistive technology might not be able to fully access information in this file. For assistance, please send e-mail to: mmwrq@cdc.gov. Type 508 Accommodation and the title of the report in the subject line of e-mail. Fourteen states (Arizona, Connecticut, Florida, Georgia, Indiana, Mississippi, New York, Ohio, Oregon, Pennsylvania, Texas, Virginia, Washington, and West Virginia) have reported investigations of multiple schoolchildren who have developed rashes. This report summarizes the investigation by state and local health departments of these rashes, which have occurred during October 2001 through February 2002, and provides examples for four states. Preliminary findings indicate that further investigation is needed to determine whether a common etiology for these rashes exists.

UNITED STATES

The first reported incident occurred October 4, 2001, in Indiana, followed by cases in Virginia that began November 20. Subsequent cases of rashes began in late January and occurred as recently as February 21. Rashes have been reported primarily from elementary schools but also among students in a few middle and high schools. The number of affected students in each state ranges from <10 to approximately 600. A few teachers and school staff have been affected, but rarely parents or siblings. Characteristics of the rashes vary, but onset has generally been acute, typically with maculopapular erythematous lesions---possibly in a reticulated pattern---on the face, neck, hands, or arms; duration

of the rash varied but in most reports it was highly pruritic. The rashes were not attributed to a defined environmental exposure or infectious agent. Children with rashes were afebrile and usually had no other associated signs or symptoms. The rashes lasted from a few hours to 2 weeks and appeared to be self-limiting. Secondary transmission has not been reported, but in-school "sympathy" cases have reportedly occurred. Diagnoses by clinicians who have examined children have included viral exanthem, contact or atopic dermatitis, eczema, chemical exposure, impetigo, and poison ivy. Approximately 40 serum samples collected in four states have been PCR or IgM negative for parvovirus B19 (1); 22 nasal swab samples have been negative for enterovirus. Environmental assessments have not identified environmental causes.

CASE REPORTS

Indiana. During October 4--November 2, 2001, rashes appeared among 18 third-grade students in an elementary school of 390 students; one substitute teacher also developed rash. No rashes among family members were reported. The rash most often began on the face, then spread to the upper extremities; most rashes occurred on exposed skin. Clinical signs---including reddish welt-type itchy rash on face and upper extremities, swollen eyes, and smooth pink cheeks---degrees of coloration, and prominence of rash varied among the children. Diagnoses in the few children examined by family physicians varied

and included contact dermatitis, chemical exposure, impetigo, and poison ivy. Because parvovirus B19 infection was diagnosed in one third-grade student on August 30, 2001, the Indiana State Department of Health collected serum specimens from four students with rashes to assess whether they had parvovirus B19 infection. All specimens tested negative for the presence of IgM antibodies. Laboratory data analysis, interviews, a building survey, and examination of the children did not identify a cause for the rashes.

Pennsylvania. The initial report of rash occurred on January 31, 2002, among 54 elementary school students who had contact dermatitis diagnosed by a local health-care provider. To date, approximately 575 cases of rashes have been reported to the Pennsylvania Department of Health; 58 schools and child-care centers have reported cases (range: one--168 cases per facility). Most cases are in elementary and middle school students, with female cases outnumbering males. The rash has been characterized as bright-red, itchy or burning, and macular, occasionally with an urticarial or papular component. The rash may be evanescent, or remain for as long as 2 weeks; recurrent cases have been reported. There have been no other associated symptoms. Among the 54 students reported initially, serologies for parvovirus B19 were drawn on 13 cases; all were negative for IgM. PCR for parvovirus B19 was negative for 10 cases; results are pending for the remainder. Another health-care provider reported that results of nonserological (biopsy) specimens from his patients were consistent with viral

exanthem. Environmental investigations at five schools have not yet identified an environmental source of the rashes.

These investigations have included sampling for dust mite and cockroach allergens, solvents and cleaners, and fungal or bacterial culture growth. Air and surface cultures are still pending.

Oregon. During February 2002, outbreaks of rashes of acute onset and short duration occurred among students in two Oregon schools. Starting February 4, rashes were reported in 53 children and 11 adults in an elementary school of 589 students in southwestern Oregon; 54 (84%) were female. The rash, which appeared on cheeks and arms, was itchy and had a sunburned appearance but no systemic symptoms. A panel of dermatologists who examined 28 of the affected children reported that the rash resembled fifth disease but that several characteristics were not compatible with that diagnosis.

Testing for parvovirus in two children was negative. Extensive questioning and environmental inspection did not uncover a source of the rash. Beginning February 21, rashes were reported by 84 children and seven adults in a middle school of 314 students in northwestern Oregon; 67 (74%) affected persons were female. No known links existed between the two schools. Rashes were characterized in a variety of ways, including eczema, and as a sunburned, itchy rash on face, arms, neck, and back; no other symptoms were reported. Tests for parvovirus in six persons were negative. An environmental evaluation of the school found no explanation for the rash. In both schools, rash improved in several children when they left school but recurred when

they returned to school.

Connecticut. On February 20, the Connecticut Department of Public Health was notified of nine

elementary schoolchildren with rashes. On February 21, an additional 16 children were identified with a similar rash. The children,

all fourth-graders, represented four classrooms in a school of 253 students and 12 classrooms. The acute rash appeared on

the trunk and extremities and was characterized by erythema and pruritis. The children were afebrile and had no other

symptoms. The illness lasted 24--72 hours. A dermatologist who examined three children attributed the rashes to an allergic reaction

to an environmental exposure. Rashes were not reported among parents or siblings of affected children. The local health

director and the state Environmental Epidemiology Program are collaborating to identify potential environmental causes. The

school was closed for 1 day to clean the classrooms, check air-handling units, and replace air filters.

PUBLIC HEALTH RESPONSE

CDC is working with state and local health and education agencies in these investigations to determine if affected

children within and between schools have developed rash as a result of a common etiology. CDC is systematically

compiling information about 1) date of onset and duration of rash; 2) settings of and circumstances surrounding the rash's

appearance; 3) the number, age, and sex of affected persons; 4) the appearance and

characteristics of the rash; 5) additional signs or symptoms, diagnoses, and treatments; and 6) investigational methods used (e.g., interviews or questionnaires, biologic sampling, and environmental sampling). To facilitate the collection of standardized information, CDC has developed and distributed to health departments a document with suggested approaches for investigating reports of rashes among groups of schoolchildren. In addition, CDC requests that dermatologists and other health-care providers who have examined affected children share their clinical observations, diagnoses, and photographs with a CDC dermatologist (bdt1@cdc.gov).

This information will help CDC assess whether affected children within and between schools developed rash caused by a common etiology. Local health and school officials with information about rashes among groups of schoolchildren in their jurisdiction are asked to report this information to their state health department.

Reported by: M Cartter, MD, P Mshar, Connecticut Dept of Public Health. H Messersmith, Indiana State Dept of Health, Epidemiology Resource Center. K Southwick, MD, K Hedberg, MD, Oregon Health Div. Y Chilcoat, Jackson County Public Health Dept, Medford, Oregon.

N Nunley, J Hersh, MEd, K Nalluswami, MD, M Moll, MD, K Waller, MD, Pennsylvania Dept of Health, Bureau of Epidemiology.

R Moodispaugh, R Swiger, Harrison-Clarksburg Health Dept Clarksburg, West Virginia. C Rubin, DVM, Div of Environmental Hazards and Health Effects, National Center for Environmental Health; A Tepper, PhD, B Lushniak, MD, Div of Surveillance, Hazard Evaluation, and

Field Studies, National Institute for Occupational Safety and Health; N Khetsuriani, MD, Div of Viral and Rickettsial Diseases, National Center for Infectious Diseases; L Kolbe, PhD, Div of Adolescent and School Health, National Center for Chronic Disease Prevention and Health Promotion; N Smith, MPH, EIS Officer, CDC.

Editorial Note:

With 53 million young people attending 117,000 schools each school day in the United States, it is expected that rashes from a wide range of causes will be observed. Environmental factors or infectious agents can cause rashes among groups of school-aged children. Rashes caused by infectious agents usually are preceded or accompanied by symptoms such as headache or fever. However, in these reports, none of the children showed signs of systemic illness, and the rash appeared to be self-limiting.

Potential environmental causes of rashes include biologic contaminants (e.g., bacteria and fungi), chemical agents (e.g., cleaning products and pesticide residues), physical agents (e.g., fiberglass), insects (e.g., biting flies and moths), and allergens

(e.g., dust mites) (2--4). If one of these environmental causes is suspected, appropriate environmental experts should be consulted.

The most commonly identified viral agent associated with rashes in school-aged children is parvovirus B19, which causes erythema infectiosum (i.e., fifth disease). Fifth disease is a mild rash illness

characterized by a "slapped-cheek" rash on the face and a lacy red rash on the trunk and limbs, which may itch; it usually resolves within 7--10 days. Low-grade fever, malaise, or upper respiratory symptoms usually precede the rash. Other manifestations of parvovirus B19 infection include arthritis and arthralgia (especially in adults), transient crisis of aplastic anemia (in persons with certain hematologic disorders such as sickle-cell anemia), neutropenia, and thrombocytopenia. In pregnant women, parvovirus B19 infection may be associated with miscarriage or nonimmune hydrops fetalis (1).

Public health response to rashes of unknown etiology involves an epidemiologic investigation that includes consultation with facilities and maintenance staff familiar with the physical plant, examination of the rash by a dermatologist, and, when appropriate, collection and analysis of biologic specimens. To date, reports from states do not document a common cause or demonstrate that all children are experiencing the same rash. State and local health departments, in collaboration with CDC, continue to investigate these and other reports of rashes among groups of schoolchildren.

Acknowledgements

This report is based on data contributed by C McRill, MD, K Komatsu, MPH, W Humble, MPH, Arizona Dept of Health Svcs; L Sands, DO, MPH, Maricopa County Dept of Public Health, Phoenix, Arizona. T Wegrzyn, MPH, J Hadler, MD, Connecticut Dept of Public Health. S Wiersma, MD, Florida Dept of Health. P Blake, MD, S Lance-Parker, PhD, C

Morin, MD, Georgia Dept of Human Resources, Div of Public Health. R Teclaw, DVM, M Wilkinson, Indiana State Dept of Health. M Currier, MD, S Slavinski, DVM, Mississippi Dept of Health. H Mackley, MPH, B Asante, MD, New York City Dept of Health. M Kohn, MD, E DeBess, DVM, L Davidoff, MD, M Scott, M Heumann, MA, Oregon Health Div; G Stevens, V Barbour, MSN, J Baures, J Manwaring, B Thomas, G Chakarun, Jackson County Public Health Dept, Medford; R Parlier, M Jaqua, M Breedlove, MBA, Yamhill County Health Dept, McMinnville, Oregon. J Rankin, Jr., DVM, C Coventon, MD, Pennsylvania Dept of Health, Bureau of Epidemiology. F Sassano, Bucks County Dept of Health, Doylestown; J Maher, MD, E Walls, Chester County Health Dept, Westchester; C Baysinger, M Supplee, MS, Montgomery County Human Svcs Center, Norristown; J Jahre, MD, St. Luke's Hospital Network, Bethlehem; N Sykes, MD, Jefferson Medical College, Philadelphia, Pennsylvania. R Stroube, MD, E Barrett, DMD, S Jenkins, VMD, Virginia Health Dept; J Florance MD, A Ansher MD, L Estrada MPH, B Bradshaw, Prince William Health Dept, Manassas, Virginia. J Hofmann, MD, J Van Eenwyk, PhD, Washington Dept of Health. L Haddy, MA, D Bixler, MD, West Virginia Dept of Health and Human Resources, Bur for Public Health; P Gordon, MD, Harrison-Clarksburg Health Dept Clarksburg, West Virginia. J Perdue, Texas Dept of Health; and other state and local health and education departments and schools. A Henderson, PhD, Div of Environmental Hazards and Health Effects, National Center for

Environmental Health. A Adija, MD, K

Griffith, MD, EIS officers, CDC.

References

Brown KE. Human parvovirus B19 epidemiology and clinical manifestations. In: Anderson LJ, Young NS, eds. Human parvovirus B19 (monographs in virology. vol 20). Basel, Switzerland: Karger 1997:42--60.

Environmental Protection Agency. Indoor air pollution: an introduction for health professionals. Washington, DC: Environmental Protection Agency, 1994. Available at <http://www.epa.gov/iaq/pubs/hpguide.htm>. Accessed February 2002.

CDC. Moth-associated dermatitis---Cozumel, Mexico. MMWR 1990;39:219--20.

CDC. Rash illness associated with gypsy moth caterpillars---Pennsylvania. MMWR 1982;31:169--70.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services. References to non-CDC sites on the Internet are provided as a service to MMWR readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of pages found at these sites.

Disclaimer

All MMWR HTML versions of articles are electronic conversions from ASCII text into HTML. This conversion may have resulted in character translation or format errors in the HTML version.

Users should not rely on this HTML document, but are referred to the electronic PDF version and/or

the original MMWR paper copy for the official text, figures, and tables.

An original paper copy of this issue can be obtained from the Superintendent of Documents,

U.S. Government Printing Office (GPO), Washington, DC 20402-9371; telephone: (202) 512-1800.

Contact GPO for current prices.

****Questions or messages regarding errors in formatting should be addressed to mmwrq@cdc.gov.**

Page converted: 2/28/2002

The first reported incident occurred October 4, 2001, in Indiana, followed by cases in Virginia that began November

20. Subsequent cases of rashes began in late January and occurred as recently as February 21. Rashes have been reported

primarily from elementary schools but also among students in a few middle and high schools. The number of affected students in

each state ranges from <10 to approximately 600. A few teachers and school staff have been affected, but rarely parents or siblings.

Characteristics of the rashes vary, but onset has generally been acute, typically with maculopapular erythematous

lesions---possibly in a reticulated pattern---on the face, neck, hands, or arms; duration of the rash varied but in most reports it

was highly pruritic. The rashes were not attributed to a defined environmental exposure or infectious agent. Children with rashes were afebrile and usually had no other associated signs or symptoms. The rashes lasted from a few hours to 2 weeks and appeared to be self-limiting. Secondary transmission has not been reported, but in-school "sympathy" cases have reportedly occurred. Diagnoses by clinicians who have examined children have included viral exanthem, contact or atopic dermatitis, eczema, chemical exposure, impetigo, and poison ivy. Approximately 40 serum samples collected in four states have been PCR or IgM negative for parvovirus B19 (1); 22 nasal swab samples have been negative for enterovirus. Environmental assessments have not identified environmental causes.

CASE REPORTS

Indiana. During October 4--November 2, 2001, rashes appeared among 18 third-grade students in an elementary school of 390 students; one substitute teacher also developed rash. No rashes among family members were reported. The rash most often began on the face, then spread to the upper extremities; most rashes occurred on exposed skin. Clinical signs---including reddish welt-type itchy rash on face and upper extremities, swollen eyes, and smooth pink cheeks---degrees of coloration, and prominence of rash varied among the children. Diagnoses in the few children examined by family physicians varied and included contact dermatitis, chemical exposure, impetigo, and poison ivy. Because

parvovirus B19 infection was diagnosed

in one third-grade student on August 30, 2001, the Indiana State Department of Health collected serum specimens from

four students with rashes to assess whether they had parvovirus B19 infection. All specimens tested negative for the presence

of IgM antibodies. Laboratory data analysis, interviews, a building survey, and examination of the children did not identify

a cause for the rashes.

Pennsylvania. The initial report of rash occurred on January 31, 2002, among 54 elementary school students who

had contact dermatitis diagnosed by a local health-care provider. To date, approximately 575 cases of rashes have been reported

to the Pennsylvania Department of Health; 58 schools and child-care centers have reported cases (range: one--168 cases

per facility). Most cases are in elementary and middle school students, with female cases outnumbering males. The rash has

been characterized as bright-red, itchy or burning, and macular, occasionally with an urticarial or papular component. The

rash may be evanescent, or remain for as long as 2 weeks; recurrent cases have been reported. There have been no other

associated symptoms. Among the 54 students reported initially, serologies for parvovirus B19 were drawn on 13 cases; all were

negative for IgM. PCR for parvovirus B19 was negative for 10 cases; results are pending for the remainder. Another

health-care provider reported that results of nonserological (biopsy) specimens from his patients were consistent with viral

exanthem. Environmental investigations at five schools have not yet identified an

environmental source of the rashes.

These investigations have included sampling for dust mite and cockroach allergens, solvents and cleaners, and fungal or

bacterial culture growth. Air and surface cultures are still pending.

Oregon. During February 2002, outbreaks of rashes of acute onset and short duration occurred among students in

two Oregon schools. Starting February 4, rashes were reported in 53 children and 11 adults in an elementary school of

589 students in southwestern Oregon; 54 (84%) were female. The rash, which appeared on cheeks and arms, was itchy and had

a sunburned appearance but no systemic symptoms. A panel of dermatologists who examined 28 of the affected

children reported that the rash resembled fifth disease but that several characteristics were not compatible with that diagnosis.

Testing for parvovirus in two children was negative. Extensive questioning and environmental inspection did not uncover a source

of the rash. Beginning February 21, rashes were reported by 84 children and seven adults in a middle school of 314 students

in northwestern Oregon; 67 (74%) affected persons were female. No known links existed between the two schools. Rashes

were characterized in a variety of ways, including eczema, and as a sunburned, itchy rash on face, arms, neck, and back; no

other symptoms were reported. Tests for parvovirus in six persons were negative. An environmental evaluation of the school

found no explanation for the rash. In both schools, rash improved in several children when they left school but recurred when

they returned to school.

Connecticut. On February 20, the Connecticut Department of Public Health was notified of nine elementary schoolchildren with rashes. On February 21, an additional 16 children were identified with a similar rash. The children, all fourth-graders, represented four classrooms in a school of 253 students and 12 classrooms. The acute rash appeared on the trunk and extremities and was characterized by erythema and pruritis. The children were afebrile and had no other symptoms. The illness lasted 24--72 hours. A dermatologist who examined three children attributed the rashes to an allergic reaction to an environmental exposure. Rashes were not reported among parents or siblings of affected children. The local health director and the state Environmental Epidemiology Program are collaborating to identify potential environmental causes. The school was closed for 1 day to clean the classrooms, check air-handling units, and replace air filters.

PUBLIC HEALTH RESPONSE

CDC is working with state and local health and education agencies in these investigations to determine if affected children within and between schools have developed rash as a result of a common etiology. CDC is systematically compiling information about 1) date of onset and duration of rash; 2) settings of and circumstances surrounding the rash's appearance; 3) the number, age, and sex of affected persons; 4) the appearance and characteristics of the rash; 5) additional signs

or symptoms, diagnoses, and treatments; and 6) investigational methods used (e.g., interviews or questionnaires, biologic sampling, and environmental sampling). To facilitate the collection of standardized information, CDC has developed and distributed to health departments a document with suggested approaches for investigating reports of rashes among groups of schoolchildren. In addition, CDC requests that dermatologists and other health-care providers who have examined affected children share their clinical observations, diagnoses, and photographs with a CDC dermatologist (bdt1@cdc.gov).

This information will help CDC assess whether affected children within and between schools developed rash caused by a common etiology. Local health and school officials with information about rashes among groups of schoolchildren in their jurisdiction are asked to report this information to their state health department.

Reported by: M Cartter, MD, P Mshar, Connecticut Dept of Public Health. H Messersmith, Indiana State Dept of Health, Epidemiology Resource Center. K Southwick, MD, K Hedberg, MD, Oregon Health Div. Y Chilcoat, Jackson County Public Health Dept, Medford, Oregon.

N Nunley, J Hersh, MEd, K Nalluswami, MD, M Moll, MD, K Waller, MD, Pennsylvania Dept of Health, Bureau of Epidemiology.

R Moodispaugh, R Swiger, Harrison-Clarksburg Health Dept Clarksburg, West Virginia. C Rubin, DVM, Div of Environmental Hazards

and Health Effects, National Center for Environmental Health; A Tepper, PhD, B Lushniak, MD, Div of Surveillance, Hazard Evaluation, and

Field Studies, National Institute for Occupational Safety and Health; N Khetsuriani, MD,

Div of Viral and Rickettsial Diseases, National Center

for Infectious Diseases; L Kolbe, PhD, Div of Adolescent and School Health, National Center for Chronic Disease Prevention and Health

Promotion; N Smith, MPH, EIS Officer, CDC.

Editorial Note:

With 53 million young people attending 117,000 schools each school day in the United States, it is

expected that rashes from a wide range of causes will be observed. Environmental factors or infectious agents can cause rashes

among groups of school-aged children. Rashes caused by infectious agents usually are preceded or accompanied by symptoms such

as headache or fever. However, in these reports, none of the children showed signs of systemic illness, and the rash appeared

to be self-limiting.

Potential environmental causes of rashes include biologic contaminants (e.g., bacteria and fungi), chemical agents

(e.g., cleaning products and pesticide residues), physical agents (e.g., fiberglass), insects (e.g., biting flies and moths), and

allergens

(e.g., dust mites) (2--4). If one of these environmental causes is suspected, appropriate environmental experts should

be consulted.

The most commonly identified viral agent associated with rashes in school-aged children is parvovirus B19, which

causes erythema infectiosum (i.e., fifth disease). Fifth disease is a mild rash illness characterized by a "slapped-cheek" rash on the

face and a lacy red rash on the trunk and limbs, which may itch; it usually resolves within 7--10 days. Low-grade fever, malaise, or upper respiratory symptoms usually precede the rash. Other manifestations of parvovirus B19 infection include arthritis and arthralgia (especially in adults), transient crisis of aplastic anemia (in persons with certain hematologic disorders such as sickle-cell anemia), neutropenia, and thrombocytopenia. In pregnant women, parvovirus B19 infection may be associated with miscarriage or nonimmune hydrops fetalis (1).

Public health response to rashes of unknown etiology involves an epidemiologic investigation that includes consultation with facilities and maintenance staff familiar with the physical plant, examination of the rash by a dermatologist, and, when appropriate, collection and analysis of biologic specimens. To date, reports from states do not document a common cause or demonstrate that all children are experiencing the same rash. State and local health departments, in collaboration with CDC, continue to investigate these and other reports of rashes among groups of schoolchildren.

Acknowledgements

This report is based on data contributed by C McRill, MD, K Komatsu, MPH, W Humble, MPH, Arizona Dept of Health Svcs; L Sands, DO, MPH, Maricopa County Dept of Public Health, Phoenix, Arizona. T Wegrzyn, MPH, J Hadler, MD, Connecticut Dept of Public Health. S Wiersma, MD, Florida Dept of Health. P Blake, MD, S Lance-Parker, PhD, C Morin, MD, Georgia Dept of Human Resources, Div of Public Health. R Teclaw, DVM, M

Wilkinson, Indiana State Dept

of Health. M Currier, MD, S Slavinski, DVM, Mississippi Dept of Health. H Mackley, MPH, B Asante, MD, New York

City Dept of Health. M Kohn, MD, E DeBess, DVM, L Davidoff, MD, M Scott, M Heumann, MA, Oregon Health Div;

G Stevens, V Barbour, MSN, J Baures, J Manwaring, B Thomas, G Chakarun, Jackson County Public Health Dept, Medford;

R Parlier, M Jaqua, M Breedlove, MBA, Yamhill County Health Dept, McMinnville, Oregon. J Rankin, Jr., DVM,

C Coventon, MD, Pennsylvania Dept of Health, Bureau of Epidemiology. F Sassano, Bucks County Dept of

Health, Doylestown; J Maher, MD, E Walls, Chester County Health Dept, Westchester; C Baysinger, M Supplee, MS,

Montgomery County Human Svcs Center, Norristown; J Jahre, MD, St. Luke's Hospital Network, Bethlehem; N Sykes, MD,

Jefferson Medical College, Philadelphia, Pennsylvania. R Stroube, MD, E Barrett, DMD, S Jenkins, VMD, Virginia Health Dept;

J Florance MD, A Ansher MD, L Estrada MPH, B Bradshaw, Prince William Health Dept, Manassas, Virginia. J

Hofmann, MD, J Van Eenwyk, PhD, Washington Dept of Health. L Haddy, MA, D Bixler, MD, West Virginia Dept of Health

and Human Resources, Bur for Public Health; P Gordon, MD, Harrison-Clarksburg Health Dept Clarksburg, West Virginia.

J Perdue, Texas Dept of Health; and other state and local health and education departments and schools. A Henderson,

PhD, Div of Environmental Hazards and Health Effects, National Center for Environmental Health. A Adija, MD, K

Griffith, MD, EIS officers, CDC.

References

Brown KE. Human parvovirus B19 epidemiology and clinical manifestations. In: Anderson LJ, Young NS, eds. Human parvovirus B19

(monographs in virology. vol 20). Basel, Switzerland: Karger 1997:42--60.

Environmental Protection Agency. Indoor air pollution: an introduction for health professionals. Washington, DC: Environmental

Protection Agency, 1994. Available at <http://www.epa.gov/iaq/pubs/hpguide.htm>.

Accessed February 2002.

CDC. Moth-associated dermatitis---Cozumel, Mexico. MMWR

1990;39:219--20.

CDC. Rash illness associated with gypsy moth caterpillars---Pennsylvania. MMWR

1982;31:169--70.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of

Health and Human Services. References to non-CDC sites on the Internet are

provided as a service to MMWR readers and do not constitute or imply

endorsement of these organizations or their programs by CDC or the U.S.

Department of Health and Human Services. CDC is not responsible for the content of pages found at these sites.

Disclaimer

All MMWR HTML versions of articles are electronic conversions from ASCII text

into HTML. This conversion may have resulted in character translation or format errors in the HTML version.

Users should not rely on this HTML document, but are referred to the electronic PDF version and/or

the original MMWR paper copy for the official text, figures, and tables.

An original paper copy of this issue can be obtained from the Superintendent of Documents,

U.S. Government Printing Office (GPO), Washington, DC 20402-9371; telephone: (202) 512-1800.

Contact GPO for current prices.

****Questions or messages regarding errors in formatting should be addressed to mmwrq@cdc.gov.**

Page converted: 2/28/2002

Indiana. During October 4--November 2, 2001, rashes appeared among 18 third-grade students in an elementary school

of 390 students; one substitute teacher also developed rash. No rashes among family members were reported. The rash

most often began on the face, then spread to the upper extremities; most rashes occurred on exposed skin. Clinical

signs---including reddish welt-type itchy rash on face and upper extremities, swollen eyes, and smooth pink cheeks---degrees of coloration,

and prominence of rash varied among the children. Diagnoses in the few children examined by family physicians varied

and included contact dermatitis, chemical exposure, impetigo, and poison ivy. Because parvovirus B19 infection was diagnosed

in one third-grade student on August 30, 2001, the Indiana State Department of Health

collected serum specimens from four students with rashes to assess whether they had parvovirus B19 infection. All specimens tested negative for the presence of IgM antibodies. Laboratory data analysis, interviews, a building survey, and examination of the children did not identify a cause for the rashes.

Pennsylvania. The initial report of rash occurred on January 31, 2002, among 54 elementary school students who had contact dermatitis diagnosed by a local health-care provider. To date, approximately 575 cases of rashes have been reported to the Pennsylvania Department of Health; 58 schools and child-care centers have reported cases (range: one--168 cases per facility). Most cases are in elementary and middle school students, with female cases outnumbering males. The rash has been characterized as bright-red, itchy or burning, and macular, occasionally with an urticarial or papular component. The rash may be evanescent, or remain for as long as 2 weeks; recurrent cases have been reported. There have been no other associated symptoms. Among the 54 students reported initially, serologies for parvovirus B19 were drawn on 13 cases; all were negative for IgM. PCR for parvovirus B19 was negative for 10 cases; results are pending for the remainder. Another health-care provider reported that results of nonserological (biopsy) specimens from his patients were consistent with viral exanthem. Environmental investigations at five schools have not yet identified an environmental source of the rashes.

These investigations have included sampling for dust mite and cockroach allergens,

solvents and cleaners, and fungal or

bacterial culture growth. Air and surface cultures are still pending.

Oregon. During February 2002, outbreaks of rashes of acute onset and short duration occurred among students in

two Oregon schools. Starting February 4, rashes were reported in 53 children and 11 adults in an elementary school of

589 students in southwestern Oregon; 54 (84%) were female. The rash, which appeared on cheeks and arms, was itchy and had

a sunburned appearance but no systemic symptoms. A panel of dermatologists who examined 28 of the affected

children reported that the rash resembled fifth disease but that several characteristics were not compatible with that diagnosis.

Testing for parvovirus in two children was negative. Extensive questioning and environmental inspection did not uncover a source

of the rash. Beginning February 21, rashes were reported by 84 children and seven adults in a middle school of 314 students

in northwestern Oregon; 67 (74%) affected persons were female. No known links existed between the two schools. Rashes

were characterized in a variety of ways, including eczema, and as a sunburned, itchy rash on face, arms, neck, and back; no

other symptoms were reported. Tests for parvovirus in six persons were negative. An environmental evaluation of the school

found no explanation for the rash. In both schools, rash improved in several children when they left school but recurred when

they returned to school.

Connecticut. On February 20, the Connecticut Department of Public Health was notified of nine

elementary schoolchildren with rashes. On February 21, an additional 16 children were identified with a similar rash. The children, all fourth-graders, represented four classrooms in a school of 253 students and 12 classrooms. The acute rash appeared on the trunk and extremities and was characterized by erythema and pruritis. The children were afebrile and had no other symptoms. The illness lasted 24--72 hours. A dermatologist who examined three children attributed the rashes to an allergic reaction to an environmental exposure. Rashes were not reported among parents or siblings of affected children. The local health director and the state Environmental Epidemiology Program are collaborating to identify potential environmental causes. The school was closed for 1 day to clean the classrooms, check air-handling units, and replace air filters.

PUBLIC HEALTH RESPONSE

CDC is working with state and local health and education agencies in these investigations to determine if affected children within and between schools have developed rash as a result of a common etiology. CDC is systematically compiling information about 1) date of onset and duration of rash; 2) settings of and circumstances surrounding the rash's appearance; 3) the number, age, and sex of affected persons; 4) the appearance and characteristics of the rash; 5) additional signs or symptoms, diagnoses, and treatments; and 6) investigational methods used (e.g., interviews or questionnaires,

biologic sampling, and environmental sampling). To facilitate the collection of standardized information, CDC has developed and distributed to health departments a document with suggested approaches for investigating reports of rashes among groups of schoolchildren. In addition, CDC requests that dermatologists and other health-care providers who have examined affected children share their clinical observations, diagnoses, and photographs with a CDC dermatologist

(bdt1@cdc.gov).

This information will help CDC assess whether affected children within and between schools developed rash caused by a common etiology. Local health and school officials with information about rashes among groups of schoolchildren in their jurisdiction are asked to report this information to their state health department.

Reported by: M Cartter, MD, P Mshar, Connecticut Dept of Public Health. H Messersmith, Indiana State Dept of Health, Epidemiology Resource Center. K Southwick, MD, K Hedberg, MD, Oregon Health Div. Y Chilcoat, Jackson County Public Health Dept, Medford, Oregon.

N Nunley, J Hersh, MEd, K Nalluswami, MD, M Moll, MD, K Waller, MD, Pennsylvania Dept of Health, Bureau of Epidemiology.

R Moodispaugh, R Swiger, Harrison-Clarksburg Health Dept Clarksburg, West Virginia. C Rubin, DVM, Div of Environmental Hazards

and Health Effects, National Center for Environmental Health; A Tepper, PhD, B Lushniak, MD, Div of Surveillance, Hazard Evaluation, and

Field Studies, National Institute for Occupational Safety and Health; N Khetsuriani, MD, Div of Viral and Rickettsial Diseases, National Center

for Infectious Diseases; L Kolbe, PhD, Div of Adolescent and School Health, National

Center for Chronic Disease Prevention and Health

Promotion; N Smith, MPH, EIS Officer, CDC.

Editorial Note:

With 53 million young people attending 117,000 schools each school day in the United States, it is

expected that rashes from a wide range of causes will be observed. Environmental factors or infectious agents can cause rashes

among groups of school-aged children. Rashes caused by infectious agents usually are preceded or accompanied by symptoms such

as headache or fever. However, in these reports, none of the children showed signs of systemic illness, and the rash appeared

to be self-limiting.

Potential environmental causes of rashes include biologic contaminants (e.g., bacteria and fungi), chemical agents

(e.g., cleaning products and pesticide residues), physical agents (e.g., fiberglass), insects (e.g., biting flies and moths), and

allergens

(e.g., dust mites) (2--4). If one of these environmental causes is suspected, appropriate environmental experts should

be consulted.

The most commonly identified viral agent associated with rashes in school-aged children is parvovirus B19, which

causes erythema infectiosum (i.e., fifth disease). Fifth disease is a mild rash illness characterized by a "slapped-cheek" rash on the

face and a lacy red rash on the trunk and limbs, which may itch; it usually resolves within 7--10 days. Low-grade fever, malaise,

or upper respiratory symptoms usually precede the rash. Other manifestations of parvovirus B19 infection include arthritis and arthralgia (especially in adults), transient crisis of aplastic anemia (in persons with certain hematologic disorders such as sickle-cell anemia), neutropenia, and thrombocytopenia. In pregnant women, parvovirus B19 infection may be associated with miscarriage or nonimmune hydrops fetalis (1).

Public health response to rashes of unknown etiology involves an epidemiologic investigation that includes consultation with facilities and maintenance staff familiar with the physical plant, examination of the rash by a dermatologist, and, when appropriate, collection and analysis of biologic specimens. To date, reports from states do not document a common cause or demonstrate that all children are experiencing the same rash. State and local health departments, in collaboration with CDC, continue to investigate these and other reports of rashes among groups of schoolchildren.

Acknowledgements

This report is based on data contributed by C McRill, MD, K Komatsu, MPH, W Humble, MPH, Arizona Dept of Health Svcs; L Sands, DO, MPH, Maricopa County Dept of Public Health, Phoenix, Arizona. T Wegrzyn, MPH, J Hadler, MD, Connecticut Dept of Public Health. S Wiersma, MD, Florida Dept of Health. P Blake, MD, S Lance-Parker, PhD, C Morin, MD, Georgia Dept of Human Resources, Div of Public Health. R Teclaw, DVM, M Wilkinson, Indiana State Dept of Health. M Currier, MD, S Slavinski, DVM, Mississippi Dept of Health. H Mackley, MPH,

B Asante, MD, New York

City Dept of Health. M Kohn, MD, E DeBess, DVM, L Davidoff, MD, M Scott, M Heumann, MA, Oregon Health Div;

G Stevens, V Barbour, MSN, J Baures, J Manwaring, B Thomas, G Chakarun, Jackson County Public Health Dept, Medford;

R Parlier, M Jaqua, M Breedlove, MBA, Yamhill County Health Dept, McMinnville, Oregon. J Rankin, Jr., DVM,

C Coventon, MD, Pennsylvania Dept of Health, Bureau of Epidemiology. F Sassano, Bucks County Dept of

Health, Doylestown; J Maher, MD, E Walls, Chester County Health Dept, Westchester; C Baysinger, M Supplee, MS,

Montgomery County Human Svcs Center, Norristown; J Jahre, MD, St. Luke's Hospital Network, Bethlehem; N Sykes, MD,

Jefferson Medical College, Philadelphia, Pennsylvania. R Stroube, MD, E Barrett, DMD, S Jenkins, VMD, Virginia Health Dept;

J Florance MD, A Ansher MD, L Estrada MPH, B Bradshaw, Prince William Health Dept, Manassas, Virginia. J

Hofmann, MD, J Van Eenwyk, PhD, Washington Dept of Health. L Haddy, MA, D Bixler, MD, West Virginia Dept of Health

and Human Resources, Bur for Public Health; P Gordon, MD, Harrison-Clarksburg Health Dept Clarksburg, West Virginia.

J Perdue, Texas Dept of Health; and other state and local health and education departments and schools. A Henderson,

PhD, Div of Environmental Hazards and Health Effects, National Center for Environmental Health. A Adija, MD, K

Griffith, MD, EIS officers, CDC.

References

Brown KE. Human parvovirus B19 epidemiology and clinical manifestations. In: Anderson LJ, Young NS, eds. Human parvovirus B19 (monographs in virology. vol 20). Basel, Switzerland: Karger 1997:42--60.

Environmental Protection Agency. Indoor air pollution: an introduction for health professionals. Washington, DC: Environmental Protection Agency, 1994. Available at <http://www.epa.gov/iaq/pubs/hpguide.htm>. Accessed February 2002.

CDC. Moth-associated dermatitis---Cozumel, Mexico. MMWR 1990;39:219--20.

CDC. Rash illness associated with gypsy moth caterpillars---Pennsylvania. MMWR 1982;31:169--70.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services. References to non-CDC sites on the Internet are provided as a service to MMWR readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of pages found at these sites.

Disclaimer

All MMWR HTML versions of articles are electronic conversions from ASCII text into HTML. This conversion may have resulted in character translation or format errors in the HTML version.

Users should not rely on this HTML document, but are referred to the electronic PDF version and/or

the original MMWR paper copy for the official text, figures, and tables.

An original paper copy of this issue can be obtained from the Superintendent of Documents,

U.S. Government Printing Office (GPO), Washington, DC 20402-9371; telephone: (202) 512-1800.

Contact GPO for current prices.

****Questions or messages regarding errors in formatting should be addressed to mmwrq@cdc.gov.**

Page converted: 2/28/2002

CDC is working with state and local health and education agencies in these investigations to determine if affected children within and between schools have developed rash as a result of a common etiology. CDC is systematically compiling information about 1) date of onset and duration of rash; 2) settings of and circumstances surrounding the rash's appearance; 3) the number, age, and sex of affected persons; 4) the appearance and characteristics of the rash; 5) additional signs or symptoms, diagnoses, and treatments; and 6) investigational methods used (e.g., interviews or questionnaires, biologic sampling, and environmental sampling). To facilitate the collection of standardized information, CDC has developed and distributed to health departments a document with suggested approaches for investigating reports of rashes among groups of schoolchildren. In addition, CDC requests that dermatologists and other health-care

providers who have examined

affected children share their clinical observations, diagnoses, and photographs with a CDC dermatologist

(bdt1@cdc.gov).

This information will help CDC assess whether affected children within and between schools developed rash caused by a

common etiology. Local health and school officials with information about rashes among groups of schoolchildren in their

jurisdiction are asked to report this information to their state health department.

Reported by: M Cartter, MD, P Mshar, Connecticut Dept of Public Health. H Messersmith, Indiana State Dept of Health,

Epidemiology Resource Center. K Southwick, MD, K Hedberg, MD, Oregon Health Div. Y Chilcoat, Jackson County Public Health Dept, Medford, Oregon.

N Nunley, J Hersh, MEd, K Nalluswami, MD, M Moll, MD, K Waller, MD, Pennsylvania Dept of Health, Bureau of Epidemiology.

R Moodispaugh, R Swiger, Harrison-Clarksburg Health Dept Clarksburg, West Virginia. C Rubin, DVM, Div of Environmental Hazards

and Health Effects, National Center for Environmental Health; A Tepper, PhD, B Lushniak, MD, Div of Surveillance, Hazard Evaluation, and

Field Studies, National Institute for Occupational Safety and Health; N Khetsuriani, MD, Div of Viral and Rickettsial Diseases, National Center

for Infectious Diseases; L Kolbe, PhD, Div of Adolescent and School Health, National Center for Chronic Disease Prevention and Health

Promotion; N Smith, MPH, EIS Officer, CDC.

Editorial Note:

With 53 million young people attending 117,000 schools each school day in the United States, it is

expected that rashes from a wide range of causes will be observed. Environmental factors or infectious agents can cause rashes among groups of school-aged children. Rashes caused by infectious agents usually are preceded or accompanied by symptoms such as headache or fever. However, in these reports, none of the children showed signs of systemic illness, and the rash appeared to be self-limiting.

Potential environmental causes of rashes include biologic contaminants (e.g., bacteria and fungi), chemical agents (e.g., cleaning products and pesticide residues), physical agents (e.g., fiberglass), insects (e.g., biting flies and moths), and allergens

(e.g., dust mites) (2--4). If one of these environmental causes is suspected, appropriate environmental experts should be consulted.

The most commonly identified viral agent associated with rashes in school-aged children is parvovirus B19, which causes erythema infectiosum (i.e., fifth disease). Fifth disease is a mild rash illness characterized by a "slapped-cheek" rash on the face and a lacy red rash on the trunk and limbs, which may itch; it usually resolves within 7--10 days. Low-grade fever, malaise, or upper respiratory symptoms usually precede the rash. Other manifestations of parvovirus B19 infection include arthritis and arthralgia (especially in adults), transient crisis of aplastic anemia (in persons with certain hematologic disorders such as sickle-cell anemia), neutropenia, and thrombocytopenia. In pregnant women,

parvovirus B19 infection may be associated with miscarriage or nonimmune hydrops fetalis (1).

Public health response to rashes of unknown etiology involves an epidemiologic investigation that includes consultation with facilities and maintenance staff familiar with the physical plant, examination of the rash by a dermatologist, and, when appropriate, collection and analysis of biologic specimens. To date, reports from states do not document a common cause or demonstrate that all children are experiencing the same rash. State and local health departments, in collaboration with CDC, continue to investigate these and other reports of rashes among groups of schoolchildren.

Acknowledgements

This report is based on data contributed by C McRill, MD, K Komatsu, MPH, W Humble, MPH, Arizona Dept of Health Svcs; L Sands, DO, MPH, Maricopa County Dept of Public Health, Phoenix, Arizona. T Wegrzyn, MPH, J Hadler, MD, Connecticut Dept of Public Health. S Wiersma, MD, Florida Dept of Health. P Blake, MD, S Lance-Parker, PhD, C Morin, MD, Georgia Dept of Human Resources, Div of Public Health. R Teclaw, DVM, M Wilkinson, Indiana State Dept of Health. M Currier, MD, S Slavinski, DVM, Mississippi Dept of Health. H Mackley, MPH, B Asante, MD, New York City Dept of Health. M Kohn, MD, E DeBess, DVM, L Davidoff, MD, M Scott, M Heumann, MA, Oregon Health Div; G Stevens, V Barbour, MSN, J Baures, J Manwaring, B Thomas, G Chakarun, Jackson County Public Health Dept, Medford;

R Parlier, M Jaqua, M Breedlove, MBA, Yamhill County Health Dept, McMinnville, Oregon.
J Rankin, Jr., DVM,
C Coventon, MD, Pennsylvania Dept of Health, Bureau of Epidemiology. F Sassano,
Bucks County Dept of
Health, Doylestown; J Maher, MD, E Walls, Chester County Health Dept, Westchester; C
Baysinger, M Supplee, MS,
Montgomery County Human Svcs Center, Norristown; J Jahre, MD, St. Luke's Hospital
Network, Bethlehem; N Sykes, MD,
Jefferson Medical College, Philadelphia, Pennsylvania. R Stroube, MD, E Barrett, DMD, S
Jenkins, VMD, Virginia Health Dept;
J Florance MD, A Ansher MD, L Estrada MPH, B Bradshaw, Prince William Health Dept,
Manassas, Virginia. J
Hofmann, MD, J Van Eenwyk, PhD, Washington Dept of Health. L Haddy, MA, D Bixler,
MD, West Virginia Dept of Health
and Human Resources, Bur for Public Health; P Gordon, MD, Harrison-Clarksburg Health
Dept Clarksburg, West Virginia.
J Perdue, Texas Dept of Health; and other state and local health and education
departments and schools. A Henderson,
PhD, Div of Environmental Hazards and Health Effects, National Center for
Environmental Health. A Adija, MD, K
Griffith, MD, EIS officers, CDC.

References

Brown KE. Human parvovirus B19 epidemiology and clinical manifestations. In:
Anderson LJ, Young NS, eds. Human parvovirus B19
(monographs in virology. vol 20). Basel, Switzerland: Karger 1997:42--60.

Environmental Protection Agency. Indoor air pollution: an introduction for health
professionals. Washington, DC: Environmental

Protection Agency, 1994. Available at <http://www.epa.gov/iaq/pubs/hpguide.htm>. Accessed February 2002.

CDC. Moth-associated dermatitis---Cozumel, Mexico. MMWR 1990;39:219--20.

CDC. Rash illness associated with gypsy moth caterpillars---Pennsylvania. MMWR 1982;31:169--70.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services. References to non-CDC sites on the Internet are provided as a service to MMWR readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of pages found at these sites.

Disclaimer

All MMWR HTML versions of articles are electronic conversions from ASCII text into HTML. This conversion may have resulted in character translation or format errors in the HTML version.

Users should not rely on this HTML document, but are referred to the electronic PDF version and/or

the original MMWR paper copy for the official text, figures, and tables.

An original paper copy of this issue can be obtained from the Superintendent of Documents,

U.S. Government Printing Office (GPO), Washington, DC 20402-9371; telephone: (202) 512-1800.

Contact GPO for current prices.

****Questions or messages regarding errors in formatting should be addressed to**
mmwrq@cdc.gov.

Page converted: 2/28/2002

With 53 million young people attending 117,000 schools each school day in the United States, it is

expected that rashes from a wide range of causes will be observed. Environmental factors or infectious agents can cause rashes

among groups of school-aged children. Rashes caused by infectious agents usually are preceded or accompanied by symptoms such

as headache or fever. However, in these reports, none of the children showed signs of systemic illness, and the rash appeared

to be self-limiting.

Potential environmental causes of rashes include biologic contaminants (e.g., bacteria and fungi), chemical agents

(e.g., cleaning products and pesticide residues), physical agents (e.g., fiberglass), insects (e.g., biting flies and moths), and

allergens

(e.g., dust mites) (2--4). If one of these environmental causes is suspected, appropriate environmental experts should

be consulted.

The most commonly identified viral agent associated with rashes in school-aged children is parvovirus B19, which

causes erythema infectiosum (i.e., fifth disease). Fifth disease is a mild rash illness characterized by a "slapped-cheek" rash on the face and a lacy red rash on the trunk and limbs, which may itch; it usually resolves within 7--10 days. Low-grade fever, malaise, or upper respiratory symptoms usually precede the rash. Other manifestations of parvovirus B19 infection include arthritis and arthralgia (especially in adults), transient crisis of aplastic anemia (in persons with certain hematologic disorders such as sickle-cell anemia), neutropenia, and thrombocytopenia. In pregnant women, parvovirus B19 infection may be associated with miscarriage or nonimmune hydrops fetalis (1).

Public health response to rashes of unknown etiology involves an epidemiologic investigation that includes consultation with facilities and maintenance staff familiar with the physical plant, examination of the rash by a dermatologist, and, when appropriate, collection and analysis of biologic specimens. To date, reports from states do not document a common cause or demonstrate that all children are experiencing the same rash. State and local health departments, in collaboration with CDC, continue to investigate these and other reports of rashes among groups of schoolchildren.

Acknowledgements

This report is based on data contributed by C McRill, MD, K Komatsu, MPH, W Humble, MPH, Arizona Dept of Health Svcs; L Sands, DO, MPH, Maricopa County Dept of Public Health, Phoenix, Arizona. T Wegrzyn, MPH, J Hadler, MD, Connecticut Dept of Public Health. S Wiersma, MD, Florida Dept of Health. P Blake,

MD, S Lance-Parker, PhD, C

Morin, MD, Georgia Dept of Human Resources, Div of Public Health. R Teclaw, DVM, M Wilkinson, Indiana State Dept

of Health. M Currier, MD, S Slavinski, DVM, Mississippi Dept of Health. H Mackley, MPH, B Asante, MD, New York

City Dept of Health. M Kohn, MD, E DeBess, DVM, L Davidoff, MD, M Scott, M Heumann, MA, Oregon Health Div;

G Stevens, V Barbour, MSN, J Baures, J Manwaring, B Thomas, G Chakarun, Jackson County Public Health Dept, Medford;

R Parlier, M Jaqua, M Breedlove, MBA, Yamhill County Health Dept, McMinnville, Oregon. J Rankin, Jr., DVM,

C Coventon, MD, Pennsylvania Dept of Health, Bureau of Epidemiology. F Sassano, Bucks County Dept of

Health, Doylestown; J Maher, MD, E Walls, Chester County Health Dept, Westchester; C Baysinger, M Supplee, MS,

Montgomery County Human Svcs Center, Norristown; J Jahre, MD, St. Luke's Hospital Network, Bethlehem; N Sykes, MD,

Jefferson Medical College, Philadelphia, Pennsylvania. R Stroube, MD, E Barrett, DMD, S Jenkins, VMD, Virginia Health Dept;

J Florance MD, A Ansher MD, L Estrada MPH, B Bradshaw, Prince William Health Dept, Manassas, Virginia. J

Hofmann, MD, J Van Eenwyk, PhD, Washington Dept of Health. L Haddy, MA, D Bixler, MD, West Virginia Dept of Health

and Human Resources, Bur for Public Health; P Gordon, MD, Harrison-Clarksburg Health Dept Clarksburg, West Virginia.

J Perdue, Texas Dept of Health; and other state and local health and education departments and schools. A Henderson,

PhD, Div of Environmental Hazards and Health Effects, National Center for Environmental Health. A Adija, MD, K Griffith, MD, EIS officers, CDC.

References

Brown KE. Human parvovirus B19 epidemiology and clinical manifestations. In: Anderson LJ, Young NS, eds. Human parvovirus B19 (monographs in virology. vol 20). Basel, Switzerland: Karger 1997:42--60.

Environmental Protection Agency. Indoor air pollution: an introduction for health professionals. Washington, DC: Environmental Protection Agency, 1994. Available at <http://www.epa.gov/iaq/pubs/hpguide.htm>. Accessed February 2002.

CDC. Moth-associated dermatitis---Cozumel, Mexico. MMWR 1990;39:219--20.

CDC. Rash illness associated with gypsy moth caterpillars---Pennsylvania. MMWR 1982;31:169--70.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services. References to non-CDC sites on the Internet are provided as a service to MMWR readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of pages found at these sites.

Disclaimer

All MMWR HTML versions of articles are electronic conversions from ASCII text into HTML. This conversion may have resulted in character translation or format errors in the HTML version.

Users should not rely on this HTML document, but are referred to the electronic PDF version and/or

the original MMWR paper copy for the official text, figures, and tables.

An original paper copy of this issue can be obtained from the Superintendent of Documents,

U.S. Government Printing Office (GPO), Washington, DC 20402-9371; telephone: (202) 512-1800.

Contact GPO for current prices.

****Questions or messages regarding errors in formatting should be addressed to mmwrq@cdc.gov.**

Page converted: 2/28/2002

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of

Health and Human Services. References to non-CDC sites on the Internet are provided as a service to MMWR readers and do not constitute or imply

endorsement of these organizations or their programs by CDC or the U.S.

Department of Health and Human Services. CDC is not responsible for the content of pages found at these sites.

Disclaimer

All MMWR HTML versions of articles are electronic conversions from ASCII text into HTML. This conversion may have resulted in character translation or format errors in the HTML version.

Users should not rely on this HTML document, but are referred to the electronic PDF version and/or

the original MMWR paper copy for the official text, figures, and tables.

An original paper copy of this issue can be obtained from the Superintendent of Documents,

U.S. Government Printing Office (GPO), Washington, DC 20402-9371; telephone: (202) 512-1800.

Contact GPO for current prices.

****Questions or messages regarding errors in formatting should be addressed to mmwrq@cdc.gov.**

Page converted: 2/28/2002

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of

Health and Human Services. References to non-CDC sites on the Internet are provided as a service to MMWR readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S.

Department of Health and Human Services. CDC is not responsible for the content of pages found at these sites.

Disclaimer

All MMWR HTML versions of articles are electronic conversions from ASCII text into HTML. This conversion may have resulted in character translation or format errors

in the HTML version.

Users should not rely on this HTML document, but are referred to the electronic PDF version and/or

the original MMWR paper copy for the official text, figures, and tables.

An original paper copy of this issue can be obtained from the Superintendent of Documents,

U.S. Government Printing Office (GPO), Washington, DC 20402-9371; telephone: (202) 512-1800.

Contact GPO for current prices. Page converted: 2/28/2002

[HOME](#) |

[ABOUT MMWR](#) |

[MMWR SEARCH](#) |

[DOWNLOADS](#) |

[RSS](#)

|

[CONTACT](#)

[POLICY](#) |

[DISCLAIMER](#) |

[ACCESSIBILITY](#)

Morbidity and Mortality Weekly Report

Centers for Disease Control and Prevention

1600 Clifton Rd, MailStop E-90, Atlanta, GA

30333, U.S.A

